Anti-inflammatory Effect of *Ampelocissus Indica* (L.) Planch (Chembravalli) in Wistar Rats

Nithya V S (✉ nithyasarangi3@gmail.com)

P G Scholar, Department of Dravyaguna vijnanam, Govt. Ayurveda College Thiruvananthapuram
<https://orcid.org/0000-0002-2064-2620>

Indulekha V C

Associate professor, Department of Dravyaguna vijnanam, Govt. Ayurveda College
Thiruvananthapuram

Helen Antony

professor, Department of Biochemistry, University of Kerala, Kariavattom, Thiruvananthapuram

Research

Keywords: Chembravalli, *Ampelocissus indica* (L.)Planch, anti inflammatory, Ayurveda, Hortus
Malabaricus, Paw edema

Posted Date: August 11th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-54571/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Abstract

Background

Ampelocissus indica (L.) Planch, *Vitaceae* locally known as *Chembravalli*, is medicinal plant used for inflammatory skin ailments and documented in Hortus Malabaricus, and traditional Ayurveda books.

Objective

To conduct phyto-chemical and acute toxicity of root of *A.indica* and to evaluate its anti-inflammatory activity.

Materials and methods

Phyto-chemical profiling of root of *A.indica* was done as per standard ICMR methods and acute toxicity as per OECD-425. Decoction of *A.indica* (AI-8.64 ml/kg, AI-4.32 ml/kg) and ethanolic extract (AIE-500 mg/kg) were orally given to rats with Carrageenan induced rat paw edema and also in other rats with Cotton pellet induced granuloma; paw volume, weight of granuloma, histopathology, serum TNF α , IL1 β were assessed.

Results

Oral administration of decoction AI-8.64 ml/kg, AI-4.32 ml/kg, and AIE-500 mg/kg significantly reduced inflammation by reducing paw volume, serum TNF α , IL1 β . AI-8.64 ml/kg significantly reduced weight of granuloma and serum TNF α , IL1 β . No acute toxicity was found upto 2000 mg/kg. Flavonoids, phenols, tannins, saponins, and steroids found in *A.indica* extract.

Conclusion

The present finding suggested that *A.indica* has a protective effect on experimental inflammation in rats and phytochemical profile will be useful for further studies

1. Background

Inflammation is the immune system's response to harmful stimuli, such as pathogens, damaged cells, toxic compounds, or irradiation, and acts by removing injurious stimuli and initiating the healing process. Inflammation is therefore a defense mechanism that is vital to health. Usually, during acute inflammatory responses, cellular and molecular events and interactions efficiently minimize impending injury or infection. However, uncontrolled acute inflammation may become chronic, contributing to a variety of

chronic inflammatory diseases(1). The early phase of acute inflammation involves the cellular influx associated with the release of mediators such as histamine and serotonin, which are primarily released from mast cells, followed by the production of bradykinin and prostaglandins. During an inflammatory response, several proinflammatory mediators are released, including interleukin-1 beta (IL-1 β), tumor necrosis factor alpha (TNF- α) and cyclooxygenase- 2 (COX-2). These mediators have important role in the initiation and amplification of inflammatory processes(2). The NF- κ B transcription factor plays important roles in inflammatory, immune response, survival, and apoptosis processes. This pathway regulates pro-inflammatory cytokine production and inflammatory cell recruitment, which contribute to the inflammatory response(1). Current anti-inflammatory medications; steroidal and nonsteroidal anti-inflammatory drugs (NSAIDs) having several adverse effects. This creates an emerging interest in natural herbal remedies and dietary supplement for inflammation(3).

According to the WHO report, about 70–80% of the world's population relies on nonconventional medicine mainly from herbal sources in their primary health care (4). The need of the hour is therefore to explore such knowledge of traditional medicines for the benefit of humanity. *A.indica*(Al) is locally known as *Chembravalli* documented in Hortus Malabaricus(5), and some Keraliya traditional books like *Chikitsamanjari*(6), *Arogyakalpadrumam*(7), *Yogamrutam*(8) and *Vaidyatarakam*(9) prescribed for inflammatory skin ailments, wound healing. Anti oxidant, anti diabetic and diuretic properties of the plant and its fractions has been explored earlier (10, 11). Since the traditional knowledge claims the cure of inflammatory skin ailments and wounds; the present study intended to see the anti-inflammatory effect of *A.indica* in Wistar rats

Carrageenan-induced paw edema is one of the most popular and highly sensitive and reproducible test for new non-steroidal anti-inflammatory drugs(12). The cotton pellet granuloma method has been widely employed to access the transudative, exudative and proliferative components of sub acute inflammation(13).

2. Materials And Methods

2.1. Plant material.: The root tuber of the *A.indica* was collected from its natural habitat in Chippancira (Kerala state, India) (Fig1) The plant was authenticated by Dr. Mathew Dan, Senior Scientist & Head Plant Genetic Resource Division Jawaharlal Nehru Tropical Botanical Garden Palod, Thiruvananthapuram, Kerala, India.(Fig 2) and the voucher specimen was deposited at the department of Dravyaguna vijnana of Govt Ayurveda College Thiruvananthapuram, Kerala, India.

2.2 Method of preparation of decoction: Fresh root tuber of *A. indica* was cleaned, dried in shade. 48 g of coarsely powdered *A.indica* was boiled with 768ml water and reduced to 96 ml (as per Ayurveda pharmacopeia of India(14).

2.3 Method of preparation of ethanolic extract: 100g of coarsely powdered tubers of *A.indica* was taken in a conical flask and refluxed for one hour in 98% 1000ml ethanol. The extract was filtered and

evaporated on a rotary evaporator under reduced pressure to obtain the viscous residue and evaporated to dryness over a water bath.

2.4 Biological activity assays

Animals and preparation of test samples for bioassay: Female Wistar rats (150 ± 50 g) were purchased from the Animal house of Sree Chitra Tirunal Institute for Medical Sciences and Technology Thiruvananthapuram and housed under standard condition (12h light / 12h dark cycle at $25 \pm 2^\circ\text{C}$) with free access to standard laboratory food and water.

Carrageenan purchased from Sigma Corporation (USA) was suspended in physiological saline at a concentration of 1%. Antibodies for ELISA and FITC conjugated antibodies were purchased from Abcam, UK. ELISA plates were purchased from NUNC, Denmark.

2.5 Acute toxicity of *A.indica* as per OECD guidelines 425

Acute toxicity of *A.indica* was evaluated as per fixed single dose procedure (limit test) by the Organization for Economic corporation and Development-OECD guidelines 425(15). Five healthy nulliparous female Wistar albino rats weighing 150g to 200 g were selected and were deprived of feed overnight and 3 h after the administration of decoction. 2000 mg/kg of decoction was administered orally and were observed individually for mortality and toxic symptoms at 10 min, 30 min, 1 h, 2 h, 4 h and 6 h and once daily thereafter for 14 days.

2.6 Establishment of carrageenan-induced hind paw edema model and drug administration

The rats were randomly divided into 6 groups ($n=5$), namely normal, Carrageenan control, indomethacin 5mg/kg (standard drug), AI-8.64ml/kg (normal dose of *A.indica* decoction), AI-4.32ml/kg (half dose of decoction of *A.indica*) and AIE -500mg/kg (ethanolic extract of *A.indica*). The decoction AI- 8.64ml/kg, AI- 4.32ml/kg and the ethanolic extract AIE-500mg/kg, indomethacin -5mg/kg, were administered orally an hour prior to the carrageenan injection. The rats in control groups were on distilled water. Carrageenan (0.1mL of 1% solution in normal saline) was injected subcutaneously into plantar fascia of the right hind paw. Measured paw volume at time points, i.e., 0, 1, 2, 3, 4 and 5 hr after carrageenan challenge using a Plethysmometer(16). Blood samples were collected 5 hour after induction of inflammation and centrifuged immediately at 3000 rpm for 10 min. The serum was transferred and stored at -20°C

2.7 Cotton pellet induced granuloma model and drug administration

Rats after overnight fasting, sterile cotton pellet (20 ± 1 mg) was implanted subcutaneously in the inter-scapular region and after the implantation the wound was closed by interrupted suture. The procedure was done under anaesthesia using ketamine hydrochloride injection (40mg/kg i.p). Animals were given the test and standard drug orally once a daily for 7 consecutive days from the day of implantation. On the 8th day, the rats were sacrificed and wet cotton pellets with granuloma were dissected out. Then, they

were dried in hot air oven at 60°C for 24 h and again the dry weight has determined(16). Blood was extracted via cardiac puncture and centrifuged at 3500 rpm for 15 minutes at 4°C. Serum was frozen at -17°C until analyzed. The skin area around granulomatous tissue were taken out and stored in 10% neutral-buffered formaldehyde.

2.8 Analysis of IL-1 β , TNF- α

TNF- α , IL-1 β levels in serum were quantified using ELISA (Engvall and Perlman, 1971) method (17). (0.5ml of o-dianisidine in methanol(1%) + 21ml of citrate phosphate buffer (PH5.0) + 2.4 μ l 30% H₂O₂) was used as a substrate and the absorbance of the colored HRP product was measured spectrophotometrically at 490nm by an automated microplate reader (Thermo Multiskan Spectrum).

2.9 Histopathological analysis: Skin tissue slices were embedded in paraffin and thin sections at 4mm thickness was made using a semi-automated microtome. The sections were stained with Hematoxylin and Eosin, and observed under light microscope to identify the inflammatory changes. Images were captured at 10 and 40 magnifications using DP71 digital camera system Histopathology was evaluated qualitatively with help of a qualified veterinary pathologist.

2.10 Phytochemical profile of *A.indica*

The physical and physicochemical parameters such as moisture content, volatile oil, ash values, fiber content, sugar content, extractive values and qualitative analysis of alkaloid, tannins, flavanoids, phenols, saponins, steroids and HPTLC fingerprint were done as per the standard procedure mentioned in Ayurveda Pharmacopoeia of India(14). Heavy metals content (copper, zinc, iron, cadmium and lead) also measured using Atomic absorption spectrometry (Thermo-scientific, USA)

2.11 Statistical analyses: comparison of paw volume at different time was done by repeated measure of ANOVA with Post hoc multiple comparison tests. Analysis of granuloma weight was by ANOVA and Tukey's multiple comparison. The TNF α and IL-1 β were analyzed by One-way ANOVA.

3. Results

3.1 Plant material

3.2 Acute toxicity study: Throughout the observation period neither incidence of mortality nor animals found in a moribund condition. Body weight gain and animal behavior were normal for 14days. Thus decoction of *A.indica* is safe up to the dose of 2000 mg/kg, in tested rats.

3.3 Carrageenan induced paw edema model: three hours after carrageenan injection, carrageenan control group showed marked paw odema, but the drug treated groups; indomethacin-5mg/kg, AI-8.64ml/kg, AI-4.32ml/kg and AIE-500mg/kg; demonstrated significant amelioration on edema comparing with the

control group ($p < 0.01$) as illustrated in Fig 3 and Table 1. The edema reduction continued up to five hours.

Table 1: post hoc multiple comparison test of mean of paw edema in different groups

group	Subset				
	1	2	3	4	5
Normal control	11.80				
Indomethacin5mg/kg		12.72			
AI8.64ml/kg		12.81	12.81		
AI4.32ml/kg			12.90	12.90	
AIE 500mg/kg				13.05	
Carrageenan control					13.44
Sig.	1.000	0.642	0.599	0.157	1.000

Data represents the post hoc multiple comparison test of mean of paw edema in different groups. Induction with Carrageenan showed obvious increase in paw volume at 1% significance in group Carrageenan control compared with normal control. All the drug treated group (indomethacin5mg/kg, AI-8.64ml/kg, AI-4.32ml/kg and AIE-500mg/kg) showed significant reduction in paw volume when compared with control group. However highest reduction was found in group treated with indomethacin5mg/kg and AI- 8.64ml/kg.

3.4 Cotton pellet induced granuloma model: There was significant reduction of weight of granuloma in indomethacin5mg/kg, AI8.64ml/kg when compared with control group as illustrated in Fig 4. No significant reduction in AI.4.32ml/kg and AIE 500mg/kg.

3.5 Analysis of IL-1 β , TNF- α

In Carrageenan induced paw edema the levels of IL-1 β , TNF α in the treated groups; indomethacin5mg/kg, AI-8.64ml/kg, AI-4.32ml/kg and AIE-500mg/kg group were significantly reduced in comparison to the control group ($p < 0.05$) depicted in table 2. Highest reduction was found in group treated with indomethacin5mg/kg and AI 8.64ml/kg.

In cotton pellet induced granuloma rats also the levels of IL-1 β , TNF α in the treated groups; Indomethacin5mg/kg, AI-8.64ml/kg, AI-4.32ml/kg and AIE-500mg/kg showed significant ($p < .05$) reduction when compared with cotton pellet control group is shown in table 2. Highest reduction was found in group treated with indomethacin-5mg/kg and AI-8.64ml/kg.

Table 2 Serum biochemical analysis

Groups	Paw edema		Cotton pellet granuloma	
	IL-1 β	TNF- α	IL-1 β	TNF- α
Normal control	0.082 $\pm$ 0.003	0.092 $\pm$ 0.003	0.0871 $\pm$ 0.003	0.0953 $\pm$ 0.003
Carrageenan control & cotton pellet control	2.29 $\pm$ 0.085 ^a	1.66 $\pm$ 0.061 ^a	0.2973 $\pm$ 0.011 ^a	0.3372 $\pm$ 0.012 ^a
AI4.32ml/kg	1.24 $\pm$ 0.046 ^{a,b}	1.1275 $\pm$ 0.042 ^{a,b}	0.2860 $\pm$ 0.010 ^{a,b}	0.3085 $\pm$ 0.011 ^{a,b}
AI8.64ml/kg	1.21 $\pm$ 0.045 ^{a,b}	0.9246 $\pm$ 0.034 ^{a,b}	0.1999 $\pm$ 0.007 ^{a,b}	0.2501 $\pm$ 0.009 ^{a,b}
AIE 500mg/kg	1.57 $\pm$ 0.0588 ^{a,b}	1.07 $\pm$ 0.039 ^{a,b}	0.2870 $\pm$ 0.010 ^{a,b}	0.2993 $\pm$ 0.011 ^{a,b}
Indomethacin 5mg/kg	0.747 $\pm$ 0.0278 ^{a,b}	0.719 $\pm$ 0.026 ^{a,b}	0.1814 $\pm$ 0.006 ^{a,b}	0.2030 $\pm$ 0.007 ^{a,b}

Values expressed as the average of 6 values $\pm$ SEM in each group, a-The statistical difference compared with normal control at (p<0.05). b- Statistical difference compared with carrageenan control at (p<0.05).

3.6 Histopathological analysis: Microscopic sections of skin as depicted in Fig 5, shows variable degree of sub acute inflammation often predominated by histiocytes. Occasional giant cells were seen indicating a granulomatous reaction found in control group which was absent in treated groups. The extent and severity of lesions could not be graded.

Histopathological analysis of skin tissue around the granuloma. A- normal rat skin (magnification 10X), B- cotton pellet control(magnification 40X),C- indomethacin5mg/kg(magnification 10X), D- AI8.64ml/kg (magnification 40X), E- AI 4.32ml/kg (magnification 10X), F AIE500mg/kg magnification 40X)

3.7 Phyto-chemical analysis : Phyto-chemical analysis of *A.indica* showed presence of alkaloid, tannins, flavanoids, phenols, saponins, steroids and other physio-chemical parameters depicted in table 3 and HPTLC fingerprint was obtained in solvent system Chloroform : Methanol: Aceton(5:3:2) shown in Fig 6. The heavy metals analysed were within limits as shown in table 4

Table 3 Physico chemical evaluation of root of *A.indica*

Si.no	Experiment	Root tuber
1	Foreign matter (%)	Nil
2	Moisture content (%)	10
3	Volatile oil (%)	Nil
4	Water soluble extractive (%)	7.09
5	Alcohol soluble extractive (%)	1.53
6	Total ash (%)	0.0308
7	Water insoluble ash (%)	0.01155
8	Acid insoluble ash (%)	0.00689
9	Total sugar (%)	4.60
	Reducing sugar (%)	3.81
10	Fibre content (%)	11.47%

Table 4 Atomic absorption spectroscopy (in parts per millionppm)

Heavy metal	<i>A.indica</i>	Maximum concentarion in ppm	Maximum permissible limits
Copper	0.1763ppm		
Zinc	0.2030ppm	20	
Cadmium	0.0240ppm	0.3	
Iron	3.0374ppm	27.4	
Lead	0.1234ppm	10	

4. Discussion

In this study, we elucidated the anti-inflammatory potential of *A.indica*. Inflammation was described as "the succession of changes which occurs in a living tissue when it is injured provided that the injury is not of such a degree as to at once destroy its structure and vitality" or "the reaction to injury of the living microcirculation and related tissues(18).

The carrageenan-induced paw edema in rats, with the advantage of high stability, convenience in operation, short time period and obvious manifestations, is a classical model to estimate the anti-inflammatory activity of natural products(19). Therefore, we applied this model to investigate the acute anti-inflammatory effect. Carrageenans are polysaccharides that induce inflammatory responses and clinical symptoms such as edema, erythema and hyperalgesia appeared immediately after the injection of carrageenan(20). The development of these symptoms resulted from the action of pro-inflammatory agents such as bradykinin, histamine, complement and reactive oxygen, which can be generated in situ at the site of insult or by infiltrating cells like Neutrophils also inflammatory cytokines, such as IL-1 β , IL-6, TNF- α , and inflammatory proteins and enzymes. These molecules can potentially serve as biomarkers for diseases diagnosis, prognosis, and therapeutic decision making(1).

Carrageenan evokes biphasic edema, the first phase of which is mediated by the release of histamine and serotonin from mast cells and the second phase of which involves neutrophil infiltration and the release of prostaglandin E2, cytokines such as IL-1 β , IL-6, IL-10, and TNF- α which can be attributed to the action of inducible cyclooxygenase (COX- 2)(21). The second phase is continued until 5 h after the induction of inflammation.

Indomethacine was the reference drug for carrageenan-induced paw edema model in this study. After 4-5 hours; all the drug treated groups produced significant reduction in paw volume ($p < .01$), when compared with carrageenan control group. Indomethacin 5mg/kg treated group and AI- 8.64ml/kg group showed high reduction in paw volume, also the AI-4.32ml/kg and AIE-500mg/kg group has significant reduction in paw volume when compared with control group. So based on the above reports, it can be inferred that the anti-inflammatory effect of *A.indica* may be due to the inhibition of the enzyme cyclooxygenase leading to the inhibition of prostaglandin synthesis.

The cotton pellet granuloma method has been widely employed to access the transudative, exudative and proliferative components of subacute inflammation(13). In this study the Indomethacin 5mg/kg group and AI-8.64ml/kg group elicited significant reduction on the weight of granuloma at $p < .01$, when compared with cotton pellet control group. AI-4.32ml/kg and AIE-500mg/kg were not elicited significant reduction on the weight of granuloma.

TNF- α and IL-1 β are pro-inflammatory cytokines that are involved in the interaction with various target cells as well as diverse immunological functions. These cytokines also mediate immunity and inflammation. Large amounts of TNF- α and IL-1 β are released by macrophages(22). The drug treated groups significantly reduced ($p < .01$) TNF- α , IL-1 β level in rat serum when compared with control group. Indomethacin 5mg/kg and AI-8.64ml/kg treated group showed high reduction in serum TNF- α , and IL-1 β than AI-4.32ml/kg and AIE500mg/kg even though they are significant when compared with the control group. Reduction of cytokine can be attributed to the anti-inflammatory activity.

Histopathological analysis was done on the skin tissue around the granuloma. All the groups showed signs of inflammation when compared with normal control. Occasional giant cells were seen indicating a

granulomatous reaction found in control group which was absent in treated groups. The extent and severity of lesions could not be graded.

In acute oral toxicity test, no mortality recorded even at the highest dose level administered. This proves that drug is safe up to the dose 2000mg/kg.

Authentication and phytochemical profiling of root of *A.indica* was carried out as a part of standardization of herbal drug. In preliminary phytochemical screening, the value of water soluble extractive is higher than the alcohol soluble extractive; which may be due to the presence of more water soluble principles and starch content of tuber. The qualitative analysis of the sample revealed the presence of flavonoids, phenols, alkaloids, tannins, saponins, which indicates that it is a potent drug. Heavy metal contents were within the permissible limit, indicating good collection and storage practice of the *A.indica* sample. The phytochemical parameters can be used for future drug evaluation.

The phytochemical analysis of *A.indica* revealed the presence of flavonoids. Flavonoids are reported as antioxidants and scavengers of free radicals(19). The antioxidant activity of *A. indica* is proven(10). So the flavonoids present in the drug *A.indica* may be the main phytochemical responsible for the observed anti-inflammatory property of the drug.

5. Conclusion

The decoction and extract of *A.indica* have anti-inflammatory effects and the probable mechanism is down regulation of cytokines such as IL-1 β , and TNF- α . The phytochemicals present in *A.indica* such as flavonoids, steroids and tannins may also modulate by its antioxidant property . The acute toxicity study proved that the drug is safe up to the dose 2000mg/kg.

6. Declarations

Ethics approval

Animal welfare and experimental procedures were carried out strictly in accordance with the Guide for the Care and Use of Laboratory Animals and approved by Institutional Animal Ethics Committee and CPCSEA committee (2015/GO/Re/S/18/CPCSEA 12/06/2018) in Government Ayurveda College Thiruvananthapuram.

Consent for publication

Not applicable

Availability of data and materials

All data generated or analysed during this study are included in this article

Competing interest

The authors declare that there are no competing interest.

Funding

Kerala State Council for Science, Technology and Environment

Authors contribution

Nithya V S – conducted the entire study

Indulekha V C – guide ship for the entire study

Helen Antony – biochemical analysis

Acknowledgment

With profound indebtedness, I am thankful to

Dr. Jollykutty Eapen M.D (Ay), former DAME, former Principal, Govt. Ayurveda College, Thiruvananthapuram for institutional support

Dr. M.A Shajahan MD (Ay), former Professor and HOD of Department of Dravyaguna vijnanam, Govt. Ayurveda College, Thiruvananthapuram for institutional support

Dr. A. Shahul Hameed MD (Ay), Professor and HOD of the Department of Dravyaguna vijnanam, Govt. Ayurveda College, Thiruvananthapuram for institutional support

Dr. Deepa. M. S MD (Ay), Associate Professor, Department of Dravyaguna vijnanam, Govt. Ayurveda College, Thiruvananthapuram for institutional support

Dr. Mathew Dan, Senior Scientist & Head Plant Genetic Resource Division JNTBGRI Palod, **Mr. Jayalal** staff of JNTBGRI Palod for authentication of plant material

Mrs. Monisha, Phd scholar Dept. of Biochemistry, Kerala University for biochemical analysis

Dr. Anilkumar scientist, Sree Chitra Tirunal Institute for Medical Sciences and Technology, Thiruvananthapuram for histopathological analysis

Prof. Somasekharan professor and HOD Dept. of statistics, Kerala University and **Mr.Nidhi P Ramesh**, Assistant Professor, Dept. of statistics, Mar Athanasius College, Ernakulam, for statistical analysis

Dr. Meena C V, Mr. Mahadevan and Mrs. Rosamma. M.P Research Officers, Drug Standardization Unit Govt. Ayurveda college, Thiruvananthapuram for phytochemical analysis.

References

1. Chen L, Deng H, Cui H, Fang J, Zuo Z, Deng J, et al. Inflammatory responses and inflammation-associated diseases in organs. *Oncotarget*. 2017;9(6):7204-18.
2. Coura CO, Souza RB, Rodrigues JAG, Vanderlei EdSO, de Araújo IWF, Ribeiro NA, et al. Mechanisms involved in the anti-inflammatory action of a polysulfated fraction from *Gracilaria cornea* in rats. *PloS one*. 2015;10(3).
3. Jitta SR, Daram P, Gourishetti K, Misra CS, Polu PR, Shah A, et al. *Terminalia tomentosa* Bark Ameliorates Inflammation and Arthritis in Carrageenan Induced Inflammatory Model and Freund's Adjuvant-Induced Arthritis Model in Rats. *Journal of Toxicology*. 2019 2019/01/17;2019:7898914.
4. Amdekar S, Roy P, Singh V, Kumar A, Singh R, Sharma P. Anti-Inflammatory Activity of *Lactobacillus* on Carrageenan-Induced Paw Edema in Male Wistar Rats. *International Journal of Inflammation*. 2012 2012/02/22;2012:752015.
5. Rheed V. *Horthus Malabaricus*. Thiruvananthapuram: University of Kerala; 2003. p. 23.
6. Namboothiri DS. *Chikitsamanjari*. 12 ed: Vidyarambham Publishers; 2015.
7. varier KR. *Arogyakalpadrumam*. Thrissur: Sahithibooks; 2003. p. 106.
8. Namboothiri DS. *Yogamrutham*. 5 ed: Vidyarambham Publishers 2010.
9. Narayanan CN. *Vaidyatharakam* 2ed. Kottayam: Dr.M N Sasidharan; 1974.
10. Sasikumar P, Prabha B, Reshmitha T, Veluthoor S, Pradeep A, Rohit K, et al. Comparison of antidiabetic potential of (+) and (-)-hopeaphenol, a pair of enantiomers isolated from *Ampelocissus indica* (L.) and *Vateria indica* Linn., with respect to inhibition of digestive enzymes and induction of glucose uptake in L6 myotubes. *RSC Advances*. 2016;6(80):77075-82.
11. Sunilson JA, Venkatenarayanan R, Thangathirupathi A, Saraswathi A, Praveen M, Anithagnanakumari A. Diuretic activity of the extract of *Vitis indica*. *Ancient science of life*. 2004;23(4):20.
12. Dzoyem J, McGaw L, Kuete V, Bakowsky U. Anti-inflammatory and anti-nociceptive activities of African medicinal spices and vegetables. *Medicinal spices and vegetables from Africa*: Elsevier; 2017. p. 239-70.
13. Ashok P, Koti B, Thippeswamy A, Tikare V, Dabadi P, Viswanathaswamy A. Evaluation of antiinflammatory activity of *Centratherum anthelminticum* (L) Kuntze seed. *Indian J Pharm Sci*. 2010;72(6):697.
14. Welfare GoIMoHaF. *The Ayurveda Pharmacopoeia Of India*. In: Ayush Do, editor. New Delhi 2007.
15. Guideline OO. 425: acute oral toxicity—up-and-down procedure. *OECD Guidelines for the Testing of Chemicals*. 2001;2:12-6.
16. Misra AK, Varma SK, Kumar R. Anti-inflammatory Effect of an Extract of *Agave americana* on Experimental Animals. *Pharmacognosy research*. 2018;10(1):104.

17. Engvall E, Perlman P. Enzymelinked immunosorbent Assay(ELISA). Quantitative assay of immunoglobulin G. Immunochemistry. 1971.
18. Neville A PUNCHARD CJW, Ian Adcock. The Journal of Inflammation Journal of Inflammation 2004.
19. Alqasoumi SI, Soliman G, Awaad AS, Donia A. Anti-inflammatory activity, safety and protective effects of *Leptadenia pyrotechnica*, *Haloxylon salicornicum* and *Ochradenus baccatus* in ulcerative colitis. *Phytopharmacology*. 2012;2(1):58-71.
20. Morris CJ. Carrageenan-induced paw edema in the rat and mouse. *Inflammation protocols*: Springer; 2003. p. 115-21.
21. Coura CO, Souza RB, Rodrigues JAG, Vanderlei EdSO, de Araújo IWF, Ribeiro NA, et al. Mechanisms involved in the anti-inflammatory action of a polysulfated fraction from *Gracilaria cornea* in rats. *PloS one*. 2015;10(3):e0119319.
22. Park SM, Lee TH, Zhao R, Kim YS, Jung JY, Park CA, et al. Amelioration of inflammatory responses by Socheongryong-Tang, a traditional herbal medicine, in RAW 264.7 cells and rats. *International journal of molecular medicine*. 2018;41(5):2771-83.

Figures

Figure 1

Histopathology samples

Figure 2

box plot diagram of analysis of granuloma weight

Figure 3

Line diagram of analysis of paw volume

Figure 4

Authenticated sample of *A.indica*

Figure 5

Root tubers of *A.indica*

Figure 6

HPTLC densitogram