

# CD83 Expression Regulates Antibody Production in Response to Influenza A Virus Infection

**Madhav Akauliya**

Hallym University College of Medicine

**Avishekh Gautam**

Hallym University College of Medicine

**Sony Maharjan**

Hallym University College of Medicine

**Byoung Kwon Park**

Hallym University College of Medicine

**Jinsoo Kim**

Hallym University College of Medicine

**Hyung-Joo Kwon** (✉ [hjookwon@hallym.ac.kr](mailto:hjookwon@hallym.ac.kr))

Hallym University College of Medicine

---

## Research

**Keywords:** Antibody production, B cells, CD83, Influenza A virus, Peritoneal cavity

**Posted Date:** August 13th, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-55905/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published on December 10th, 2020. See the published version at <https://doi.org/10.1186/s12985-020-01465-0>.

# Abstract

**Background:** CD83 is known to regulate lymphocyte maturation, activation, homeostasis, and antibody response to immunization and infection. While CD83 has a major part in B cell function, its role in influenza A virus infection has not yet been investigated.

**Methods:** We investigated the role of CD83 using C57BL/6J wild type mice and CD83 knockout (KO) mice after intraperitoneal administration of the influenza A/WSN/1933 virus. We analyzed cells of the peritoneal cavity, splenocytes, and cells of the bone marrow with FACS to investigate CD83 expression and cell population change in response to the virus infection. ELISA was performed with sera and peritoneal cavity fluids to detect A/WSN/1933 virus-specific IgG and the subclasses of IgG.

**Results:** FACS analysis data showed a transient but distinct induction of CD83 expression in the peritoneal B cells of wild type mice. CD83 KO mice exhibited a delayed recovery of B cells in the bone marrow after influenza virus infection and overall, a smaller T cell population compared to wild type mice. The peritoneal cavity and serum of the wild type mice contained a high titer of IgG within 14 days after infection, whereas the CD83 KO mice had a very low titer of IgG.

**Conclusions:** These results show the importance of CD83 in lymphocytes homeostasis and antibody production during influenza A virus infection.

## Background

CD83, an evolutionary conserved member of immunoglobulin, is a highly glycosylated type 1 transmembrane glycoprotein composed of 175 amino acids in mice [1] and 186 amino acids in humans [2]. CD83 consists of a variable extracellular Ig-like domain, transmembrane (TM) domain, and intracellular C-terminal cytoplasmic domain sharing 63% homology of amino acids between mice and humans [3]. CD83 is expressed in mature dendritic cells (DCs) and is an activation marker for DCs [3]. However, CD83 is also expressed in natural killer cells [4], activated macrophages [5, 6], neutrophils [7, 8], and activated T cells and B cells [2, 9–11]. CD83 is also expressed on thymic epithelial cells, where it contributes to the selective development and maturation of CD4<sup>+</sup> T cells [12].

The TM domain of CD83 promotes and stabilizes the cell surface expression of major histocompatibility complex II (MHC II) and CD86 on DCs by inhibiting their association with membrane-bound E3 ubiquitin ligase MARCH1 thereby affecting the activation of CD4<sup>+</sup> T cells and B cells [13]. CD83 is required for the differentiation and stability of regulatory T (Treg) cells [14]. Moreover, CD83 also promotes the expansion and antigenic specificity of CD8<sup>+</sup> T cells [15].

Immature B cells beyond the pre-B cell stage, after acquiring functional B cell receptors (BCRs), express a low level of CD83, which has an important role in B cell activation, maturation, homeostasis, effective functioning, longevity, and germinal center response [16–18]. CD83 also reduces the sensitivity of BCRs

preventing the overstimulation of B cell [19]. Furthermore, the CD83 antibody in the human system was shown to make B cells unresponsive to antigens, along with a reduced response to CD4<sup>+</sup> T cells [20].

Because CD83 regulates the development and function of various immune cells, it also modulates the immune response during infections. Infection by different viruses leads to the degradation of membrane CD83 in dendritic cells, interfering with the maturation and immune response of DCs [21–24]. For example, upon hepatitis B virus infection in humans, DCs exhibit a reduced capacity for antigen presentation, cytokine production, phagocytosis, and migration [22]. Previous studies have shown that CD83 also affects B cell function during infection. Mice constitutively expressing CD83 under MHC I promoter (CD83Tg) showed reduced production of antigen-specific antibody production upon infection with *Leishmania major* and *Trypanosoma cruzi* [11]. Similarly, *in vitro* stimulation of CD83Tg B cells with LPS resulted in enhanced interleukin-10 (IL-10) production along with diminished Ig secretion [25].

Although CD83 is important for B cell function, its role in influenza A virus infection has not yet been investigated. Previously, we showed that intraperitoneal infection with the influenza A/WSN/1933 virus caused a transient but substantial depletion of B cells in the peritoneal cavity [26]. Because B cell function is important to combat virus infection, we investigated the modulation of the B cell and T cell population at different time points after influenza A virus infection and confirmed the requirement of CD83 in the virus-specific antibody production using CD83 knockout (KO) mice.

## Methods

### Cell line and viruses

Madin-Darby Canine Kidney (MDCK) cell lines used for this study were purchased from the American Type Culture Collection (ATCC, Manassas, VA, USA) and grown on Minimum Essential Medium (MEM) with 10% fetal bovine serum (FBS), 100 µg/ml streptomycin and 100 U/ml penicillin. The influenza A virus, A/WSN/1933 (H1N1), was obtained from Professor Man-Seong Park (Korea University, Seoul, Korea). The virus was amplified using specific-pathogen-free (SPF) embryonated eggs followed by infecting the MDCK cell lines. MDCK cells ( $2 \times 10^5$ /well) were cultured in 6-well plates using MEM media containing 10% FBS at 37°C overnight in a CO<sub>2</sub> incubator. After the overnight culture, the cells were washed with PBS, and each well was infected with the influenza A virus at MOI 0.01 in MEM media containing 1 µg/ml L-tosylamide-2-phenylethyl chloromethyl ketone (TPCK)-treated trypsin and then incubated at 37°C in CO<sub>2</sub> incubator. After 1 h incubation, the supernatants were removed and then cultured for 3 days in MEM media containing 0.3% BSA at 37°C in a CO<sub>2</sub> incubator. The virus culture supernatants were collected and centrifuged at 2,000 rpm for 10 min at 4°C to remove the cell debris. The quantification of the amplified viruses was performed by plaque assay. MDCK cells ( $6 \times 10^5$ /well) were cultured in 6-well plates. The cells were cultured, washed with PBS, and infected with the amplified influenza A virus culture supernatants after a ten-fold serial dilution. After 1 h incubation, the supernatants were removed and overlaid with 2 ml of DMEM/F12 media containing 2 mM glutamine, 4%

BSA, 10 mM HEPES, 2.5% sodium bicarbonate, 50 mg/ml DEAE dextran, 1 µg/ml L-tosylamide-2-phenylethyl chloromethyl ketone (TPCK)-treated trypsin, 100 U/ml penicillin, 100 mg/ml streptomycin and 0.6% immunodiffusion-grade agar. After 72 h incubation, the plates were stained with crystal violet (0.1% crystal violet in 20% methanol) for 1 h. The plaques were counted to determine the virus titers. The entire process for the virus preparation and cell culture procedure was conducted in a biosafety level 2 laboratory.

## Mice

Eight-week-old female C57BL/6J mice were purchased from NARA Biotech, Inc. (Seoul, Korea), and C57BL/6J CD83 KO mice (B6.129S4-*Cd83<sup>tm1Tft</sup>*/J, JAX stock #017703) were purchased from The Jackson Laboratory (Bar Harbor, ME, USA). The mice were maintained at the Animal Center of Hallym University under specific pathogen-free conditions (20–25°C, 40–45% humidity, 12 h light/dark cycle, and food and water access, *ad libitum*). Mice experiments were performed in a biosafety level 2 facility in the Hallym Clinical and Translational Science Institute.

## Virus infection

A/WSN/1933 virus inactivation was performed by 254 nm UV light exposure with 1,500 mW/s/cm<sup>2</sup> UV for 15 min from a height of 5 cm. The virus inactivation was verified by a plaque assay with MDCK cells [27]. The live A/WSN/1933 virus or UV-inactivated A/WSN/1933 virus was intraperitoneally infected at a dose of  $5 \times 10^6$  pfu per mouse.

## FACS analysis

C57BL/6J wild type or C57BL/6J CD83 KO mice were infected intraperitoneally with  $5 \times 10^6$  pfu of A/WSN/1933 virus. After the virus infection, cells of the peritoneal cavity, splenocytes, and cells of the bone marrow were harvested in RPMI 1640 media containing 5% FBS. The cells were centrifuged at 1,200 rpm and 4°C for 5 minutes, and the pellets were treated with RBC lysis buffer (20 mM Tris HCl and 140 mM NH<sub>4</sub>Cl) for 5 min. After washing with RPMI media, the cells were washed with FACS buffer (1% FBS in PBS) and transferred into Falcon tubes and then treated with 10 µg/ml anti-FcγRII/III antibody (Catalogue No: 553142, BD Biosciences, San Jose, CA, USA). To analyze lymphoid population, PerCP Cy5.5-conjugated anti-CD3 antibody (Catalog No: 551163, BD Biosciences), Pacific blue ef450-conjugated anti-CD4 antibody (Catalog No: 558107, BD Biosciences), FITC-conjugated anti-CD8 antibody (Catalog No: 553031, BD Biosciences), APC-Cy7-conjugated anti-CD19 antibody (Catalog No: 557655, BD Biosciences), PE-conjugated anti-CD23 antibody (Catalog No: 12-0232-81, eBioscience, San Diego, CA, USA), and APC-conjugated anti-CD83 antibody (Catalog No: 558208, BD Biosciences) were used.

## ELISA

For the detection of A/WSN/1933 virus-specific IgG, the subclasses of IgG, ninety-six well Immuno plates (Nunc™, Roskilde, Denmark) were coated with the A/WSN/1933 virus. After overnight incubation at 4°C,

the plates were washed three times with PBST (0.1% Tween-20 in PBS) and blocked with 1% BSA for 1 h and then loaded with three-fold dilutions of sera and peritoneal supernatants in PBST. After incubation at room temperature for 2 h, the plates were washed three times with PBST and goat anti-mouse IgG/IgG1/IgG2a/IgG2b/IgG3 antibodies labeled with Horseradish peroxidase (Southern Biotechnology Associates, Inc. Birmingham, AL, USA) with dilutions 1:500 were used to bind total IgG and each subclasses antibody. The plates were incubated for 1 h at room temperature, and then, TMB substrate solution was added (Kirkegaard and Perry Laboratories, Gaithersburg, MD, USA). The absorbance was measured by a Spectra Max 250 microplate reader (Molecular Devices, Sunnyvale, CA, USA) at 450 nm

## Statistical analysis

Results are shown as the mean  $\pm$  standard deviation. The statistical significance of the differences between the two samples was evaluated using Student's t-test with  $P < 0.05$  as the threshold for statistical significance.

## Ethical approval

Mice experimental procedures were undertaken following the guidelines of Laboratory Animals of the National Veterinary Research and Quarantine Service of Korea. Animal use and related experimental procedures were approved by the Institutional Animal Care and Use Committee of Hallym University (Permit Number: Hallym R2/2018-25). Mice were anesthetized with 3% to 5% isoflurane (Pharmaceutical, Seoul, Korea) inhalation to minimize any pain. The mice were sacrificed by CO<sub>2</sub> inhalation after the experiments and euthanized by CO<sub>2</sub> inhalation if they lost 25% of their baseline adult body weight or if they revealed evidence of debilitation, pain or distress such as a hunched posture, rough hair coat, reduced food consumption, emaciation, inactivity, ambulation difficulty, and respiratory problems, and all efforts were made to limit suffering.

# Results

## CD83 expression in peritoneal B cells upon influenza A virus infection

B cells perform a vital function in combating infection through the production of antibodies and cytokines and antigen presentation to CD4<sup>+</sup> and CD8<sup>+</sup> T cells [28]. Because CD83 has an important role in the development and function of B cells and T cells [17], we investigated the role of CD83 in B cell function during influenza A virus infection in mice. To determine the effect of influenza A virus on CD83 expression in peritoneal cells, C57BL/6J wild type mice were intraperitoneally injected with  $5 \times 10^6$  plaque forming units (pfu)/mouse of live or UV-inactivated A/WSN/1933 virus. FACS analysis results showed that the expression of CD83 on CD19<sup>+</sup> peritoneal B cells was drastically increased at 3 h and then restored to basal level at 24 h after infection with live A/WSN/1933 virus (Fig. 1b). The expression of CD83 on CD19<sup>+</sup> B cells was also increased at 3 h in the mice infected with UV-inactivated WSN virus; however, the levels of CD83 expression at 6 h were substantially lower than those of live virus-infected

mice (Fig. 1a). CD83 expression on CD19<sup>+</sup> B cells in mice infected with UV-inactivated virus was also restored to basal level at 24 h. The expression of CD83 on CD19<sup>+</sup> B cells from bone marrow and the spleen was not affected by live or UV-inactivated A/WSN/1933 virus-infection (Fig. 1b). However, CD3<sup>+</sup> T cells from the peritoneum cavity, bone marrow, and spleen showed no difference in expression of CD83 upon infection with either live or UV-inactivated virus (Fig. 1c). These results show that infection of influenza A virus leads to a transient but distinct increase of CD83 expression in the surface of peritoneal B cells.

CD83 is required for the restoration of lymphocyte populations in the bone marrow after intraperitoneal challenge with influenza A virus

Previously, we have reported that intraperitoneal inoculation with the A/WSN/1933 virus induced a substantial but transient B cell depletion at 12 h post-infection, which was partially recovered at 7 days with a gradual restoration to normal levels by 14 days [26]. To investigate the effect of CD83 expression in the lymphocyte population and its restoration pattern in the peritoneal fluid, bone marrow, and spleen, wild type mice and CD83 KO mice were intraperitoneally injected with the live A/WSN/1933 virus at a dose of  $5 \times 10^6$  pfu/mouse. Peritoneal cells, bone marrow cells, and splenocytes were harvested at 5, 7, and 14 days post-infection, and cell populations were analyzed by flow cytometry. In the wild type mice, depletion of CD19<sup>+</sup> B cells was observed at 5 days post-infection in the peritoneal cavity (Figure 2a) and bone marrow (Figure 2b), followed by a partial recovery at 7 days and restoration to near-normal levels at 14 days post-infection as previously described [26]. In CD83 KO mice, transient depletion and then recovery to the normal level were observed in CD19<sup>+</sup> B cells of the peritoneal cavity similar to wild type mice (Fig. 2a). However, the CD19<sup>+</sup> B cells in bone marrow exhibited an incomplete recovery in the CD83 KO mice until 14 days post-infection, suggesting a delay in the restoration process (Fig. 2b). On the other hand, the peritoneal cells exhibited an increment in CD3<sup>+</sup> T cells for both the wild type and CD83 KO mice after infection although the amounts of T cells at 7 and 15 days post-infection were higher in the wild type mice than in the CD83 KO mice (Fig. 2a). Furthermore, the basal levels of CD3<sup>+</sup> T cells in the peritoneal cavity of the CD83 KO mice were much lower (about 50%) compared to those of wild type mice (Fig. 2a). The T cell numbers of the bone marrow declined at day 5 and slowly recovered at day 7 and 14 in the wild type mice and CD83 KO mice (Fig. 2b). In the spleen, the CD19<sup>+</sup> B cells and CD3<sup>+</sup> T cells were decreased at 5 days post-infection, followed by a partial recovery at 7 days and restoration to near-normal levels at 14 days post-infection with no prominent difference between the wild type mice and CD83 KO mice (Fig. 2c). Collectively, these results show the requirement of CD83 for the homeostasis of T cells in the peritoneal cavity fluids and for the restoration of B cells in the bone marrow following influenza A virus infection.

**Influenza A virus triggers an altered behavior by the T cell subpopulation in the peritoneal cavity of CD83 knockout mice.**

B cell populations are composed of subclasses B-1 and B-2 cells. B-1 cells, unlike the classical B-2 cells, are characterized by a low level of CD23 expression and participates in the antibody response during vaccination and infection [29]. Therefore, we analyzed the B-1 and B-2 cells in the peritoneal cavity fluids after intraperitoneal infection of live A/WSN/1933 virus. The depletion of both the B-1 and B-2 cells was observed at 5 days post-infection, followed by partial recovery at 7 days in both wild type mice and CD83 KO mice (Fig. 3a). While B-2 cells had a full restoration of their cell numbers at day 14, B-1 cells were unable to recover to a normal level. However, there was no significant difference between the wild type and CD83 KO mice (Fig. 3a). Next, we checked the T cell populations and found that after the virus inoculation, the CD8<sup>+</sup> T cell number underwent continuous expansion as observed at day 5, 7, and 14 in both the wild type (3 fold) and CD83 KO (5 fold) mice (Fig. 3b). In contrast, CD4<sup>+</sup> T cells exhibited an altered behavior between the wild type and CD83 KO mice. As observed earlier, the T cell population was smaller in the CD83 KO mice [30]. In the wild type mice, the CD4<sup>+</sup> T cell number was reduced at day 5 post-infection and later was restored to the normal level by day 14. However, the CD4<sup>+</sup> T cell number mildly increased after infection in the CD83 KO mice; however, the total number of cells in the CD83 KO mice was still much lower compared to the wild type mice (Fig. 3b). Overall, these results suggest that CD83 loss could lead to differential properties, especially in the T cell population in the peritoneal cavity.

### **Effect of CD83 expression on antibody production after intraperitoneal challenge with influenza A virus**

Upon infection, some of the naïve B cells are converted to plasma cells that produce antibodies against the antigenic components of invading pathogens, which has an essential role in the clearance of pathogens [31]. Antibody production by B cells could be CD4<sup>+</sup> T cell-dependent or independent [32]. Because the loss of CD83 affected the B cell population and CD4<sup>+</sup> T cell population after infection, we investigated the effect of CD83 expression on antibody production in response to virus infection by measuring the level of virus-specific IgG and its subclasses in the peritoneal cavity fluids and serum at 5, 7 and 14 days post-infection. There were no detectable virus-specific IgG in the PBS control mice. A negligible amount of the virus-specific IgG was produced in the peritoneal cavity fluids and serum at day 5 and 7 post-infection, which later increased to a considerable value by day 14 in the wild type mice. Notably, IgG2b and IgG3 were the major IgG isotypes in the peritoneal cavity fluids (Fig. 4a), whereas IgG2b was the most prominent in the serum (Fig. 4b). Conversely, the CD83 KO mice failed to produce virus-specific antibodies comparable to the wild type mice (Fig. 4). Although IgG levels were increased at day 14 in the CD83 KO mice, the amount was significantly lower compared to the wild type mice both in the peritoneal cavity fluids (Fig. 4a) and serum (Fig. 4b). Taken together, these results suggest that CD83 has an important role in B cell response to influenza A virus infection with a direct influence on the production of virus-specific antibodies.

## **Discussion**

Although CD83 is a key component of the immune system that can modulate T cell and B cell function, no studies involving CD83 and influenza virus have been carried out so far. Recently, we demonstrated

that intraperitoneal injection of influenza A/WSN/1933 virus induced an efficient immune response and robust virus-specific antibody production by B cells that sufficiently provides cross-protection against other strains of influenza A virus [26]. Here, we investigated the effect of CD83 on B and T cells during influenza virus infection using C57BL/6J wild type mice in parallel with CD83 KO mice. We found that CD83 expression is required for the optimal production of virus-specific antibodies during influenza virus infection.

B cells have a crucial role in combating virus infection via antigen presentation to the T cells and the production of high-affinity antibodies against virus antigens [33]. To explore the function of CD83 in B cells, we examined the expression of CD83 on the cell surface of B cells after intraperitoneal administration of influenza A virus on wild type mice. Influenza A virus infection induced the cell surface expression of CD83 on peritoneal B cells, reaching a plateau within 3 h, without affecting the B cells in the spleen and bone marrow. The expression of CD83 returned to the basal level within 24 h. Although CD83 also has an important role in the development and functioning of T cells [12, 15], influenza A infection did not significantly affect the CD83 expression level on the T cells in our experimental settings. These initial results suggest that CD83 mainly functions through B cells during the early stages of influenza A infection like priming or activation of the B cells.

Previously, we demonstrated that intraperitoneal administration of influenza A virus triggered an increase of T cells and a transient loss of B cells and macrophages, which recovered after 2 weeks post-infection in the peritoneal cavity [26]. Through experiments with CD83 KO mice in this study, it was found that without CD83, the B cell population was recovered in the peritoneal cavity fluids and spleen similarly to wild type, but the B cell recovery in the bone marrow was prominently delayed. Importantly, the T cell population was smaller in the peritoneal cavity, bone marrow, and spleen in the CD83KO mice with the highest difference in the peritoneal cavity. B cells can activate T cells via antigen presentation on the MHC molecules, and the activated helper T cells can activate the B cells to enter the germinal center reaction where the activated B cells expand exponentially and later differentiate to memory cells or plasma cells [34, 35]. CD83 is known to stabilize the MHC II molecule on the cell surface by inhibiting its interaction with ubiquitin ligase. Thus, it is plausible that in the absence of CD83, the expansion of CD4<sup>+</sup> T cells in the peritoneal cavity is impaired, resulting in the delayed activation of B cells and lesser IgG production after influenza A virus infection. Antibodies that are derived from B cells in response to influenza virus are crucial in preventing mortality and morbidity, providing a full recovery, and producing memory cells for future immunity [31]. In this study, mice infected with influenza A virus produced a high amount of IgG after 14 days as measured in the peritoneal cavity fluids and serum. Concomitantly, with a reduced population of CD4<sup>+</sup> T cells, CD83 KO mice had very low levels of virus-specific antibodies. Previous studies showed that the surface expression of CD83 on B cells is indirectly proportional to the amount of IgG production. Kretschmer *et al.* showed that CD83Tg overexpressing B cells produce less Igs, whereas CD83 deficient B cells produced higher levels after T-cell dependent and independent immunization [25]. In our experimental conditions, the virus infection induced CD83 expression only transiently, and the expression level returned to the basal level within 24 h post-infection. Therefore, CD83

is likely to affect the activation and proliferation of B cells. The lower number of the CD4<sup>+</sup> T cell population in the CD83 KO mice [12] could also have affected the antibody production efficacy after virus infection. Further studies are required to dissect the precise role of CD83 on B cells and the interaction of CD83<sup>+</sup> B cells and CD4<sup>+</sup> T cells in antibody production. Moreover, the role of CD83 on macrophages and dendritic cells against influenza virus infection needs to be determined in future studies.

Gordon et al. [36] showed that IgG2 deficiency was associated with an increased risk of severe H1N1 infection in obese, pregnant, and immunosuppressed cases leading to the severe pandemic 2009 influenza A infection. Another study reported the effect of age on IgG subtypes after influenza vaccination and found that IgG1 and much more IgG3 were elevated in the serum. The higher level of IgG3 production was found to be positively correlated with the level of tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) and interleukin-6 (IL-6) [37] because IL-6 is the major cytokine produced during influenza infection [38]. In response to the intraperitoneal infection of the influenza A/WSN/1933 virus in C57BL/6J wild type mice, IgG2b and IgG3 were the dominant IgG in the peritoneal cavity fluids whereas IgG2b was the major IgG in the serum. In the CD83 KO mice, both in the peritoneal cavity fluids and serum, IgG was found to be decreased in comparison to the wild type mice. In contrast to our study, previous reports showed that IgG1 was the major subclass in the serum of C57BL/6J mice, while IgG2a was the major one in BALB/c mice infected with the A/Queensland/6/72 (H3N2) influenza virus [39]. Another study showed serum IgG2a as prominent in CBA/CaH mice during intraperitoneal inoculation of A/Queensland/6/72 (H3N2) [40]. Altogether, it is likely that the prevalence of IgG subclass depends on various factors like the type of virus strain, route of inoculation, days of measurement, and types of experimental mice undertaken for the study.

## Conclusions

This work showed the important role of CD83 in B and T cell homeostasis and antibody production during influenza A virus infection. These results could provide important insights into future studies that are directed towards antibody response and vaccine production for influenza viruses.

## Abbreviations

DC Dendritic cell

ELISA Enzyme-linked immunosorbent assay

FACS *Fluorescence-activated cell sorting*

FBS Fetal bovine serum

KO Knockout

MDCK Madin-Darby Canine Kidney

MEM Minimum Essential Medium

pfu Plaque forming unit

## Declarations

### Ethics approval and consent to participate

Animal use and related experimental procedures were approved by the Institutional Animal Care and Use Committee of Hallym University (Permit Number: Hallym R2/2018-25).

### Consent for publication

Not applicable. No personal data is published in this study.

### Availability of data and materials

Not applicable

### Competing interests.

We do not have any conflict of interests.

### Funding

This research was supported by grants from the National Research Foundation (2020R1A2B5B02001806) funded by the Ministry of Science and ICT in the Republic of Korea.

### Authors' Contributions

H.J.K. designed research; M.A., A.G., B.K.P., J.K. performed experiments; H.J.K., M.A., A.G., S.M. analyzed data; H.J.K., M.A., S.M. wrote the manuscript. All authors read and approved the final manuscript.

### Acknowledgements

Not applicable

### Authors' information

<sup>1</sup>Department of Microbiology, College of Medicine, Hallym University, Chuncheon 24252, Republic of Korea

<sup>2</sup>Institute of Medical Science, College of Medicine, Hallym University, Chuncheon 24252, Republic of Korea


## References

1. Twist CJ, Beier DR, Disteché CM, Edelhoff S, Tedder TF. The mouse Cd83 gene: structure, domain organization, and chromosome localization. *Immunogenetics*. 1998;48:383–93.
2. Zhou LJ, Schwarting R, Smith HM, Tedder TF. A novel cell-surface molecule expressed by human interdigitating reticulum cells, Langerhans cells, and activated lymphocytes is a new member of the Ig superfamily. *J Immunol*. 1992;149:735–42.
3. Zhou LJ, Tedder TF. Human blood dendritic cells selectively express CD83, a member of the immunoglobulin superfamily. *J Immunol*. 1995;154:3821–35.
4. Mailliard RB, Alber SM, Shen H, Watkins SC, Kirkwood JM, Herberman RB, Kalinski P. IL-18-induced CD83 + CCR7 + NK helper cells. *J Exp Med*. 2005;202:941–53.
5. Cao W, Lee SH, Lu J. CD83 is preformed inside monocytes, macrophages and dendritic cells, but it is only stably expressed on activated dendritic cells. *Biochem J*. 2005;385 Pt1:85–93.
6. Nicod LP, Joudrier S, Isler P, Spiliopoulos A, Pache JC. Upregulation of CD40, CD80, CD83 or CD86 on alveolar macrophages after lung transplantation. *J Heart Lung Transplant*. 2005;24:1067–75.
7. Yamashiro S, Wang JM, Yang D, Gong WH, Kamohara H, Yoshimura T. Expression of CCR6 and CD83 by cytokine-activated human neutrophils. *Blood*. 2000;96:3958–63.
8. Iking-Konert C, Wagner C, Deneffle B, Hug F, Schneider M, Andrassy K, Hansch GM. Up-regulation of the dendritic cell marker CD83 on polymorphonuclear neutrophils (PMN): divergent expression in acute bacterial infections and chronic inflammatory disease. *Clin Exp Immunol*. 2002;130:501–8.
9. Reinwald S, Wiethe C, Westendorf AM, Breloer M, Probst-Kepper M, Fleischer B, Steinkasserer A, Buer J, Hansen W. CD83 expression in CD4 + T cells modulates inflammation and autoimmunity. *J Immunol*. 2008;180:5890–7.
10. Wolenski M, Cramer SO, Ehrlich S, Steeg C, Grossschupff G, Tenner-Racz K, Racz P, Fleischer B, von Bonin A. Expression of CD83 in the murine immune system. *Med Microbiol Immunol*. 2003;192:189–92.
11. Breloer M, Kretschmer B, Lüthje K, Ehrlich S, Ritter U, Bickert T, Steeg C, Fillatreau S, Hoehlig K, Lampropoulou V, Fleischer B. CD83 is a regulator of murine B cell function in vivo. *Eur J Immunol*. 2007;37:634–48.
12. Fujimoto Y, Tu L, Miller AS, Bock C, Fujimoto M, Doyle C, Steeber DA, Tedder TF. CD83 expression influences CD4 + T cell development in the thymus. *Cell*. 2002;108:755–67.
13. Tze LE, Horikawa K, Domaschütz H, Howard DR, Roots CM, Rigby RJ, Way DA, Ohmura-Hoshino M, Ishido S, Andoniou CE, Degli-Esposti MA, Goodnow CC. CD83 increases MHC II and CD86 on dendritic cells by opposing IL-10-driven MARCH1-mediated ubiquitination and degradation. *J Exp Med*. 2011;208:149–65.
14. Doebbele M, Koenig C, Krzyzak L, Seitz C, Wild A, Ulas T, Baßler K, Kopelyanskiy D, Butterhof A, Kuhnt C, Kreiser S, Stich L, Zinser E, Knippertz I, Wirtz S, Riegel C, Hoffmann P, Edinger M, Nitschke L, Winkler T, Schultze JL, Steinkasserer A, Lechmann M. CD83 expression is essential for Treg cell differentiation and stability. *JCI Insight*. 2018;3:e99712.

15. Hirano N, Butler MO, Xia Z, Ansén S, von Bergwelt-Baildon MS, Neuberger D, Freeman GJ, Nadler LM. Engagement of CD83 ligand induces prolonged expansion of CD8 + T cells and preferential enrichment for antigen specificity. *Blood*. 2006;107:1528–36.
16. Prazma CM, Yazawa N, Fujimoto Y, Fujimoto M, Tedder TF. CD83 expression is a sensitive marker of activation required for B cell and CD4 + T cell longevity in vivo. *J Immunol*. 2007;179:4550–62.
17. Breloer M, Fleischer B. CD83 regulates lymphocyte maturation, activation and homeostasis. *Trends Immunol*. 2008;29:186–94.
18. Krzyzak L, Seitz C, Ubat A, Hutzler S, Ostalecki C, Gläsner J, Hiergeist A, Gessner A, Winkler TH, Steinkasserer A, Nitschke L. CD83 Modulates B Cell Activation and Germinal Center Responses. *J Immunol*. 2016;196:3581–94.
19. Cerutti A, Cols M, Puga I. Marginal zone B cells: virtues of innate-like antibody-producing lymphocytes. *Nat Rev Immunol*. 2013;13:118–32.
20. Wong KY, Baron R, Seldon TA, Jones ML, Rice AM, Munster DJ. CD83 Antibody Inhibits Human B Cell Responses to Antigen as well as Dendritic Cell-Mediated CD4 T Cell Responses. *J Immunol*. 2018;200:3383–96.
21. Kummer M, Turza NM, Muhl-Zurbes P, Lechmann M, Boutell C, Coffin RS, Everett RD, Steinkasserer A, Prechtel AT. Herpes simplex virus type 1 induces CD83 degradation in mature dendritic cells with immediate-early kinetics via the cellular proteasome. *J Virol*. 2007;81:6326–38.
22. Yonejima A, Mizukoshi E, Tamai T, Nakagawa H, Kitahara M, Yamashita T, Arai K, Terashima T, Iida N, Fushimi K, Okada H, Yamashita T, Sakai Y, Honda M, Kaneko S. Characteristics of Impaired Dendritic Cell Function in Patients With Hepatitis B Virus Infection. *Hepatology*. 2019;70:25–39.
23. Salio M, Cella M, Suter M, Lanzavecchia A. Inhibition of dendritic cell maturation by herpes simplex virus. *Eur J Immunol*. 1999;29:3245–53.
24. Morrow G, Slobedman B, Cunningham AL, Abendroth A. Varicella-zoster virus productively infects mature dendritic cells and alters their immune function. *J Virol*. 2003;77:4950–9.
25. Kretschmer B, Lüthje K, Guse AH, Ehrlich S, Koch-Nolte F, Haag F, Fleischer B, Breloer M. CD83 modulates B cell function in vitro: increased IL-10 and reduced Ig secretion by CD83Tg B cells. *PLoS One*. 2007;2:e755.
26. Gautam A, Park BK, Kim TH, Akauliya M, Kim D, Maharjan S, Park J, Kim J, Lee H, Park MS, Lee Y, Kwon HJ. Peritoneal Cells Mediate Immune Responses and Cross-Protection Against Influenza A Virus. *Front Immunol*. 2019;10:1160.
27. Rhee JW, Kim D, Park BK, Kwon S, Cho S, Lee I, Park MS, Seo JN, Kim YS, Choi HS, Lee Y, Kwon HJ. Immunization with a hemagglutinin-derived synthetic peptide formulated with a CpG-DNA-liposome complex induced protection against lethal influenza virus infection in mice. *PLoS One*. 2012;7:e48750.
28. Shen P, Fillatreau S. Antibody-independent functions of B cells: a focus on cytokines. *Nat Rev Immunol*. 2015;15:441–51.
29. Haas KM. B-1 lymphocytes in mice and nonhuman primates. *Ann N Y Acad Sci*. 2015;1362:98–109.


30. Ju X, Silveira PA, Hsu WH, Elgundi Z, Alingcastre R, Verma ND, Fromm PD, Hsu JL, Bryant C, Li Z, Kupresanin F, Lo TH, Clarke C, Lee K, McGuire H, Fazekas de St Groth B, Larsen SR, Gibson J, Bradstock KF, Clark GJ, Hart DN. The Analysis of CD83 Expression on Human Immune Cells Identifies a Unique CD83+-Activated T Cell Population. *J Immunol.* 2016;197:4613–25.
31. Gerhard W, Mozdzanowska K, Furchner M, Washko G, Maiese K. Role of the B-cell response in recovery of mice from primary influenza virus infection. *Immunol Rev.* 1997;159:95–103.
32. Mozdzanowska K, Furchner M, Zharikova D, Feng J, Gerhard W. Roles of CD4 + T-cell-independent and -dependent antibody responses in the control of influenza virus infection: evidence for noncognate CD4 + T-cell activities that enhance the therapeutic activity of antiviral antibodies. *J Virol.* 2005;79:5943–51.
33. Lam JH, Baumgarth N. The Multifaceted B Cell Response to Influenza Virus. *J Immunol.* 2019;202:351–9.
34. Hong S, Zhang Z, Liu H, Tian M, Zhu X, Zhang Z, Wang W, Zhou X, Zhang F, Ge Q, Zhu B, Tang H, Hua Z, Hou B. B Cells Are the Dominant Antigen-Presenting Cells that Activate Naive CD4(+) T Cells upon Immunization with a Virus-Derived Nanoparticle Antigen. *Immunity.* 2018;49:695–708.e4.
35. Zabel F, Fettelschoss A, Vogel M, Johansen P, Kündig TM, Bachmann MF. Distinct T helper cell dependence of memory B-cell proliferation versus plasma cell differentiation. *Immunology.* 2017;150:329–42.
36. Gordon CL, Johnson PD, Permezel M, Holmes NE, Gutteridge G, McDonald CF, Eisen DP, Stewardson AJ, Edington J, Charles PG, Crinis N, Black MJ, Torresi J, Grayson ML. Association between severe pandemic 2009 influenza A (H1N1) virus infection and immunoglobulin G(2) subclass deficiency. *Clin Infect Dis.* 2010;50:672–8.
37. Frasca D, Diaz A, Romero M, Mendez NV, Landin AM, Blomberg BB. Effects of age on H1N1-specific serum IgG1 and IgG3 levels evaluated during the 2011–2012 influenza vaccine season. *Immun Ageing.* 2013;10:14.
38. Liu S, Yan R, Chen B, Pan Q, Chen Y, Hong J, Zhang L, Liu W, Wang S, Chen JL. Influenza Virus-Induced Robust Expression of SOCS3 Contributes to Excessive Production of IL-6. *Front Immunol.* 2019;10:1843.
39. Hocart MJ, Mackenzie JS, Stewart GA. The immunoglobulin G subclass responses of mice to influenza A virus: the effect of mouse strain, and the neutralizing abilities of individual protein A-purified subclass antibodies. *J Gen Virol.* 1989;70 Pt 9:2439–48.
40. Hocart MJ, Mackenzie JS, Stewart GA. The IgG subclass responses to influenza virus haemagglutinin in the mouse: effect of route of inoculation. *J Gen Virol.* 1989;70(Pt4):809–18.

## Figures




**Figure 1**

CD83 expression in peritoneal cells, bone marrow cells, and splenocytes after intraperitoneal injection of A/WSN/1933 virus. Cells were collected at 1, 3, 6, and 24 h after intraperitoneal injection of live A/WSN/1933 virus (WSN) or UV-inactivated A/WSN/1933 virus (UV-WSN) in C57BL/6J mice, and CD83 expression was analyzed by FACS. a CD83 expression in CD19<sup>+</sup> B cells of the peritoneal cavity after infection. b, c CD83 expression in CD19<sup>+</sup> B cells (b) and CD3<sup>+</sup> T cells (c) of the peritoneal cavity cells, bone marrow cells, and splenocytes after infection.




**Figure 2**

Population of lymphoid cells in the peritoneal cavity fluid, bone marrow, and spleen after intraperitoneal injection of A/WSN/1933 virus in C57BL/6J wild type mice and CD83 KO mice. a-c Wild type mice and CD83 KO mice (n=5/group) were injected intraperitoneally with  $5 \times 10^6$  pfu of A/WSN/1933 virus. Peritoneal cells, bone marrow cells, and splenocytes were harvested at 5, 7, and 14 days after infection. Cells were enumerated and stained with fluorescence-conjugated antibodies to be analyzed by FACS. Total cells of peritoneal cavity fluid (a), bone marrow (b), and spleen (c) were sorted into CD19<sup>+</sup> B cells and CD3<sup>+</sup> T cells for all experimental groups of mice. The left panel displays the FACS profile with the percentage of the subpopulation in each quadrant. The right panel displays the absolute number of CD19<sup>+</sup> B cells and CD3<sup>+</sup> T cells in each subgroup.



**Figure 3**

Analysis of B cells and T cells in the peritoneal cavity fluid after intraperitoneal injection of A/WSN/1933 virus in C57BL/6J wild type mice and CD83 KO mice. Wild type mice and CD83 KO mice (n=5/group) were injected intraperitoneally with  $5 \times 10^6$  pfu of A/WSN/1933 virus. Peritoneal cells were harvested at 5, 7, and 14 days after infection. a CD19<sup>+</sup> B cells were gated and further sorted into B-1 cells (CD19<sup>+</sup>, CD23<sup>-</sup>) and B-2 cells (CD19<sup>+</sup>, CD23<sup>+</sup>). b CD3<sup>+</sup> T cells were analyzed specifically for the presence of the CD4<sup>+</sup> and CD8<sup>+</sup> T cell populations. The bar diagram on the right side showed the absolute number of cells in each subgroup.



**Figure 4**

A/WSN/1933 virus-specific antibody production by C57BL/6J wild type mice and CD83 KO mice after intraperitoneal injection of A/WSN/1933 virus. Peritoneal cavity fluids and sera were collected at 5, 7, and 14 days after intraperitoneal injection of live A/WSN/1933 virus in mice (n=5/group) with a dose of  $5 \times 10^6$  pfu/mice. The amount of total IgG and IgG subclasses were quantified by ELISA. a A/WSN/1933 virus-specific antibody production in the peritoneal cavity fluid. b A/WSN/1933 virus-specific antibody production in sera.