Upregulated CDKN2A expression may be an independent protective factor in Luminal-like breast cancer

Yier Qiu (✉ qiuyierhello@163.com)

Yinzhou People's Hospital <https://orcid.org/0000-0002-3721-6644>

Guowen Lu

Yinzhou People's Hospital

Yingjie Wu

Yinzhou People's Hospital

Research

Keywords: CDKN2A, breast cancer, Prognosis, TCGA, GSEA

Posted Date: November 20th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-56061/v2>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Previous studies revealed that CDKN2A (cyclin-dependent kinase inhibitor 2A) functioned as a tumour suppressor in various types of malignant tumours. The aim of the study was to clarify the value of CDKN2A expression in the prognosis of breast cancer.

Method: Using the Cancer Genome Atlas (TCGA) database, we compared CDKN2A mRNA levels between breast cancer tissues and normal tissues and analyzed the relationship between clinical features and CDKN2A expression with the Wilcox test and the Kruskal-Wallis test. Kaplan-Meier and Cox analyses were performed to determine the correlation between CDKN2A expression and breast cancer prognosis. Gene set enrichment analysis (GSEA) was performed using the TCGA data set.

Results: We first found that CDKN2A expression was markedly higher in breast cancer tissues than in normal tissues using the TCGA database ($P=0.000$). In addition, CDKN2A mRNA expression in breast cancer was positively correlated with age ($P=0.018$), histological types ($P=0.028$), ER status ($P=0.000$), PR status ($P=0.000$) and molecular subtypes ($P=0.000$). Kaplan-Meier analysis showed that increased CDKN2A expression was associated with increased survival time in breast cancer patients ($P=0.000$), especially in Luminal-like subtype. Univariate and multivariate Cox analyses indicated that CDKN2A expression was an independent prognostic biomarker for breast cancer ($P=0.037$). GSEA suggested that pathways involving cell adhesion molecules (CAMs), cytokine-receptor interactions, cytosolic DNA sensing, the cell cycle, and killer cell-mediated cytotoxicity were differentially enriched in the CDKN2A-high expression group.

Conclusion: Our research demonstrated that high CDKN2A mRNA expression may be an independent protective factor for improved prognosis in Luminal-like breast cancer. Additionally, the signaling pathways related to CAMs, cytokine-receptor interactions, cytosolic DNA sensing, the cell cycle, and killer cell-mediated cytotoxicity regulated by CDKN2A mRNA expression should be further studied.

Background

Breast cancer (BC) has the highest incidence of any cancer not only in China (38.9% of total cases) but also worldwide (30% of total cases) and is also second most common female malignancy worldwide[1,2]. Based on the expression of different molecular biomarkers, breast cancer patients are grouped into four molecular subtypes: luminal A, luminal B, HER-2 enriched, and triple-negative BC(TNBC) [3-5]. Accordingly, dependent on molecular subtypes, there are numerous standard protocols and emerging treatments for breast cancer patients in the last few years[6,7]. However, treatment resistance, tumour metastasis and recurrence in breast cancer seriously influence patient survival and quality of life[8]. Therefore, it remains important to detect effective prognostic biomarkers and therapeutic targets for breast cancer patients.

Indeed, recent studies focus on an ideal target for breast cancer molecular targeting therapy. CDKN2A, which was situated on chromosome 9p21 and encoded the tumour suppressor protein p16[9], played a

vital role in the occurrence of melanoma in the earliest stage[10]. The function and importance of CDKN2A in different types of cancer have been frequently inferred. Steven et al. demonstrated that CDKN2A served as a tumour suppressor in mice and prevented plexiform neurofibroma progression[11]. Moreover, another study showed that increased CDKN2A immunostaining indicated increased survival time in cats with oral squamous cell carcinomas[12]. In addition, Ganjoo's study noted that CDKN2A deletion was associated with poor prognosis in histological subtypes of sarcoma[13]. Similarly, by inhibiting CDKN2A/p16, C-terminal binding protein-2 promoted proliferation and migration, as well as lower p16 expression was predictive of lower survival rate in BC patients[14]. In contrast, Lee et al. suggested that elevated CDKN2A expression in breast cancer predicted poor survival outcomes[15].

Given the controversy surrounding CDKN2A, we used TCGA data to identify the role of CDKN2A expression in breast cancer. Furthermore, GSEA was in order to recognize signalling pathways activated in breast cancer about the mechanism of CDKN2A. In our present study, we revealed that increased CDKN2A expression correlated with increased survival time in breast cancer. GSEA showed that signalling pathways associated with cell adhesion molecules (CAMs), the cell cycle, cytokine-receptor interactions, DNA replication, killer cell-mediated cytotoxicity and P53 were enriched in breast cancer patients with high CDKN2A expression.

Methods

Data mining

We downloaded CDKN2A expression data and corresponding clinical information from TCGA database (<https://portal.gdc.cancer.gov/>)[16].

GSEA

GSEA is a powerful processor that catches on several cancer-related datasets and reveals many common biological pathways[17]. In our study, GSEA was implemented to distinguish signalling pathways activated in breast cancer between high and low CDKN2A groups. Gene sets with a normal P-value <0.05 and a false discovery rate (FDR) <0.25 were considered to be significantly enriched.

Statistical analysis

Most statistical analyses were conducted using R (v.3.6.0) (a free and widely used statistical programming based on the R language environment, the NES reference line and the HR value). Differential expression of CDKN2A between normal tissues and tumour tissues in breast cancer patients was evaluated by Wilcox test. The relationship between clinicopathological parameters and CDKN2A mRNA expression was analysed with Wilcox test and Kruskal-Wallis test. Patients were assigned into two groups (high and low CDKN2A level) according to median value. Chi-square test was used to analyze associations of CDKN2A expression with clinicopathological data clinicopathological characteristics through IBM SPSS 24.0 (SPSS Inc, Chicago, USA) software. Survival analysis was displayed by Kaplan-

Meier method and log-rank test. Univariate and multivariate analysis was implemented to distinguish prognostic factors with the Cox proportional hazard regression test. P-values of < 0.05 were regarded as statistically significant.

Results

Differential expression of CDKN2A in breast cancer patients based on TCGA data

First, the CDKN2A expression data of 122 normal tissues and 1,066 tumor tissues from breast cancer patients were downloaded from TCGA in March 2019. We detected that CDKN2A levels were markedly higher in tumor tissues ($P= 0.000$) (Figure 1A). Besides, we compared CDKN2A mRNA levels in normal tissues and matched tumor tissues in breast cancer patients. Similarly, CDKN2A levels were higher in the tumor tissues ($P= 0.000$) (Figure 1B), which implies that CDKN2A may play a relevant role in breast cancer.

Clinicopathological parameters

As displayed in Table 1, we analysed Clinicopathological parameters of 832 female breast cancer patients according to clinical data downloaded from TCGA after excluding incomplete clinical prognostic information. The median age of confirmed diagnosis was 58 years old. Most of pathologic types (73.9%, $n= 615$) were of infiltrating ductal carcinoma, 19.6% ($n= 163$) were infiltrating lobular carcinoma, least were mixed histology ($n= 25$) and another specific carcinoma ($n= 29$). Moreover, clinical stage I of breast cancer was found in 143 patients (17.19%), stage II was found in 482 patients (57.93%), stage III was found in 193 patients (23.20%), and stage IV was found in 14 patients (1.68%). Besides, there were four different subtypes of breast cancer, including Luminal A (73.6%, $n= 613$), Luminal B (7.7%, $n= 64$), HER2-enriched (17.1%, $n= 142$) and Triple negative (1.6%, $n= 13$). The mean and median follow-up times after surgery were 3.2 and 2.3 years respectively. During the 23 years of follow-up in this survey, 109 (13.1%) patients died.

Table 1. Clinicopathological parameters of the breast cancer patients.

Clinicopathological parameters	N
Age	
≤58	416
>58	416
Histological type	N
Infiltrating ductal carcinoma	615
Infiltrating lobular carcinoma	163
Mixed histology	25
Another specific carcinoma	29
Tumor classification	
T1	220
T2	486
T3	104
T4	22
Lymphatic classification	
N0	396
N1	281
N2	105
N3	50
Distant classification	

M0	818
M1	14
Clinical stage	
I	143
II	482
III	193
IV	14
Estrogen receptor	
Positive	666
Negative	166
Progesterone receptor	
Positive	583
Negative	249
HER2	
Positive	77
Negative	755
Molecular subtypes	
Luminal A	613
Luminal B	64

Triple negative	142
HER2-enriched	13
Survival status	
Death	109
Survival	723

CDKN2A expression in breast cancer tissues according to clinical characteristics

We further investigated CDKN2A expression between tumour tissues and normal tissues in breast cancer patients based on clinical characteristics (Figure 2). Our results revealed that there were differences in CDKN2A expression among four molecular subtypes ($P= 0.000$) [Figure 2I]. In addition, we found that as the clinical stage increased, CDKN2A expression decreased inversely ($P= 0.491$). However, no differences were observed according to TNM stage, distant metastasis and HER2 status ($P= 0.917$, $P= 0.458$, $P= 0.5$ and $P= 0.579$ respectively).

CDKN2A expression is associated with clinicopathological features

To evaluate the clinical significance of CDKN2A, we analysed the associations between CDKN2A expression and the clinicopathological characteristics by chi-square test. CDKN2A expression was correlated with age ($P= 0.018$), histological types ($P= 0.028$), ER status ($P= 0.000$), PR status ($P= 0.000$) and molecular subtypes ($P= 0.000$), respectively. However, no significant correlation was detected in TNM stage and HER2 status (Table 2).

Table 2. Relationships of CDKN2A expression with clinical features based on TGCA data.

Variable		CDKN2A (n)		
		low	high	<i>P</i> ¹
		n=416	n=416	
Age (years)	≤58	191	225	0.018
	>58	225	191	
Histological types	Infiltrating ductal carcinoma	315	300	0.028
	Infiltrating lobular carcinoma	69	94	
	Mixed histology	18	7	
	Other specific carcinoma	14	15	
T classification	T1	113	107	0.924
	T2	239	247	
	T3	52	52	
	T4	12	10	
Lymphatic metastasis	N0	195	201	0.051
	N1	135	146	
	N2	65	40	
	N3	21	29	
Distant metastasis	M0	409	409	1.000
	M1	7	7	
Clinical stage	I	69	74	0.831

	II	238	244	
	III	102	91	
	IV	7	7	
<hr/>				
ER	Positive	364	302	0.000
	Negative	52	114	
<hr/>				
HR	Positive	316	267	0.000
	Negative	100	149	
<hr/>				
HER2	Positive	41	36	0.550
	Negative	375	380	
<hr/>				
Molecular subtypes	Luminal A	334	279	0.000
	Luminal B	35	29	
	Triple negative	41	101	
	HER2-enriched	6	7	
<hr/>				

The bold number is on behalf of the P-values with significant differences. ¹PValue was calculated by χ^2 test. ER: estrogen receptor; PR: progesterone receptor; HER2: human epidermal growth factor receptor 2.

High CDKN2A expression is an independent protective factor for OS

Survival analysis showed that high CDKN2A expression was positively associated with overall survival (OS) (P=0.000, Figure 3A). We also found that increased CDKN2A expression was related to superior overall survival, in spite of age, lymphatic metastasis and HER2 status. Beyond that, high CDKN2A expression was correlated with superior OS, especially in breast cancer patients with T2-T3 classification, clinical stage II-III and luminal-like subtype but without distant metastasis (Figure 3B-P).

Univariate Cox regression analysis showed that CDKN2A expression, age, TNM stage and HER2 status were significantly associated with OS (Table 3). Moreover, multivariate analysis confirmed that CDKN2A expression and age were independent predictors of OS in breast cancer patients (Table 3).

Table 3. Univariate and multivariate analyses in breast cancer patients

Variable	Univariate analysis			Multivariate analysis		
	HR	95% CI	Pvalue	HR	95% CI	Pvalue
CDKN2A	0.701	0.584-0.842	0.000	0.976	0.953-0.998	0.037
Age	1.033	1.018-1.049	0.000	1.030	1.014-1.046	0.000
T	1.624	1.280-2.059	0.000	1.260	0.964-1.645	0.091
N	1.624	1.334-1.978	0.000	1.267	0.999-1.607	0.051
M	3.251	1.961-5.391	0.000	1.885	0.925-3.842	0.081
STAGE	0.571	0.419-1.049	0.000	0.704	0.482-1.028	0.069
ER	1.067	0.697-1.632	0.765			
PR	1.225	0.831-1.806	0.304			
HER-2	0.501	0.312-0.811	0.004	0.743	0.441-1.253	0.265
Molecular types	1.143	0.939-1.391	0.182			

HR: Hazard ratio; CI: Confidence interval. The bold number is on behalf of P-values with significant differences.

CDKN2A-related signaling pathway

We implemented GSEA to recognize the signalling pathways activated in breast cancer by comparing the low and high CDKN2A expression datasets. GSEA showed significant differences (FDR<0.25, NOM P-value <0.05) in the enrichment of the MSigDB Collection pathways. As shown in Table 4, we chose the signalling pathways that were most significantly enriched in breast cancer patients with high CDKN2A

expression, including pathways related to bladder cancer, CAMs, the cell cycle, cytokine-receptor interactions, cytosolic DNA sensing, DNA replication, killer cell-mediated cytotoxicity and P53 (Figure 4).

Table 4. Gene sets enriched in the high CDKN2A expression.

Gene set name	NES	NOM p-val	FDR q-val
BLADDER_CANCER	1.857	0.006	0.108
DNA_REPLICATION	1.853	0.013	0.094
CELL_ADHESION_MOLECULES_CAMS	1.763	0.032	0.062
CELL_CYCLE	1.780	0.040	0.058
CYTOKINE_CYTOKINE_RECEPTOR_INTERACTION	1.656	0.042	0.086
NATURAL_KILLER_CELL_MEDIATED_CYTOTOXICITY	1.634	0.043	0.085
P53_SIGNALING_PATHWAY	1.602	0.047	0.093
CYTOSOLIC_DNA_SENSING_PATHWAY	1.556	0.048	0.114

NES: normalized enrichment score; NOM: nominal; FDR: false discovery rate. Gene sets with NOM p-val<0.05 and FDR q-val<0.25 are regarded as significant.

Discussion

Breast cancer (BC) accounts for approximately 2.1 million new cases and 62,000 deaths per year, with high mortality and morbidity rates in women worldwide[1]. Meanwhile, owing to gene expression profiles of breast cancer, there were four molecular subtypes as well as each requires different treatment[18]. Despite substantial research efforts devoted to exploring effective biomarkers for breast cancer, few basic research results have been applied in the clinical field. As a consequence, it is badly in need of recognizing truly effective biomarkers and therapeutic targets for breast cancer. Recent studies have reported that CDKN2A functions as a tumour suppressor in different types of carcinoma, such as melanoma[10], ovarian cancer[19], lung cancer[20], atypical neurofibroma[11], malignant mesothelioma[21] and bladder cancer[22]. Several studies also showed that decreased CDKN2A expression correlated to poor prognosis in osteosarcoma[23], soft tissue sarcoma[13], and oral squamous cell carcinoma[24].

With respect to the roles of CDKN2A expression in breast cancer, Feng et al found that KDM2B promoted triple-negative breast cancer cell proliferation by suppressing CDKN2A/p16 expression[25]. Fu et al. also showed that C-terminal binding protein-2 motivated proliferation and migration in breast cancer via the inhibition of CDKN2A/p16[14]. The above results suggest that reduced CDKN2A expression contributes to poor outcomes in breast cancer. However, contrasting results showed that increased expression of CDKN2A/p16 was associated with poor prognosis in breast cancer [15].

Therefore, there still exists controversy regarding the prognostic value of CDKN2A in BC, and signalling pathways related to the regulatory mechanism have not yet been determined. In our study, we first found that CDKN2A expression was markedly elevated in BC tissues than in normal tissues using TCGA database, which implied that CDKN2A may play its part in the tumorigenesis of BC. Thus, we further analysed the relationship between CDKN2A expression and clinical features of BC patients. The results showed that the CDKN2A expression was positively related to age, ER status, PR status and molecular subtypes in breast cancer. Then, the Kaplan-Meier curves for OS also showed that reduced CDKN2A expression correlated to decreased survival time, especially in Luminal-like subtype, a finding that was in contrast to the findings of Lee's study[15] but was consistent with the previously identified roles of CDKN2A in breast cancer[14,25,23]. Then Univariate and multivariate Cox analyses indicated CDKN2A expression and age were prognostic biomarkers for BC patients. Additionally, to identify the potential signalling pathways, we implemented GSEA comparing the low and high CDKN2A expression datasets and observed that pathways related to bladder cancer, CAMs, the cell cycle, cytokine-receptor interactions, cytosolic DNA sensing, DNA replication, killer cell-mediated cytotoxicity and P53 were enriched in breast cancer. A previous study showed that loss of CDKN2A negatively regulated with the cell cycle and CDK4/6/RB pathway which was shown to promote breast cancer cell proliferation in Luminal-like breast cancer patients [26]. While the CDK4/6 inhibitor was preferentially effective in inhibiting the cell proliferation of ER+ cancer cells in preclinical studies[27]. For the sake of perspective, CDKN2A overexpression may inhibit breast cancer cells growth which should confirmed in vivo and in vitro experiment. Meanwhile, another study pointed out that cancer gene therapy mediated by CDKN2A was beneficial for the induction of cell death in non-small-cell lung carcinoma[20]. In addition, we found CDKN2A mRNA expression was associated with bladder cancer, the cell cycle and DNA replication[28-30]. However, it is the first time that the results in this paper showing correlations between CDKN2A expression and signalling pathways involving CAMs, cytokine-receptor interactions, cytosolic DNA sensing and killer cell-mediated cytotoxicity are reported, as well as biological mechanism needs to be further clarified.

Nonetheless, there were several limitations to our study that should be noted. First, more large-scale studies should be required to further improve the reliability of our findings. Second, the clinical information from the TCGA database was not comprehensive, which not included the latest molecular classification of breast cancer. Finally, we did not explore the exactly underlying molecular mechanisms in this study, which will be elucidated in future researches.

Conclusions

In conclusion, upregulated CDKN2A expression may be an effective prognostic biomarker in luminal-like breast cancer suggesting longer survival times. However, we should focus on the association between luminal-like subtypes and CDKN2A to inform the current controversial results. Additionally, signalling pathways related to CAMs, cytokine-receptor interactions, cytosolic DNA sensing and killer cell-mediated cytotoxicity may be pivotal pathways regulated by CDKN2A in breast cancer, and these relationships should be further clarified.

Abbreviations

CDKN2A: cyclin dependent kinase inhibitor 2A; TGCA: The Cancer Genome Atlas; GSEA: Gene Set Enrichment Analysis; OS: overall survival; HR: Hazard ratio; CI: Confidence interval; NES: normalized enrichment score; NOM: nominal; FDR: false discovery rate

Declarations

Acknowledgements

None.

Authors' contributions

YEQ and YJW participated in the study design. WJX and GWL contributed to data collection and analysis. All authors were involved in the writing of the article. YEQ critically reviewed the manuscript. All authors have read and approved the final manuscript.

Funding

None.

Availability of data and materials

The datasets analyzed during the current study are available in the TGCA (<https://portal.gdc.cancer.gov/>).

Ethics approval and consent to participate

None.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Siegel RL, Miller KD, Jemal A (2018) Cancer statistics, 2018. CA Cancer J Clin 68 (1):7-30. doi:10.3322/caac.21442
2. Feng RM, Zong YN, Cao SM, Xu RH (2019) Current cancer situation in China: good or bad news from the 2018 Global Cancer Statistics? Cancer Commun (Lond) 39 (1):22. doi:10.1186/s40880-019-0368-6

3. Charles M. Perou TSr, Michael B. Eisen, Matt van de Rijnß, Stefanie S. Jeffreyk, Christian A. Rees, Jonathan R. Pollackð, Douglas T. Rossð, Hilde Johnsen, Lars A. Akslen, Eystein Flugel, Alexander Pergamenschikov, Cheryl Williams, Shirley X. Zhuß, Per E. Lønning, Anne-Lise Børresen-Dale, Patrick O. Brown & David Botstein (2000) Molecular portraits of human breast tumours. *Nature* 406 (6797):747-752
4. Therese Sørliea CMP, Robert Tibshiranie, Turid Aasf, SG, Hilde Johnsenb, Trevor Hastiee, Michael B. Eisenh, Matt van de Rijni, Stefanie S. Jeffreyj, Thor Thorsenk, Hanne Quistl, JCM, Patrick O. Brownm, David Botsteinc, Per Eystein Lønningg, and Anne-Lise Børresen-Daleb, n (2001) Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications. *Proc Natl Acad Sci US A* 98 (19):10869-10874.
5. Goutsouliak K, Veeraraghavan J, Sethunath V, De Angelis C, Osborne CK, Rimawi MF, Schiff R (2019) Towards personalized treatment for early stage HER2-positive breast cancer. *Nat Rev Clin Oncol*. doi:10.1038/s41571-019-0299-9
6. Lehmann BD, Bauer JA, Chen X, Sanders ME, Chakravarthy AB, Shyr Y, Pietenpol JA (2011) Identification of human triple-negative breast cancer subtypes and preclinical models for selection of targeted therapies. *J Clin Invest* 121 (7):2750-2767. doi:10.1172/JCI45014
7. Basu A, Ramamoorthi G, Jia Y, Faughn J, Wiener D, Awshah S, Kodumudi K, Czerniecki BJ (2019) Immunotherapy in breast cancer: Current status and future directions. *Adv Cancer Res* 143:295-349. doi:10.1016/bs.acr.2019.03.006
8. Harbeck N, Penault-Llorca F, Cortes J, Gnant M, Houssami N, Poortmans P, Ruddy K, Tsang J, Cardoso F (2019) Breast cancer. *Nat Rev Dis Primers* 5 (1):66. doi:10.1038/s41572-019-0111-2
9. Andre L. Reed JC, Paul Cairns, William H. Westra, Richard M. Jones, Wayne Koch, Steven Ahrendt, Yolanda Eby, Duane Sewell, Homaira Nawroz, Jiri Bartek and David Sidransky (1996) High Frequency of p16 (CDKN2/MTS-1/INK4A) Inactivation in Head and Neck Squamous Cell Carcinoma. *CANCER RESEARCH* 56:3630-3633
10. William D. Foulkes TYF, * Pamela M. Pollock,t, Haywardt aNK (1997) THE CDKN2A(pl6) GENE AND HUMAN CANCER. *Molecular Medicine* 3 (1):15-20
11. Rhodes SD, He Y, Smith A, Jiang L, Lu Q, Mund J, Li X, Bessler W, Qian S, Dyer W, Sandusky GE, Horvai AE, Armstrong AE, Clapp DW (2019) Cdkn2a (Arf) loss drives NF1-associated atypical neurofibroma and malignant transformation. *Hum Mol Genet* 28 (16):2752-2762. doi:10.1093/hmg/ddz095
12. Munday JS, He Y, Aberdein D, Klobukowska HJ (2019) Increased p16(CDKN2A), but not p53, immunostaining is predictive of longer survival time in cats with oral squamous cell carcinomas. *Vet J* 248:64-70. doi:10.1016/j.tvjl.2019.04.007
13. Bui NQ, Przybyl J, Trabucco SE, Frampton G, Hastie T, van de Rijn M, Ganjoo KN (2019) A clinico-genomic analysis of soft tissue sarcoma patients reveals CDKN2A deletion as a biomarker for poor prognosis. *Clin Sarcoma Res* 9:12. doi:10.1186/s13569-019-0122-5

14. Xiaojing Yang¹ YS, Hongling Li¹, Yuhui Shao¹, Depeng Zhao², Weiwei Yu¹, Jie Fu¹ (2017) C-terminal binding protein-2 promotes cell proliferation and migration in breast cancer via suppression of p16INK4A. *Oncotarget* 8 (16):26154-26168
15. Pare R, Shin JS, Lee CS (2016) Increased expression of senescence markers p14(ARF) and p16(INK4a) in breast cancer is associated with an increased risk of disease recurrence and poor survival outcome. *Histopathology* 69 (3):479-491. doi:10.1111/his.12948
16. Weisenberger DJ (2014) Characterizing DNA methylation alterations from The Cancer Genome Atlas. *J Clin Invest* 124 (1):17-23. doi:10.1172/JCI69740
17. A. Subramanian PT, V.K. Mootha, S. Mukherjee, B.L. Ebert, M.A. Gillette, et al. (2005) Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profile. *Proc Natl Acad Sci* 102:15545–15550
18. Nadia Harbeck FP-L, Javier Cortes, Michael Gnant, Nehmat Houssami⁶, Philip Poortmans, Kathryn Ruddy, Janice Tsang and Fatima Cardoso (2019) Breast cancer. *NATURE REVIEWS | DISEASE PRIMERS* 5 (66):1-31
19. Xia L, Zhang W, Gao L (2019) Clinical and prognostic effects of CDKN2A, CDKN2B and CDH13 promoter methylation in ovarian cancer: a study using meta-analysis and TCGA data. *Biomarkers* 24 (7):700-711. doi:10.1080/1354750X.2019.1652685
20. Xande JG, Dias AP, Tamura RE, Cruz MC, Brito B, Ferreira RA, Strauss BE, Costanzi-Strauss E (2019) Bicistronic transfer of CDKN2A and p53 culminates in collaborative killing of human lung cancer cells in vitro and in vivo. *Gene Therapy*. doi:10.1038/s41434-019-0096-1
21. Kukuyan AM, Sementino E, Kadariya Y, Menges CW, Cheung M, Tan Y, Cai KQ, Slifker MJ, Peri S, Klein-Szanto AJ, Rauscher FJ, 3rd, Testa JR (2019) Inactivation of Bap1 Cooperates with Losses of Nf2 and Cdkn2a to Drive the Development of Pleural Malignant Mesothelioma in Conditional Mouse Models. *Cancer Res* 79 (16):4113-4123. doi:10.1158/0008-5472.CAN-18-4093
22. Garcia-Perdomo HA, Usubillaga-Velasquez JP, Zapata-Copete JA, Reis LO (2019) Mutations in CDKN2A and the FGFR3 genes on bladder cancer diagnosis: a systematic review and meta-analysis. *World J Urol* 37 (10):2001-2007. doi:10.1007/s00345-019-02779-7
23. Zhang H, Mao JS, Hu WF (2019) Functional Genetic Single-Nucleotide Polymorphisms (SNPs) in Cyclin-Dependent Kinase Inhibitor 2A/B (CDKN2A/B) Locus Are Associated with Risk and Prognosis of Osteosarcoma in Chinese Populations. *Med Sci Monit* 25:1307-1313. doi:10.12659/MSM.915001
24. Padhi SS, Roy S, Kar M, Saha A, Roy S, Adhya A, Baisakh M, Banerjee B (2017) Role of CDKN2A/p16 expression in the prognostication of oral squamous cell carcinoma. *Oral Oncol* 73:27-35. doi:10.1016/j.oraloncology.2017.07.030
25. Zheng Q, Fan H, Meng Z, Yuan L, Liu C, Peng Y, Zhao W, Wang L, Li J, Feng J (2018) Histone demethylase KDM2B promotes triple negative breast cancer proliferation by suppressing p15INK4B, p16INK4A, and p57KIP2 transcription. *Acta Biochim Biophys Sin (Shanghai)* 50 (9):897-904. doi:10.1093/abbs/gmy084

26. Araki K, Miyoshi Y (2018) Mechanism of resistance to endocrine therapy in breast cancer: the important role of PI3K/Akt/mTOR in estrogen receptor-positive, HER2-negative breast cancer. *Breast Cancer* 25 (4):392-401. doi:10.1007/s12282-017-0812-x
27. Richard S Finn JD, Dylan Conklin, Ondrej Kalous, David J Cohen, Amrita J Desai, Charles Ginther, Mohammad Atefi, Isan Chen, Camilla Fowst, Gerret Los and Dennis J Slamon (2009) PD 0332991, a selective cyclin D kinase 4/6 inhibitor, preferentially inhibits proliferation of luminal estrogen receptor-positive human breast cancer cell lines in vitro. *Breast Cancer Research* 11 (5):1-13. doi:10.1186/bcr2419)
28. Worst TS, Weis CA, Stohr R, Bertz S, Eckstein M, Otto W, Breyer J, Hartmann A, Bolenz C, Wirtz RM, Erben P (2018) CDKN2A as transcriptomic marker for muscle-invasive bladder cancer risk stratification and therapy decision-making. *Sci Rep* 8 (1):14383. doi:10.1038/s41598-018-32569-x
29. Pal A, Potjer TP, Thomsen SK, Ng HJ, Barrett A, Scharfmann R, James TJ, Bishop DT, Karpe F, Godsland IF, Vasen HF, Newton-Bishop J, Pijl H, McCarthy MI, Gloyn AL (2016) Loss-of-Function Mutations in the Cell-Cycle Control Gene CDKN2A Impact on Glucose Homeostasis in Humans. *Diabetes* 65 (2):527-533. doi:10.2337/db15-0602
30. Gadhikar MA, Zhang J, Shen L, Rao X, Wang J, Zhao M, Kalu NN, Johnson FM, Byers LA, Heymach J, Hittelman WN, Udayakumar D, Pandita RK, Pandita TK, Pickering CR, Redwood AB, Piwnica-Worms H, Schlacher K, Frederick MJ, Myers JN (2018) CDKN2A/p16 Deletion in Head and Neck Cancer Cells Is Associated with CDK2 Activation, Replication Stress, and Vulnerability to CHK1 Inhibition. *Cancer Res* 78 (3):781-797. doi:10.1158/0008-5472.CAN-17-2802

Figures

Figure 1

CDKN2A expression in breast cancer patients based on TCGA data. A: Scatter gram represents CDKN2A expressions from normal to tumor tissues, respectively; B: A dot plot with connecting lines represents CDKN2A expressions from normal to matched tumor tissues. P-values of < 0.05 were considered significant.

Figure 2

CDKN2A expression on the basis of TCGA data. A-I: CDKN2A expression was compared according to patient age, T classification, lymph node, distant metastasis and clinical stage, respectively.

Figure 3

Survival curves on TCGA data. (A): Comparisons of the overall survival duration between the low and high CDKN2A expression; (B-C): Independent of patient age; (D-E): Independent of T classifications; (F-G): Independent of lymphatic metastasis; (H): Independent of distant metastasis; (I-J): Independent of clinical stage; (K-P): Independent of ER status, PR status, HER2 and molecular subtypes.

Figure 4

Enrichment plots from GSEA. Different pathways including bladder cancer (A), cell adhesion molecules cams (B), cell cycle (C), cytokine receptor interaction (D), cytosolic DNA sensing cell cycle (E), DNA replication (F), killer cell mediated cytotoxicity (G) and P53 signaling (H) are enriched in breast cancer patients with high CDKN2A expression according to GSEA. (I): Sum of 8 enriched pathways in breast cancer patients with high CDKN2A expression.