

Single blinded semi-field evaluation of MAÏA® topical repellent ointment compared to unformulated 20% DEET against *Anopheles gambiae* s.s., *Anopheles arabiensis* and *Aedes aegypti* in Tanzania

Emmanuel Mbuba (✉ embuba@ihi.or.tz)

Ifakara Health Institute <https://orcid.org/0000-0003-0157-1780>

Olukayode Odufuwa

Ifakara Health Institute

Frank Tenywa

Ifakara Health Institute

Rose Philipo

Ifakara Health Institute

Mgeni Tambwe

Ifakara Health Institute

Johnson Kyeba Swai

Ifakara Health Institute

Jason Moore

Ifakara Health Institute

Sarah Moore

Ifakara Health Institute

Research

Keywords: Malaria, mosquito, topical repellent, ointment, protective efficacy, complete protection time, CPT, DEET

Posted Date: September 29th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-56100/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Malaria Journal on January 6th, 2021. See the published version at <https://doi.org/10.1186/s12936-020-03461-9>.

Abstract

Background: N,N-Diethyl-3-methylbenzamide (DEET) topical mosquito repellents are effective personal protection tools. However, DEET-based repellents tend to have low consumer acceptability because they are cosmetically unappealing. More attractive formulations are needed to encourage regular user compliance. This study evaluated the protective efficacy and protection duration of a new topical repellent ointment containing 15% DEET, MAÏA[®] compared to 20% DEET in ethanol using malaria and dengue mosquito vectors in Bagamoyo Tanzania.

Methods: Fully balanced 3x3 Latin square design studies were conducted in large semi-field chambers using laboratory strains of *Anopheles gambiae* s.s., *An. arabiensis* and *Aedes aegypti*. Human volunteers applied either MAÏA[®] ointment, 20% DEET or ethanol to their lower limbs six hours before the start of tests. Approximately 100 mosquitoes per strain per replicate were released inside each chamber, with 25 mosquitoes released at regular intervals during the collection period to maintain adequate biting pressure throughout the test. Volunteers recaptured mosquitoes landing on their lower limbs for six hours over a period of six to 12-hours post-application of repellents. Data analysis was conducted using mixed-effects logistic regression.

Results: The protective efficacy of MAÏA[®] and 20% DEET was not different for each of the mosquito strains: 95.9% vs 97.4% against *An. gambiae* (OR=1.53 [95% CI: 0.93–2.51] p=0.091); 96.8% vs 97.2% against *An. arabiensis* (OR =1.08 [95% CI: 0.66 –1.77] P=0.757); 93.1% vs 94.6% against *Ae. aegypti* (OR=0.76 [95% CI: 0.20-2.80] p=0.675). Average complete protection time (CPT) of MAÏA[®] and that of DEET was similar for each of the mosquito strains: 571.6 minutes (95% CI: 558.3-584.8) vs 575.0 minutes (95% CI: 562.1-587.9) against *An. gambiae*; 585.6 minutes (95% CI: 571.4-599.8) vs 580.9 minutes (95% CI: 571.1-590.7) against *An. arabiensis*; 444.1 minutes (95% CI: 401.8-486.5) vs 436.9 minutes (95% CI: 405.2-468.5) against *Ae. aegypti*.

Conclusions: MAÏA[®] repellent ointment provides complete protection for 9 hours against both *An. gambiae* and *An. arabiensis*, and 7 hours against *Ae. aegypti* similar to 20% DEET (in ethanol). MAÏA[®] repellent ointment can be recommended to be used as a tool for prevention of outdoor biting mosquitoes in tropical locations as it protects for more than 6 hours.

Background

The use of insecticide-treated bed nets (ITNs) and indoor residual sprays (IRS) has almost halved malaria burden throughout sub-Saharan Africa [1, 2]. However, a considerable proportion of malaria transmission occurs outside of sleeping hours when ITNs are not in use [3]. Topical mosquito repellents are one of the oldest interventions used to prevent contact between humans and mosquito vectors [4, 5], and may be useful for people who spend their time outdoors in the evening or overnight for occupational or social activities [6-9]. Some studies have shown that people who use topical repellents consistently in addition to ITNs are protected from malaria [10, 11]. However, their utility as a disease prevention tool is limited by

poor user compliance [12]. Therefore, deployment of topical mosquito repellents for malaria prevention is not recommended by the World Health Organisation (WHO) as an intervention with public health value but maybe beneficial as an intervention to provide personal protection against malaria [13, 14].

Topical repellents must be applied daily or even several times a day, and poor user compliance is a major limitation to the effectiveness of topical repellents [15-17]. Common reasons for poor compliance include forgetting to apply the repellent [15, 18], poor acceptability including unpleasant smell or greasy “feel” on the skin [18] and the perception that repellents are poisonous [15]. DEET is a common active ingredient of most of the topical mosquito repellents present in the market today. Over 200 million persons use DEET-based repellents every year with the market growing annually due to increases in vector borne diseases such as Dengue and Zika [19]. DEET has an excellent safety profile and is safe for use among children and pregnant women [20, 21].

According to Carroll *et al*, “a good repellent ointment should be effective against target vector strain, easy to apply and has a nice odour and the residual feeling after application” [22]. To obtain better compliance, consideration of human customs and behaviour that may encourage consistent use of repellents is essential. In a study conducted in Burkina Faso, it was observed that about 91% of children under 5–years werewashed in the evening and 80% of them receive ointment on their skin after bathing and before bedtime (Traoré *et al*; *manuscript under preparation*). During this time, mosquitoes are actively host-seeking and interacting with humans outdoors [23-25]. Therefore, a well-formulated topical mosquito repellent with skin softening properties for daily use after bathing may improve user compliance as well as protect against vector-borne diseases such as dengue and malaria when people are outdoors in the early evening.

The Maïa Africa SAS [26], a company based in Burkina Faso worked in collaboration with local mothers to develop MAÏA[®] repellent ointment that is formulated with , shea butter, cotton oil, beeswax, fragrance, and 15% DEET. Shea butter-based ointments are widely used for skin softening purposes [27, 28] and MAÏA[®] repellent is designed to repel mosquitoes in addition to softening the human skin. This study evaluated the protective efficacy and duration of repellency of MAÏA[®] repellent ointment comparing with the gold standard, unformulated 20% DEET against two species of *Anopheles* i.e. *An. gambiae* and *An. arabiensis* and *Aedesaegypti* mosquitoes under semi-field system conditions in Bagamoyo Tanzania. The semi-field evaluation of topical repellents generates data that are comparable to a full-field evaluation [29].

Methods

Study area

This study was conducted under ambient conditions in a semi-field system measuring 29 x 21 meters built from a fabricated greenhouse frame modified to make two compartments with a central corridor and an opaque polyethylene roof for rain protection (Fig. 1) [29]. The semi-field system (SFS) is located at 6°

8° S, 30° 37' E at the Ifakara Health Institute in Bagamoyo district in Tanzania. Bagamoyo district experiences annual rainfall between 800 mm and 1000 mm, temperatures between 22°C and 33°C, and mean relative humidity of 73%. This evaluation followed the WHO Guidelines for efficacy testing of topical repellents [30].

Study design

This study was divided into two parts to accommodate the circadian rhythm of mosquitoes investigated. *An. gambiae* s.s. and *An. arabiensis* were tested at night and *Ae. aegypti* in the early morning. A pilot study for the first six hours after repellent application was conducted and found that a longer testing period was required. The final study tested the repellent from six to and twelve hours post-application.

A study for *Anopheles*

In the pilot study of *Anopheles* strains, repellents were applied at 17:45 hours and mosquito collection was conducted between 18:00 hours to 00:00 hours (Fig. 2A). In the final study, repellents were applied at 17:45 hours and mosquito collection was conducted between 00:00 hours to 06:00 hours (i.e. 6-12 hours after the application of repellents) (Fig. 2B).

Both the pilot and final study consisted of two fully balanced (3x3) Latin squares (LS) design conducted in two chambers of the SFS simultaneously over nine nights using six volunteers. In each LS, three volunteers rotated sequentially between the three mosquito collection positions each day inside the chamber and swapped repellents after every 3-days. After nine days of the study period, each volunteer had tested each of the repellents at each of the three mosquito collection positions inside the SFS chamber three times. The study flow plan for *An. gambiae* s.s. and *An. arabiensis* is shown in (Fig. 2).

A study for *Aedes*

For *Ae. aegypti* repellents were applied at 05:50 hours in the pilot study and at 23:45 hours in the final study. Mosquito collection was conducted from 06:00 hours to 10:00 hours in both the pilot and final study. The pilot study consisted of two fully balanced (3x3) LS conducted in two chambers of the semi-field system simultaneously over nine nights using six volunteers. In the final study, one fully balanced (3x3) LS was conducted in one chamber over nine nights using three additional volunteers. In both studies, volunteers rotated sequentially between three collection positions each day inside the chamber of the SFS and switched repellents after every 3-days. After the nine-days study, each volunteer had tested each of the repellents at each of the three mosquito collection positions inside the SFS chamber three times. The study flow plan for *Ae. aegypti* is shown in (Fig. 3).

Mosquito strains

This study used nulliparous female laboratory-reared mosquitoes, aged 5-7 days old, sugar starved and reactive to human odour. Mosquito strains used were pyrethroid-resistant (20% mortality with pyrethroid) *An.arabiensis* (Kingani, colonized Tanzania 2006), fully pyrethroid susceptible *An. gambiae* (Kisumu,

colonized Kenya 1975) and fully pyrethroid susceptible *Ae. aegypti* (Bagamoyo, colonized Tanzania 2018). Mosquitoes were reared following the MR4 guideline [31]. Before the experiment, *An. arabiensis* were lightly marked by placing them in a cup coated with a fluorescent dye to make them distinguishable from the morphologically identical *An. gambiae*. By very lightly marking the mosquitoes there is no significant effect on their fitness nor host preference [32]. Mosquitoes were then sugar-starved for 8 hours. About 30 minutes before the experiment, 100 female mosquitoes that were responsive to human odour were selected and transported in boxes to the SFS chambers to acclimatize with the ambient environmental conditions.

Repellents tested

MAÏA® ointment and 20% DEET were shipped to Ifakara Health Institute (IHI) Vector Control Product Testing Unit (VCPTU) in plastic jars by MAÏA Africa SAS [26]. The amount of MAÏA® repellent ointment and DEET 97% (reference number 26028, lot number 2436308) received at IHI was 600 and 200 ml respectively. After the products were received, they were stored the same day at room temperature between 25° and 29°C until used in the experiment. The 20% DEET in ethanol (V/V) was prepared in-house before the experiment.

Volunteers

Nine male volunteers between the age of 24 to 30 years were recruited after signing informed consent forms written in Swahili. All volunteers were tested for malaria parasite infection using SD BIOLINE Malaria Ag P.f [33] rapid malaria diagnostic kits before participating in the study and once per week during the study period as part of IHI health and safety procedures. Only male volunteers were recruited for cultural reasons as women would be expected to work overnight with men.

Allocation of volunteers

At the beginning of the study (pilot study), six male volunteers were assigned into two groups of 3 volunteers. One group was assigned to chamber '1' and the other in chamber '2' of the SFS (Fig. 1). Inside each chamber, three mosquito recapture positions were marked each 9 meters from the mosquito releasing point (Fig. 4). Each volunteer was randomly assigned to one of the three mosquito recapture positions inside one chamber and rotated between positions nightly. After nine days of experiments, it was discovered that the complete protection time of both repellents was above six hours. Therefore, the final study of 12hours recapture period was set up with an additional three volunteers recruited to run the *Ae aegypti* which was conducted immediately after the *Anopheles* experiment.

Application of repellents

Volunteers wore shorts and washed their lower limbs using water without soap before starting the experiment and they also wore closed shoes and a mesh bug jacket to ensure mosquitoes have access to the lower limbs only. Volunteers were non-smokers, and were requested to not drink alcohol or use

perfumed soaps or ointment during the entire study period. We calculated a lower limb-skin surface area for each volunteer using the following formula at the beginning of the study; $\text{Area} = \frac{1}{2} (K+A) \times L$

Where “L” represents the leg length between the knee and the ankle and “K” represents the circumference at the knee and “A” represents the circumference of the ankle area. The average lower limb skin-surface area of volunteers was 1259.2 cm². The amount of repellents used on the day of experiment was measured using Ohaus CS200 weighing scale (Ohaus Corporation, USA). The average amount of MAÏA® repellent ointment applied per limb was 2.52 grams corresponding to a target dose of 2 mg/cm² (approximately 0.3 mg of DEET per cm²). The average amount of 20% DEET in ethanol applied per limb was 2.4g corresponding to 1.9mg/cm² (approximately 0.38 mg of DEET per cm²). All volunteers applied repellents using latex-gloved hands to minimize absorption onto the hands. Empty plastic cups that contained MAÏA® ointment were weighed to determine the amount left in the cups after application. Repellents for the *Anopheles* study were applied at 17:45 hours and for the *Aedes* study at 05:50 hours during the pilot study, while during the final study repellents were applied at 17:45 hours for the *Anopheles* study and at 23:45 hours for the *Aedes* study. In the final study, all participants rested with their trousers rolled up to prevent abrasion of the repellents after repellent application.

Duration of the study

Six-hour pilot test

During the first 9 days, 50 *An. arabiensis* and 50 *An. gambiae* were released in each of the SFS chambers and testing was conducted for 6 hours. During the 6 hours recapture period, no mosquitoes were recaptured by volunteers who applied MAÏA® and 20% DEET. Therefore, we extended the recapture period to 12-hours in order to confidently determine the duration of complete protection of MAÏA® repellent ointment and 20% DEET.

Twelve-hour test

During the LS with 12-hours of recapture period, the same volunteers, chambers and rotation schedule was used for *Anopheles* experiment: 100 *An. arabiensis* and 100 *An. gambiae* were released (25 per release) starting at 00:00 hours. Three additional volunteers were recruited following study procedure and were assigned to the *Ae. aegypti* experiments in which 100 *Ae. aegypti* were released with 25 released every hour between 06:00 and 10:00hours.

Mosquito recapture

Volunteers recorded the time of a first mosquito recapture of each experiment (*Anopheles* or *Aedes*) and placed in a separate cup labelled with the time of recapture, volunteer’s code, position and repellent (treatment) code. Volunteers collected subsequent mosquitoes that landed for a 6-hour collection period up to 12-hours post-application of repellent with cups labelled with repellent code, position and hour of collection. Cups were changed after every hour. At the end of recapture time, mosquitoes were killed by

refrigeration at 4°C for about 40 minutes and then sorted to species level. If fewer than 50% of mosquitoes were recaptured by a negative control volunteer, the data were discarded and the replicate was repeated. Temperature, relative humidity and wind speed were recorded on the day at the beginning of the experiment.

Data management and statistical analysis

Data were recorded in paper forms and then double entered and cleaned in Microsoft Excel 2016. Data analyses were performed in Stata 15.1 (Stata Corp, USA). Descriptive analyses of mosquitoes recaptured by repellents was performed. The mean complete protection time (CPT) of each repellent for each mosquito strain was estimated using the Kaplan-Meier survival analysis. The protective efficacy (PE) was established for the data collected up to 12-hours and calculated using this formula; $P = ((C - T)/C) * 100$; where 'P' represents the percentage protection, 'C' represents the number of mosquitoes recaptured on the negative control (ethanol) and 'T' represents the number of mosquitoes recaptured on volunteer's lower limbs treated with either MAIA[®] or 20% DEET.

Statistical analysis was performed using a mixed-effects binary logistic regression to compare the protective efficacy between MAIA[®] and 20% DEET (as the reference in the statistical model). Several models were tested using the proportion of recaptured mosquitoes as the outcome variable and repellent type (treatments), volunteer, position of the volunteer and time of recapture as fixed effects and day of test as random effects. The best-fit model was determined using the Akaike's Information Criterion (AIC) and the model with the smallest AIC value was selected. In the *Anopheles* data, repellent and time after application were fixed effects and the day of test was a random effect. In the *Aedes* data, the type of repellent, time of recapture and position of the volunteer were fixed effects and the day of test was a random effect.

Results

General test conditions

Environmental conditions for *Anopheles* experiment temperature was 26.5°C (95% CI: 26.4–26.6) temperature, 82.96% (95% CI: 82.4–83.5) relative humidity, and 0.36 m/s (95% CI: 0.3–0.4) wind speed, and for *Ae. aegypti* experiment, temperature was 24.6°C (95% CI: 24.5–24.7), relative humidity 59.45% (95% CI: 55.7–63.2), and wind speed 0.00 m/s (95% CI: 0.0–0.0). All tests were conducted with recapture rate in the negative control arm exceeding 50%.

Descriptive analysis

The percentage recapture for *An. gambiae* was 69.9% (1258/1800) and that of *An. arabiensis* was 75.4% (1358/1800) during the 12 hour tests. The percentage recapture for *Ae. aegypti* was 88.9% (800/900) (Table 1). The geometric mean (GM) hourly mosquito landings are shown in Table 1. The GM hourly landings on volunteers who applied absolute ethanol was 9.46 (95% CI: 8.4 – 10.6) for *An. gambiae*, 10.2

(95% CI: 9.0–11.5) for *An. arabiensis*, and 10.0 (95% CI: 8.3–12.1) for *Ae. aegypti*. There was no variability between volunteers in terms of mosquito attractiveness but most users preferred using MAÏA® repellent ointment than 20% DEET (in ethanol).

The protective efficacy of MAÏA® and 20% DEET

For the recapture period, 6-12 hours post-application, both MAÏA® and 20% DEET provided greater than 93% protective efficacy against all strains (Table 1). There was no significant difference in the protective efficacy (PE) between the MAÏA® repellent ointment and the unformulated 20% DEET (in ethanol) over the 12-hour test period for any of the mosquito strains tested. For *An. gambiae* s.s., MAÏA® PE was 95.9% (95% CI: 95.4 - 96.3) and 20% DEET, PE was 97.4% (95% CI: 97.1-97.6), OR= 1.53 [95% CI: 0.93 – 2.51] p=0.091. For *An. arabiensis*, MAÏA® PE was 96.8% (95% CI: 96.3 - 97.3) and 20% DEET, PE was 97.2% (95% CI: 96.9-97.4), OR =1.08 [95% CI: 0.57 – 2.04] p=0.757. For *Ae. aegypti*, MAÏA® PE was 93.1% (95% CI: 92.2- 94.1) and 20% DEET, PE was 94.6% (95% CI: 93.8-95.4), OR=0.76 [95% CI: 0.20 – 2.80] p=0.675 (Table 1).

Complete protection time (minutes) of MAÏA® and 20% DEET

MAÏA® repellent ointment and unformulated 20% DEET had similar complete protection time (CPT) exceeding 9 hours (Table 1, Fig. 5). The average CPT (minutes) of MAÏA® was 571.6 (95% CI: 558.3-584.8) and 20% DEET was 575.0 (95% CI: 562.1-587.9) against *An. gambiae* s.s.. Average CPT (minutes) of MAÏA® was 585.6 (95% CI: 571.4-599.8) and 20% DEET was 580.9 (95% CI: 571.1-590.7) for *An. arabiensis*. Average CPT (minutes) was 444.1 (95% CI: 401.8-486.5) and 20% DEET was 436.9 (95% CI: 405.2-468.5) for *Ae. aegypti* (Fig. 5).

Table 1: The percentage (%) recapture, geometric mean of hourly mosquito landings, percentage protection, mean complete protection time (CPT) and odds ratio between MAÏA® repellent ointment 20% DEET and absolute ethanol.

Test systems (mosquito strain)	Test items (repellents)	Percentage recapture	Geometric mean hourly landings	Percentage protection (PE) and 95% CI	Odds Ratio	P value	95% Confidence Interval	Mean CPT in minutes (95% CI)
Susceptible <i>An. gambiae</i> s.s.	20% DEET	2.5% (31/1258)	1.21 (1.05 - 1.40)	97.4% (97.1 - 97.6)	1	-	-	575.0 (562.1 - 587.9)
	MAÏA® Ointment	3.9% (49/1258)	1.57 (1.28 - 1.94)	95.9% (95.4 - 96.3)	1.53	0.091	0.93 - 2.51	571.6 (558.3 - 584.8)
	Absolute Ethanol	93.6% (1178/1258)	9.46 (8.43 - 10.60)	-	62.63	<0.0000	40.04 - 97.97	-
Resistant <i>An. arabiensis</i>	20% DEET	2.7% (36/1357)	1.40 (1.15 - 1.71)	97.2% (96.9 - 97.4)	1	-	-	580.9 (571.1 - 590.7)
	MAÏA® Ointment	3% (40/1357)	1.52 (1.18 - 1.96)	96.8% (96.3 - 97.3)	1.08	0.757	0.66 - 1.77	585.6 (571.4 - 599.8)
	Absolute Ethanol	94.4% (1281/1357)	10.17 (9.01 - 11.48)	-	58.93	<0.0000	38.72 - 89.71	-
<i>Ae. aegypti</i>	20% DEET	4.8% (38/800)	3.36 (2.35 - 4.82)	94.6% (93.8 - 95.4)	1	-	-	436.9 (405.2 - 468.5)
	MAÏA® Ointment	6% (48/800)	2.47 (1.7 - 3.5)	93.1% (92.2 - 94.1)	0.76	0.675	0.20 - 2.80	444.1 (401.8 - 486.5)
	Absolute Ethanol	89.3% (714/800)	10.0 (8.30 - 12.12)	-	14.87	<0.0000	6.99 - 31.65	-

Discussion

DEET-based topical mosquito repellents provide personal protection against numerous arthropods of medical importance including malaria vectors, such as *An. gambiae* and *An. arabiensis* and the dengue vector *Aedes aegypti* [34]. The evaluation of the protective efficacy and duration of protection of MAÏA® repellent ointment compared to a standard unformulated 20% DEET in ethanol under semi-field conditions was done in the SFS. Sangoro *et al*, demonstrated that a semi-field evaluation of topical mosquito repellent gives similar results to field studies but is far safer as only disease-free laboratory-reared mosquitoes are used [29]. This is especially important in the case of *Ae. aegypti* evaluations in areas of active dengue transmission such as Tanzania [35]. Moreover, semi-field environments allow volunteers to be accessible to a known number of mosquitoes of known age, physiological status, and avidity, and minimize heterogeneity in the data allowing a more precise estimation of true repellent efficacy. The study is also conservative as it used young, never blood-fed mosquitoes raised under optimal conditions to maximize body size and then sugar starved, which are less repelled by DEET than older or smaller mosquitoes [36, 37].

The study results demonstrated that MAÏA® repellent ointment is comparable to unformulated 20% DEET in terms of mean repellency over 12 hours as well as the duration of CPT against both malaria and dengue vectors. MAÏA® repellent ointment and 20% DEET had CPT of more than 9-hours against *An. gambiae* and *An. arabiensis* and more than 7-hours for *Ae. aegypti* post-application of the repellents. These results are similar to another study which evaluated the effectiveness of MAÏA® repellent ointment

under field conditions in Burkina Faso in which the authors concluded that MAÏA® and 20% DEET are comparable in terms of duration of CPT against *Anopheles* strain and *Ae. aegypti* (Guelbeogo *et al.*, *Manuscript under preparation*).

The study results indicated that MAÏA® repellent ointment is an effective mosquito repellent suitable for the use even under high mosquito biting pressure. According to Goodyer *et al.*, an ideal mosquito repellent should provide CPT greater than 6-hours under the highest mosquito biting pressure [38]. The present study fulfilled this characteristic by demonstrating the mean CPT above 9-hours against *An. gambiae* and *An. arabiensis* and above 7-hours against *Ae. aegypti* for both MAÏA® and 20% DEET with an average of 10 mosquito landings per strain per hour in the control, 30 landings per hour in total.

User compliance is a major limitation of most topical insect repellents [15-17] and since topical repellents are applied to the skin, most users prefer insect repellents which are cosmetically pleasant in terms of odour and feel on the skin; in addition to providing protection from biting insects [39, 40]. In this study, all volunteers preferred to use MAÏA® repellent ointment compared to 20% DEET because those who applied MAÏA® repellent reported that the ointment felt better on their skin. This observation confirms that some repellent users may be influenced by product characteristics such as texture, skin feel, and odour [40, 41]. DEET tends to damage some plastics [42] which is a disadvantage in an area where plastic footwear is commonly worn. An additional benefit observed by the users was that MAÏA® repellent ointment did not damage plastics when intentionally applied to a plastic watch strap. Therefore, the use of MAÏA® repellent ointment may be a suitable alternative to a less cosmetically appealing DEET-based formulations. However, more studies are required to specifically assess user acceptability of MAÏA® repellent ointment compared to other formulated products available on the market and whether this may improve compliance.

Study limitations

After the initial nine days of the experiment, the study was extended for another nine days to confidently determine the CPT of MAÏA® repellent ointment and 20% DEET. This led to the addition of volunteers to work exclusively on the *Aedes* experiment conducted in one chamber of the SFS and only one Latin square design conducted. Therefore, the mean CPT of MAÏA® repellent ointment and 20% DEET was achieved using different and fewer (3) volunteers in the *Aedes* experiment compared to six volunteers and two SFS chambers used simultaneously in the *Anopheles* experiments. Ideally, the experiment would use more volunteers to capture the repellent efficacy against a wide range of people [43]. The study did not assess the efficacy of the repellent against nuisance mosquitoes such as *Culex quinquefasciatus*, which may also be important when assessing the consumer acceptance of repellents. This study did not involve female participants although DEET is generally more protective against women than men due to their smaller size that makes them less attractive to mosquitoes [44].

Conclusion

In conclusion, MAÏA® repellent ointment is comparable to unformulated 20% DEET under high biting pressure. Therefore, it may be recommended for use in disease-endemic areas as it protects for more than 6 hours. It is a cosmetically appealing mosquito bite protection tool that also nourishes and moisturizes the skin, which may improve consumer acceptability and fit into daily life if used every evening after bathing.

Abbreviations

AIC: Aikaike's Information Criterion

CI: Confidence Interval

CPT: Complete Protection Time

DEET: N,N-Diethyl-3-methylbenzamide

GM: Geometric mean

HQ: Headquarter

IHI: Ifakara Health Institute

IRB: Institutional Review Board

LS: Latin square

NIMR: National Institute for Medical Research

PE: Protective Efficacy

SFS: Semi field system

USA: United States of America

V/V: Volume/Volume

VCPTU: Vector Control Product Testing Unit

WHO: World Health Organisation

Declarations

Ethics approval and consent to participate

This study was approved by the Institutional Review Board of Ifakara Health Institute with approved certificate number: IHI/IRB/No: 38 – 2019. Also, the National Institute for Medical Research ethical

clearance committee approved this study with certificate number NIMR/HQ/R.8a/Vol. IX/ 3445. All volunteers involved in this study were recruited after signing a written informed consent form and were informed that they were free to leave the study at any time without any repercussion.

Consent for publication

Permission to publish these findings was granted by the National Institute of Medical Research, Tanzania.

Availability of data and materials

Data are available upon reasonable request to the corresponding author.

Competing interests

All authors work in the Vector Control Product Testing Unit (VCPTU) in Bagamoyo Tanzania and conduct vector control product evaluations for a number of companies.

Funding

The evaluation was commissioned by Maïa Africa SAS and funded by the Roddenberry Foundation (USA). The funder had no influence on study design, data collection, analysis, and the contents of this manuscript or the decision to publish.

Author's contributions

EM, OGO, MM, FT, KS, JM, SM designed the study. RP provided logistical support. EM wrote the study protocol and supervised the experiment. EM, SM conducted data analysis and wrote the manuscript. All authors reviewed the manuscript and gave permission to publish.

Acknowledgements

Authors thank study volunteers who tirelessly devoted their time and effort to this study and the Vector Control Product Testing Unit (VCPTU) management for providing mosquitoes used in this study. Authors thank the MAÏA Africa SAS for providing repellents used in this study.

References

1. Lengeler C: **Insecticide-treated bed nets and curtains for preventing malaria.** *Cochrane Database Syst Rev* 2004: Cd000363.
2. Bhatt S, Weiss DJ, Mappin B, Dalrymple U, Cameron E, Bisanzio D, Smith DL, Moyes CL, Tatem AJ, Lynch M, et al: **Coverage and system efficiencies of insecticide-treated nets in Africa from 2000 to 2017.** *Elife* 2015, **4**.

3. Sherrard-Smith E, Skarp JE, Beale AD, Fornadel C, Norris LC, Moore SJ, Mihreteab S, Charlwood JD, Bhatt S, Winskill P, et al: **Mosquito feeding behavior and how it influences residual malaria transmission across Africa.** *Proc Natl Acad Sci U S A* 2019, **116**:15086-15095.
4. Moore S: **History of Insect Repellents. Insect Repellents: Principles, Methods and Uses. Edited by: Debboun M, Frances SP, Strickman D.** Boca Raton: CRC Press, Taylor and Francis Group; 2006.
5. Katz TM, Miller JH, Hebert AA: **Insect repellents: historical perspectives and new developments.** *J Am Acad Dermatol* 2008, **58**:865-871.
6. Monroe A, Asamoah O, Lam Y, Koenker H, Psychas P, Lynch M, Ricotta E, Hornston S, Berman A, Harvey SA: **Outdoor-sleeping and other night-time activities in northern Ghana: implications for residual transmission and malaria prevention.** *Malar J* 2015, **14**:35.
7. Monroe A, Mihayo K, Okumu F, Finda M, Moore S, Koenker H, Lynch M, Haji K, Abbas F, Ali A, et al: **Human behaviour and residual malaria transmission in Zanzibar: findings from in-depth interviews and direct observation of community events.** *Malar J* 2019, **18**:220.
8. Moshi IR, Manderson L, Ngowo HS, Mlacha YP, Okumu FO, Mnyone LL: **Outdoor malaria transmission risks and social life: a qualitative study in South-Eastern Tanzania.** *Malar J* 2018, **17**:397.
9. Gryseels C, Durnez L, Gerrets R, Uk S, Suon S, Set S, Phoeuk P, Sluydts V, Heng S, Sochantha T, et al: **Re-imagining malaria: heterogeneity of human and mosquito behaviour in relation to residual malaria transmission in Cambodia.** *Malaria Journal* 2015, **14**:165.
10. Rowland M, Downey G, Rab A, Freeman T, Mohammad N, Rehman H, Durrani N, Reyburn H, Curtis C, Lines J, Fayaz M: **DEET mosquito repellent provides personal protection against malaria: a household randomized trial in an Afghan refugee camp in Pakistan.** *Trop Med Int Health* 2004, **9**:335-342.
11. Hill N, Lenglet A, Arnéz AM, Carneiro I: **Plant based insect repellent and insecticide treated bed nets to protect against malaria in areas of early evening biting vectors: double blind randomised placebo controlled clinical trial in the Bolivian Amazon.** 2007, **335**:1023.
12. Maia MF, Kliner M, Richardson M, Lengeler C, Moore SJ: **Mosquito repellents for malaria prevention.** *Cochrane Database of Systematic Reviews* 2018.
13. WHO: *Guidelines for malaria vector control.* Geneva: World Health Organization; 2019.
14. Wilson AL, Chen-Hussey V, Logan JG, Lindsay SW: **Are topical insect repellents effective against malaria in endemic populations? A systematic review and meta-analysis.** *Malaria Journal* 2014, **13**:446.
15. Gryseels C, Uk S, Sluydts V, Durnez L, Phoeuk P, Suon S, Set S, Heng S, Siv S, Gerrets R, et al: **Factors influencing the use of topical repellents: implications for the effectiveness of malaria elimination strategies.** *Scientific Reports* 2015, **5**:16847.
16. Maia MF, Kliner M, Richardson M, Lengeler C, Moore SJ: **Mosquito repellents for malaria prevention.** *The Cochrane database of systematic reviews* 2018, **2**:CD011595-CD011595.

17. Sluydts V, Durnez L, Heng S, Gryseels C, Canier L, Kim S, Van Roey K, Kerkhof K, Khim N, Mao S, et al: **Efficacy of topical mosquito repellent (picaridin) plus long-lasting insecticidal nets versus long-lasting insecticidal nets alone for control of malaria: a cluster randomised controlled trial.** 2016, **16**:1169-1177.
18. Sangoro O, Turner E, Simfukwe E, Miller JE, Moore SJ: **A cluster-randomized controlled trial to assess the effectiveness of using 15% DEET topical repellent with long-lasting insecticidal nets (LLINs) compared to a placebo lotion on malaria transmission.** *Malar J* 2014, **13**:324.
19. Analytics A: **Global Mosquito Repellent Market - Analysis By Product Type (Coils, Liquid Vaporizers, Sprays, Mats, Creams and Oil, Others), By Distribution Channel, By Region, By Country (2019 Edition): Opportunities and Forecast (2019-2024).** 2019.
20. McGready R, Hamilton KA, Simpson JA, Cho T, Luxemburger C, Edwards R, Looareesuwan S, White NJ, Nosten F, Lindsay S, et al: **Safety of the insect repellent N, N-diethyl-M-toluamide (DEET) in pregnancy.** 2001, **65**:285-289.
21. Koren G, Matsui D, Bailey B: **DEET-based insect repellents: safety implications for children and pregnant and lactating women.** *Cmaj* 2003, **169**:209-212.
22. Carroll SP: **Evaluation of topical insect repellents and factors that affect their performance.** In *Insect repellents : principles, methods, and uses.* CRC Press: Boca Raton 2007: 245 -261
23. Finda MF, Moshi IR, Monroe A, Limwagu AJ, Nyoni AP, Swai JK, Ngowo HS, Minja EG, Toe LP, Kaindoa EW, et al: **Linking human behaviours and malaria vector biting risk in south-eastern Tanzania.** *PLoS One* 2019, **14**:e0217414.
24. Maia MF, Kreppel K, Mbeyela E, Roman D, Mayagaya V, Lobo NF, Ross A, Moore SJ: **A crossover study to evaluate the diversion of malaria vectors in a community with incomplete coverage of spatial repellents in the Kilombero Valley, Tanzania.** *Parasit Vectors* 2016, **9**:451.
25. Musiime AK, Smith DL, Kilama M, Rek J, Arinaitwe E, Nankabirwa JI, Kanya MR, Conrad MD, Dorsey G, Akol AM, et al: **Impact of vector control interventions on malaria transmission intensity, outdoor vector biting rates and Anopheles mosquito species composition in Tororo, Uganda.** *Malar J* 2019, **18**:445.
26. **MAIA Africa** [<https://www.maia-africa.com/en/>]. Accessed 20 May 2020
27. Hon KL, Tsang YC, Pong NH, Lee VW, Luk NM, Chow CM, Leung TF: **Patient acceptability, efficacy, and skin biophysiology of a cream and cleanser containing lipid complex with shea butter extract versus a ceramide product for eczema.** *Hong Kong Med J* 2015, **21**:417-425.
28. Lin TK, Zhong L, Santiago JL: **Anti-Inflammatory and Skin Barrier Repair Effects of Topical Application of Some Plant Oils.** *Int J Mol Sci* 2017, **19**.
29. Sangoro O, Lweitojera D, Simfukwe E, Ngonyani H, Mbeyela E, Lugiko D, Kihonda J, Maia M, Moore S: **Use of a semi-field system to evaluate the efficacy of topical repellents under user conditions provides a disease exposure free technique comparable with field data.** *Malaria Journal* 2014, **13**:159.

30. WHOPEs: **GUIDELINES FOR EFFICACY TESTING OF MOSQUITO REPELLENTS FOR HUMAN SKIN.** Geneva: World Health Organization; 2009.
31. MR4: *Methods in Anopheles Research Manual*. 4 edn2014.
32. Saddler A, Kreppel KS, Chitnis N, Smith TA, Denz A, Moore JD, Tambwe MM, Moore SJ: **The development and evaluation of a self-marking unit to estimate malaria vector survival and dispersal distance.** *Malar J* 2019, **18**:441.
33. **SD Bioline Malaria Ag Pf** [<https://www.alere.com/en/home/product-details/sd-bioline-malaria-ag-pf-hrp-2.html>]. Accessed 20 May 2020.
34. Alpern JD, Dunlop SJ, Dolan BJ, Stauffer WM, Boulware DR: **Personal Protection Measures Against Mosquitoes, Ticks, and Other Arthropods.** *Med Clin North Am* 2016, **100**:303-316.
35. Mwanyika G, Mboera LEG, Rugarabamu S, Lutwama J, Sindato C, Paweska JT, Misinzo G: **Co-circulation of Dengue Virus Serotypes 1 and 3 during the 2019 epidemic in Dar es Salaam, Tanzania.** bioRxiv 2019:763003.
36. Xue RD, Barnard DR, Schreck CE: **Influence of body size and age of Aedes albopictus on human host attack rates and the repellency of deet.** *J Am Mosq Control Assoc* 1995, **11**:50-53.
37. Mulatier M, Porciani A, Nadalin L, Ahoua Alou LP, Chandre F, Pennetier C, Dormont L, Cohuet A: **DEET Efficacy Increases With Age in the Vector Mosquitoes Anopheles gambiae s.s. and Aedes albopictus (Diptera: Culicidae).** *J Med Entomol* 2018, **55**:1542-1548.
38. Goodyer L, Lawrence J, Feller CA. **Marketing for Efficacy, Acceptability, Safety and Health.** In: Debboun M, Frances SP, Strickman D, editors. *Insect Repellents Handbook*. CRC Press: Boca Raton; 2014: 253-263.
39. Debboun M, Frances SP, Strickman D: *Insect repellents handbook*. CRC Press; 2014.
40. Debboun M, Frances SP, Strickman D: *Insect repellents: principles, methods, and uses*. CRC press; 2006.
41. Frances SP, Debboun M: **User acceptability: public perceptions of insect repellents.** In: Debboun M, Frances SP, Strickman D, editors. *Insect Repellents. Principles, Methods, and Uses*. CRC Press: Boca Raton; 2007:397-404
42. Armed Forces Pest Management Board. **Technical Guide No. 36: Personal Protective Measures Against Insects and Other Arthropods of Military Significance.** Walter Reed Army Medical Center, Washington DC, 2015.
43. Rutledge LC, Gupta RK: **Variation in the protection periods of repellents on individual human subjects: an analytical review.** *J Am Mosq Control Assoc* 1999, **15**:348-355.
44. Gilbert IH, Gouck HK, Smith N: **Attractiveness of Men and Women to Aedes aegypti and Relative Protection Time Obtained with Deet.** *The Florida Entomologist* 1966, **49**:53-66.

Figures



Figure 1

A semi-field system (SFS) A semi-field system (SFS) with 29 x 21 meters built from a fabricated greenhouse frame modified to make two chambers with a central corridor and an opaque polyethylene roof for rain protection.




Figure 2

Study flow for *Anopheles*. Study flow for *An.s gambiae s.s.* and *An. arabiensis* conducted for 6 hours and 12-hours of recapture period. Two semi-field system (SFS) chambers (1 & 2) were used and two fully balanced Latin square (LS) design (3x3) were conducted simultaneously using six human volunteers. Mosquito recapture started immediately after application of repellent in a pilot study (Figure 2A) and started 6-hours post-application of repellents in a final study (Figure 2B). Total mosquitoes of each strains released were 50 (25 per release) and 100 (25 per release) in a pilot and final study, respectively.




Figure 3

Study flow for *Aedes aegypti*. Study flow plan for *Ae. aegypti* experiment conducted for 6 hours and 12 hours of recapture period. Two semi-field system (SFS) chambers (1 & 2) were used simultaneously and two fully balanced Latin square (LS) design (3x3) were conducted using six human volunteers in a pilot study. Mosquito recapture started immediately after application of repellent in a pilot study (Figure 3A) and started 6 hours post-application of repellents in a final study (Figure 3B). Total female *Ae.s aegypti* released were 50 (25 per release) and 100 (25 per release) in the pilot and final study, respectively.




Figure 4

An experimental set up in a semi-field system. A schematic diagram of a semi-field system showing two chambers (A & B), three mosquito collection positions and one mosquito releasing point inside each chamber. The distance between releasing point and mosquito collection positions was nine meters.




Figure 5

Complete Protection Time (CPT) of MAIA® repellent ointment and 20% DEET. The CPT of MAIA® repellent ointment and 20% DEET against laboratory-reared mosquito strains. (A) Probability of no *An. gambiae* landing on lower limbs of volunteers treated with MAIA® repellent ointment (green) and 20% DEET (red). (B) Probability of no *An. arabiensis* landing on lower limbs of volunteers treated with MAIA® repellent ointment (green) and 20% DEET (red). (C) The probability of no *Ae. aegypti* landing on lower limbs of volunteers treated with MAIA® repellent ointment (green) and 20% DEET (red).