Quantum Fourier Transform Is the Building Block for Creating Entanglement

Mario Mastriani (✉ mmastria@fiu.edu)

Florida International University

Research Article

Keywords: quantum Fourier transform (QFT), quantum computing, quantum communication, future quantum Internet

Posted Date: June 1st, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-563987/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Scientific Reports on November 15th, 2021. See the published version at <https://doi.org/10.1038/s41598-021-01745-x>.

Quantum Fourier Transform is the building block for creating entanglement

Mario Mastriani

*Knight Foundation School of Computing & Information Sciences,
Florida International University, 11200 S.W. 8th Street, Miami, FL 33199, USA*

This study demonstrates entanglement can be exclusively constituted by quantum Fourier transform (QFT) blocks. A bridge between entanglement and QFT will allow incorporating a spectral analysis to the already traditional temporal approach of entanglement, which will result in the development of new more performant, and fault-tolerant protocols to be used in quantum computing as well as quantum communication, with particular emphasis in the future quantum Internet.

ORCID Id: 0000-0002-5627-3935

mmastria@fiu.edu

Introduction. -Jean-Baptiste Joseph Fourier (Auxerre, France, 21 March 1768, Paris, 16 May 1830) was a French mathematician and physicist, a disciple of Joseph-Louis Lagrange (Turin, Italy, 25 January 1736, Paris, 10 April 1813), known for his work on the decomposition of periodic functions into convergent trigonometric series called Fourier series, a method with which he managed to solve the heat equation. The projection of his work in the two centuries following his death on areas as diverse as electricity, optics, electronics, and so on, culminated during the twentieth century in the creation of the famous Discrete Fourier Transform¹, Fast Fourier Transform², and Quantum Fourier Transform³ (QFT), where the latter constitutes a key piece within Quantum Information Processing⁴ for the case of those quantum algorithms that require a phase estimation⁵.

On the other hand, entanglement⁶⁻⁸, so reviled by Albert Einstein, Boris Podolsky, and Nathan Rosen in their 1935 paper⁹, has become the cornerstone of Quantum Computing⁴ and Quantum Communication¹⁰, in particular, in communications protocols such as quantum teleportation¹¹, quantum secret sharing¹², quantum key distribution¹³, quantum secure direct communication¹⁴, and quantum repeaters¹⁵, with a marked commitment to the future quantum Internet¹⁶⁻²⁰.

The union of both entities, i.e. QFT, and entanglement, seems at first something quite strange, at least in the way it is presented in this work, where the first becomes a basal element for the creation of the second, however, the approach that will be presented then it will allow access to the hidden face of the entanglement, its spectral face.

QFT is constituted by an important family of quantum operations over the ring $\mathbb{Z}_2^n$. The n -qubit QFT makes a coherent mapping from an input state or qubit string $|x\rangle = |x_1 \text{ K } x_n\rangle$ to an output state or qubit string $|y\rangle = |y_1 \text{ K } y_n\rangle$ in the computational basis²¹ as follows:

$$|y\rangle \text{ a } \frac{1}{\sqrt{2^n}} \sum_{v=0}^{2^n-1} \omega_{2^n}^{u \cdot v} |x\rangle, \quad u = 0, 1, 2, \text{ K }, 2^n - 1, \quad (1)$$

where $\omega_{2^n} = e^{i2\pi/2^n}$ is the 2^n root of unity, while the inverse QFT is:

$$|x\rangle \text{ a } \frac{1}{\sqrt{2^n}} \sum_{u=0}^{2^n-1} \omega_{2^n}^{-v \cdot u} |y\rangle, \quad v = 0, 1, 2, \text{ K }, 2^n - 1. \quad (2)$$

The Hadamard matrix H is equivalent to the 1-qubit QFT and its inverse^{22, 23},

$$F_{2^1} = H = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} = H^{-1} = F_{2^1}^{-1} \in \mathbb{C}^{2^1 \times 2^1}. \quad (3)$$

That is, for the 1-qubit QFT all its components are equivalent. Instead, for the 2-qubit QFT, the same does not happen, since

$$F_{2^2} = \frac{1}{2} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & i & -1 & -i \\ 1 & -1 & 1 & -1 \\ 1 & -i & -1 & i \end{bmatrix}, \quad (4a) \quad \text{and} \quad F_{2^2}^{-1} = \frac{1}{2} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & -i & -1 & i \\ 1 & -1 & 1 & -1 \\ 1 & i & -1 & -i \end{bmatrix}, \quad (4b)$$

are different: $F_{2^2} \neq F_{2^2}^{-1}$, where $F_{2^2} \wedge F_{2^2}^{-1} \in \mathbb{F}^{2^2 \times 2^2}$. On the other hand, the Feynman's gate⁴ (also known as *Controlled-X*, CNOT, or CX gate), as well as its flipped version are respectively:

$$CNOT = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}, \quad (5a) \quad \text{and} \quad CNOT_{flipped} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}, \quad (5b)$$

where the difference between them consists in that in Eq.(5a) the upper qubit is the control qubit, while the lower qubit is the target qubit. Instead, in the version of Eq.(5b) it is exactly the opposite, being: $CNOT_{flipped} = (H \otimes H) \times CNOT \times (H \otimes H)$, “ $\times$ ” the matrix product, and “ $\otimes$ ” the Kronecker product⁴.

Multiplying both F_{2^2} by itself and $F_{2^2}^{-1}$ by itself, both multiplications result equal to the $CNOT_{flipped}$ gate of Eq.(5b): $F_{2^2} \times F_{2^2} = F_{2^2}^{-1} \times F_{2^2}^{-1} = CNOT_{flipped}$. This can be easily verified by multiplying $CNOT_{flipped}$ by itself, and $F_{2^2} \times F_{2^2}$ by $F_{2^2}^{-1} \times F_{2^2}^{-1}$ and regrouping,

$$\begin{aligned} CNOT_{flipped} \times CNOT_{flipped} &= (F_{2^2} \times F_{2^2}) \times (F_{2^2}^{-1} \times F_{2^2}^{-1}) \\ &= F_{2^2} \times (F_{2^2} \times F_{2^2}^{-1}) \times F_{2^2}^{-1} = F_{2^2} \times I \times F_{2^2}^{-1} = F_{2^2} \times F_{2^2}^{-1} = I_{2^2 \times 2^2}. \end{aligned} \quad (6)$$

However,

$$\sqrt{CNOT} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & (1+i)/2 & (1-i)/2 \\ 0 & 0 & (1-i)/2 & (1+i)/2 \end{bmatrix}, \quad (7a) \quad \text{and} \quad \sqrt{CNOT_{flipped}} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & (1+i)/2 & 0 & (1-i)/2 \\ 0 & 0 & 1 & 0 \\ 0 & (1-i)/2 & 0 & (1+i)/2 \end{bmatrix}, \quad (7b)$$

therefore, $\sqrt{CNOT} \neq F_{2^2}$ and $\sqrt{CNOT_{flipped}} \neq F_{2^2}$. Finally, the $CNOT$ gate is equal to the flipped version of the multiplication of QFT F_{2^2} by itself,

$$(H \otimes H) \times (F_{2^2} \times F_{2^2}) \times (H \otimes H) = (H \otimes H) \times (F_{2^2}^{-1} \times F_{2^2}^{-1}) \times (H \otimes H) = CNOT. \quad (8)$$

Equation (8) is fundamental in the creation of the entanglement for two or more qubits, as well as in all the applications that require it, as is the case of quantum teleportation¹¹.

Bell states.-Pauli's matrices⁴ can be expressed in terms of the so-named Hadamard rotation gates²⁴ or the general unitary operator $U(\theta, \varphi, \lambda) = \begin{bmatrix} \cos(\theta/2) & -e^{i\lambda} \sin(\theta/2) \\ e^{i\varphi} \sin(\theta/2) & e^{i(\lambda+\varphi)} \cos(\theta/2) \end{bmatrix}$ as follows:

$$I = H_I H_I = H_{III} H_{III} = H_{II} H_{IV} = H_{IV} H_{II}, \quad (9a)$$

$$X = H_{III} H_{II} = H_{II} H_I = H_I H_{IV} = H_{IV} H_{III}, \quad (9b)$$

$$Y = i H_{III} H_I = i H_{II} H_{II} = -i H_{IV} H_{IV} = -i H_I H_{III}, \text{ and} \quad (9c)$$

$$Z = -H_{II} H_{III} = H_I H_{II} = -H_{III} H_{IV} = H_{IV} H_I, \quad (9d)$$

where I is a 2×2 identity matrix, $i = \sqrt{-1}$, $H_I = H = U(\pi/2, 0, \pi)$ of Eq.(3), while

$$H_{II} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} = U(\pi/2, 0, 0), \quad H_{III} = \frac{1}{\sqrt{2}} \begin{bmatrix} -1 & 1 \\ 1 & 1 \end{bmatrix} = U(5\pi/2, \pi, 0), \quad \text{and} \quad H_{IV} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix} = U(\pi/2, \pi, \pi). \quad (10)$$

The equivalence of Eq.(9b) can be used to represent the four Bell states:

$$|\beta_{00}\rangle = |\Phi^+\rangle = 1/\sqrt{2}(|00\rangle + |11\rangle) = CNOT(H \otimes I)|00\rangle, \quad (11a)$$

$$|\beta_{01}\rangle = |\Psi^+\rangle = 1/\sqrt{2}(|01\rangle + |10\rangle) = CNOT(H \otimes I)|01\rangle = CNOT(H \otimes I)|0\rangle X|0\rangle, \quad (11b)$$

$$|\beta_{10}\rangle = |\Phi^-\rangle = 1/\sqrt{2}(|00\rangle - |11\rangle) = CNOT(H \otimes I)|10\rangle = CNOT(H \otimes I)X|0\rangle|0\rangle, \quad \text{and} \quad (11c)$$

$$|\beta_{11}\rangle = |\Psi^-\rangle = 1/\sqrt{2}(|01\rangle - |10\rangle) = CNOT(H \otimes I)|11\rangle = CNOT(H \otimes I)X|0\rangle X|0\rangle, \quad (11d)$$

with $|0\rangle = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, and $|1\rangle = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$. That is, replacing in Eq.(11) the equivalence corresponding to X of Eq.(9b),

H of Eq.(3), I of Eq.(9a), and $CNOT$ of Eq.(8), it is possible to implement the four Bell states exclusively basing on QFT blocks. Without losing generality, Fig. 1(a, and c) represents the implementation of the $|\beta_{00}\rangle = |\Phi^+\rangle$ Bell state in terms of its two original versions (direct and flipped), while Fig. 1(c, and d) constitute their respective counterparts based exclusively on QFT blocks, confirming that these blocks are all that is needed to create entanglement while revealing its spectral nature.

FIG. 1. Representation of the $|\beta_{00}\rangle = |\Phi^+\rangle$ Bell state in terms of QFT: a) original version based on H and $CNOT$ gates, b) its representations in terms of one $QFT_{2^1 \times 2^1}$ and two flipped $QFT_{2^2 \times 2^2}$, c) original version with one H and one flipped $CNOT$ gates, and d) its representations in terms of one $QFT_{2^1 \times 2^1}$ and two $QFT_{2^2 \times 2^2}$.

N -qubits Greenberger–Horne–Zeilinger (GHZ_N) states.- This family of configurations is the most commonly used in practice when it comes to entanglement between three or more particles^{4, 6-8}, being its general form as follows:

$$|GHZ_N\rangle = 1/\sqrt{2}(|0\rangle^{\otimes N} + |1\rangle^{\otimes N}). \quad (12)$$

Without loss of generality, in this study only $|GHZ_3\rangle$ and $|GHZ_4\rangle$ are implemented in terms of QFT blocks, where:

$$|GHZ_3\rangle = 1/\sqrt{2}(|000\rangle + |111\rangle) = (I_{2 \times 2} \otimes CNOT)(CNOT \otimes I_{2 \times 2})(H \otimes I_{4 \times 4})|000\rangle, \quad \text{and} \quad (13)$$

$$|GHZ_4\rangle = 1/\sqrt{2}(|0000\rangle + |1111\rangle) = (I_{4 \times 4} \otimes CNOT)(I_{2 \times 2} \otimes CNOT \otimes I_{2 \times 2})(CNOT \otimes I_{4 \times 4})(H \otimes I_{8 \times 8})|0000\rangle. \quad (14)$$

Equations (13) and (14) are graphically represented in Figures 2(a) and 2(d), respectively. Figures 2(b) shows $|GHZ_3\rangle$ with one $QFT_{2^1 \times 2^1}$ and four flipped $QFT_{2^2 \times 2^2}$, while Fig. 2(c) represents it thanks to one $QFT_{2^1 \times 2^1}$ and two flipped $QFT_{2^3 \times 2^3}$. Finally, Fig. 2(e) shows $|GHZ_4\rangle$ with one $QFT_{2^1 \times 2^1}$ and six flipped $QFT_{2^2 \times 2^2}$, while Fig. 2(f) represents it thanks to one $QFT_{2^1 \times 2^1}$ and two flipped $QFT_{2^4 \times 2^4}$.

Equivalences between Fig. 2(a, b, c) as well as between Fig. 2(d, e, f), show again the spectral nature of the entanglement even for the case of more than two particles entangled at the same time. In fact, the equivalences present in Fig. 2 between $(N-1)$ pairs of flipped QFT $_{2^2 \times 2^2}$ and two flipped QFT $_{2^N \times 2^N}$, show that the equivalence of Fig. 1 is not a simple coincidence for a particular case like a Bell state, but actually, the entanglement in all its manifestations has a spectral nature, where the QFT is the essential instrument for a spectral tomography of it. It only remains to project this equivalence, in perhaps the most conspicuous application of entanglement, quantum teleportation¹¹.

FIG. 2. Implementations of $|GHZ_3\rangle$ and $|GHZ_4\rangle$: a) original version of $|GHZ_3\rangle$ in terms of one H and two CNOT gates, b) $|GHZ_3\rangle$ thanks to one QFT $_{2^1 \times 2^1}$ and four flipped QFT $_{2^2 \times 2^2}$, c) $|GHZ_3\rangle$ based on one QFT $_{2^1 \times 2^1}$ and two flipped QFT $_{2^3 \times 2^3}$, d) original version of $|GHZ_4\rangle$ in terms of one H and three CNOT gates, e) $|GHZ_4\rangle$ thanks to one QFT $_{2^1 \times 2^1}$ and six flipped QFT $_{2^2 \times 2^2}$, f) $|GHZ_4\rangle$ based on one QFT $_{2^1 \times 2^1}$ and two flipped QFT $_{2^4 \times 2^4}$.

Teleportation.—This protocol¹¹ is implemented in three different ways in Fig. 3, where a qubit $|\psi\rangle$ to be teleported is prepared and introduced in the upper qubit on the left of the protocol. A Bell state like that of Fig. 1 is distributed between Alice and Bob. Subsequently, a module applied in the two upper qubits and constituted by a CNOT gate, an H gate, and two quantum measurement blocks (QuMe) constitute what in practice is known as a Bell State Measurement (BSM) module^{4,6-8}. The double lines at the output of each QuMe convey classical information from Alice to Bob in the form of two classical disambiguation or control bits. For this reason, this means of transport is known as a classic channel of disambiguation, control, or simply as an auxiliary channel.

A 2-qubits Controlled-Z gate or simply CZ gate can be constructed from QFT blocks according to the equivalence of Eq.(9d), or in terms of two H and one CNOT (Controlled-X or CX) gates from the following identity:

$$CZ = (I_{2 \times 2} \otimes H)CX(I_{2 \times 2} \otimes H) \quad (15)$$

As can be seen in Fig. 3(c), except for the QuMe blocks, everything else in this protocol is representable using QFT blocks. This extends, with identical results, to all other protocols that are based on entanglement such as quantum secret sharing¹², quantum key distribution¹³, quantum secure direct communication¹⁴, and quantum repeaters¹⁵, and that are used in quantum Internet¹⁶.

FIG. 3. Quantum teleportation protocol: a) the original implementation, b) Controlled-Z is replaced with its equivalent in terms of one CX and two H gates, and c) a version based exclusively on QFT blocks, where QuMe means *quantum measurement*, and the double lines represent classical information obtained after each QuMe.

Quantum Spectral Analysis.—A time decomposition, based on Fig. 4, is developed according to the intervention of each QFT block in the creation of the $|\beta_{00}\rangle = |\Phi^+\rangle$ Bell state. This analysis begins with the creation of a flipped $\text{QFT}_{2^2 \times 2^2}$ in terms of a $\text{QFT}_{2^2 \times 2^2}$ and four Hadamard (H) gates, where the last ones are used for flipping it as follows,

$$\begin{aligned}
 F_{2^2, \text{flipped}} &= (H \otimes H) F_{2^2} (H \otimes H) \\
 &= \frac{1}{2} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{bmatrix} \frac{1}{2} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & i & -1 & -i \\ 1 & -1 & 1 & -1 \\ 1 & -i & -1 & i \end{bmatrix} \frac{1}{2} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & i & -i \\ 1 & -1 & -i & i \end{bmatrix}. \quad (16)
 \end{aligned}$$

FIG. 4. Timeline of the $|\beta_{00}\rangle = |\Phi^+\rangle$ Bell state of Fig. 1(b) based on QFT blocks.

Next, the complete timeline is developed according to Fig. 4, starting at t_1 , where $|\psi(t_1)\rangle$ is the wave-function in that instant,

$$|\psi(t_1)\rangle = |0\rangle \otimes |0\rangle = |00\rangle = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} = |q_u(t_1)\rangle \otimes |q_l(t_1)\rangle, \quad (17)$$

$|q_u(t_1)\rangle = |0\rangle$ is the upper qubit in Fig. 4, and $|q_l(t_1)\rangle = |0\rangle$ is the lower qubit in that figure. The qubits obtained $|q_u\rangle$ and $|q_l\rangle$ at time t_1 are *completely independent*⁶, and are used as inputs to the next step, which is made up of an H gate in $|q_u\rangle$ and an identity matrix in $|q_l\rangle$,

$$\begin{aligned} |\psi(t_2)\rangle &= (H \otimes I)|\psi(t_1)\rangle = (H \otimes I)|00\rangle = \begin{bmatrix} 1/\sqrt{2} & 0 & 1/\sqrt{2} & 0 \\ 0 & 1/\sqrt{2} & 0 & 1/\sqrt{2} \\ 1/\sqrt{2} & 0 & -1/\sqrt{2} & 0 \\ 0 & 1/\sqrt{2} & 0 & -1/\sqrt{2} \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 1/\sqrt{2} \\ 0 \\ 1/\sqrt{2} \\ 0 \end{bmatrix} \\ &= (H|0\rangle) \otimes |0\rangle = |+\rangle \otimes |0\rangle = |q_u(t_2)\rangle \otimes |q_l(t_2)\rangle, \end{aligned} \quad (18)$$

where $|+\rangle = \begin{bmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{bmatrix}$, $|q_u(t_2)\rangle = |+\rangle$, and $|q_l(t_2)\rangle = |0\rangle$. As in the previous case, Eq.(18) shows us that both qubits obtained at time t_2 are also completely independent⁶. Instead, in the next step yields,

$$|\psi(t_3)\rangle = F_{2^2, \text{flipped}} |\psi(t_2)\rangle = \frac{1}{2} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & i & -i \\ 1 & -1 & -i & i \end{bmatrix} \begin{bmatrix} 1/\sqrt{2} \\ 0 \\ 1/\sqrt{2} \\ 0 \end{bmatrix} = \begin{bmatrix} 1/\sqrt{2} \\ 0 \\ (1+i)/2\sqrt{2} \\ (1-i)/2\sqrt{2} \end{bmatrix} \neq |q_u(t_3)\rangle \otimes |q_l(t_3)\rangle, \quad (19)$$

where Eq.(19) indicates that $|q_u\rangle$ and $|q_l\rangle$ at time t_3 are *classically correlated*⁶, i.e. although this case does not result in entanglement, $|\psi(t_3)\rangle$ cannot be factored. In this intermediate instance, between both flipped $\text{QFT}_{2^2 \times 2^2}$, the impossibility of factoring can be observed, since there are no values of $|q_u(t_3)\rangle$ and $|q_l(t_3)\rangle$, such that $|\psi(t_3)\rangle = |q_u(t_3)\rangle \otimes |q_l(t_3)\rangle$. Undoubtedly, this constitutes advance respect to the literature on the subject in force to date, which associated the aforementioned impossibility with the intervention of the CNOT gate, as a whole, in an exclusive way, or with an inappropriate coupling between the individual contributions of H and CNOT gates. In consequence, this analysis makes explicit an intermediate instance to the one already known for the non-separability and indistinguishability of the states during entanglement, which is exclusively the responsibility of a particular characteristic of the Discrete Fourier Transform¹ (DFT) and that is inherited by the QFT³. This characteristic refers to the fact that the DFT is a dense matrix, i.e. all its elements are different from zero, since, they are the N roots of the unit or twiddle factors¹, which when they are multiplied by the input vector produce an output vector where each of its elements represents a mixture or weighted sum of the incoming vector. Finally, the intervention of the second flipped $\text{QFT}_{2^2 \times 2^2}$ allows obtaining the wave-function at the time t_4 ,

$$|\psi(t_4)\rangle = F_{2^2, \text{flipped}} |\psi(t_3)\rangle = \frac{1}{2} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & i & -i \\ 1 & -1 & -i & i \end{bmatrix} \begin{bmatrix} 1/\sqrt{2} \\ 0 \\ (1+i)/2\sqrt{2} \\ (1-i)/2\sqrt{2} \end{bmatrix} = \begin{bmatrix} 1/\sqrt{2} \\ 0 \\ 0 \\ 1/\sqrt{2} \end{bmatrix} \neq |q_u(t_4)\rangle \otimes |q_l(t_4)\rangle, \quad (20)$$

where, as in the previous case, it is impossible to decompose $|\psi(t_4)\rangle$ into two independent states $|q_u\rangle$ and $|q_l\rangle$, that is, $|\psi(t_4)\rangle$ is not factorable. This gives rise to a very particular state of null spin called *entanglement*⁶.

The four density matrices associated with every wave-function of Fig. 4 are the following:

$$\rho(t_1) = |\psi(t_1)\rangle\langle\psi^*(t_1)| = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad (21)$$

where $(\bullet)^*$ is the complex conjugate of $(\bullet)$,

$$\rho(t_2) = |\psi(t_2)\rangle\langle\psi^*(t_2)| = \begin{bmatrix} 1/\sqrt{2} \\ 0 \\ 1/\sqrt{2} \\ 0 \end{bmatrix} \begin{bmatrix} 1/\sqrt{2} & 0 & 1/\sqrt{2} & 0 \end{bmatrix} = \begin{bmatrix} 1/2 & 0 & 1/2 & 0 \\ 0 & 0 & 0 & 0 \\ 1/2 & 0 & 1/2 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad (22)$$

$$\begin{aligned} \rho(t_3) = |\psi(t_3)\rangle\langle\psi^*(t_3)| &= \begin{bmatrix} 1/\sqrt{2} \\ 0 \\ (1+i)/2\sqrt{2} \\ (1-i)/2\sqrt{2} \end{bmatrix} \begin{bmatrix} 1/\sqrt{2} & 0 & (1+i)/2\sqrt{2} & (1-i)/2\sqrt{2} \end{bmatrix} \\ &= \begin{bmatrix} 1/2 & 0 & (1+i)/4 & (1-i)/4 \\ 0 & 0 & 0 & 0 \\ (1+i)/4 & 0 & i/4 & 1/4 \\ (1-i)/4 & 0 & 1/4 & -i/4 \end{bmatrix}, \text{ and} \end{aligned} \quad (23)$$

$$\rho(t_4) = |\psi(t_4)\rangle\langle\psi^*(t_4)| = \begin{bmatrix} 1/\sqrt{2} \\ 0 \\ 0 \\ 1/\sqrt{2} \end{bmatrix} \begin{bmatrix} 1/\sqrt{2} & 0 & 0 & 1/\sqrt{2} \end{bmatrix} = \begin{bmatrix} 1/2 & 0 & 0 & 1/2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1/2 & 0 & 0 & 1/2 \end{bmatrix}. \quad (24)$$

The four density matrices can be seen in Table 1, where only $\rho(t_3)$ has an imaginary part. On the other hand, comparing the 3D bars of the density matrices at t_2 and t_4 , it can be seen that the consecutive action of both flipped QFT_{2²×2²} has a stretching effect as far as the locations of the bars are concerned. This shows that both flipped QFT_{2²×2²} are the architect of a bad copy of wave-function $|\psi(t_2)\rangle$ of Eq. (18), that is to say,

$$U|\psi(t_2)\rangle = U(|+\rangle|0\rangle) \neq |+\rangle|+\rangle, \quad (25)$$

where $U = \text{flipped QFT}_{2^2 \times 2^2} \times \text{flipped QFT}_{2^2 \times 2^2}$, being the true result generated by both,

$$U|\psi(t_2)\rangle = U(|+\rangle|0\rangle) = |\beta_{00}\rangle. \quad (26)$$

This shows that entanglement is the result of an inadequate copy by a very inefficient copy machine embodied by both flipped $QFT_{2^2 \times 2^2}$. The aforementioned stretching effect added to the unification of the entanglement's own wave-function triggers its most conspicuous characteristic, that is to say, the impossibility of factoring the wave function of Eq.(20).

Table 1. Density matrices of the four wave-functions of Fig. 4.

Instant	Real part of the density matrices	Imaginary part of the density matrices
t_1		—
t_2		—
t_3

Moreover, given two subsystems (A, and B) that interact with each other, their density matrices treated individually are,

$$\rho^A = \rho^B = \frac{1}{2}(|0\rangle\langle 0| + |1\rangle\langle 1|) = \frac{1}{2}I = \frac{1}{2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad (21)$$

and their von Neumann entropies are,

$$S^A = S^B = -\text{tr} \left[\rho^A \log(\rho^A) \right] = -\text{tr} \left[\rho^B \log(\rho^B) \right] = -\text{tr} \left[\frac{1}{2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \log \left(\frac{1}{2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \right] = 1, \quad (22)$$

where $\text{tr}(\bullet)$ is the *trace of the square matrix* ($\bullet$), and $\log(\bullet)$ is *logarithm base 2 of* ($\bullet$). In the same way, for a composed system, the entropy is,

$$S^{A \cup B} = -\text{tr} \left[\rho^{A \cup B} \log(\rho^{A \cup B}) \right]. \quad (23)$$

$S^{A \cup B}$ depends on the degree of correlation (completely independent, classically correlated, and entangled) between both subsystems. Besides, in the classical and the quantum worlds, the correlations between the subsystems are those established by the additional information. In the case of composite quantum systems, the mutual information $S^{A \cap B}$ is introduced to quantify that additional information, allowing us to obtain the degree of correlation between both subsystems⁶,

$$S^{A \cap B} = S^A + S^B - S^{A \cup B} \geq 0. \quad (24)$$

Therefore, the entropy of the composite system $S^{A \cap B}$ indicates that the uncertainty of a state $\rho^{A \cup B}$ is less than the two subsystems S^A and S^B added together.

Table 2. Entropies in terms of the degree of correlations between both subsystems.

Degree of correlation between both subsystems	$S_{A \cup B}$	$S_{A \cap B}$	Graphic of sets

completely independent	2	0	
classically correlated	1	1	
entangled	0	2	

Table 2 shows entropies in terms of the degree of correlations between both subsystems, in such a way that when $S^{A \cup B} = 2$, the entropy of the composite system $S^{A \cap B} = S^A + S^B - S^{A \cup B} = 1 + 1 - 2 = 0$, which means that both subsystems do not have mutual information, and this null degree of correlation corresponds to the case of Eq.(18) of Fig. 4 at time t_2 , where $|\psi(t_2)\rangle$ is factored into $|q_u(t_2)\rangle = |+\rangle$, and $|q_l(t_2)\rangle = |0\rangle$, that is, both subsystems are completely independent. Instead, when $S^{A \cup B} = 1$, the entropy of the composite system $S^{A \cap B} = S^A + S^B - S^{A \cup B} = 1 + 1 - 1 = 1$, this case corresponds to Eq.(19) at time t_3 of Fig. 4, where both subsystems are classically correlated, i.e., $|\psi(t_3)\rangle$ cannot be factored in terms of $|q_u(t_3)\rangle$ and $|q_l(t_3)\rangle$. Although both subsystems (A, and B) share information and are not separable, they do not give rise to entanglement. Finally, if $S^{A \cup B} = 0$, the entropy of the composite system $S^{A \cap B} = S^A + S^B - S^{A \cup B} = 1 + 1 - 0 = 2$, that is to say, the mutual information between both subsystems is maxima. It is about the presence of entanglement corresponding to Eq.(20) at time t_4 of Fig. 4, where, as in the previous case, it is impossible to decompose $|\psi(t_4)\rangle$ into two independent states $|q_u\rangle$ and $|q_l\rangle$, that is, $|\psi(t_4)\rangle$ is not factorable.

Conclusions.-This study demonstrated the existing relationship between the Feynman⁴ gate, known as Controlled-X, CNOT, or CX, with a pair of flipped QFT $2^2 \times 2^2$. This, added to the already known equivalence between the Hadamard gate (H), and one QFT $2^1 \times 2^1$ matrix, gives rise to the creation of entanglement based exclusively on QFT blocks. This equivalence is extended to the creation of entanglement between more than two particles, as is the case of the states $|GHZ_3\rangle$, and $|GHZ_4\rangle$.

A representation of the famous quantum teleportation protocol¹¹ based exclusively on QFT^{3,4} blocks is achieved, which highlights a clear projection of the study carried out here on the future quantum Internet¹⁶⁻²⁰.

The decomposition of the configuration for the creation of the entanglement in QFT blocks allows, through the timeline of Fig. 4, to perform an internal tomography of the entanglement, revealing, for the first time in the literature, the three degrees of correlation between particles⁶, that is, completely independent, classically correlated, and entanglement, from a single configuration.

In the same process mentioned previously, it becomes evident as never before that entanglement arises from a defective copy starring both flipped QFT $2^2 \times 2^2$.

Finally, the spectral analysis of all quantum computing and quantum communication protocols, added to the traditional temporal analysis present in the literature, will allow a better understanding of the inner nature of the entanglement, so that this new approach can help to create new and more efficient algorithms and fault tolerant protocols.

Acknowledgements.-M.M. thanks the staff of the Knight Foundation School of Computing and Information Sciences at Florida International University for all their help and support.

Competing Interests.-M.M. declares that he has no competing interests.

References

1. Briggs, W. L. & van Emden, H. The DFT: An Owner's Manual for the Discrete Fourier Transform (SIAM, Philadelphia, 1995).
2. Jain, A. K. Fundamentals of Digital Image Processing (Englewood Cliffs, N.J., 1989).
3. Weinstein, Y. S. *et al.* Implementation of the Quantum Fourier Transform, *Phys Rev Lett.* 86, 1889 (2001).
4. Nielsen, M. A. & Chuang, I. L. Quantum Computation and Quantum Information (Cambridge University Press, N.Y., 2004).
5. Shor, P. W. Polynomial-Time Algorithms for Prime Factorization and Discrete Logarithms on a Quantum Computer, *SIAM J. Comput.*, 26:5, 1484–1509 (1997).
6. Audretsch, J. Entangled Systems: New Directions in Quantum Physics (Wiley-VCH Verlag GmbH & Co., Weinheim, 2007).
7. Jaeger, G. Entanglement, Information, and the Interpretation of Quantum Mechanics. (Springer-Verlag, Berlin, 2009).
8. Horodecki, R. *et al.* Quantum entanglement, *Rev. Mod. Phys.*, 81:2, 865-942 (2009).
9. Einstein, A., Podolsky, B. & Rosen, N. Can Quantum-Mechanical Description of Physical Reality Be Considered Complete? *Physical Review.* 47:10, 777–780 (1935).
10. Cariolaro, G. Quantum Communications: Signals and Communication Technology (Springer, N.Y., 2015).
11. Bennett, C. H. *et al.* Teleporting an unknown quantum state via dual classical and Einstein-Podolsky-Rosen channels, *Phys. Rev. Lett.* 70:13, 1895-1899 (1993).
12. Joy, D. *et al.* Implementation of quantum secret sharing and quantum binary voting protocol in the IBM quantum computer, *Quantum Inf Process.* 19:33 (2020).
13. Ekert, A. K. Quantum cryptography based on Bell's theorem, *Phys. Rev. Lett.* 67:6, 661–663 (1991).
14. Zhou, Z. *et al.* Measurement-device-independent quantum secure direct communication, *Sci Chin Phys Mech Astron.* 63:3, 230362 (2020).
15. Ruihong, Q. & Ying, M. Research Progress Of Quantum Repeaters, *IOP J Phys Conf Ser.* 1237:052032 (2019).
16. Caleffi, M. *et al.* The Rise of the Quantum Internet, *Computer.* 53:06, 67-72 (2020).
17. Cacciapuoti, A. S. *et al.* The Quantum Internet: Networking Challenges in Distributed Quantum Computing, *IEEE Network*, 34:1, 137-143 (2020).
18. Gyongyosi, L. & Imre, S. Entanglement Accessibility Measures for the Quantum Internet, *Quant. Info. Proc.* 19:115 (2020).
19. Gyongyosi, L. & Imre, S. Entanglement Access Control for the Quantum Internet, arXiv:quant-ph/1905.00256 (2019).
20. Gyongyosi, L. & Imre, S. Opportunistic Entanglement Distribution for the Quantum Internet, arxiv:quant-ph/1905.00258 (2019).
21. Browne, D. E. Efficient classical simulation of the quantum Fourier transform, *New J. Phys.* 9:146 (2007).
22. Rieffel, E. & Polak, W. Quantum Computing: A Gentle Introduction (The MIT Press, Cambridge, Massachusetts, 2011).
23. Fijany, A. & Williams, C. P. Quantum Wavelet Transforms: Fast Algorithms and Complete Circuits, arxiv:quant-ph/9809004 (1998).
24. Gruska, J. Quantum computing (McGraw-Hill, N.Y., 1999-2005).

Figures

Figure 1

See the Manuscript Files section for the complete figure caption.

Figure 2

See the Manuscript Files section for the complete figure caption.