

The Beneficial Effects of Square Dance on Musculoskeletal System in Early Postmenopausal Chinese Women: A Cross-sectional Study.

Jie Sun

Affiliated Hospital of Nantong University

Chen Yao

Affiliated Hospital of Nantong University

Ziping Wang

Affiliated Hospital of Nantong University

Jiancheng Wu

Affiliated Hospital of Nantong University

Bo Zhang

Affiliated Hospital of Nantong University

Zhenyu Zhou

Affiliated Hospital of Nantong University

Fan Liu

Affiliated Hospital of Nantong University

Yafeng Zhang (✉ fcius2004@hotmail.com)

Affiliated Hospital of Nantong University <https://orcid.org/0000-0002-6823-9961>

Research article

Keywords: postmenopausal women; square dance; BMD; musculoskeletal system

Posted Date: August 13th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-56419/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: This study was set to investigate the beneficial effects of square dance on musculoskeletal system of early postmenopausal Chinese women.

Methods: Chinese postmenopausal women, who had been without menstruation for 1 to 10 years from the onset of menopause were recruited from community centers for this study. A standardized structured face-to-face interview was performed to collect demographic information, life styles, personal medical history, diet and menstrual status. Subjects who had been practicing regular square dance without participated in other sports activities for more than 2 years and over 4 hours per week (usually more than 45 minutes per time and more than 5 times per week) were assigned to square dance group. Those postmenopausal women who had not participated in regular exercises (no more than 0.5 hour per week) were recruited as the sedentary control group. Bone mineral density (BMD) of spine, total hip and femoral neck was measured by using dual-energy X-ray absorptiometry (DXA). Low limb muscle strength was measured for the non-dominant leg, body flexibility was measured by a simple trunk bend-and-reach test, and body balance was evaluated using a single-stance test for the non-dominant leg. Independent two-tailed Student's t-test was used for data analysis.

Results: 152 subjects from community centers were selected for this study and divided into square dance group (n=74) and control group (n=78). The square dance subjects had higher lumbar spine BMD (P=0.01) and total hip BMD (P=0.02) than control subjects, but there was no significant difference of femoral neck BMD (P=0.48) between these two groups. Functional testing indicated that square dance subjects had higher low limb muscle strength (P<0.01) and longer single-stance time (P=0.02) than the control subject, but there was no significant difference in trunk bend-and-reach (P=0.12) between these two groups.

Conclusion: Our results show that postmenopausal Chinese women can get beneficial effects, like higher BMD, stronger low limb muscle and improved body balance ability on musculoskeletal system by participating in square dance regularly.

Introduction

Aging of population has become an extremely tough challenge that the whole world must face in the twenty-first century. On mainland China, the residents who are 60 years old and above account for over 13% of the total population according to the latest national population census report. The age-related skeletal problems, such as osteoporosis and osteoarthritis will become more and more common in China. The structure and function of human locomotor system deteriorate with advancing age, such as bone loss, obesity reduction of muscle mass or strength and decreased coordination or balance ability, which may cause impaired physical function, higher risk of fall or fractures, reduced life quality and even higher mobility(1–3). All these factors that are more predominant in the female population after the onset of menopause. In women, almost 20% of the overall bone loss can happen and accelerate after the onset of

menopause and up to half of them suffer fragility fractures(4, 5). It is not clear whether these changes are just an effect of estrogen deficiency, or whether it is associated with malnutrition, vascular changes, or deterioration of neuromuscular function. Whatever the etiology, many studies have been performed to investigate prevention strategies including drugs, lifestyle adjustment or physical exercises(6–8).

More and more studies have reported physical exercises have multiple benefits on the locomotor system in postmenopausal women, including the prevention of muscle weakness and bone loss, thereby reducing fracture risks. The type of exercises included aerobic, stretching, resistance and balance. Different exercises focus on different effects and some exercises are span multiple categories like Yoga, Pilates and Tai Chi Chun (TCC)(9, 10). Square dance, also known as Dama square dance, is a unique phenomenon in China and is gaining popularity with middle-aged or older women due to increased awareness of health and exercise(11, 12). Every morning and evening, Chinese aunties, also called Dama in Mandarin, gather on open areas such as playgrounds, parks, streets or courts and dance according to loud music for hours. Every Chinese people or foreigners who went to China may see them. Young people always hate them because of the noisy music. However, elder women delight in it and seem never get tired, which is so interesting.

Square dance is classified as a moderate intensity exercise with all sorts of styles which can be tangos, waltzes, line-dancing, rock n' roll style jigging or just free-style bumping and grinding. In a sense, square dance is a compound exercise like TCC, while it's relatively simple and low-impact, with a shorter learning curve. Maybe these explain why square dance is becoming more popular than TCC with over 100 million participants in China(13). Published studies have reported that TCC was associated with reduced bone loss, lower fracture risks, improved cardiopulmonary function and better mental health in early postmenopausal women or other aged populations(14–16). However, the similar research about square dance is rare and limited(17, 18). Therefore, we eager to know whether or not early postmenopausal Chinese women can benefit from regular square dancing, in terms of musculoskeletal system. The study performed a horizontal comparison to explore the effects of square dance on bone mineral density (BMD), muscle strength, body flexibility and balance ability in early postmenopausal Chinese women.

Methods

Subjects and study design

A cross-sectional study was conducted to compare the postmenopausal women who participant in square dance regularly and those who without this daily fitness in terms of musculoskeletal system. Chinese postmenopausal women, aged between 50 to 60 years, who had been without menstruation for 1 to 10 years from the onset of menopause were recruited from community centers in Nantong city, Jiangsu province, China. All subjects agreed to take part in the study and signed the informed consent before the measurement. The study protocol was approved by the Clinical Research Ethics Committee of the Affiliated Hospital of Nantong University.

A standardized structured face-to-face interview was performed to collect demographic information, life styles, personal medical history, diet and menstrual status. A short version of the Health Habits and History Questionnaire (HHHQ) was used during this procedure(19). Subjects who had been practicing regular square dance without participated in other sports activities for more than 2 years and over 4 hours per week (usually more than 45 minutes per time and more than 5 times per week) were assigned to square dance group. Those postmenopausal women who had not participated in regular exercises (no more than 0.5 hour per week) were recruited as the sedentary control group. Exclusive criteria were as the follows: subjects who had injuries affecting the body function tests, subjects who were practicing other sports activities besides square dance, subjects who were receiving hormone replacement therapy, calcium supplementation or drug treatment affecting bone metabolism, subjects who had some conditions such as hypo- or hyperparathyroidism, hypo- or hyperthyroidism, renal or liver disease.

Outcome Evaluations

Bone mineral density (BMD)

Standard BMD measurements were performed by using dual-energy X-ray absorptiometry (DXA) with Hologic QDR 4500 bone densitometers (Hologic, Inc. Waltham, MA). The proximal femur and lumbar spine vertebrae were evaluated. Calibration was performed daily on a lumbar spine phantom, and the coefficient of variation was 0.7%.

Low Limb Muscle Strength, Body Flexibility And Balance Ability

Low limb muscle strength was measured for the non-dominant leg with Biodex System 3Pro isokinetic muscle strength testing instrument produced by USA. The measurement method was the same as previously reported(20). Measurements were repeated three times and the average value was used for evaluation. Body flexibility was measured by a simple trunk bend-and-reach test. The subjects stand on the measuring board with no gap between the feet, keeping the knees straight. The lowest level their fingers could reach after trunk bending forwards and downwards was recorded. Measurements were repeated three times and the average value was used for data analysis. Body balance was evaluated using a single-stance test for the non-dominant leg. Measurement was repeated three times with a time interval of 30 seconds between each measurement. The average time was used for evaluation.

Statistics

Data was presented as mean \pm SD. Independent two-tailed Student's t-test was used for data analysis. P-value of less than 0.05 was regarded as statistically significant, all the analyses were performed with SPSS software (version 13.0, Chicago, USA).

Results

Finally, a total of 152 postmenopausal women were evaluated in this study, with 74 subjects in the square dance group and 78 subjects in the sedentary control group. There was no significant difference

about demographic characteristics, dietary calcium intake and daily activity between two groups. The details of the comparison were showed in Table 1.

Table 1
Demographic characteristics, dietary calcium intake and daily activity between the square dancing and sedentary control groups.

	Square dancing group N = 74	Control group N = 78	P Value
Age (years)	56.2 ± 3.4	55.4 ± 2.8	0.325
Height (cm)	155.2 ± 4.8	156.1 ± 5.7	0.681
Weight (kg)	55.5 ± 6.3	55.2 ± 5.8	0.837
BMI (kg/m ²)	23.0 ± 3.1	22.7 ± 3.2	0.812
Age at menopause	50.8 ± 3.5	50.3 ± 3.4	0.562
Years since menopause	5.4 ± 3.2	5.1 ± 3.5	0.880
Calcium intake (mg/week)	6052 ± 516	6134 ± 580	0.662
Outdoor walking (h/week)	10.5 ± 3.2	9.8 ± 4.5	0.762

The comparison of BMD between two groups was summarized in Table 2. Subjects in the square dance group had higher lumbar spine BMD and total hip BMD than those in the control group (0.874 ± 0.120 vs 0.812 ± 0.118 , $p < 0.05$; 0.788 ± 0.104 vs 0.744 ± 0.112 , $p < 0.05$; respectively). There was no significant difference of femoral neck BMD between two groups (0.704 ± 0.098 vs 0.688 ± 0.100 , $p = 0.28$). Subjects in the square dance group had higher strength of hip extension (226.5 ± 22.6 N vs 191.8 ± 26.1 N, $P < 0.01$), knee extension (202.4 ± 19.6 N vs 173.5 ± 23.5 N, $P < 0.01$) and ankle dorsal flexion (124.8 ± 21.3 N vs 191.8 ± 26.1 N, $P < 0.01$), Single-stance time was significantly longer in square dance group than in control group (116.5 ± 40.2 s vs 70.8 ± 58.4 s, $P < 0.05$). No significant difference was found in trunk bend-and-reach between these two groups (-2.4 ± 8.8 cm vs 1.8 ± 8.0 cm, $P = 0.12$). The details of results of functional tests were summarized in Table 3.

Table 2
Comparison of BMD between square dancing and sedentary control groups.

	Square dancing group N = 74	Control group N = 78	P Value
Spine BMD	0.874 ± 0.120	0.812 ± 0.118	0.01
Femoral neck BMD	0.704 ± 0.098	0.688 ± 0.100	0.48
Total hip BMD	0.788 ± 0.104	0.742 ± 0.112	0.02

Table 3
Comparison of functional test between square dancing and sedentary control groups.

	Square dancing group N = 74	Control group N = 78	P Value
Hip extension strength (N)	226.5 ± 22.6	191.8 ± 26.1	< 0.01
Knee extension strength (N)	202.4 ± 19.6	173.5 ± 23.5	< 0.01
Ankle flexion strength (N)	124.8 ± 21.3	102.4 ± 20.4	< 0.01
Trunk bend-and-reach (cm)	-2.4 ± 8.8	1.8 ± 8.0	0.12
Single-stance time (s)	116.5 ± 40.2	70.8 ± 58.4	0.02

Discussion

The results of this study showed that subjects who participated in regular square dance had higher lumbar spine BMD and total hip BMD than sedentary control subjects. In addition, the function tests indicated higher strength of low limb muscles and better balance ability in square dance group. These results suggested that regular square dance may be beneficial for the locomotor system in early postmenopausal women.

Different from western countries, those high-impact or difficult exercises, such as running, jumping, swimming or weight lifting, may not be preferred by postmenopausal Chinese women. They prefer simple, low-impact and free-style exercises rather than go to the fitness center. As a unique Chinese exercise, TCC has been recommended to elder Chinese subjects for hundreds of years. TCC may be classified as a multicomponent exercise involving low velocity of muscle contraction and neuromuscular coordination. Square dance is something like TCC and has been becoming more popular all over China than TCC. It is still unclear whether square dance has similar benefits to multiple systems like TCC and our study provided some reference value in the field.

The results of this study showed that square dance was correlated with high BMD density of lumbar spine and total hip except femoral neck. Square dance is a relatively low-impact exercise with small size of joint movement and does not apply much bending force to femoral neck, which is different from those high-impact exercises such as running or jumping(21). Higher low limb muscle strength and longer single-stance time were detected in subjects of square dance group, which suggested that square dance was associated with better muscle function and body balance. Improved muscle function and balance ability should be considered as important as increased BMD, because they can help prevent unexpected fall, decreasing risk of fragility fractures. On the other hand, the results showed no significant difference about body flexibility between two groups. Square dance is relatively skill-free without any special style and subjects may perform any actions they like easily on a comfortable state. They do not actively extend their muscles, ligaments or joints to a large range, which therefore have limited effects on the body flexibility.

Several limitations of our study exist and deserve mention. Firstly, we presented a pilot cross-sectional analysis with a limited sample size, which influenced the evidence level of this study. Secondly, due to the obvious diversity of the square dance, we have no accurate data about special intensity or style of every subjects, increasing the heterogeneity of our study. Thirdly, all the data of square dance was collected by self-report, it may have personal bias. Lifestyle factors, different styles or intensities of square dance, and comprehensive dietary patterns containing calcium intake should be included in the finally evaluation. We have been performing a prospective randomized case-control study including more subjects to investigate the effect the square dance on musculoskeletal system in early postmenopausal Chinese women and the outcomes will be reported in the near future.

Conclusion

In summary, this cross-sectional study showed postmenopausal Chinese women who participate in square dance regularly had higher BMD, stronger low limb muscle and improved body balance ability. As a simple, free-style and multicomponent exercise like TCC, square dance may have a protective effect on postmenopausal deterioration of musculoskeletal system and is worth promoting. Further longitudinal research should be carried out to investigate the effect of square dance on bone health in early postmenopausal Chinese women.

List Of Abbreviations

BMD: Bone mineral density; DXA: dual-energy X-ray absorptiometry; TCC: Tai Chi Chun; HHHQ: Health Habits and History Questionnaire

Declarations

Ethics approval and consent to participate

This study was approved by the Clinical Research Ethics Committee of The Affiliated Hospital of Nantong University. The study was conducted according to the ethical principles stated in the Declaration of Helsinki.

Consent for publication

All the participants in this study have given their informed consent for the article to be published.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the National Natural Science Foundation of China (No. 81501913), Natural Science Foundation of Jiangsu province (No. BK20151275) and Postgraduate Research & Practice Innovation Program of Jiangsu Province (NO. SJCX19_0865).

Authors' contributions

Chen Yao, Jie Sun, Fan Liu and Yafeng Zhang conceived and designed the study. Jie Sun, Bo Zhang, Ziping Wang and Jiancheng Wu performed the DXA, functional test and collected the rude data. Yafeng Zhang and Zhenyu Zhou performed the statistical analysis. Chen Yao and Yafeng Zhang wrote the manuscript. All authors approved the final version to be published.

Acknowledgements

The authors would like to thank all the participants for their contributions to this study.

References

1. Gomez-Cabello A, Ara I, Gonzalez-Aguero A, Casajus JA, Vicente-Rodriguez G. Effects of training on bone mass in older adults: a systematic review. *Sports Med.* 2012;42(4):301-25.
2. Krist L, Dimeo F, Keil T. Can progressive resistance training twice a week improve mobility, muscle strength, and quality of life in very elderly nursing-home residents with impaired mobility? A pilot study. *Clin Interv Aging.* 2013;8:443-8.
3. Fien S, Henwood T, Climstein M, Keogh JW. Feasibility and benefits of group-based exercise in residential aged care adults: a pilot study for the GrACE programme. *PeerJ.* 2016;4:e2018.
4. Compston JE, McClung MR, Leslie WD. Osteoporosis. *Lancet.* 2019;393(10169):364-76.

5. Kemmler W, Haberle L, von Stengel S. Effects of exercise on fracture reduction in older adults: a systematic review and meta-analysis. *Osteoporos Int.* 2013;24(7):1937-50.
6. Hoke M, Omar NB, Amburgy JW, Self DM, Schnell A, Morgan S, et al. Impact of exercise on bone mineral density, fall prevention, and vertebral fragility fractures in postmenopausal osteoporotic women. *J Clin Neurosci.* 2020;76:261-3.
7. Lems WF, Dreinhofer KE, Bischoff-Ferrari H, Blauth M, Czerwinski E, da Silva J, et al. EULAR/EFORT recommendations for management of patients older than 50 years with a fragility fracture and prevention of subsequent fractures. *Ann Rheum Dis.* 2017;76(5):802-10.
8. Wang H, Dai X, Wu J, Wu X, Nie X. Influence of urban green open space on residents' physical activity in China. *BMC Public Health.* 2019;19(1):1093.
9. Benedetti MG, Furlini G, Zati A, Letizia Mauro G. The Effectiveness of Physical Exercise on Bone Density in Osteoporotic Patients. *Biomed Res Int.* 2018;2018:4840531.
10. Xie BC, Chen SC, Wu JH, Wang QH, Zhou CH, Xu DH. The effects of Tai Chi exercise on bone mineral density in postmenopausal women. *J Sports Med Phys Fitness.* 2019;59(6):1093-4.
11. Zhou L. Music is not our enemy, but noise should be regulated: thoughts on shooting/conflicts related to Dama square dance in China. *Res Q Exerc Sport.* 2014;85(3):279-81.
12. Peng F, Yan H, Sharma M, Liu Y, Lu Y, Zhu S, et al. Exploring factors influencing whether residents participate in square dancing using social cognitive theory: A cross-sectional survey in Chongqing, China. *Medicine (Baltimore).* 2020;99(4):e18685.
13. Xiao J, Hilton A. An Investigation of Soundscape Factors Influencing Perceptions of Square Dancing in Urban Streets: A Case Study in a County Level City in China. *Int J Environ Res Public Health.* 2019;16(5).
14. Jia X, Jiang C, Tao J, Li Y, Zhou Y, Chen LD. Effects of core strength training combined with Tai Chi Chuan for the musculoskeletal system and cardiopulmonary function in older adults: A study protocol for a randomized controlled trial. *Medicine (Baltimore).* 2018;97(35):e12024.
15. Wayne PM, Kiel DP, Buring JE, Connors EM, Bonato P, Yeh GY, et al. Impact of Tai Chi exercise on multiple fracture-related risk factors in post-menopausal osteopenic women: a pilot pragmatic, randomized trial. *BMC Complement Altern Med.* 2012;12:7.
16. Guo C, Xiang G, Xie L, Liu Z, Zhang X, Wu Q, et al. Effects of Tai Chi training on the physical and mental health status in patients with chronic obstructive pulmonary disease: a systematic review and meta-analysis. *J Thorac Dis.* 2020;12(3):504-21.
17. Wang S, Yin H, Meng X, Shang B, Meng Q, Zheng L, et al. Effects of Chinese square dancing on older adults with mild cognitive impairment. *Geriatr Nurs.* 2019.
18. Zhang X, Ni X, Chen P. Study about the effects of different fitness sports on cognitive function and emotion of the aged. *Cell Biochem Biophys.* 2014;70(3):1591-6.
19. Boeckner LS, Pullen CH, Walker SN, Abbott GW, Block T. Use and reliability of the World Wide Web version of the Block Health Habits and History Questionnaire with older rural women. *J Nutr Educ Behav.* 2002;34 Suppl 1:S20-4.

20. Song QH, Zhang QH, Xu RM, Ma M, Zhao XP, Shen GQ, et al. Effect of Tai-chi exercise on lower limb muscle strength, bone mineral density and balance function of elderly women. *Int J Clin Exp Med*. 2014;7(6):1569-76.
21. Montgomery GJ, Abt G, Dobson CA, Evans WJ, Aye M, Ditroilo M. A 12-month continuous and intermittent high-impact exercise intervention and its effects on bone mineral density in early postmenopausal women: a feasibility randomized controlled trial. *J Sports Med Phys Fitness*. 2020;60(5):770-8.