Quality of Life in Women With Postmenopausal Osteoporosis - Pilot Studies in The Łódź Province

Bogumiła Górczewska (✉ bogumila.gorczevska@gmail.com)

CKD ŁÓDŹ <https://orcid.org/0000-0001-6820-7401>

Elżbieta Jakubowska-Pietkiewicz

Department of Peadiatrics, Pathology Neoanathology and Metabolic Desease

Research

Keywords: QUALEFFO-41, VAS, postmenopausal osteoporosis, women, quality of live

Posted Date: September 28th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-56428/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Osteoporosis is a social disease and epidemic of the 21st century, and is also a growing problem in the ageing Polish society. The aim of the study was to assess the quality of life (QoL) in women with postmenopausal osteoporosis.

Material and methods The study was an outpatient, questionnaire-based poll, carried out from VI.2018 to V.2019. A group of 198 women, aged 72.31 ± 8.59 years with diagnosed postmenopausal osteoporosis in two Osteoporosis Clinics in the city of Łódź (Poland) were included. The paper uses: the Polish version of the QUALEFFO 41 quality of life scale, the VAS scale and a survey of my own authorship. The data were analysed using multiple linear regression.

Results The average result of QUALEFFO-41 was 40.26 ± 16.92 points. Analysing individual QUALEFFO-41 domains the lowest quality of life was found in the mental function domain (mental function 48.49 ± 18.06 points). The visual analogue scale of pain was 4.87 ± 2.39 points. The VAS scale positively correlated with all QUALEFFO 41 domains. Marital status, education and financial situation had a significant statistical impact on the quality of life ($p < 0.05$). Multiple regression step-by-step analysis showed that pain expressed on the VAS scale, pain awakening from sleep, taking sleeping pills and older age of respondents deteriorate the quality of life.

Conclusion The analysis showed that Polish women with postmenopausal osteoporosis have an average quality of life. In Poland the assessment of quality of life of chronically ill patients and routine evaluation of pain severity on the VAS scale is not common.

Background

The World Health Organization reports that the number of 60 or more years-old people over represents more than 11% of the world's population and is projected to increase to about 22% by 2050 [1]. As life expectancy increases, the number of cases of osteoporosis continues to increase. This statement is disturbing for countries that have an ageing population, including Poland. It is estimated that in Poland, 30–40% of postmenopausal women are struggling with osteoporosis [2].

The greatest complication of osteoporosis is bone fractures, which are associated with pain and cause a reduction in physical fitness and social functions, which are an aspect of health-related quality of life (HRQoL). Kanis et. al. assume that due to demographic changes in the world's population, the annual number of bone fractures in people with osteoporosis will increase from 3.5 million in 2010 to 4.5 million in 2025 – what gives us an increase of 28% [3]. In a recently published study of Glinkowski W. et al. it has been demonstrated that femoral neck fractures in people with osteoporosis over 50 years of age are more and more common in the Polish population [4]. In many people, bone fracture results in loss of independence, increased pain and deterioration of health-related quality of life, which may affect the mental condition of the patient and cause limitations in social life.

The ICUROS results revealed that the quality of life substantially deteriorates immediately after the bone fracture and only about 18 months after the bone fracture, the mental condition improves, but still remains lower than before the fracture [5]. Moreover, bone fracture is associated with the development of other harmful complications such as disability, depression, cardiovascular disease, stroke and increased mortality. The increasing number of bone fractures in the course of osteoporosis, especially in developed countries, is a serious health and economic problem [6]. It is estimated that in Poland the costs of osteoporosis treatment (i.e. The costs in the first year after the fracture, costs in subsequent years after the fracture and the cost of pharmacological intervention, including administration and monitoring) will increase by 27% by 2025 [7]. By contrast, the costs for all countries in the European Union will increase by 25% (i.e. from € 37. 4 billion in 2010 to € 46. 8 billion in 2025) [7].

Moreover, Curtis E. M. et al, (2020) point out that osteoporosis, compared to other non-communicable diseases, is rarely adequately funded by the government and healthcare providers, but unfortunately, this trend is also observed in Poland. Analysing individual EU countries, there are 4.3 densitometric devices (DXA-dual-energy X-ray absorptiometers) per million inhabitants in Poland, fewer only in Lithuania 3.4 (DXA/million), Luxembourg 2. 0 (DXA/million) and Bulgaria 1. 2 (DXA/million) [8]. On the other hand, the largest number of DXA examination devices per one million inhabitants in the EU, with as many as 53 in Belgium, shows the scale of the problem of difficulties in access to diagnostics and appropriate monitoring and treatment of this disease (over the last 5 years, the rates of osteoporosis treatment in the whole EU have decreased significantly) [8].

It should be remembered that as the age increases, sleep problems, affecting the quality of life in the elderly, also increase [9]. Sleep problems are more common in women than in men and are related to menopause. Polo-Cantola reported that among women aged 50–64 years 25% of them had sleep issues and 15% of them had serious sleep disorders, which significantly affected their quality of life [10]. There are many detailed questionnaires available to measure the quality of life in people with osteoporosis [11], but none of them is as specific as QUALEFFO-41 [12].

Restrictions related to old age (e.g: difficulties in moving around, loneliness, sleep disorders), chronic pain, difficulties in accessing diagnostics (DXA) [8] and increased risk of bone fractures, are just some of the issues faced by people with osteoporosis every day. A great challenge is to maintain a good mental state and high quality of life. Therefore, the main objective of our study was the general quality of life assessment (QoL) among female respondents with osteoporosis, depending on their sociodemographic situation and pain assessed on the VAS scale (regardless of fracture history).

Materials And Method

The survey was conducted from June 2018 until December 2018. The patients in two Osteoporosis Clinics in Łódź were examined. The study involved 198 postmenopausal women aged 72.31 ± 8.59 (range 51-90 years). The main criterion for the selection of patients was postmenopausal osteoporosis

diagnosed in the patient's history by the physician treating them in the Osteoporosis Outpatient Clinic according to ICD 10 - M81.0.

Other criteria for including women in the study group were:

- consent to participate in the study;
- full completion of the AIS and SLWS questionnaire;
- no hospitalization during the last six months in the hospital;

Women excluded from the study:

- women with active cancer, with malignant bone metastases;
- women who fail to complete the AIS and SLWS questionnaires in full;
- women with secondary osteoporosis;
- women treated with glucocorticoids;
- women with currently broken bones;
- women hospitalized in the last 6 months in a hospital;

The research method was a diagnostic survey and the tools were as follows:

1. The quality of life was assessed on the basis of the Polish version of the detailed questionnaire - QUALEFFO-41, which consists of 41 questions, in which we distinguish 5 main domains: pain (5 questions), physical functions (17 questions), social functions (7 questions), general health perception (3 questions) and mental functions (9 questions) [13]. In our study the field of physical functions was divided into: physical activities (4 questions), homework (5 questions) and mobility (8 questions). The analysis of the result was performed according to the algorithm proposed by the International Osteoporosis Foundation on a scale from 0 to 100. In interpreting the results of the questionnaire the principle was applied that the more points the worse the quality of life. When filling in the questionnaire, one should choose only one answer for each question. The results of each domain and the total QUALEFFO 41 result were analysed.
2. The VAS scale for assessing the severity of pain during the last week.
3. Author's questionnaire, consisting of questions about socio-demographic data (e.g. age, marital status, place of residence, financial situation).

The vast majority of study participants had abnormal body weight, average BMI was $25.71 \pm 4.73 \text{ kg/m}^2$. Moreover, the study group presented the average body weight of $64.5 \pm 11.66 \text{ kg}$ and the average height was $158.6 \pm 6.52 \text{ cm}$. Almost half of the respondents (47%, $n=93$) were married, 36% ($n=72$) were widows, 10% ($n=19$) divorced, and 7.0% ($n=14$) had never been married. Most of the respondents had secondary education (35%, $n=70$) and higher education (25%, $n=50$), only 21% ($n=41$) had primary education, and vocational education 19% ($n=37$). In the studied group, osteoporosis was diagnosed at the age of 61.82 ± 9.74 on average. On the other hand, the duration of osteoporosis among the respondents was on average 10.7 ± 8.53 years. In addition, 29% of respondents ($n=57$) stated that osteoporosis occurred in the

immediate family, of which 56% (n=32) in mothers and 31% (n=18) in the sisters of the surveyed women. Characteristics of the studied group in Table 1.

The respondents slept 6.56 h (about 7 h) on average. Sleeping pills were taken by 51% (n=101) of women in the group. Sleep problems caused by pain were reported by 49% (n=98) of respondents.

Ethical issues

The survey was voluntary and was conducted according to the principles of human research specified in the Helsinki Declaration. The respondents signed an informed consent form and were informed that the study would be anonymous, according to the currently applicable regulations and the provisions of the GDPR. Before filling the questionnaires, the patients were informed about the purposefulness of the study and instructed how to complete the questionnaires correctly. Each person qualified for the examination provided answers in the office, in which they stayed together with the surveyor. Some patients asked the subject to read aloud the questions, which they then answered verbally, and which was circled in questionnaires. The duration of the survey was adjusted to the individual capabilities of the examined women. The study was approved by the Bioethics Committee of the Medical University of Lodz (Resolution No. RNN / 215 / 18KE of June 12.2018).

Statistical analysis

The results obtained from questionnaires were statistically analyzed. The values of the analyzed parameters were provided with the use of: average, median, standard deviation, minimum and maximum values. The Shapiro-Wilk test was used to test the normality of the quantitative variable distribution. Correlation coefficients between quantitative variables were calculated using the Spearman or Pearson correlation, accordingly to the distribution of the variable. Comparisons quantitative variables were made with the use of the Student's t-test, ANOVA or Mann-Whitney U test, correspondingly to the distribution of the variable. Data were analyzed using multiple linear regression. The adopted level of statistical significance was $p < 0.05$. The static analysis was performed with the use of the Statistica 13 PL program (StatSoft, Tulusa, OK, USA).

Results

The surveyed women evaluated their quality of life at the average level and scored an average result of 40.26 ± 16.92 (MEDIANA 39), according to the QUALEFFO 41 scale.

Detailed analysis of QUALEFFO 41 questionnaire shows that the surveyed women reported the least number of problems in the domain of the activities of daily living, on average 24.58 ± 19.49 points (MEDIANA 25) and locomotion functions on average 34.56 ± 24.80 points (MEDIANA 29). The domain of pain amounted to 42.27 ± 24.47 (MEDIANA 40) on average and was rated at the same level as the domain of housework 37.27 ± 26.64 (MEDIANA 40). Social functions (leisure, social activities) were assessed on average at 44.29 ± 23.34 (MEDIANA 46), while general health perception at 44.61 ± 28.34

(MEDIANA 42). The respondents assessed mental function as the worst, with the average value of 48.49 ± 18.06 (MEDIANA 50). The domain activities of daily living, was rated twice as high as the mental function domain.

Then the socio-demographic features were statistically analysed. It has been statistically significant that education, marital status and financial situation have an impact on the quality of life (Table 1). People with secondary and higher education had a higher quality of life, compared to people with primary education. In addition, the respondents, who assessed their financial situation as very good and divorced persons assessed their quality of life as the highest. The relationship between BMI, place of residence and quality of life was not statistically significant.

The respondents who had any bone fracture in their history rated their quality of life lower on the QUALEFFO 41 scale (42.05 ± 17.31 points, $n = 137$), compared to persons without fractures (36.25 ± 15.40 points, $n = 61$). The greatest loss of QoL was observed in patients with a hip fracture (51.88 ± 18.90 pts, $p = 0.00534$) followed by an ankle fracture (47.85 ± 16.53 pts, $p = 0.08168$), vertebral fracture (44.38 ± 15.41 pts, $p = 0.06346$) and upper limb fracture (40.02 ± 18.05 pts, $p = 0.95804$). The reliability index - Cronbach's alpha Cronbach in all QUALEFFO-41 total domains was high and amounted to 0.90. Cronbach's alpha internal value for the individual domains was: pain 0.87, physical functions 0.86, social functions 0.87, general health perception 0.90 and mental functions 0.89.

The reliability index - Cronbach's alpha in all QUALEFFO-41 total domains was high and amounted to 0.90. Cronbach's alpha internal value for the individual domains was: pain 0.87, physical functions 0.86, social functions 0.87, general health perception 0.90 and mental functions 0.89.

On the VAS scale, patients rated the pain as mild. The average level obtained on the VAS scale is 4.87 ± 2.39 points. A statistically significant ($p < 0.05$), high positive correlation ($r = 0.5419$) between pain assessed in the VAS scale and quality of life in the QUALEFFO-41 scale was demonstrated. The relation between VAS scale and domains of QUALEFFO-41 scale was analysed statistically and positive correlation was obtained in all domains. The results of this correlation were: pain $r = 0.5877$, activities of daily living $r = 0.4871$, jobs around the house $r = 0.5609$, mobility $r = 0.5639$, leisure and social activities $r = 0.2935$, general health perception $r = 0.1406$ and mental function $r = 0.1369$.

In order to estimate the predictors of the quality of life, a variable regression analysis was implemented. The parameters tested by univariable analysis were next analysed in a multivariable logistic regression analysis using AIC-based backward selection to drop insignificant terms from the model. The model turned out to be significant ($F = 40.019$; $p < 0.000$), and all the predictors explained together 45.33% of the dependent variable (the rest of the variation, i.e. 54.67% is explained by other factors not included in our model).

The quality of life was substantially affected by four factors: age ($\beta = 0.31$; $t = 5.49$; $p = 0.000$), pain awakening at night ($\beta = 0.14$; $t = 2.32$; $p = 0.021$), occurrence of pain ($\beta = 0.35$; $t = 5.69$; $p = 0.000$) and taking sleeping pills ($\beta = 0.15$; $t = 2.42$; $p = 0.016$).

Interpreting the estimated value of the individual parameters' assessment, we can state that with each year of life, the quality of life of the women surveyed deteriorates by 0.61% on the QUALEFFO 41 scale. On the other hand, the occurrence of painful ailments that cause nocturnal arousals in patients devolve the quality of life by 4.9%. However, an increase in the VAS scale by one point deteriorates the quality of life by 2.4% and the use of sleeping pills by 4.9%.

Discussion

The survey showed that the quality of life of the respondents was at an average level. The results obtained are similar to those previously reported by Baczyk G. et al. for women with osteoporosis tested in Wielkopolska [Poland] evaluated in the QUALEFFO 41 scale [14]. Analyzing the individual QUALEFFO-41 domains, we found the lowest quality of life in the mental function domain, which reveals, among others the emotional impact of the diagnosis made and evaluates the mental state of the patient. This result is consistent with the previous study (2016) in the Polish population, which indicated that women with osteoporosis suffer from low quality of life, especially in the domain of mental function, general health perception and social function [15]. It is suggested that the quality of life assessed on the QUALEFFO 41 scale differs between continents and ethnic groups. In most analyses, QoL was lowest in Asia and Europe and highest in North America and Oceania [16]. Further research is undoubtedly needed to understand this phenomenon.

The study shows that marital status, education and financial situation have a significant statistical impact on the quality of life. We have shown that women who were divorced assessed their mental state more positively.

Another discovery in this study is that the patient's level of education is also a factor determining the quality of life. In our survey, secondary and higher education was a protective factor against poorer quality of life. Similar relations were described by de Oliveira Ferreira N et al. and Abourazzak, et. al. [17, 18]. This can be explained by the fact that women with higher education usually look for more information about their illness (they broaden their knowledge) [18], so that they take supplements more often (e.g. vitamin D3, calcium preparations) or introduce hormone replacement therapy [19].

We have also shown that the financial situation affects the quality of life in women with postmenopausal osteoporosis. It is suggested that a higher level of wealth provides access to better medical services, e.g. DXA tests [20]. Other authors confirm our results that well-paid work improves the quality of life assessed in the QUALEFFO 41 scale in women with postmenopausal osteoporosis [17]. The literature reports show that the factors significantly related to quality of life were body mass index (BMI), race, school education, age at menopause and implementation of hormone therapy [17]. Osteoporosis reduces the patient's quality of life by affecting all spheres of the patient's life. This shows that further interventions are needed to improve psychological support and education of patients and to improve long-term treatment of osteoporosis.

Research conducted by Janiszewska M. et. al. among 292 women aged 51–83 years in Lublin (Poland, 2014) showed that better educated women living in cities with very good or good social and welfare conditions showed a much higher level of knowledge about osteoporosis prevention [2]. In addition, the same authors concluded that also women undergoing bone densitometric testing do not have sufficient knowledge about osteoporosis prevention. The low level of women's knowledge of prevention, diagnosis and treatment of osteoporosis additionally causes stress, triggers negative emotions and leads to a decrease in their quality of life.

The results of this study showed that the overall (total) QoL among women with a bone fracture in the survey was lower than among respondents without fractures. Bączyk et. al. Lesnyak O. et al, Gold et al showed the same dependence [14, 21, 22]. Interestingly, we showed significantly, in terms of statistics, that women with a femoral neck fracture more negatively assessed the quality of life than people with a fracture in other places. A similar relationship has been recently observed (2020) in patients over 50 years of age with osteoporosis in Russia, where the quality of life of patients with femoral neck fracture has been shown to be lower than that of the spine, humerus and ankle [21]. Hagino and et al, after a year-long observation of women with osteoporosis and the femoral neck fracture, concluded that the loss of QoL did not improve to the level from before the bone fracture and was more severe than after the wrist fracture [23]. However, so far, vertebral fracture has been reported as one of the factors contributing to the worse quality of life in women with postmenopausal osteoporosis [15].

On the other hand, it should be noted that regardless of the presence of bone fractures, women with osteoporosis show a reduced quality of life in different QUALEFFO-41 domains. This is confirmed by reports of, among others Hopman et al. (2019) [24], Ciubean et al. [25] and Bianchi et. al [26]. This may be related to increased chest kyphosis, decreased lean muscle mass and generalized muscle weakness in these patients [27]. Therefore, the authors suggest that HRQOL should be tested before any bone fracture to develop appropriate interventions for all stages of the disease [24]. It would therefore be appropriate to widen this study group in order to analyse this issue more closely.

This study indicates that one of the factors related to the perception of general health is sleep. Poor quality of sleep (sleep disorders) may be associated with osteoporosis [28], by affecting the bone microarchitecture [29]. It is known that a good night's sleep is a key factor in maintaining quality of life. [30] In the current study, we have noted that taking sleeping pills deteriorates quality of life. These observations can be explained by an earlier study, which suggests that the use of sleeping pills in women may be associated with worse bone health and should be considered when reviewing older patients [29].

Moreover, awakening from sleep through pain affects the mental state of the patients. The existing literature describes that women in Sweden, after a compression fracture of their spine, complained about sleep problems at night, woke up with severe pain, and many of them took sleeping pills for better rest [31]. We have not shown a statistically significant relationship between sleep length and QUALEFFO 41. The literature (2020) describes that short sleep time (less or equal to 5 h) was associated with lower BMD and higher risk of osteoporosis [32]. Ochs-Balcom HM et al in a group of 11 084 postmenopausal women

observed that short sleep time (less than or equal to 5 h) was associated with lower BMD and higher risk of osteoporosis.

However, other authors emphasize that both short and long sleep time in middle aged persons and elderly people is associated with significantly increased risk of osteoporosis, and an appropriate sleep time may help in delaying or preventing osteoporosis [33, 34]. However, discussions on the appropriate length of sleep in people with osteoporosis are still ongoing.

Pain deserves special attention in this study because it was associated with poor quality of life. There was a good correlation between VAS and QUALEFFO-41 in all domains. A similar relationship in women with osteoporosis in Norway was observed by Stanghelle Brita et al. [35]. The authors suggest that pain intensity negatively affects HRQoL in women with osteoporosis with or without vertebral fracture [36]. Moreover, chronic pain contributes to unbalanced gait, which increases the risk of falls in older people [37]. And Albayrak et. al. found no statistically significant differences in total VAS and QUALEFFO-41 results ($p > 0.05$) [38]. It would be advisable to assess the pain at the beginning of therapy and then monitor its level during treatment.

With the extension of life, a deterioration in quality of life was observed. These results are similar to those of other tests using QUALEFFO 41. In many studies, the authors stressed that the occurrence of bone fracture and age significantly affect HRQoL in people with osteoporosis [15, 22]. Only appropriate counselling interventions, support and psychological care for the elderly can help patients to develop effective strategies for the admission and management of the disease [39]. In addition, improving medical care in Poland would require the implementation of a national fracture prevention programme. In our work we used the Polish version of the QUALEFFO 41 questionnaire. Alpha Cronbach's mirroring of the reliability of the Polish version of QUALEFFO-41 for all domains was satisfactory and falls within the scope of internal consistency. Our Cronbach's alpha values are similar to those presented for the Polish population in the study Bączek et. al [14].

Our study has some limitations that should be mentioned. The results of the study concern women with postmenopausal osteoporosis from one city, it would be advisable to conduct the study in another city to verify the results. After the extension of the examination time, the same respondents could be examined at the turn of two or three years, analysing the different stages of treatment of osteoporosis and fracture history. After increasing the research group, it would be advisable to analyse the factors influencing the hygiene and quality of sleep of the respondents. In addition, an analysis could be made according to the current bone mineral density.

To sum up, the quality of life of women with postmenopausal osteoporosis in the Łódź Province is at an average level. The unfavourable trend observed in this study, corresponding to the deterioration of mental condition, is associated with the occurrence of pain, which should be considered a serious health problem. Moreover, factors affecting the quality of life have been shown: awakening pain during sleep and taking sleeping pills. It would be advisable to introduce new methods of psychological assistance,

sleep testing (hypersomnia), implementation of more effective methods (pharmacotherapy, physiotherapy) in reducing pain and improving sleep quality in this group of patients.

Conclusion

Long-term medical care in patients with osteoporosis should also include an assessment of their quality of life, which could indicate areas for wider specialist intervention.

List Of Abbreviations

QoL - quality of life assessment

BMI - body mass index

Declarations

Ethics approval and consent to participate - The study was approved by the Bioethics Committee of the Medical University of Lodz (Resolution No. RNN / 215 / 18KE of June 12.2018).

Consent for publication - 'Not applicable'

Availability of data and materials - 'Not applicable'

Competing interests - 'Not applicable'

Funding - 'Not applicable'

Authors' contributions – 50 % BG and 50% EJP

References

1. Ageing and Life Course. [(accessed on 18 June 2019)]; Available online: <http://www.who.int/ageing/about/facts/en/>
2. Janiszewska M, Firlej E, Żołnierczuk-Kieliszek D, Dziedzic M. (2016). Knowledge about osteoporosis prevention among women screened by bone densitometry. *Menopause review*, 15(2), 96–103. <https://doi.org/10.5114/pm.2016.61192>
3. Kanis J. A., Cooper C., Rizzoli R., Reginster J. Y., Scientific Advisory Board of the European Society for Clinical and Economic Aspects of Osteoporosis (ESCEO) and the Committees of Scientific Advisors and National Societies of the International Osteoporosis Foundation (IOF) (2019). European guidance for the diagnosis and management of osteoporosis in postmenopausal

- women. *Osteoporosis international* : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA, 30(1), 3–44. <https://doi.org/10.1007/s00198-018-4704-5>
4. Glinkowski W, Narloch J, Krasuski K, Śliwczyński A. The Increase of Osteoporotic Hip Fractures and Associated One-Year Mortality in Poland: 2008-2015. *J Clin Med*. 2019;8(9):1487. <https://doi.org/10.3390/jcm8091487>
 5. Svedbom A, Borgström B, Hernlund E, Ström O, Alekna V, Bianchi ML, Clark P, Curiel MD, Dimai HP, Jürisson M, Kallikorm R, Lember M, Lesnyak O, McCloskey E, Sanders KM, Silverman S, Solodovnikov A, Tamulaitiene M, Thomas T, Toropectsova N, Uusküla A, Tosteson ANA, Jönsson B, Kanis JA. (2018) Quality of life for up to 18 months after low-energy hip, vertebral, and distal forearm fractures-results from the ICUROS. *Osteoporos Int*. 29:557–566.
 6. [Johnell O](#), [Kanis JA](#). (2006) An estimate of the worldwide prevalence and disability associated with osteoporotic fractures. *Osteoporos Int*. 17(12):1726-33. <https://doi.org/10.1007/s00198-006-0172-4>
 7. Hernlund, E., Svedbom, A., Ivergård, M., Compston, J., Cooper, C., Stenmark, J., McCloskey, E. V., Jönsson, B., & Kanis, J. A. (2013). Osteoporosis in the European Union: medical management, epidemiology and economic burden. A report prepared in collaboration with the International Osteoporosis Foundation (IOF) and the European Federation of Pharmaceutical Industry Associations (EFPIA). *Archives of osteoporosis*, 8(1-2), 136. <https://doi.org/10.1007/s11657-013-0136-1>
 8. Curtis, E. M., Woolford, S., Holmes, C., Cooper, C., & Harvey, N. C. (2020). General and Specific Considerations as to why Osteoporosis-Related Care Is Often Suboptimal. *Current osteoporosis reports*, 18(1), 38–46. <https://doi.org/10.1007/s11914-020-00566-7>
 9. Andenæs, R., Småstuen, M. C., Misvær, N., Ribu, L., Vistad, I., & Helseth, S. (2020). Associations between menopausal hormone therapy and sleep disturbance in women during the menopausal transition and post-menopause: data from the Norwegian prescription database and the HUNT study. *BMC women's health*, 20(1), 64. <https://doi.org/10.1186/s12905-020-00916-8>
 10. Polo-Kantola P. Sleep problems in midlife and beyond. 2011; 68(3):224–232. <https://doi.org/10.1016/j.maturitas.2010.12.009>.
 11. Madureira, M. M., Ciconelli, R. M., & Pereira, R. M. (2012). Quality of life measurements in patients with osteoporosis and fractures. *Clinics (Sao Paulo, Brazil)*, 67(11), 1315–1320. [https://doi.org/10.6061/clinics/2012\(11\)16](https://doi.org/10.6061/clinics/2012(11)16)
 12. Koevska V, Nikolikj-Dimitrova E, Mitrevska B, Gjeracaroska-Savevska C, Gocevska M, Kalcovska B. Effect of Exercises on Quality of Life in Patients with Postmenopausal Osteoporosis - Randomized

Trial. Open Access Maced J Med Sci. 2019;7(7):1160–1165. Published 2019 Apr 14.
<https://doi.org/10.3889/oamjms.2019.271>

13. Bączyk G, Opala T, Kleka P. Quality of life in postmenopausal women with reduced bone mineral density: psychometric evaluation of the Polish version of QUALEFFO-41. Arch Med Sci. 2011;7(3):476–485. <https://doi.org/10.5114/aoms.2011.23415>
14. Bączyk G, Opala T, Klejewski A, Tesarska B. Evaluation of the functioning of women with osteoporosis with vertebral fractures and without vertebral fractures using the Polish version of Qualeffo-41. Chir Narzadow Ruchu Ortop Pol. 2010 May-Jun;75(3):172-7.
<https://www.ncbi.nlm.nih.gov/pubmed/21038636>
15. Bączyk G, Samborski W, Jaracz K. Evaluation of the quality of life of postmenopausal osteoporotic and osteopenic women with or without fractures. Arch Med Sci. 2016;12(4):819–827.
<https://doi.org/10.5114/aoms.2015.55012>
16. van Schoor, N.M., Yu, H., Bobula, J. P. Lips. Cross-geographic region differences in quality of life in women with and without vertebral fracture. Osteoporos Int 20, 1759–1766 (2009).
<https://doi.org/10.1007/s00198-009-0853-x>
17. de Oliveira Ferreira N, Arthuso M, da Silva R, et al. Quality of life in women with postmenopausal osteoporosis: correlation between QUALEFFO 41 and SF-36. Maturitas 2009; 62:85–90.
<https://doi.org/10.1016/j.maturitas.2008.10.012>
18. Abourazzak, F.E., Allali, F., Rostom, S. et al. Factors influencing quality of life in Moroccan postmenopausal women with osteoporotic vertebral fracture assessed by ECOS 16 questionnaire. Health Qual Life Outcomes 7, 23 (2009). <https://doi.org/10.1186/1477-7525-7-23>
19. Pluskiewicz W., Adamczyk P., Czekajło A., Grzeszczak W., Drozdowska B. Influence of education, marital status, occupation, and the place of living on skeletal status, fracture prevalence, and the course and effectiveness of osteoporotic therapy in women in the RAC-OST-POL Study. Bone Miner. Metab. 2014;32:89–95. <https://doi.org/10.1007/s00774-013-0471-8>
20. Holmberg, T., Möller, S., Rothmann, M. et al. Socioeconomic status and risk of osteoporotic fractures and the use of DXA scans: data from the Danish population-based ROSE study. Osteoporos Int 30, 343–353 (2019). <https://doi.org/10.1007/s00198-018-4768-2>
21. Lesnyak O., Svedbom A., Belova K., Dobrovolskaya O., Ershova O., Golubev G., Grebenshikov V., Ivanov S., Kochish A., Menshikova L., Nikitinskaya O., Nurligayanov R., Solodovniko, Toropectsova N., Varavko J., Zotkin E., Borgstrom F. Kanis J. A. (2020). Quality of life after fragility fracture in the Russian Federation: results from the Russian arm of the International Cost and Utility Related to Osteoporotic Fractures Study (ICUROS). Archives of osteoporosis, 15(1), 37.
<https://doi.org/10.1007/s11657-020-0699-6>

22. Gold, T., Williams, S. A., Weiss, R. J., Wang, Y., Watkins, C., Carroll, J., Middleton, C., & Silverman, S. (2019). Impact of fractures on quality of life in patients with osteoporosis: a US cross-sectional survey. *Journal of drug assessment*, 8(1), 175–183.
<https://doi.org/10.1080/21556660.2019.1677674>
23. Hagino H, Nakamura T, Fujiwara S, Oeki M, Okano T, Teshima R. Sequential change in quality of life for patients with incident clinical fractures: a prospective study. *Osteoporos Int*. 2009;20(5):695–702. . <https://doi.org/1007/s00198-008-0761-5>
24. Hopman WM, Berger C, Joseph L, Morin SN, Towheed T, Anastassiades T, Adachi JD, Hanley DA, Prior JC, Goltzman D; CaMos Research Group. Longitudinal assessment of health-related quality of life in osteoporosis: data from the population-based Canadian Multicentre Osteoporosis Study. *Osteoporos Int*.2019 Aug;30(8):1635-1644. <https://doi.org/10.1007/s00198-019-05000-y>
25. Ciubean, A. D., Ungur, R. A., Irsay, L., Ciortea, V. M., Borda, I. M., Onac, I., Vesa, S. C., & Buzoianu, A. D. (2018). Health-related quality of life in Romanian postmenopausal women with osteoporosis and fragility fractures. *Clinical interventions in aging*, 13, 2465–2472.
<https://doi.org/10.2147/CIA.S190440>
26. Bianchi, M. L., Orsini, M. R., Saraifoger, S., Ortolani, S., Radaelli, G., & Betti, S. (2005). Quality of life in post-menopausal osteoporosis. *Health and quality of life outcomes*, 3, 78.
<https://doi.org/10.1186/1477-7525-3-78>
27. Miyakoshi, N., Kudo, D., Hongo, M. et al. Comparison of spinal alignment, muscular strength, and quality of life between women with postmenopausal osteoporosis and healthy volunteers. *Osteoporos Int* 28, 3153–3160 (2017). <https://doi.org/10.1007/s00198-017-4184-z>
28. Sasaki N, Fujiwara S, Yamashita H, Ozono R, Teramen K, Kihara Y. Impact of sleep on osteoporosis: sleep quality is associated with bone stiffness index. *Sleep Med*.2016; 25:73-77.
<https://doi.org/10.1016/j.sleep.2016.06.029>.
29. Bevilacqua, G., Denison, H. J., Laskou, F., Jameson, K. A., Ward, K. A., Cooper, C., & Dennison, E. M. Self-reported Sleep Quality and Bone Outcomes in Older Adults: Findings from the Hertfordshire Cohort Study. *Calcif Tissue Int*. 2020; 106(5), 455–464. <https://doi.org/10.1007/s00223-020-00657-8>
30. Andenæs, R., Småstuen, M. C., Misvær, N., Ribu, L., Vistad, I., & Helseth, S. (2020). Associations between menopausal hormone therapy and sleep disturbance in women during the menopausal transition and post-menopause: data from the Norwegian prescription database and the HUNT study. *BMC women's health*, 20(1), 64. <https://doi.org/10.1186/s12905-020-00916-8>
31. Svensson HK, Olofsson EH, Karlsson J, Hansson T, Olsson LE. A painful, never ending story: older women's experiences of living with an osteoporotic vertebral compression fracture. *Osteoporos Int*. 2016;27(5):1729- <https://doi.org/10.1007/s00198-015-3445-y>

32. Ochs-Balcom HM, Hovey KM, Andrews C, Cauley JA, Hale L, Li W, Bea JW, Sarto GE, Stefanick ML, Stone KL, Watts NB, Zaslavsky O, Wactawski-Wende J . Short Sleeps Associated With Low Bone Mineral Density and Osteoporosis in the Women's Health Initiative. *J Bone Miner Res*. 2020 Feb;35(2):261-268. <https://doi.org/10.1002/jbmr.3879>.
33. Wang D, Ruan W, Peng Y, Li W. Sleepduration and the risk of osteoporosis among middle-aged and elderly adults: a dose-response meta-analysis. *Osteoporos Int*. 2018 Aug;29(8):1689-1695. <https://doi.org/10.1007/s00198-018-4487-8>.
34. Moradi S, Shab-Bidar S, Alizadeh S, Djafarian K . Association between sleep duration and osteoporosis risk in middle-aged and elderly women: A systematic review and meta-analysis of observational studies. 2017; 69:199-206. <https://doi.org/10.1016/j.metabol.2017.01.027>.
35. Stanghelle B, Bentzen H, Giangregorio L, Pripp AH, Bergland A. Associations between health-related quality of life, physical function and pain in older women with osteoporosis and vertebral fracture. *BMC geriatrics* 19,1 298. 4 Nov. 2019, <https://doi.org/10.1186/s12877-019-1268-y>
36. Hubscher M, Vogt L, Schmidt K, Fink M, Banzer W. Perceived pain, fear of falling and physical function in women with osteoporosis. *Gait & posture*. 2010;32(3):383–5. <https://doi.org/10.1016/j.gaitpost.2010.06.018>
37. Pain Is Associated With Poor Balance in Community-Dwelling Older Adults: A Systematic Review and Meta-analysis Tatsuya H, T, Okubo Y Sturnieks DL Stephen R. *Journal of the American Medical Directors Association*,2020, Volume 21, Issue 5, 597 - 603.e8 <https://doi.org/10.1016/j.jamda.2020.02.011>
38. Albayrak I, Aydogmus M, Ozerbil OM, Levendoglu F The association between bone mineral density, quality of life, quality of sleep and fatigue. *Acta Clin Belg*. 2016 Apr;71(2):92-8. <https://doi.org/10.1179/2295333715Y.0000000061>.)
39. Bianchi ML, Orsini MR, Saraifoger S, Ortolani S, Radaelli G, Betti S. Quality of life in post-menopausal osteoporosis. *Health Qual Life Outcomes*. 2005;3:78. <https://www.ncbi.nlm.nih.gov/pubmed/16321148>

Table

Table 1 Characteristics of the study population. Relations QUALEFFO 41

		Liczebność [n]	QUALEFFO 41[pkt ±]**	ANOVA (p-value)
BMI*	Underweight	13	44.99±24.38	p=0.25157
	Normal body weight	78	37.49±15.98	
	Overweight	75	41.24±16.37	
	Obesity	32	42.82±16.63	
Education	None	41	53.36±15.47	p=0.00000
	Completed primary	37	41.24±15.18	
	Completed secondary	70	35.04±15.88	
	Completed tertiary	50	36.11±15.18	
Place residence	Village	17	47.61±14.16	p=0.13686
	City of above 50.000 inhabitants	26	35.74±18.69	
	City 50 000- 100.000 inhabitants	112	38.79±15.36	
	City of over 100.000 inhabitants	43	40.76±17.23	
Living conditions	Very good	17	35.81±14.13	p=0.00000
	Good	105	36.03±14.40	
	Unsatisfactory	51	42.72±18.61	
	Very bad	25	56.08±15.19	
Marital status	Never married	14	37.91±22.79	p=0.00261
	Married	93	39.45±16.04	
	Widowed	72	44.74±16.99	
	Divorced	19	29.03±9.08	
Any fractures	Yes	137	42.05±17.31	p=0.02554
	No	61	36.25±15.40	
Pain weaknes up durning sleep	Yes	98	47.28±15.13	p=0.00000
	No	100	33.39±15.77	
Pain durning the day	Yes	156	43.15±16.40	p=0.00000
	No	42	29.56±14.45	

*BMI - Body Mass Index (underweight: under 18.5 kg/m², normal weight: 18.5kg/m² to 25kg/m², overweight: 25kg/m² to 30kg/m², obese: over 30kg/m²).

** The quality of live is assessed on a scale of 0 to 100, which 0 indicating the highest quality of live and 100 indicating the lovest quality of live.