

Synthesis of Nickel-Chitosan Nanoparticles for Controlling Blast Diseases in Asian Rice

Ramalingam Parthasarathy

Indian Institute of Science

Chelliah Jayabaskaran

Indian Institute of Science

Appu Manikandan

Zhejiang Gongshang University

Sathiyarayanan Anusuya (✉ anusathsar.rajesh@gmail.com)

St Joseph's College Tiruchirappalli

Short Report

Keywords: Ni-Ch nanoparticles, Protein profile, Seed germination, Oryza sativa, Pyricularia oryzae, Rice blast

Posted Date: January 31st, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-564322/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Rice blast disease is a major challenge for rice production and consumption worldwide. To explore a control measure of rice blast alternative to fungicides, nickel-chitosan nanoparticle (Ni-Ch NPs) was prepared using nickel chloride and investigated for growth improvement and inhibition of *Pyricularia oryzae* (blast pathogen). SEM, TEM and DLS analyses exhibited smooth and spherical shaped nanoparticles with size range from 20–70 nm. Zeta potential revealed the colloidal stability of nanoparticles with PDI of 0.22. EDX spectroscopy corroborated the presence of nickel (14.05%) in Ni-Ch NPs. Application of Ni-Ch NPs to paddy seeds showed a significant increase in germination and growth attributes in terms of shoot and root length and number of lateral roots over control. Further, the nanoparticle treatment to plants under glasshouse condition demonstrated a remarkable improvement in plant growth condition showing no toxic effect. Protein profiling of treated plants revealed new polypeptides (rubisco units) which enlightened the enhanced photosynthetic rate. In addition, Asian rice exhibited reduced blast symptoms on leaves treated with nanoparticle over control under glasshouse condition while displaying 64 % mycelia inhibition in Petri plates. All these results suggest that nickel-chitosan nanoparticles could be exploited as a plant growth promoter cohort in controlling rice blast disease.

1. Introduction

Rice blast disease as a result of *Pyricularia oryzae* is solicitude in tackling worldwide subsistence unpredictably accountable for roughly 30% food grain manufacturing drop globally- equivalent of nourishing 60 million populations. This forfeiture escalates the universal grain cost and diminishes customer well-being and food certainty [1]. Distinctive strategies had been developed to boom the productivity and conquer demanding situations of controlling rice blast diseases. Though rice blast may be curtailed through the use of fungicides and resistant cultivars, development of latest pathogenic races hostility to commercially obtainable antifungal agent is an enduring problem [2]. To overcome this issue, nanomaterials are predominantly used in crop protection as an alternate to fungicides owing to their homogenous size, larger surface volume ratio, quirky optical properties and ecologically secure [3]. Over the last few years, metal nanoparticles including Ag, Au, ZnO, nano-sized TiO₂, Al₂O₃, Fe₃O₄, CeO₂ and Cu NPs are reported to exert positive effects on plant growth [4–7]. These metallic nanoparticles had been shown to penetrate the biological tissue inducing physiological and biochemical modifications in plants [8, 9]. Conversely, different studies provided conflicting evidence about the positive nexuses of nanomaterials on grain sprouting and vegetative maturation in limiting environmental conditions [10–14].

Nickel (Ni), an essential plant micronutrient was proclaimed to regulate seed germination in higher plants including cereal grains and legumes which is less toxic [15]. It activates enzymes such as urease and glycosidase-I required for nitrogen metabolism and assimilation [16]. It has been shown that nickel gets transported easily into plant system inducing physiological changes in plants [17, 18]. Gajewska and Sklodowska [19] described the use of nickel salt as fertilizer for promoting plant growth. Nowadays biopolymers such as chitosan, cellulose, alginate and pectin has fascinated researchers for synthesizing metal nanoparticles which acts as a good support preventing agglomeration and proven to be a better stabilizing agent as they allow better particle size control [20, 21].

Chitosan, a non-toxic, biocompatible and biodegradable natural polymer has the capability to chelate numerous metal ions such as Cu²⁺ and Zn²⁺ which can make an ideal composite based on the concept of nanotechnology. As well, chitosan have been shown to boost the yield and protein content emphatically in numerous plants likely maize, mulberry grains, broad bean and corn [22]. In particular, metal based chitosan nanoparticles (Cu-chitosan, ZnO-chitosan, Ag-chitosan) demonstrated an outstanding antimicrobial activity against phytopathogens accompanying plant growth promotion [23–26]. However, only few investigations have been concentrated on the antifungal activity of nickel nanoparticles [27]. In this context, we hereby describe primarily, the synthesis of nickel-chitosan nanoparticles and assessed the effect of nanoparticles on growth improvement in Asian rice and control of *P. oryzae* causing rice blast disease.

2. Materials And Methods

2.1. Synthesis of Nickel-Chitosan nanoparticles

In brief, 100 ml of 0.1% (w/v) chitosan solution was processed employing (0.1%, w/v) acetic acid. The mixture was precipitated with 10N NaOH and centrifuged at 10,000g for 30 min. The supernatant was discarded and immensely washed with distilled water to eliminate excess sodium hydroxide.

For the preparation of nanoparticles, 0.01% (w/v) of 25 ml nickel chloride solution was added drop wise to the prepared chitosan solution and stirred continuously for 12h at 70° C using a magnetic stirrer. To the resultant complex, 0.5 M NaOH was added drop wise to remove excess Cl₂ ions from the nanoparticles. The composite was finally centrifuged at 10,000 g for 10 min, concentrated under vacuum and stored at 4° C.

2.1. Physico-Chemical Characterization of Ni-Ch NPs

The optical measurement of Ni-Ch NPs was carried out by UV-visible spectrophotometer (Varian Cary 100) ranging from 200–500 nm to authenticate the formation of nanoparticles. FTIR spectroscopic studies were documented in the span of 4000 – 400 cm⁻¹ on Spectrum RX I to authenticate the functional groups and bonding of chitosan to nickel nanoparticles. The mean particle size, distribution and polydispersity index (PDI) of Ni-Ch NPs was carried out by Dynamic Light Scattering (DLS). The stability of the nanoparticle in suspension was determined by measuring the zeta potential of nanoparticles (Zeta sizer Nano ZS, Malvern). SEM analysis was being done to identify the dimension, shape and surface arrangement of Ni-Ch NPs. EDX spectroscopy analysis was performed to determine the presence of elemental nickel in the nanoparticle. The size and morphology of nickel-chitosan nanoparticle was characterized using TEM (JEOL JEM-2100 F) by drop coating of the NPs onto a carbon-coated copper grid.

2.3. *In vitro* seedling bioassay

The seeds of Asian rice (ADT-43) were obtained from Tamil Nadu Rice Research Institute, Aduthurai, TamilNadu, India. Briefly, 50 paddy seeds were disinfected with 0.001% (w/v) sodium hypochlorite for 10 minutes and washed several times with distilled water. The disinfected seeds were placed in Petri plates holding filter paper with 5ml of 0.1% (w/v) Ni-Ch NPs (treated) and 5ml of distilled water serves as control. Seed germination was recorded and the Petri dishes were kept in a growth chamber at $28 \pm 2^\circ \text{C}$ for 10 days. Each experiment was repeated three times. The shoot length, root length and number of lateral roots per plant were measured after 10 days of germination. Seedling vigour index was calculated using the formula as described by Abdul-Baki & Anderson [28]: Seed vigor index = (germination %) x (seedling length)

2.4. Inhibitory effect of Ni-Ch NPs against *Pyricularia oryzae*

The fungus *Pyricularia oryzae* was obtained from Tamil Nadu Rice Research Institute, Aduthurai, India. *In vitro* antifungal assay against *P. oryzae* was evaluated by the inhibition of mycelia growth (%). The *P. oryzae* mycelia bit (5.0 mm) was placed at the centre of Potato Dextrose broth (PDA) plates containing 0.1% (w/v) Ni-Ch NPs as treated. The PDA plate without nanoparticles served as control. The plates were maintained at 28°C for 7 days and monitored for mycelial growth. The percent inhibition of mycelial growth was measured. Each experiment was replicated thrice.

2.5. Effect of Ni-Ch NPs on vegetative growth and rice blast control under glasshouse environment

Ni-Ch NP treated paddy seeds were placed in containers (27-cm width; 26-cm vertex) holding soil + manure, inoculated with spores of *Pyricularia oryzae* (1×10^5 spores ml^{-1}) under glasshouse condition. Plants grown in soil with spores of *P. oryzae* (1×10^5 spores ml^{-1}) alone served as control. Growth attributes with regard to shoot length, root length and number of leaves was monitored regularly in treated and control plants. For all experiment, 50 plants were used and repeated thrice. Blast symptom including lesions was observed on leaves of control and treated plants. Additionally, disease severity (DS) and percent efficacy of disease control (PEDC) were assessed as mentioned by Chester [29] and Wheeler [30].

DS = Sum of all individual disease rating \times 100/total number of plant assessed \times maximum rating.

PEDC = Disease severity in control - disease severity in treatment/ disease severity in control \times 100

2.6. Effect of Ni-Ch NPs on Protein profiling

The control and treated (Ni-Ch NP + pathogen) leaves of paddy were collected on 25th day after treatment. Leaves (1g / 2ml) were homogenized with sodium acetate buffer (0.02 M, pH 5.8) and centrifuged at 10,000 G for 10 min at 4°C . From the supernatant obtained, protein content was evaluated according to Bradford method [31].

The samples (20 μg) were denatured for 3 min in a boiling water bath with sodium dodecyl sulphate (SDS) containing sample buffer and separated ahead with marker proteins following the procedure of Laemmli [32]. Localization of polypeptides on PAGE was completed by staining with Coomassie Brilliant Blue.

2.7. Statistical analysis

All the statistics were subjected to two-manner analysis of variance. The significance of variations among treatment approach was as compared by means of Duncan's multiple range tests at $P < 0.05$.

3. Results And Discussion

Metal and chitosan based nanoparticles promoting plant growth and protection have been known for years finding its application in modern agriculture [33–36]. Naturally occurring polymers for managing rice blast disease are secure to environment and acts as an effective alternative to chemical fungicides. The utility of nano-composite which include micronutrients complements the consumption and utility of minerals in cereal crops [37]. As well, chitosan acts as a carrier to obtain new polymeric material in order to prevent the undesirable nutrient loss to soil and can be modified easily without affecting its innate capabilities [38, 39]. The present study herein strongly demonstrated the control of rice blast by promoting the growth of paddy plants.

3.1. Characterization of nickel-chitosan NPs

The successful formation of nanoparticles was confirmed by a peak at 240 nm using UV-visible spectrophotometer (Fig. 1). FTIR analysis was done to track the interaction between nickel and chitosan nanoparticles. As depicted in the Fig. 2, FTIR spectra exhibited the characteristic peaks at 3431cm^{-1} (asymmetrical stretching frequency due to overlap of OH and $-\text{NH}_2$); 2924cm^{-1} and 2865cm^{-1} ($-\text{C}-\text{H}$ stretching vibration); 1633cm^{-1} ($-\text{C}=\text{O}$ amide group); 1472cm^{-1} , 1412cm^{-1} ($-\text{CH}_2-\text{N}$ stretching with combined with 1378cm^{-1} ($-\text{N}-\text{H}$ stretching) and 1186cm^{-1} (skeleton stretching frequencies of skeleton C-O and $-\text{C}-\text{O}-\text{C}$); The band at 889cm^{-1} stretching frequency shows $-\text{C}-\text{O}-\text{C}$ bridge as well as glycosidic linkage. The peak at 1420cm^{-1} and 1072cm^{-1} indicates the $-\text{CH}_2$ bending and $-\text{C}-\text{O}-\text{C}$ stretching frequency which confirmed the glycosidic linkage formed between nickel and chitosan nanoparticle [40].

DLS analyses revealed the average size and narrow distribution of nanoparticle as 89.81 nm with a polydispersity index of 0.22 (Fig. 3a). As shown in Fig. 3b, the zeta potential of nickel-chitosan nanoparticle was found to possess negative surface charge (-17.8 mv) which implicated the colloidal stability of nanoparticles. Chitosan, a positively charged amino polysaccharide in acidic condition can form strong chemical and physical bonds with metal ions insinuating the stability of nanoparticles [9, 23].

SEM study revealed the well dispersed and spherical shape of nanoparticles ranging from 30–50 nm (Fig. 4a). The EDX analysis inferred the elemental composition (14.05%) of the nanocomposite by exhibiting very strong signals at nickel-rich complex for nickel ions along with other weak signals displaying carbon and oxygen. The weak signals were attributed to the chitosan polymer surrounding nickel nanoparticle complex (Fig. 4b). The TEM micrograph

disclosed the spherical shape and smooth surface of the discrete nanoparticles without any aggregation (Fig. 5). The mean size of the nickel-chitosan nanoparticle as observed by TEM was 30–40 nm. It was obvious that chitosan as a dispersing agent interconnect superficially to nickel-chitosan nanoparticles thereby modifying the morphology, structure and magnetic properties [41]. This is in accordance with Hajipour and Abolfathi [42] who prepared uniform size, spherical shaped nickel nanoparticles supported on to the chitosan surface. The dissimilarity in dimensions among TEM and DLS study was because of the techniques intricated in preparation of samples [43].

3.2. Effect of Ni-Ch NPs on seed germination and growth of paddy

To check out the plant growth stimulating activity of Ni-Ch nanoparticles in paddy, the development criterion for instance shoot length, root length, number of sideward roots and seedling vigour index was calculated. Germination studies of Ni-Ch NP on paddy seeds revealed 100% germination while control showed only 82% germination. Plant growth promoting activity was found to be higher in Ni-Ch NP treatment (Fig. 6a). As depicted in Table 1, a significant increase in shoot (54%) and root length (64%) was noticed in NP applied seedlings as compared to control. Seed treatment of nanoparticles was reported to enhance the seed vigor and seedling development in many crops [44]. The Ni-Ch NP gets imbibed into the seeds and transported along with the photosynthates. Tan et al. [45] reported higher root number, root length and plant fresh weight of the *Oryza sativa* treated with uniconazole encapsulated silica nanoparticles. Besides increased shoot length, the growth rate was higher in root tissue following exposure to NP exhibiting enhancement in lateral root formation compared to control. Root elongation is a more sensitive indicator to study the phytotoxicity of engineered nanoparticles as reported by Tamsah and Joner [46]. At the root plane, rhizodermis lateral root junctions may anticipate easy access to NMs, largely near the root tip, although upper parts are impermeable since impregnation of suberin on outer layer [47]. Further, the negatively charged nanoparticles are translocated directly into root central cylinder and promote their migration ascending towards the aerial part of plant [48, 49].

Table 1
Effect of Ni-Ch NPs on germination, shoot length, root length, number of lateral roots and seedling vigor index of paddy. ^{a,b} and ^c show significance at $P \leq 0.05$.

Treatment	Germination (%)	Mean Root length (cm)	Mean Shoot length (cm)	Mean Seedling length (cm)	Seedling vigor index	Number of lateral roots
Control	82 ± 0.4 ^a	8 ± 0.5 ^b	5.5 ± 0.2 ^a	13.5 ± 0.3 ^a	1305	8 ± 0.5 ^a
Ni-Ch NPs	100 ± 0.1 ^c	12.5 ± 0.9 ^c	10.1 ± 0.05 ^c	22.5 ± 0.2 ^b	2054	22 ± 0.5 ^c

It is noteworthy that NP treatment promoted the growth of paddy plants under pot condition showing no toxic effects (Fig. 6b). Some of the synthesized metal nanoparticles (Iron oxide, ZnO, TiO₂, and copper) were straight away administered to soil as nanofertilizers through irrigation or foliar spray in distinct plants, together with mung bean plant, cucumber and rape [50–53]. Despite this fascinating evidence, soil amendment of nanoparticles may additionally create a risk to soil microorganisms dampening the use of metallic nanoparticles in agriculture [54, 55]. But, metal chitosan nanocomposites are said to be much less toxic due to slow release phenomenon and confirmed the ever-lasting effect in plants [52, 56, 57]. Chitosan nanoparticles had been used for controlled launch of nitrogen, phosphorus and potassium in wheat by way of foliar uptake [4]. Because of excessive nitrogen content material and low C/N ratio, chitosan will be immediately used as a fertilizer to enhance crop growth. In a few different studies, chitosan nanoparticles loaded with three triazine herbicides have shown decreased environmental effect and less toxic effects on genotype of onion plants [58]. The nanoparticle treatment displaying positive impact may additionally be because of the dimension, form, strength, chemical framework, effective electrical charge of nanoparticles [59].

3.3. Effect of Ni-Ch NPs on *Pyricularia oryzae*

The obliteration of mycelial growth was ascertained in petri plates holding 0.1 % (w/v) Ni-Ch NP after 7 days at 28° C and the percentage of inhibition were measured. No inhibition was noticed in control plates while the treated plates displayed (64%) higher growth inhibition (Fig. 7). This observation is in agreement with Pandian et al. [60] demonstrating the antifungal activity of nickel nanoparticle against *candida* and *Aspergillus* sps. Antifungal activity of metal nanoparticles is dependent on concentration, exposure time, suspension preparation procedure and the class of fungal strains [60, 61]. The inhibition of mycelial growth might be due to the electrostatic interplay between undoubtedly charged nickel ions and negatively charged microbial cellular membranes. The inhibitory effect also results from the chitosan interaction with phospholipid components of fungal membrane inflicting leakage of cell contents which ultimately results in cell death [62, 63]. Furthermore, chitosan readily binds with minor nutrients fabricating crucial elements unaccessible for normal fungal growth [64]. Therefore the direct antifungal activity of NPs could have relevant implications in controlling blast pathogen.

To investigate the virtue of Ni-Ch NPs against blast pathogen, potting experiments were conducted by inoculating the spores of *P. oryzae* under glasshouse condition. After 25 days of pathogen infection, the disease was assessed for each leaf and disease severity was calculated. Initial symptoms such as oval shaped grey color lesions were observed in control. Later it becomes dark and the entire leaves get affected while a significant control of blast disease was observed in NP treated paddy (Fig. 6a). The disease severity was higher in control (63%) compared to treated plants (21%). Percentage Efficacy of Disease Control was 23.4% in control while the treated plants showed 72% higher resistance to pathogen (Table 2). Cu-chitosan NPs have been outlined successful against early blight and *Fusarium* wilt of tomato [65]. In plants, chitosan is recognized by specific receptors available on the cellular plasma membrane together with pathogen-associated molecular pattern (PAMP) receptor. After this recognition, chitosan can spark off PAMP- brought on immunity this is capable of set off defense responses against potential fungal, bacterial, and viral pathogens [66].

Table 2
Effect of Ni-Ch NPs on paddy plants after 25 days of pathogen infection under glasshouse condition

Treatment	Disease severity (%)	Percent Efficacy of Disease Control (%)
Control	64.1 ± 0.9 ^c	33.4 ± 0.5 ^a
Treated	21 ± 0.25 ^a	72 ± 0.78 ^a

3.4. Protein profile

Protein content was found to be higher in leaves of treated plants compared to control (Fig. 8a). Metal NPs such as zinc oxide NPs increased the protein content by modulating the photosynthetic efficiency in tomato plants [67]. Also, Mehmood and Murtaza [68] reported that exposure of pea seeds to silver NPs significantly increased the carbohydrate and protein content. Multiwalled carbon nanotubes extended the number of flowers and fruits in tomato by activating the proteins important for plant bloom and development [69].

Ribulose biphosphate carboxylase/oxygenase (Rubisco), a vital photosynthetic enzyme was quantified by electrophoresis [70]. SDS-PAGE separation of protein in leaves of paddy obtained on 25th day treatment revealed polypeptides ranging from 25 kDa to 110 kDa in control (pathogen alone) as well as treated (Ni-Ch NPs + pathogen) plants (Fig. 8b). Apart from that, three new polypeptides were observed with molecular weight 15, 18 and 28 kDa in leaves of treated paddy plants. As seen in Fig. 8b, a molecular mass of 52.2 kDa was predominantly expressed in treated seedlings compared to control. RuBP oxygenase-carboxylase (rubisco), a key enzyme in photosynthesis contains two large subunits (50–55 kDa) that are associated with two small subunits (12–18 kDa) comprising up to 50% of the total soluble protein and 26% of total nitrogen found in leaf material [71]. Perdomo et al. [72] highlighted the importance of Rubisco for enhancing the photosynthetic overall performance of wheat, rice and maize at distinctive climatic conditions. Also, Rubisco and nitrogen courting rice was stated to promote leaf photosynthesis and plant growth [73].

In monocots, the degradation of rubisco is constantly preponderant after complete growth of leaf blade leading to a prompt decrease in rubisco content. But in response to NPs, rubisco tend to remain even after 25 days following exposure to NPs and pathogen which was confirmed by electrophoresis. This sincerely explains the position of NPs in enhancing the photosynthetic activity by means of mentioned expression of rubisco eventually promoting plant growth.

Conclusion

From this study it is clear that nickel chitosan nanoparticles through seed and soil amendment can be used in modern-day agriculture for increasing productiveness and as a novel remedy for rice blast disease. Further studies on biochemical and molecular aspects underneath field situations are compulsory to validate the exact mechanisms.

Declarations

Acknowledgements

The authors are grateful to Dean and the authorities of RVS Agricultural College, Tanjore, Tamilnadu, India for providing facilities to carry out the work.

Ethical Approval

Not Applicable

Consent to Participate

Not Applicable

Consent to Publish

Not Applicable

Authors Contributions

Ramalingam Parthasarathy: Conceptualization, Methodology, Writing - original

draft. **Sathiyarayanan Anusuya**: Writing- review & editing - original draft. **Appu Manikandan**: review & editing. **Chelliah Jayabaskaran**: Supervision, Project administration.

Competing Interests

The authors declare that they have no known competing financial interests

Availability of data and materials

References

- [1] Nalley, L., Tsiboe, F., Durand-Morat, A., Shew, A., Thoma, G. (2016). Economic and Environmental impact of Rice blast pathogen (*Magnaporthe oryzae*) alleviation in the United States. *Plos One*. 11 (12):e0167295. doi:10.1371/journal.pone.0167295
- [2] Asibi, A.E., Chai, Q., Coulter, J.A. (2019). Rice blast: A disease with implications for global food security. *Agronomy*. <https://doi.org/10.3390/agronomy9080451>
- [3] Sanzari, I., Leone, A., Ambrosone, A. (2019). Nanotechnology in Plant Science: To Make a Long Story Short. *Front. Bioeng. Biotechnol*, <https://doi.org/10.3389/fbioe.2019.00120>
- [4] Abdel-Aziz, H. M. M., Hasaneen, M. N. A., and Omer, A. M. (2016). Nano chitosan-NPK fertilizer enhances the growth and productivity of wheat plants grown in sandy soil. *Spanish J. Agric. Res.* 14, 1–9. doi: 10.5424/sjar/2016141-8205
- [5] Sathiyabama, M., Manikandan, A. (2018). Application of Copper-Chitosan Nanoparticles Stimulate Growth and Induce Resistance in Finger Millet (*Eleusine coracana* Gaertn.) Plants against Blast Disease. *J. agric. Food chem.* 66 (8), 1784-1790.
- [6] Ali, S., Rizwan, M., Noureen, S., Anwar, S., Ali, B., Naveed, M., Abd Allah, E.F., Alqarawi, A.A., Ahmad, P. (2019). Combined use of biochar and zinc oxide nanoparticle foliar spray improved the plant growth and decreased the cadmium accumulation in rice (*Oryza sativa* L.) plant. *Environ. Sci. Pollut. Res.* 26, 11288–11299. doi: 10.1007/s11356-019-04554-y.
- [7] Rizwan, M., Ali, S., ur Rehman, M.Z., Malik, S., Adrees, M., Qayyum, M.F., Alamri, S.A., Alyemeni, M.N., Ahmad, P. (2019). Effect of foliar applications of silicon and titanium dioxide nanoparticles on growth, oxidative stress, and cadmium accumulation by rice (*Oryza sativa*) *Acta Physiol. Plant.* 41, 35. doi: 10.1007/s11738-019-2828-7
- [8] Adili, A., Crowe, S., Beaux, M.F., Cantrell, T., Shapiro, P.J., McIlroy, D.N., Gustin, K.E. (2008). Differential cytotoxicity exhibited by silica nanowires and nanoparticles. *Nanotoxicol.* 2, 1–8.
- [9] Anusuya, S., Nibiya Banu, K. (2016). Silver-chitosan nanoparticles induced biochemical variations of chickpea (*Cicerarietinum* L.). *Biocatal. Agric. Biotechnol.* 8, 39-44.
- [10] Pokhrel, L. R., and Dubey, B. (2013). Evaluation of developmental responses of two crop plants exposed to silver and zinc oxide nanoparticles. *Sci. Total Environ.* 452–453, 321–332. doi: 10.1016/j.scitotenv.2013.02.059
- [11] Disfani, M.N., Mikhak, A., Kassaee, M.Z., Maghari, A. (2017). Effects of nano Fe/SiO₂ fertilizers on germination and growth of barley and maize. *Arch. Agron. Soil Sci.* 63, 817–826. doi: 10.1080/03650340.2016.1239016.
- [12] Shang, Y., Hasan, K., Ahammed, G.J., Li, M., Yin, H., Zhou, J. (2019). Applications of Nanotechnology in Plant Growth and Crop Protection: A Review. *Molecules.* 24(14), 2558.
- [13] Wang, Y., Lin, Y., Xu, Y., Yin, Y., Guo, H., Du, W. (2019). Divergence in response of lettuce (var. ramosa Hort.) to copper oxide nanoparticles/microparticles as potential agricultural fertilizer. *Environ. Pollut. Bioavailab.* 31, 80–84. doi: 10.1080/26395940.2019.1578187.
- [14] Du, W., Yang, J., Peng, Q., Liang, X., Mao, H. (2019). Comparison study of zinc nanoparticles and zinc sulphate on wheat growth: From toxicity and zinc biofortification. *Chemosphere.* 227, 109–116. doi: 10.1016/j.chemosphere.2019.03.168.
- [15] Fabiano, C.C., Tezotto, T., Favarin, J.L., Polacco, J.C., Mazzafera, P. (2015). Essentiality of nickel in plants: a role in plant stresses. *Front Plant Sci.* 6, 754.
- [16] Shahzad, B., Tanveera, M., Rehman, A., Cheema, S.A., Fahad, S., Rehman, S., Sharma A. (2018). Nickel; whether toxic or essential for plants and environment - A review. *Plant Physiol. Biochem.* 132, 641- 651.
- [17] Zhao, J., Shi, G., Yuan, Q. (2008). Polyamines content and physiological and biochemical responses to ladder concentration of nickel stress in *Hydrocharis*, Dubai (Bl.) Backer leaves. *Bio-Metals.* 21, 665–674.
- [18] Magaye, R., Zhao, J. (2012). Recent progress in studies of metallic nickel and nickel-based nanoparticles' genotoxicity and carcinogenicity. *Environ. Toxicol. Pharmacol.* 34, 644–650.
- [19] Gajewska, E., Skłodowska, M. (2009). Nickel-induced changes in nitrogen metabolism in wheat shoots. *J. Plant Physiol.* 166, 1034–1044.
- [20] Wei, D., Qian, W. (2008). Facile synthesis of Ag and Au nanoparticles utilizing chitosan as a mediator agent. *Colloids Surf B Biointerfaces*, 62(1), 136-142.
- [21] Ozhava, D., Kılıçaslan, N.Z., Ozkar, S. (2015). PVP-stabilized nickel(0) nanoparticles as catalyst in hydrogen generation from the methanolysis of hydrazine borane or ammonia borane. *Applied Catalysis B: Environmental.* 162, 573.

- [22] Behboudi, F., Tahmasebi, Z., Servestani, T., Kassaei, M.Z., Sanavi, S.A.M.M., Sorooshzadeh, A., Ahmadi, S.B. (2018). Evaluation of Chitosan Nanoparticles Effects on Yield and Yield Components of Barley (*Hordeum vulgare* L.) under Late Season Drought Stress. *J. Water Environ. Nanotechnol.* 3, 22-39.
- [23] Manikandan, A., Sathiyabama, M. (2015). Green synthesis of copper-chitosan nanoparticles and study of its antibacterial activity. *J. Nanomed. Nanotechnol.* 6 (1), 1.
- [24] Pereira, A.E.S., Silva, P.M., Oliveira, J.L., Oliveira, H.C., Fraceto, L.F. (2017). Chitosan nanoparticles as carrier systems for the plant growth hormone gibberellic acid. *Colloids Surf B Biointerfaces.* 150, 141-152.
- [25] Kaur, P., Duhan, J.S., Thakur, R., Chaudhury, A. (2018). Comparative pot studies of chitosan and chitosan-metal nanocomposites as nano-agrochemicals against fusarium wilt of chickpea (*Cicer arietinum* L.). *Biocatal. Agric. Biotechnol.* 14, 466-471.
- [26] Le, V.T., Bach, L.G., Pham, T.T., Le, N.T.T., Ngoc, U.T.P., Tran, D.H.N., Nguyen, D.H. (2019). Synthesis and antifungal activity of chitosan-silver nanocomposite synergize fungicide against *Phytophthora capsici*. *J. Macromol. Sci. Part A.* 56, 522–528. doi: 10.1080/10601325.2019.1586439.
- [27] Sharma, P., Sharma, A., Sharma, M., Bhalla, N., Estrela, P., Jain, A., Thakur, P., Thakur. (2017). Nanomaterial Fungicides: *In Vitro* and *In Vivo* Antimycotic Activity of Cobalt and Nickel Nanoferrites on Phytopathogenic Fungi. *Global Challenges.* 1 (9), 1700041.
- [28] Abdul-Baki, A.A., Anderson, J.D. (1973). Vigor determination in soybean by multiple criteria. *Crop Sci.* 13: 630–633.
- [29] Chester, K.S. (1950). Plant disease losses: their appraisal and interpretation. *Plant Disease Reporter Supplement.* 193, 189-362.
- [30] Wheeler, B.E.J. (1969). An introduction to plant diseases. 2nd Edition, John Wiley and Sons, London, 301.
- [31] Bradford, M.M. (1976). 'A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding'. *Anal Biochem.* 72, 248–254.
- [32] Laemmli, U. K. (1970). Cleavage of Structural Proteins during the Assembly of the Head of Bacteriophage T4. *Nature.* 227, 680-685.
- [33] Ghormade, V., Deshpande, M.V., Paknikar, K.M. (2010). Perspectives for nanobiotechnology enabled protection and nutrition of plants. *Biotechnol. Adv.* 29, 792-803.
- [34] Khot, L.R., Sankaran, S., Mari Maja, J., Ehsani, R., Schuster, E.W. (2012). Applications of nanomaterials in agricultural production and crop protection: A review. *Crop Prot.* 35, 64-70.
- [35] Yang, L., Cao, W., Rui, Y. (2017). Interactions between nanoparticles and plants: phytotoxicity and defense mechanisms. *Journal of plant interactions.* 12, 158-169.
- [36] Venkatachalam P., N. Priyanka, K. Manikandan, I. Ganeshbabu, P. Indiraarulsevi, N. Geetha, K. Muralikrishna, R.C. Bhattacharya, M. Tiwari, N. Sharma, S.V. S (2017). Enhanced plant growth promoting role of phycomolecules coated zinc oxide nanoparticles with P supplementation in cotton (*Gossypium hirsutum* L.). *Plant Physiol. Biochem.*, 110, 118-127
- [37] Jinghua, G. (2004). Synchrotron radiation, soft X-ray spectroscopy and nanomaterials. *J. Nanotechnol.* 1, 193-225.
- [38] De Rosa, M.R., Monreal, C., Schnitzer, M., Walsh, R., Sultan, Y. (2010). Nanotechnology in fertilizers. *Nat. Nanotechnol J.* 5, 91.
- [39] Divya, K., Jisha, M.S. (2018). Chitosan nanoparticles preparation and applications. *Environ. Chem. Lett.* 16, 101–112.
- [40] Kasaai, M.R. (2008). A review of several reported procedures to determine the degree of N-acetylation for chitin and chitosan using infrared spectroscopy. *Carbohydr. Polym.* 71, 497–508.
- [41] Liu, D., Ren, S., Wu, H., Zhang, Q., Wen, L. (2008). Morphology control in synthesis of nickel nanoparticles in the presence of polyvinylpyrrolidone (PVPK30). *J. Mat. Sci.* 43(6), 1974–1978.
- [42] Hajipour, A.R., Abolfathi, P. (2017). Novel triazole-modified chitosan@nickel nanoparticles: efficient and recoverable catalysts for Suzuki reaction. ***New J. Chem.*** 41, 2386-2391.
- [43] Khlebtsov, B.N., Khlebtsov, N.G. (2011). On the measurement of gold nanoparticle sizes by the dynamic light scattering method. *Colloid J.* 73, 118–127.
- [44] Arruda, S.C.C., Silva, A.L.D., Galazzi, R.M., Azevedo, R.A., Arruda, M.A.Z. (2015). Nanoparticles applied to plant science: A Review. *Talanta*, 131, 693–705.
- [45] Tan, W.M., Hou, N., Pang, S., Zhu, X.F., Li, Z.H., Wen, L.X., Duan, L.S. (2012). Improved biological effects of uniconazole using porous hollow silica nanoparticles as carriers. *Pest Manag Sci.* 68(3), 437-43.
- [46] El-Temsah, Y.S., Joner, E.J. (2012). Impact of Fe and Ag nanoparticles on seed germination and differences in bioavailability during exposure in aqueous suspension and soil. *Environ. Toxicol.* 27, 42–49.

- [47] Chichiricco, G., Poma, A. (2015). Penetration and Toxicity of Nanomaterials in Higher Plants. *Nanomaterials* (Basel). 5(2), 851–873.
- [48] Avellan, A., Schwab, F., Masion, A., Chaurand, P., Borschneck, D., Vidal V., et al. (2017). Nanoparticle uptake in plants: gold nanomaterial localized in roots of *Arabidopsis thaliana* by X-ray computed nanotomography and hyperspectral imaging. *Environ. Sci. Technol.* 51, 8682–8691.
- [49] Zhao, X., Meng, Z., Wang, Y., Chen, W., Sun, C., Cui B., et al. (2017). Pollen magnetofection for genetic modification with magnetic nanoparticles as gene carriers. *Nat. Plants.* 3, 956–964.
- [50] Gao, F., Hong, F., Liu, C., Zheng, L., Su, M., Wu, X., et al. (2006). Mechanism of nano-anatase TiO₂ on promoting photosynthetic carbon reaction of spinach: inducing complex of Rubisco-Rubisco activase. *Biol. Trace Elem. Res.* 111, 239–253. doi: 10.1385/BTER:111:1:239
- [51] Tarafdar, J.C., Raliya, R., Mahawar, H., Rathore, I. (2014). Development of zinc nanofertilizer to enhance crop production in pearl millet (*Pennisetum americanum*). *Agric. Res.* 3, 257–262. doi: 10.1007/s40003-014-0113-y
- [52] Saharan, V., Kumaraswamy, R.V., Choudhary, R.C., Kumari, S., Pal, A., Raliya, R., et al. (2016). Cu-chitosan nanoparticle mediated sustainable approach to enhance seedling growth in maize by mobilizing reserved food. *J. Agric. Food Chem.* 64, 6148–6155. doi: 10.1021/acs.jafc.6b02239
- [53] Verma, S. K., Das, A. K., Patel, M. K., Shah, A., Kumar, V., and Gantait, S. (2018). Engineered nanomaterials for plant growth and development: a perspective analysis. *Sci. Total Environ.* 630, 1413–1435. doi: 10.1016/j.scitotenv.2018.02.313
- [54] Simonin, M., Richaume, A., Guyonnet, J. P., Dubost, A., Martins, J. M. F., and Pommier, T. (2016). Titanium dioxide nanoparticles strongly impact soil microbial function by affecting archaeal nitrifiers. *Sci. Rep.* 6, 33643. doi: 10.1038/srep33643
- [55] Goncalves, M. F. M., Gomes, S. I. L., Scott-Fordsmand, J. J., and Amorim, M. J. B. (2017). Shorter lifetime of a soil invertebrate species when exposed to copper oxide nanoparticles in a full lifespan exposure test. *Sci. Rep.* 7, 1355. doi: 10.1038/s41598-017-01507-8
- [56] Wang, X.H., Du, Y.M., Liu, H. (2004). Preparation, characterization and antimicrobial activity of chitosan-Zn complex. *Carbohydr. Polym.* 56, 21–26.
- [57] Choudhary, R.C., Kumaraswamy, R.V., Kumari, S., Pal, A., Raliya, R., Biswas, P., Saharan, V. (2017). Synthesis, Characterization, and Application of Chitosan Nanomaterials Loaded with Zinc and Copper for Plant Growth and Protection. In book: Nanotechnology: 227-247.
- [58] Grillo, R., Clemente, Z., Oliveira, J. L., de Campos, E. V. R., Chalupe, V. C., Jonsson, C. M., et al. (2015). Chitosan nanoparticles loaded the herbicide paraquat: the influence of the aquatic humic substances on the colloidal stability and toxicity. *J. Hazard. Mater.* 286, 562–572. doi: 10.1016/j.jhazmat.2014.12.021
- [59] Rastogi, A., Zivcak, M., Sytar, O., Kalaji, H.M., He, X., Mbarki, S., Brestic, M. (2017). Impact of Metal and Metal Oxide Nanoparticles on Plant: A Critical Review. *Front Chem.* 5, 78.
- [60] Pandian, C.J., Palanivel, R., Dhanasekaran, S. (2016). Screening Antimicrobial Activity of Nickel Nanoparticles Synthesized Using *Ocimum sanctum* Leaf Extract. *Journal of Nanoparticles.* <http://dx.doi.org/10.1155/2016/4694367>
- [61] Gunalan, S., Sivaraj, R, Rajendran, V. (2012). Green synthesized ZnO nanoparticles against bacterial and fungal pathogens, *Progress in Natural Science: Materials International.* 22 (6), 693–700.
- [62] Rincon, J.G., Perez, J.V., Guerra-Sanchez, M.G., Hernandez-Lauzardo, A.N., Pena-Diaz, A., Velazquez-DelValle, M.G. (2010). Effect of chitosan on growth and plasma membrane properties of *Rhizopus stolonifer* (Ehrenb.:Fr.) Vuill," *Pesticide Biochem. Physiol.* 97, 275–278.
- [63] Liu, H., Du, Y., Wang, X., Sun, L. (2004). Chitosan kills bacteria through cellmembrane damage," *Int. J. Food Microbiol.* 95, 147–155.
- [64] Roller, S., Covill, N. (1999). The antifungal properties of chitosan in laboratory media and apple juice. *Int. J. Food Microbiol.* 47, 67–77.
- [65] Saharan, V., Sharma, G., Yadav, M., Choudhary, M.K., Sharma, S.S., Pal, A., Raliya, R., Biswas, P. (2015). Synthesis and *in vitro* antifungal efficacy of Cu-chitosan nanoparticles against pathogenic fungi of tomato. *Int. J. Biol. Macromol.* 75, 346–353.
- [66] Pusztahelyi, T. (2018). Chitin and chitin-related compounds in plant–fungal interactions. *Mycology*, 9, 189–201.
- [67] Faizan, M., Faraz, A., Yusuf, M., Khan, S.T., Hayat, S. (2017). Zinc oxide nanoparticle-mediated changes in photosynthetic efficiency and antioxidant system of tomato plants. *Photosynthetica.* 56, 678–686
- [68] Mehmood, A., Murtaza, G. (2017). Impact of biosynthesized silver nanoparticles on protein and carbohydrate contents in seeds of *Pisum sativum* L. *Crop Breed. Appl. Biotechnol.* 17 (4), <https://doi.org/10.1590/1984-70332017v17n4a51>
- [69] Khodakovskaya, M. V., Kim, B. S., Kim, J. N., Alimohammadi, M., Dervishi, E., Mustafa, T. et al. (2013). Carbon nanotubes as plant growth regulators: effects on tomato growth, reproductive system, and soil microbial community. *Small.* 9, 115–123. doi: 10.1002/smll.201201225
- [70] Aranjuelo, I., Pérez, P., Hernández, L., Irigoyen, J.J., Zita, G., Martínez-Carrasco, R., Sánchez-Díaz, M. (2005). **The response of nodulated alfalfa to water supply, temperature and elevated CO₂: Photosynthetic downregulation.** *Physiol. Plant*, 123, 348-358.

[71] Tabita, F.R., Hanson, T.E., Li, H., Satagopan, S., Singh, J., Chan, S. (2007). Function, structure, and evolution of the RubisCO-like proteins and their RubisCO homologs. *Microbiol. Mol. Biol. Rev.* 71, 576-599

[72] Perdomo, J.A., Bauçà, S.C., Silva, E.C., Galmés J. (2017). Rubisco and Rubisco Activase Play an Important Role in the Biochemical Limitations of Photosynthesis in Rice, Wheat, and Maize under High Temperature and Water Deficit. *Front Plant Sci.* 8, 490.

[73] Makino, A. (2003). Rubisco and nitrogen relationships in rice: Leaf photosynthesis and plant growth. *Soil sci. plant nutri.* 49, 319-327.

Figures




Figure 1

UV-Visible spectra of Ni-Ch NPs

Figure 2

FTIR spectra of Ni-Ch NPs

Figure 3

DLS (a) and Zeta potential distribution (b) of Ni-Ch NPs



Figure 4

SEM micrograph (a) and EDX analysis (b) of Ni-Ch NPs




Figure 5

TEM micrograph of Ni-Ch NPs

Figure 6

Effect of Ni-Ch NPs on growth of paddy under 'in vitro' (a) and glasshouse condition (b).

Figure 7

Inhibitory effect of Ni-Ch NPs on *Pyricularia oryzae* under 'in vitro' (a) and blast symptoms on control and treated leaves under glasshouse condition (b).

Figure 8

Protein content in control and Ni-Ch NP treated leaves of paddy plants (a); Localization of polypeptides in leaves of paddy (b) on SDS-SPAGE (10% separating gel; 4 % stacking gel; M: Standard marker proteins of different molecular weight (Bio Rad Chem. Co., USA); C: control; T: Ni-Ch NP treated).