

# A new candidate vaccine for human brucellosis based on influenza viral vectors: development of immunization schedule in guinea pig model

**Dina Bugybayeva**

Research Institute for Biological Safety Problems

**Zhailaubay Kydyrbayev**

Research Institute for Biological Safety Problems

**Nadezhda Zinina**

Research Institute for Biological Safety Problems

**Nurika Assanzhanova**

Research Institute for Biological Safety Problems

**Bolat Yespembetov**

Research Institute for Biological Safety Problems

**Yerken Kozhamkulov**

Research Institute for Biological Safety Problems

**Kunsulu Zakarya**

Research Institute for Biological Safety Problems

**Sholpan Ryskeldinova** (✉ [ryskeldinova1964@mail.ru](mailto:ryskeldinova1964@mail.ru))

Research institute for biosafety problems <https://orcid.org/0000-0002-6538-8486>

**Kaissar Tabynov**

Kazakh National Agrarian University

---

## Research Article

**Keywords:** Human brucellosis, influenza viral vectors, vaccine candidate, protection, guinea pigs, pre-existing immunity

**Posted Date:** November 14th, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-56762/v3>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published on February 16th, 2021. See the published version at <https://doi.org/10.1186/s40249-021-00801-y>.

# Abstract

**Background:** A new candidate vector vaccine against human brucellosis based on recombinant influenza viral vectors (rIVV) subtypes H5N1 expressing *Brucella* Omp16, L7/L12, Omp19 or Cu-Zn SOD proteins has been developed. This paper presents the results of the study of protection of the vaccine with various options of administering, dosing and frequency of use on guinea pigs.

**Methods:** General states of guinea pigs was assessed based on , behavior and dynamics of a guinea pig weight-gain test. The effectiveness of the new anti-brucellosis vector vaccine was determined by studying its protective effect after conjunctival, intranasal and sublingual administration in doses  $10^5$  EID<sub>50</sub>,  $10^6$  EID<sub>50</sub> and  $10^7$  EID<sub>50</sub> during prime and boost vaccinations of animals, followed by challenge with a virulent strain of *B. melitensis* 16M infection. For sake of comparison, the commercial *B. melitensis* Rev.1 vaccine was used as a control. The protective properties of vaccines were assessed by quantitation of *Brucella* colonization in organs and tissues of infected animals and compared to the control groups.

**Results:** It was observed a gradual increase in body weight of guinea pigs after prime and booster immunization with the vaccine using conjunctival, intranasal and sublingual routes of administration, as well as after using various doses of vaccine. . The most optimal way of using the vaccine has been established: double intranasal immunization of guinea pigs at a dose of  $10^6$  EID<sub>50</sub>, which provides 80% protection of guinea pigs from *B. melitensis* 16M infection ( $P < 0.05$ ), which is comparable to the results of the effectiveness of the commercial *B. melitensis* Rev.1 vaccine.

**Conclusions:** We developed effective vaccine candidate against brucellosis and developed its immunization protocol in guinea pig model. We believe that this study is a substantial step for using the vaccine for future pre-clinical and clinical trials in human.

## Background

Brucellosis is a chronic infectious disease of humans and animals, which is included in the list of quarantine diseases as a social threat factor. There are 10 known species of the causative agent of brucellosis which include those that pose a threat to human health - *B. melitensis*, *B. abortus*, *B. suis* and *B. canis* [1,2]. *B. melitensis* that causes the most severe and acute form of infection is responsible for 80–90% of human brucellosis cases [3]. Most cases of brucellosis in humans are the result of occupational exposure to the bacteria and consumption of infected dairy products [4].

Despite the fact that brucellosis is amenable to antibiotic therapy, it seriously weakens the human body and many patients require long-term recovery. In addition, due to the intracellular tropism of *Brucella*, only a very limited number of antibiotics can be used to treat this infection [5]. It is important to note that in 5–40% of cases the antibiotic therapy results in relapses of the disease, which requires protracted treatment using different combinations of antibiotics [6]. This fact, as well as the lack of a licensed anti-brucellosis vaccine for humans represent a serious problem for the brucellosis endemic areas.

Earlier, USSR and China were widely using live attenuated vaccines based on *B. abortus* 19 BA or *B. melitensis* 104M strains for human vaccination [7]. Normally, in vaccinated people these vaccines ensured a short-term immune response and were also accompanied by high reactogenicity and hypersensitivity, especially when repeated doses of the vaccine were administered.

One of the most important strategies in the development of safe and effective anti-brucellosis vaccines is the use of live genetically modified vectors - non-pathogenic microorganisms (bacteria and viruses) that produce brucellosis antigens. Nowadays, *Lactococcus Lactis* [8], *Escherichia coli* [9], *Salmonella enterica* [10] and *Semliki Forestvirus* (SFV) [11] are used as vectors for the expression of brucellosis proteins *in vivo*. It has been proved experimentally that the tested bacterial (intracellular) and viral vectors are capable of infecting a wide range of cell types and expressing brucellosis antigens within the infected cell.

Previously, we used recombinant Influenza A viral vectors of both subtypes a H5N1 for prime vaccination and H1N1 for booster vaccination that express brucellosis immunodominant outer membrane proteins (Omp) 16 and ribosomal L7/L12 in order to develop a new *Brucella abortus* vaccine (Flu-BA) against brucellosis in cattle [12,13]. Now, the Flu-BA vaccine is registered in Kazakhstan (registration certificate #RK-VP-1-3775-19 dated from January 14, 2019) and is at the stage of commercialization for vaccination of cattle against *B. abortus* infection.

The effectiveness of cattle vaccination is comparable to the results of use of the commercial *B. abortus* S19 vaccine. The use of influenza viral vectors (IVV) subtype H5N1 may serve as an additional factor in increasing the effectiveness of the vaccine for humans. The fact is that there is no pre-existing immunity to influenza virus H5N1 in human population, and, therefore IVV of the H5N1 subtype has more opportunities for replication and expression of brucellosis proteins.

In this study, we used the entire existing stock of previously obtained recombinant IVV type A of subtype H5N1 that express *Brucella* outer membrane proteins (Omp) 16 and 19, ribosomal L7/L12 or Cu-Zn superoxide dismutase (SOD) from the open reading frame of the NS1 gene at amino acid position 80 [14].

*Brucella* antigens Omp 16 and 19, L7/L12 and SOD induce a pronounced cellular response which is necessary for protection against brucellosis infection [15, 16, 17]. It should be noted that the proteins expressed by IVV are immunodominant and common (genetically similar by 95–99%) for *B. melitensis*, *B. abortus*, *B. suis*, and *B. canis* [18, 19, 20]. In this regard, we have developed an anti-brucellosis tetravalent IVV based human vaccine candidate expressing the brucellosis proteins Omp16, L7/L12, Omp19 and SOD.

This article presents the results of studying clinical observation of general states of guinea pigs and protectiveness of the anti-brucellosis human vaccine with different modes of administration, with different doses and frequency of vaccination in guinea pigs.

# Methods

## Generation of viral constructs and preparation of vaccine samples

Influenza viruses were obtained by a standard reverse genetics method using eight bi-directional plasmids pHW2000. In this study, we used IVV type A of H5N1 subtype that express *Brucella* proteins Omp16, Omp19, L7/L12 or SOD from NS1 open reading frames at amino acid position 80. A detailed procedure for the construction of the IVV has been described previously [14]. Schematic picture for IVV is presented in Figure 1. Obtained recombinant influenza vectors (Flu-NS1-80-Omp16, Flu-NS1-80-L7/L12, Flu-NS1-80-Omp19, Flu-NS1-80-Cu-Zn SOD) were then used to produce vaccine for immunization guinea pigs. Briefly, the viral vector was accumulated in 10-day chick embryos (CE) at 34°C for 48 hrs. The IVV titer was determined by hemagglutination assay based on the generally accepted method as reported previously [21].

Allantoic suspensions of IVV were clarified, purified and concentrated by ultracentrifugation and then the resulting material was sent to diafiltration and sterilizing filtration and then combined with a sterile stabilizing medium containing 12% soy peptone (Sigma-Aldrich, USA) and 6% sucrose (Sigma-Aldrich, USA). The combined material was added to 1 ml ampoules, lyophilized and stored at 2-8 °C. The lyophilized vaccine was resuspended in phosphate buffered saline (PBS) prior to vaccination of the animals.

## Bacterial strains

The virulent *B. melitensis* 16M strain was obtained from the collection of microorganisms of the Research Institute for Biological Safety Problems (RIBSP). Bacterial cells were cultured under aerobic conditions on a Brucella Agar Base nutrient medium (Sigma-Aldrich, USA) at 37 °C. All experiments with live *Brucella* cells were conducted in Biosafety level 3 facility (BSL-3) at the RIBSP. Infected animals were contained in specialized animal BSL-3 facility.

## Bioethics and groups of experimental animals

The studies were conducted in accordance with national and international regulations and guidelines for the handling and use of laboratory animals. The study protocol was approved by the RIBSP Bioethics Committee (Protocol no. 0418/04). The animals contained in cages on 12 light/12 dark cycle and were fed *ad libitum* with standard rodent diet and had no water restrictions. Animals from control and experimental groups were kept isolated from each other throughout the whole experiment. This study used conventional bred female guinea pigs weighing 250-330 g (National Center for Expertise of Drugs, Medical Products and Equipment, Kazakhstan).

## Immunization of guinea pigs

To determine the method of applying the vaccine, 5 groups (n=5 animals per group) were formed - three experimental and two control groups. The experimental groups was administered with vaccine

conjunctivally (*c.*) in a volume of 100 µl (50 µl in each eye) or intranasally (*i.n.*) in a volume of 200 µl (100 µl in each nasal cavity) or sublingually (*s.l.*) in a volume of 200 µl. The infectious titer of the IVV in the experimental vaccine samples was 6.14-6.75 log<sub>10</sub> EID<sub>50</sub>/µl. On day 21 after the prime vaccination, guinea pigs were boosted according to the immunization protocol. The animals in negative control group were injected subcutaneously (*s.c.*) with 200 µl of PBS. In the positive control group, a dose of 10<sup>5</sup> CFU/animal commercial *B. melitensis* Rev.1 vaccine (Antigen LLP, Kazakhstan) was injected once *s.c.*, according to the manufacturer's recommendations.

Further on, to determine the protective dose and the frequency of immunization, the vaccine was administered *i.n.* Three doses of the vaccine - 10<sup>5</sup> EID<sub>50</sub>, 10<sup>6</sup> EID<sub>50</sub> and 10<sup>7</sup> EID<sub>50</sub> were evaluated in six experimental groups of guinea pigs (n=5 animals per group), including control groups, after prime-boost immunization according to the protocol presented above.

In order to assess weight changes associated with the vaccine, immunized guinea pigs were clinically monitored with weekly weighing for 42 days upon both prime and boost vaccinations. The clinical assessment was evaluated based on parameters including the survival rate, general condition, behavior and dynamics of animals' weight.

### **Assessment protectiveness of the vaccine**

On the day 21 after the boost immunization, the guinea pigs (45 animals in total) from the vaccinated and control groups were *s.c.* infected with a virulent strain of *B. melitensis* 16M infection at a dose of 1.3 log<sub>10</sub> CFU/animal. Guinea pigs from positive (n = 10) control groups were infected in a similar way on day 42 after a single vaccination with *B. melitensis* Rev.1. Thirty days after infection, all guinea pigs were euthanized with CO<sub>2</sub> and lymph nodes (retropharyngeal, lower cervical and right and left inguinal), liver, spleen and bone marrow were extracted aseptically. Bacteriological examination and evaluation of the results were carried out according to the previously described method [23]. Briefly, after 10-fold serial dilution the tissue homogenates in 0.1% Triton X-100 - PBS solution were inoculated into plates with *Brucella* Base Agar (HiMedia, India) and incubated at 37°C for 2 weeks with periodical counting of the bacterial colonies during this period.

The concentration of bacteria (colony forming units (CFU)/g per tissue) in tissue samples was determined by standard colony counting. An animal was considered infected if a *Brucella* colony was found in culture of one or more organs. The results of the bacteriological study were assessed according to several parameters: a) vaccination efficacy or the number of animals (expressed in %) from which no *Brucella* colonies were isolated in any animal samples; b) generalization of the infectious process or the index of infection (the number of organs and lymph nodes of animals from which *Brucella* was isolated and which is represented in the arithmetic mean); c) the intensity/severity of the infectious process or the degree of *Brucella* colonization of organs and lymph nodes (LN) expressed in log<sub>10</sub> CFU/g of tissue.

### **Statistical Analysis**

The index of infection and *Brucella* colonization in tissues between groups was analyzed with one-way or two-way ANOVA followed by multiple comparisons of the Turkey test, Sidak or Dunnett's tests. The variance in protective efficacy of animal groups was compared by one-sided Fisher exact test.  $P < 0.05$  was considered significant. Means are reported with standard errors (SEM). Statistical analysis of all experimental data was performed using Graph Pad Prism Software Version 8.0 (Graph Pad Software Inc., La Jolla, CA, USA). The experiments have been repeated, and the results were reproducible.

## Results

Assessing general states of the guinea pigs after immunization with the vaccine

The effect of the vaccine on guinea pigs was evaluated with different ways of administering, with the use of different doses of vaccine and the frequency of use in comparison with a negative (PBS) control group.

It was found that the vaccine was safe in guinea pigs in different ways of vaccine administering - conjunctival, intranasal and sublingual, as well as in various doses in primary and secondary immunization (prime and boost). There were no animal deaths or signs of disease by the end of the observation period. In general, the condition of the animals, both in the control and in the experimental groups, was satisfactory in terms of physical activity, appetite and general health condition.

Evaluation of changes in animal body weight within 21 days after prime-boost immunization showed that the weight of animals (on average) increased in all test groups (Figure 2) by 45 - 58% and amounted to 158-175 g which was comparable to the control group - 55 % and 162 g.

No group was differed significantly in mean of animal weight from the negative (PBS) control group.

### **Protectiveness of viral vector based vaccine candidate at different ways of its administration against *B. melitensis* 16M infection**

The protective efficacy of the vaccine candidate was evaluated in guinea pigs using conjunctival (c.), intranasal (i.n.) and sublingual (s.l.) routes of administration and compared with reference *B. melitensis* Rev.1 vaccine or PBS control groups.

The protective efficacy of the vaccine was assessed by parameters as the index of infection, the efficacy of vaccination and the number of bacteria of the virulent strain of *B. melitensis* 16M obtained from organs and tissues of vaccinated and unvaccinated animals.

The results of the bacteriological study showed that with various mucosal routes of immunization, the new anti-brucellosis vaccine candidate provided protection at a level of 1.64 to 2.30  $\log_{10}$  units. In comparison with the unvaccinated control group (PBS), all vaccine samples, regardless of the route of administration, ensured protection of guinea pigs from *B. melitensis* 16M infection ( $\alpha = 0.0001 - 0.02$ ) (Table 1.)

**Table 1** Degree of protective efficacy of the vaccines by different route of administration in guinea pigs

Vaccine	Route of administration	Log <sub>10</sub> CFU/animal (mean ± SE)	*Log <sub>10</sub> protection	†Value (P)
Influenza viral vector based brucellosis vaccine candidate	<i>c.</i>	0.56±0.22	2.3	<0.003
	<i>i.n.</i>	0.06±0.04	2.8	<0.0001
	<i>s.l.</i>	1.22±0.29	1.64	<0.02
Commercial vaccine <i>B. melitensis</i> Rev.1	<i>s.c.</i>	0.48±0.18	2.38	<0.0005
Control (PBS)	<i>i.n.</i>	2.86±0.19	0.00	-

**Table note:** Degree of protective efficacy of the vaccines as evaluated by the isolation rate of *Brucella* from organs and tissues of guinea pigs challenged with the virulent strain of *B. melitensis* 16M infection

1. – conjunctivally, *i.n.* – intranasally, *s.l.* – sublingually; \* Log<sub>10</sub> protection units were obtained by subtracting the mean log<sub>10</sub> CFU of the control (PBS) group from the mean of log<sub>10</sub> CFU for the experimental group. †Compared with control group (PBS). Statistical analysis was performed using two-way ANOVA followed by Tukey’s multiple comparisons test.

Significant protection of the vaccine was observed after *i.n.* administration (2.8 log<sub>10</sub>), whereas for *s.l.* immunization, the unit of protection was 1.64 log<sub>10</sub>. In case when the vaccine administered by *c.* route the unit of its protection was 2.3 log<sub>10</sub>, which was comparable to the commercial *B. melitensis* Rev.1 vaccine results.

According to the index of infection (Fig. 3, B), significant protection compared to the control group (PBS) was achieved in the groups vaccinated *c.* ( $P < 0.01$ ; vaccination efficiency 60%) and *i.n.* ( $P < 0.002$ , vaccination efficiency 80%), as well as in animals vaccinated with *B. melitensis* Rev.1 ( $P < 0.005$ ; vaccination efficiency 80%). It should be noted that the index of infection in *i.n.* vaccinated animals was similar ( $P > 0.99$ ) to the index in animals immunized with the commercial *B. melitensis* Rev.1 vaccine.

Based on the results of these studies, the *i.n.* route of vaccine administration was applied for further research in order to determine the optimal dose for immunization.

### **Study of protective efficacy of a novel vaccine candidate at different doses and after infection with *B. melitensis***

In this study, the protectiveness of the vaccine was assessed at various doses of *i.n.* vaccine administration after prime-boost immunization, as well as after challenge of guinea pigs with the virulent *B. melitensis* 16M strain.

The evaluation of the protective efficacy of the tetravalent vector brucellosis vaccine against the infection was carried out according to the following parameters - 1) determination of the effectiveness of vaccination (the degree of complete protection against infection, expressed in percent), 2) study of the infection index, as well as by counting *Brucella* colonies in the tissues and lymph nodes of guinea pigs 30 days after infection.

Bacteriological studies of the organs and lymph nodes of infected animals showed that the tested doses of the recombinant vector vaccine, as well as the commercial vaccine, provided significant protection ( $P < 0.04$  versus the unvaccinated group) of guinea pigs against *B. melitensis* 16M infection ranging from 1.64 to 2.8 log<sub>10</sub> units. Our results showed that the highest level of protection (vaccination efficiency) against infection in guinea pigs was in the groups immunized at doses of 10<sup>6</sup> EID<sub>50</sub> and 10<sup>7</sup> EID<sub>50</sub> (80%) compared with the control group (PBS) after the challenge, in which the infection rate was 100% and these data were statistically significant ( $P < 0.04$ ). It should be noted that in the group vaccinated with *B. melitensis* Rev.1 the vaccination efficiency was also 80% (Table 2).

According to the index of infection (Fig. 4, D), a significant level of protection in comparison with the control group (level of infection - 100%) was achieved in all three tested doses and only after secondary boost vaccination ( $P < 0.0004$ ,  $P < 0.02$ ). The index of infection in the groups of animals immunized with a dose of vaccine 10<sup>6</sup> EID<sub>50</sub> and *B. melitensis* Rev.1 vaccine was comparable with each other and these groups differed significantly compared to the negative control group  $P = 0.003 - 0.001$ .

**Table 2** Rates of protection in guinea pigs after challenge with the virulent strain *B. melitensis* 16M

Immunization group (prime-boost)	Total animals	Isolation of <i>B. melitensis</i> in animals, n (%)	*Value (P)	
			(+) control	(-) control
Vector vaccine at dose of 10 <sup>5</sup> EID <sub>50</sub>	5	2 (40)	>0.05	>0.05
Vector vaccine at dose of 10 <sup>6</sup> EID <sub>50</sub>	5	1 (20)	>0.05	<0.05
Vector vaccine at dose of 10 <sup>7</sup> EID <sub>50</sub>	5	1 (20)	>0.05	<0.05
<i>B. melitensis</i> Rev.1	5	1 (20)	-	<0.05
Control (PBS)	5	5 (100)	<0.05	-

**Table note:** \*In comparison with control untreated PBS or *B. melitensis* Rev.1 groups. Statistical analysis was performed using a one-sided Fisher's exact test. P-value less than 0.05 (< 0.05), P-value higher than 0.05 (> 0.05).

It should be noted that the values of the index of infection between the *B. melitensis* Rev.1 group and the experimental groups did not have significance ( $P>0.05$ ).

## Discussion

To our knowledge, this study is the first trial conducted in guinea pigs to evaluate the protective properties of a new candidate for vector vaccine against human brucellosis. This phase in vaccine trials is an important step in making an experimental vaccine a promising candidate for further human clinical trials to determine its effectiveness. In this study, double *i.n.* immunization with a vector vaccine based on influenza viral vectors expressing the immunodominant brucellosis proteins Omp16, L7/L12, Omp19 and Cu-Zn SOD at a dose of  $10^6$  EID<sub>50</sub> ensured protection against *B. melitensis* 16M infection comparable to the effect of commercial *B. melitensis* Rev.1 vaccine.

The choice of guinea pigs as model animals for evaluation of body gain changes and vaccine candidate protection was determined by their natural resistance to influenza infection in comparison with laboratory mice. In this case, the use of a more resistant model animal seemed to be a key condition in the study of protection, since, in the long run, the vaccine is designed for humans.

The previous success of using IVV in the development of the anti-brucellosis vaccine Flu-BA for cattle [12], which is now at the stage of commercialization in Kazakhstan, served as the basis for this study. The idea of developing an anti-brucellosis human vaccine is that the high efficacy of the vaccine is achieved in cattle which naturally resistant to influenza infection (i.e., as a non-replicable viral vector), and, in our opinion, should be even more pronounced in humans. This assumption is based on the fact that humans are a natural host for the influenza virus (Influenza A), including the influenza viral vectors we use.

It should be noted that we used an influenza viral vector (IVV) of the H5N1 subtype, because there is no immune background to this type of pathogen [22] in the human population and IVV of the H5N1 subtype has a greater potential as a vaccine vector.

We began the process of creating an effective anti-brucellosis vector vaccine for humans with the formation of requirements for the developed product, production technology and methods of its application and testing in healthcare practice. To this end, we have accumulated the existing experience in the development of vector vaccines for public health, have chosen the most generally accepted requirements for such vaccines and their manufacturing technologies, the method and frequency of their use and have developed criteria for assessing the effectiveness and safety of vaccine candidates.

An analysis of the compliance of the developed vaccine with the above requirements (which are more general in nature than specific) showed that the viral vectors we selected, as well as the method for preparing and using the vaccine, correspond to them. In particular, we use non-pathogenic influenza viral vectors, the general safety of which has been confirmed by studies on guinea pigs with various ways of administering and dose of immunization.

In order to obtain influenza viral vectors (IVV), we used A/Puerto Rico/8/34 (H1N1) with a length-modified NS1-80 gene encoding 80 amino acids in the N-terminal region of the protein as the initial strain. The surface genes of hemagglutinin (HA) and neuraminidase (NA) were taken from A/chicken/Astana/6/05 strains (H5N1, with the HA cleavage site preliminary removed). The safety or attenuation of IVV is ensured by the truncated NS1 protein (interferon antagonist), which results in their limited replicative capabilities (they make one cycle of reproduction in the cell and do not leave it) [14]. It is known that the degree of IVV attenuation is directly dependent on the length of the NS1 protein [23]. We have an IVV with NS1 length in 80 amino acid. NS1-124 was used to create a veterinary brucellosis vaccine as it for use in cattle a more aggressive IVV was required. As for humans, far preferable may be IVV with NS1-80. With IVV subtype H5N1 (a pathogenic variety of influenza virus), the attenuation was additionally achieved by removing the proteolytic cleavage site in the HA protein, that is, double attenuation was performed. During repeated re-inoculation in chick embryos, IVV retained all their basic biological properties, including signs of attenuation, and did not lose the brucellosis insertion segment [14], which indicates their genetic stability. In addition, the influenza viral vectors we use are RNA-containing viruses that are limited to cytoplasmic replication, thus eliminating the risk of integration and long-term persistence.

The next important phase of our research was devoted to the study of the general safety control of the vaccine candidate at the early stage with different ways of administration and dose of use in guinea pigs. The vaccine has been found to be safe for guinea pigs when administered *c.*, *i.n.* and *s.l.* The experimental animals did not show death or signs of any disease; by the end of observation (on day 14 after the prime-boost vaccinations), the body weight gain in guinea pigs was observed both after prime and after boost immunization. At the same time, the increase in body weight of guinea pigs in the experimental groups was comparable to the control group of animals that were injected with PBS. As a result of this work, the vaccine was recognized as a safe drug and was used in the future to assess its protectiveness depending on the immunization schedule.

Further assessment of the effectiveness of the vaccine with different routes of administration to mucosal areas was determined using *c.*, *i.n.* and *s.l.* methods of vaccine immunization in prime-boost mode. Since the influenza virus has a tropism for mucosal surfaces, it was assumed that the optimal way to administer a vaccine based on an influenza viral vector would be one of the tested mucosal routes. Since *Brucella* should be considered as a mucosal pathogen penetrating mucous surfaces, the “gates” of infection are the mucosal surfaces of the nose or mouth. Consequently, mucosal vaccination is capable to generate protective responses against pathogens at the site of the infection “gate” [24]. Our bacteriological study demonstrated that significant protection of guinea pigs after challenging with virulent strain of *B. melitensis* 16M infection was achieved through *i.n.* administration of the vaccine in comparison with other methods of application.

The next important step in our study was devoted to the choice of the vaccination dose and, at the same time, the frequency of vaccination, where the study of the protection and immunogenicity of the vaccine candidate was evaluated in animals by the ability to retain bacteria in organs and lymph nodes after animal infection with standard methods. Another distinctive feature of our studies was that the vaccine

protection was assessed not only by the *Brucella* culture isolation from the tissues of vaccinated and unvaccinated animals, but also by such aspects as vaccination efficiency and infection index. It is believed that these indicators jointly provide a more complete and objective characterization of the vaccine protection. The new vaccine induced significant protection in response to *B. melitensis* 16M infection within a range of 60-80% when administered *i.n.* in a double vaccination mode for all tested doses, and it was not inferior in efficiency to *B. melitensis* Rev.1, which is currently used in veterinary practice as the most immunogenic brucellosis vaccine. The level of protection of the *B. melitensis* Rev.1 vaccine obtained in our studies corresponds to the science literature data [25]. At the same time, it was found that the new vaccine candidate does not possess protection after primary vaccination, regardless of the dose. When choosing an immunizing dose of the vaccine, it is recommended to use a vaccination dose of  $10^6$  EID<sub>50</sub>, since the protection at the  $10^6$  EID<sub>50</sub> dose (80% efficiency) was higher than  $10^5$  EID<sub>50</sub> (60% efficiency) and similar to  $10^7$  EID<sub>50</sub> (80% efficiency). The choice of an immunizing dose of  $10^6$  EID<sub>50</sub> is determined by the reduction of possible adverse effect of vaccination and the cost of the production process of the vaccine. The vaccine is targeted at a specific risk group - laboratory scientists working with the pathogen, veterinarians, slaughterhouse workers and people involved in animal care industry.

## Conclusions

The results of the study in guinea pig models show that the recombinant vector anti-brucellosis vaccine candidate is a safe product with wide dose of application and mucosal immunization techniques and its protection properties in prime-boost immunization mode are comparable to that of the commercial *B. melitensis* Rev.1 vaccine against *B. melitensis* 16M infection. We conclude the immunization schedule for our new candidate vaccine against human brucellosis – tetravalent vaccine formulation based on recombinant influenza A virus subtype H5N1 expressing *Brucella* Omp16, L7/L12, Omp19 and SOD in prime-boost intranasal immunization mode at immunization dose of  $10^6$  EID<sub>50</sub>. We could further use this recombinant vaccine vector for pre-clinical and clinical trials in humans.

## Declarations

### Ethics approval and consent to participate

Not applicable.

### Consent for publication

Not applicable.

### Availability of data and materials

All data related to the present study are available in the manuscript.

### Competing interests

No potential conflict of interest was reported by the authors.

## Funding

This work was supported by the Science Committee of the Ministry of Education and Science of the Republic of Kazakhstan under Grant No. AP05131463.

## Authors' contributions

ShR, DB, NZ and KT: conception and design of the study, or acquisition of data. DB: analyze the results, wrote the manuscript and revising it critically for important intellectual content. NA, ZhK, BY, YK and KZ: final approval of the version to be submitted. KT: revised final version of the manuscript. All authors read and approved the final manuscript.

## Acknowledgements

The authors express their gratitude to M. Sarmyкова employee of the Research Institute for Biological Safety Problems, for her assistance in the present study.

## Abbreviations

EID - embryo infectious dose

CFU - colony forming units

rIVV – recombinant influenza viral vector

Omp - outer membrane protein

PBS - phosphate-*buffered* saline

HA - hemagglutinin

NA - neuraminidase


## References

1. Głowacka P, Żakowska D, Naylor K, Niemcewicz M, Bielawska-Drózd A. Brucella - Virulence Factors, Pathogenesis and Treatment. *Pol J Microbiol* 2018; 67(2):151-161.
2. Avijgan M, Rostamnezhad M, Jahanbani-Ardakani H. Clinical and serological approach to patients with brucellosis: A common diagnostic dilemma and a worldwide perspective. *Microb Pathog.* 2019;129:125-130.
3. Pappas G, Akritidis N, Bosilkovski M & Tsianos E. *New Engl J Med* 2005; 352:2325-2336.
4. Corbel MJ. Brucellosis: an overview. *Emerg Infect Dis.* 1997; 3(2):213-21.

5. Al-Tawfiq JA. Therapeutic options for human brucellosis. *Expert Rev Anti Infect Ther* 2008; 6:109-120.
6. Ariza J, Gudiol F & Pallares R. Treatment of human brucellosis with doxycyclin plus rifampicin or doxycycline plus streptomycin. A randomized, double-blind study. *Ann Intern Med* 1992; 117:25-31.
7. Perkins SD, Smither SJ, Atkins HS. Towards a *Brucella* vaccine for humans. *FEMS Microbiol Rev*. 2010;34(3):379-394.
8. Miyoshi A, Bermúdez-Humarán LG, Ribeiro LA, et al. Heterologous expression of *Brucella abortus* GroEL heat-shock protein in *Lactococcus lactis*. *Microb Cell Fact*. 2006; 5:14.
9. Gupta VK, Radhakrishnan G, Harms J, Splitter G. Invasive *Escherichia coli* vaccines expressing *Brucella melitensis* outer membrane proteins 31 or 16 or periplasmic protein BP26 confer protection in mice challenged with *B. melitensis*. *Vaccine*. 2012; 30(27):4017-22.
10. Zhao Z, Li M, Luo D, Xing L, Wu S, Duan Y, Yang P, Wang X. Protection of mice from *Brucella* infection by immunization with attenuated *Salmonella enterica* serovar typhimurium expressing A L7/L12 and BLS fusion antigen of *Brucella*. 2009; 27:5214-19.
11. Oñate AA, Donoso G, Moraga-Cid G, Folch H, Céspedes S, Andrews E. An RNA vaccine based on recombinant Semliki Forest virus particles expressing the Cu,Zn superoxide dismutase protein of *Brucella abortus* induces protective immunity in BALB/c mice. *Infect Immun*. 2005;73(6):3294-3300.
12. Tabynov K. Influenza viral vector based *Brucella abortus* vaccine: a novel vaccine candidate for veterinary practice. *Expert Rev Vaccines*. 2016; 15(10):1237-9.
13. Tabynov K, Kydyrbayev Z, Ryskeldinova S, Yespembetov B, Zinina N, Assanzhanova N, Kozhamkulov Y, Inkarbekov D, Gotskina T, Sansyzybay A. Novel influenza virus vectors expressing *Brucella* L7/L12 or Omp16 proteins in cattle induced a strong T-cell immune response, as well as high protectiveness against *B. abortus* infection. *Vaccine*. 2014;32(18):2034-
14. Tabynov K, Sansyzybay A, Kydyrbayev Z, Yespembetov B, Ryskeldinova S, Zinina N, Assanzhanova N, Sultankulova K, Sandybayev N, Khairullin B, Kuznetsova I, Ferko B, Egorov A. Influenza viral vectors expressing the *Brucella* OMP16 or L7/L12 proteins as vaccines against *abortus* infection. *Viol J*. 2014; 11:69.
15. Hosseinabadi A, Korani M, Esmaeili D. In Silico Analysis *Brucella* OMPs and CagA for expansion of a subunit vaccine candidate versus Brucellosis. *International Journal of Medical Reviews*. 2019; 6(1):14-20.
16. Rice PA, Steitz TA. Ribosomal protein L7/L12 has a helix-turn-helix motif similar to that found in DNA-binding regulatory proteins. *Nucleic Acids Res*. 1989; 17(10):3757-3762.
17. Muñoz-Montesino C, Andrews E, Rivers R, et al. Intraspleen delivery of a DNA vaccine coding for superoxide dismutase (SOD) of *Brucella abortus* induces SOD-specific CD4+ and CD8+ T cells. *Infect Immun*. 2004; 72(4):2081-7.
18. Tibor A, Decelle B, Letesson JJ. Outer membrane proteins Omp10, Omp16, and Omp19 of *Brucella* spp are lipoproteins. *Infect Immun*. 1999; 67:4960-2.



19. Oliveira SC, Zhu Y, Splitter GA. Sequence of the rplJL operon containing L10 and L7/L12 genes from *Brucella abortus*. *Gene*. 1994; 140:137-8.
20. Bricker BJ, Tabatabai LB, Judge BA, Deyoe BL, Mayfield JE. Cloning, expression, and occurrence of the *Brucella* Cu-Zn superoxide dismutase. *Infect Immun*. 1990; 58: 2935-9.
21. Brauer R, Chen P. Influenza virus propagation in embryonated chicken eggs. *J Vis Exp*. 2015; (97):52421.
22. Hien TT, de Jong M, Farrar J. Avian influenza – a challenge to global health care structures. *N Engl J Med*. 2004; 351(23):2363-5.
23. García-Sastre A, Egorov A, Matassov D, Brandt S, Levy D, et al. Influenza A virus lacking the NS1 gene replicates in interferon-deficient systems. *Virology*. 1998; (252): 324-330.
24. Belyakov IM, Ahlers JD. What role does the route of immunization play in the generation of protective immunity against mucosal pathogens?. *J Immunol*. 2009; 183(11):6883-92.
25. Clapp B, Yang X, Thornburg T, Walters N, Pascual, W. Nasal vaccination stimulates CD8(+) T cells for potent protection against mucosal *Brucella melitensis* challenge. *Immunology and cell biology*. 2016; 94(5):496–508.

## Figures




**Figure 1**

Schematic representation of the recombinant influenza viral vector construction destined to generate a vaccine against brucellosis. (A) Full size NS1 protein virulence factor for antagonizing interferon system and (B) deletion part of NS1 replaced by Brucella immunodominant proteins NS1-Omp 16, NS1-Omp 19, NS1-L7/L12 and NS 1-Cu-Zn SOD. The blue box and red box (B) represent RNA binding site of NS1 and part of NS1 protein at amino acid position 80 for insertion brucellosis segment, respectively. The scheme is not drawn according to scale.

**A****B****Figure 2**


Dynamics of body weight change of guinea pigs on day 42 after prime-boost immunization. Animals were immunized with vaccine candidate against human brucellosis at different routes of immunization (A) and various doses (B). Statistical analysis was performed with two-way ANOVA followed by Dunnett's multiple comparisons test showed that during 42-day body weight measurement between the PBS control and vaccinated groups were not significant.  $P < 0.05$  values were considered significant.



**Figure 3**

Protective efficacy of the vaccines in guinea pigs when administered by different routes. Protective efficacy of vaccines as evaluated by the effectiveness of vaccination (A), index of infection (B) and isolation rate of *Brucella* (C) from the tissues of control and experimental groups of guinea pigs on day 30 after challenging with the virulent strain of *B. melitensis* 16M. Animals were vaccinated with the vector vaccine by prime-boost conjunctival (c.), intranasal (i.n.), sublingual (s.l.) at interval of 21 days, and with

*B. melitensis* Rev.1 by single subcutaneous (s.c.) vaccination. Guinea pigs in negative control group were injected with PBS. The challenge of animals was performed with *B. melitensis* 16M at a dose of  $1.3 \log_{10}$  CFU/animal using s.c. route. Bacteriological evaluation was assessed by the index of infection in animals (the arithmetic mean  $\pm$  standard error was given for each group; the number of organs and lymph nodes from which *Brucella* was isolated for each animal) and by counting *Brucella* colonies in tissues, where data is expressed as  $\log_{10}$  CFU/g. Statistical analysis for (B) was performed using a one-way ANOVA followed by Dunnett's multiple comparisons test, (\* $P < 0.01$ , \*\* $P < 0.002$ ) and for (C) two-way ANOVA, Tukey's multiple comparisons test. \* $P=0.04-0.01$ , \*\* $P=0.009 - 0.001$ , \*\*\* $P=0.0004 - .0002$ , \*\*\*\* $P < 0.0001$ .



**Figure 4**

Protectiveness of vaccine samples at different doses in guinea pigs. (A-C) Colonization of *B. melitensis* in organs and tissues of prime-booster vaccinated guinea pigs and (C) index of *Brucella* infection upon challenge with *B. melitensis* 16M. Guinea pigs were immunized twice i.n. 21 days apart with a new vaccine candidate at dose  $10^5$  EID<sub>50</sub>,  $10^6$  EID<sub>50</sub> and  $10^7$  EID<sub>50</sub> or a single delivery of commercial vaccine *B. melitensis* Rev.1. via s.c. immunization. Guinea pigs of positive control groups were injected with PBS. Animals challenged with virulent strain of *B. melitensis* 16M at a dose of  $1.3 \log_{10}$  CFU/animal

using s.c. route. Bacteriological evaluation was assessed by counting *Brucella* colonies in tissues, where data is expressed as log<sub>10</sub> CFU/g and the index of infection of infected animals and compared to the control groups (the arithmetic mean  $\pm$  standard error was given for each group; number of tissues from where *Brucella* was isolated for each animal). Statistical analysis for (A-C) was performed using a one-way ANOVA followed by Dunnett's multiple comparisons test and for (D) using two-way ANOVA, Tukey's multiple comparisons test.