

Prevalence, Pattern and Impact of Self Medication of Anti-infective Agents During COVID-19 Outbreak in Dhaka City

Morshed Nasir (✉ morshednasir@hotmail.com)

Holy Family Red Crescent Medical College <https://orcid.org/0000-0001-5296-7406>

Khandaker Abu Talha

Sylhet Eomen's Medical College

ASM Salauddin Chowdhury

Holy Family Red Crescent Medical College

Tahmina Zahan

Holy Family Red Crescent Medical College

Rawshan Ara Perveen

Holy Family Red Crescent Medical College

Research Article

Keywords: COVID-19, self-medication, anti-infective agents, drug dispensing, rational use of drug, prescription-only drugs

Posted Date: August 12th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-57011/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Global Journal of Medical Research on August 12th, 2020. See the published version at <https://doi.org/10.34257/GJMRBVOL20IS7PG1>.

Prevalence, Pattern and Impact of Self Medication of Anti-infective Agents During COVID-19 Outbreak in Dhaka City

Morshed Nasir¹, Khandaker Abu Talha², ASM Salauddin Chowdhury³, Tahmina Zahan⁴,
Rawshan Ara Perveens

Abstract:

Background: The outbreak of coronavirus disease-2019 is putting a massive strain on vulnerable healthcare system in low and middle-income countries like Bangladesh. Inequitable access to healthcare is further widened by the socio-economic gap and sense of insecurity during this pandemic. Self-medication is a common practice in Bangladesh as it provides a low-cost alternative for people, which involves inappropriate and injudicious use of medicines to treat self-recognized symptoms by the people. During the outbreak of COVID-19 in Dhaka city, the tendency of taking medicines by own decision was thought to be increased alarmingly due to unusual distress, caused by high self-awareness of their health and buying capacity of medication. *Objectives:* To observe the prevalence, pattern, sources and impact of self-medication practice among the respondents with high socio-economic standings and education. *Setting:* Citizens living in Dhaka city, Bangladesh during COVID-19 outbreak from April 2020 to May 2020. *Method:* Cross sectional online survey conducted on 626 citizens without having background of medical knowledge or related to any healthcare services (doctors, nurses, pharmacist, medical students) by structured questionnaires during COVID-19 outbreak. *Outcome measures:* Frequency of self-medication, commonly used antimicrobial agents, symptoms causing self-medication, sources of inspiration, relation with COVID-19 test results, economic burden involved in self-medication during outbreak. *Results:* The prevalence of self-medication amid the outbreak of COVID-19 was 88.33% and only 179 (28.59%) took medication with doctors' advice and remaining 447 (71.40%) respondents took the drugs as "self-medication" by other sources. The most frequently used anti-infective drugs during the outbreak were ivermectine (77.15%), azithromycin (54.15%), doxycycline (40.25%). The common symptoms were fever, throat pain, dry-cough and total 105 (16.77%) respondents took medications without having any symptoms. Almost 355 (85.33%) had taken medication without doing any test for COVID-19. The causes of self-medication as news of spread, effects and remedies in media channels, internet; mental stress of lockdown and isolation, insecurity and panic about scarcity of drug and healthcare support. *Conclusion:* High risks of developing antibiotic resistance, adverse drug reactions and financial loss was predictable with absence of strict regulatory enforcement to protect people and proper utilization resources during COVID-19 outbreak in Dhaka city.

Impact on Practice:

- Self-medication of anti-infective agents during COVID-19 outbreak is high due to inadequate access to the doctors, easy availability of any drugs without prescription and remedy of common symptoms.
- Most of the educated citizens having high socio-economic status in Dhaka become stressed due to scarcity of supply as informed by news media, internet and friends that provoked self-medication of anti-infective agents.
- Unapproved medications for COVID-19 like azithromycin, doxycycline, hydroxychloroquine and ivermectine were used in self-medication that added unnecessary financial burden.

- Lack of monitoring, surveillance and regulations in drug dispensing can be a major cause of resistance, organ damage, adverse drug reactions and interrupted supply chain for high demand during COVID-19 outbreak in Dhaka city.

Introduction:

According to the World Health Organization (WHO), self-medication is explained as “the selection and use of medicines by individuals (or a member of the individual’s family) to treat self-recognized or self-diagnosed conditions or symptoms. Reasons commonly adducted for indulgence in self-medication includes delayed access to healthcare centers, socio cultural belief, relatively high cost of hospital treatment, previous experience of treatment of same symptoms, easy availability of drugs, poor regulatory practice, urgency of feeling relieved, advice from friends and media. The practice of self-medication is prevalent in most parts of Bangladesh regardless of socio-economic status and level of education. While this is indisputable, the incidence of self- medication is may be higher in the low or middle-income countries without consulting with qualified health professionals¹. Although the WHO stressed that rational self-medication practice helps in the prevention and treatment of some minor pathological conditions at affordable cost², but otherwise it may cause wastage of resources, resistance to pathogens and serious health hazards with adverse drug reactions and prolonged morbidity³. In a developing country like Bangladesh, the practice of self-medication may provide an alternative for people as low-cost to avoid high cost of clinical services and many drugs dispensed over the counter (OTC) without prescription⁴. But it is gone beyond the OTC drugs and sometimes prescription-only drugs like antimicrobials, sedatives, hypnotics and strong NSAIDs also reported to be dispensed without prescription in few cases.

Moreover, the ratio between doctor and patient in Bangladesh is currently as low that places the country at second position from the bottom, among the South Asian countries, according to the WHO⁵. Besides, some individuals practice out of ignorance, poverty and generalized hypes.

Dhaka, being the capital and the major economic-hub of the country; it lags behind in the ratio between healthcare workers compared to other neighboring countries, thus hampering proper, and timely healthcare. So, practice of self-medication is almost inevitable as the country has only 6 doctors, nurses, and midwives for every 10,000 populations, according to the report of health bulletin published yearly by the Health Ministry⁶. Some studies on general tendency and pattern of self-medication practice of prescription-only drugs among students and people with or without medical knowledge showed alarming scenario in Dhaka city⁷.

The outbreak of coronavirus disease-2019 is putting a massive strain on vulnerable healthcare system in low and middle-income countries like Bangladesh. Inequitable access to healthcare is further widened by the socio-economic gap and sense of insecurity during this pandemic since the beginning of 2020⁸. Besides, the population of higher socio-economic standings are more potential to have access to better health information, medications and affordability that may lead to self-medication practice in mass. Similarly, people with higher education are reported to have much distress, probably due to high self-awareness and access to mass information network⁹.

The combat against COVID-19 is still continuing in Bangladesh, with the highest incidence rate in Dhaka city. The available data by WHO revealed that the highest AR was observed to continue in the Dhaka (2321.7/1,000,000) and was highest (9422.1/1,000,000) during April to June 2020¹⁰. As there is no approved cure for COVID-19 or a vaccine against SARS-CoV-2, the

aim of treatment is focused to manage and reduce symptoms until clinical recovery. Most people (around 80%) are asymptomatic or mild infection that can be treated at home.

As stated in the National Guideline on Case Management of COVID-19 in Bangladesh, there is no precise effective treatment for COVID-19, the mainstay of management is early diagnosis and supportive care of symptoms and optimum support for organ function in severe illness. No drug is yet recommended as chemoprophylaxis as there is no quality evidence of efficacy and safety in COVID 19. Though Patients should be managed in hospital setting; however, proper home care may also be advised with mild illness unless rapid deterioration or inability to avail hospital if necessary¹¹. But lack of rapid response, scarcity of hospital beds, absence of private practice, inadequate capacity of testing (RT-PCR), spread of un-authenticated treatment protocols are putting the citizens in dilemma of choosing medical advices. Lot of prescription-only drugs like antimicrobials (azithromycin, doxycycline), anti-parasitic (ivermectine), anti-malarial (hydroxychloroquine) are reported to become shortage in the dispensary and peripheral supply chain throughout the city. Moreover, risk of wide range of side effects (Table-I) are also of great concern for undocumented self-medication.

Table-I: List of possible side effects of anti-infective agents used in self-medication during COVID-19

	Mild / common	Severe / rare
Azithromycin	Diarrhea, nausea, abdominal pain, vomiting, headache	Prolong QT interval, Arrhythmia, Hepatic dysfunction, Myasthania
Doxycycline	loss of appetite, nausea, vomiting, diarrhea, rash, sensitivity to the sun, hives, discoloring of teeth, bloody diarrhea, stomach cramping and pain, fever, dehydration, weight loss	headache, blurry vision, double vision, vision loss, Irritation of esophagus, Anemia, Pancreatitis. pain in upper abdomen, fever, skin reactions, blisters, peeling skin, small purple spots
Hydroxychloroquine	nausea, vomiting, stomach pain or cramps, loss of appetite, weight loss, diarrhea, dizziness, spinning sensation, headache, ringing in ears, mood changes, nervousness, irritability, skin rash, itching, or hair loss.	Bone marrow depression, anemia, aplastic anemia, agranulocytosis, leukopenia, and thrombocytopenia. Hemolysis reported in individuals with glucose-6- phosphate dehydrogenase deficiency, Cardiomyopathy, cardiac failure, prolongs the QT interval, Ventricular arrhythmias, torsade de pointes
Ivermectine	tiredness, loss of energy, stomach pain, loss of appetite, nausea, vomiting, diarrhea, dizziness, sleepiness or drowsiness, itchiness, trouble breathing, swelling of throat or tongue, skin rash	Bronchospasm, hepatotoxicity, seizure

Because of commonly occurring symptoms of soreness or pain in throat, dry cough, fever, body ache, breathlessness; people started to take medicines without being diagnosed or tested for COVID-19. Besides the risk of misuse or overuse of these drugs may lead to immediate or delayed complications including adverse drug reactions (hypersensitivity, anaphylaxis), drug interactions, malfunction or destruction of vital organs such as liver, kidney; the practice of self-medication gives a deceitful sense of security and masking the correct diagnosis. Many countries are using different drugs but they are not using those as the guidelines and should only be used under the supervision of physicians on a case-by-case basis, not as a general recommendation¹².

In view of this high prevalence of self-medication of prescription-only drugs in Bangladesh and its associated adverse socio-economic impact on individual and the healthcare service system, supply chain at large. This study is therefore done to evaluate the awareness, pattern and attitude towards self-medication of anti-infective agents among the high socio-economic and educated citizens in Dhaka during the outbreak of COVID-19 without testing, diagnosis and prescription.

Ethics Approval: None of the authors has any conflict including employment, consultancies, stock ownership, honoraria, paid expert testimony, patent applications/registrations for this study.

Methodology:

Using and relying on the authors' network with people living in Dhaka city, the capital of Bangladesh; this online survey adopted a descriptive non-experimental research design to investigate the awareness and practice of self-medication conducted from April to June 2020, the period during the nationwide lockdown and up surging of number of positive COVID-19 cases. Because it was not feasible to do a community-based sampling survey during this outbreak period, we decided to collect data online by convenience sampling. A structured questionnaire was circulated to complete via clicking the link, connected to Google form. The questionnaire contained brief introduction on the background, objective, procedure, voluntary nature of participation, declaration of anonymity and confidentiality, and notes for filling in the online questionnaire.

The online questionnaire was developed and validated through face and content validity techniques, by giving the draft questionnaire to a few of the citizens with inclusion criteria at Dhaka city, to assess whether the response looks meaningful, well designed and/or a good measure of the construct to an innocent bystander. The response was used to refine and modify the questionnaire further.

The content validity was done by giving the resultant questionnaire to three independent scholars from the fields of Public Health, Pharmacology and Social Statistics to assess its appropriateness, clarity, coverage and relevance to the study.

The reliability of the validated questionnaire was ascertained by test retest method. The questionnaire was administered twice at two weeks' interval on ten respondents from Dhaka city who practiced self-medication during COVID-19 outbreak. The responses were compared and the reliability coefficient determined ($r=0.83$). The incorporated draft questionnaire was recast for ambiguity and repetitive questions were struck off.

The cross sectional populations were included with the inclusion criteria as, the adult citizens living in Dhaka city, aged 25 years or more, with education level of graduation or above, non-

medical professionals, having email address, agreed to participate voluntarily. Respondents having involvement or knowledge on medical background (medical graduates, medical practitioners, nurses, medical researchers, pharmacist) were excluded to find out the public perceptions and their responses to take medications without prescription during COVID-19 outbreak. The frequencies of response were recorded in datasheet and observed according to demographic characteristics, sources of information, clinical symptoms, status of COVID-19 test results and cost involvement. The market prices of commonly used medicines as self-medication during the outbreak were calculated to reveal the economic burden caused COVID-19 driven self-medication practice.

Results:

Total 639 participants completed the online survey questionnaire and submitted with e-mail verification. After excluding 13 respondents, of whom 10 were doctors by profession and 3 were below the minimum age limit (25 years); the final sample consisted of 626 valid participants. Among the final sample, 316 (50.47%) respondents were from the age group of 45-54 years, 346 (55.27%) were women, 312 (49.84%) held a bachelor degree and 230 (36.74%) engaged in non-civil services. Other demographic characteristics are shown in Table-II.

Table-II: Frequency distribution of respondents by their socio-demographic features

Variables	Frequency	Percentage (%)
Gender:		
Male	282	45.04%
Female	346	55.27%
Age:		
25-34	96	15.33%
35-44	156	24.92%
45-54	316	50.47%
55 +	58	9.26%
Education level:		
Undergraduate	110	17.57%
Bachelor degree	312	49.84%
Master degree	178	28.43%
Doctoral and advanced	26	4.15%
Work status:		
Students	84	13.41%
Civil service	13	2.06%
Non-civil service	230	36.74%
Self-employed / business	184	29.39%
Retired	38	6.07%
Unemployed	77	12.30%

Table-III: Frequency distribution of respondents by use of anti-infective agents

		Number of Respondents	Azithromycin		Doxycycline		Hydroxychloroquin		Ivermectin	
			N	Cost involved (USD)	N	Cost involved (USD)	N	Cost involved (USD)	n	Cost involved (USD)
RT-PCR test	Positive	132	111	490	93	227	84	420	109	160
	Negative	78	20	102	31	76	0	0	55	81
	Not done	416	208	1046	128	313	44	215	319	469
Total		626	339	1638	252	616	128	635	483	710

- Cost involved = Unit price x Daily dosage x Number of days
- Total cost converted from local BDT to USD

Out of 626 respondents, only 73 (11.66%) did not take any medication during the survey period as 12 (1.91%) were negative by rRT-PCR test for COVID-19 and 61 (9.74%) did not test. Remaining 132 respondents (21.08%) was found positive test result for COVID-19.

The most frequently used prescription-only drug among the respondents were ivermectine (77.15%), azithromycin (54.15%), montelukast (43.13%), calcium supplements (41.37%), doxycycline (40.25%) and hydroxychloroquine (20.44%) respectively. The frequency of taking the prescription-only drugs among the respondents who were not even the diagnosed by rRT-PCR positive result, were reported higher with ivermectine (76.68%) and azithromycin (50.0%) as shown in Table-III.

Figure-1: Distribution of respondents taken medication according to source of advices






Figure-2: Distribution of respondents taken medication according to symptoms



Among the respondents, only 179 (28.59%) took medication with doctors' advice and remaining 447 (71.40%) respondents took the drugs as "self-medication" by other sources like friends/family, pharmacy/dispensary and media/internet as shown in Fig-1.

Total 105 (16.77%) respondents took medications without having any symptoms. Among the remaining 521 respondents, common symptoms for which the respondents took medications were fever (37.61%), throat pain (28.79%), dry-cough (14.20%), loss of smell (9.21%), loss of taste (3.45%), body ache (4.99%) and rarely diarrhea (1.72%) respectively as shown in Fig-2.

Figure-3: Raised price of the anti-infective agents during COVID-19 outbreak



The market price of the anti-infective agents was higher than the retail price as shown in Fig-3. Azithromycin raised almost 37%, hydroxychloroquine 18%, doxycycline 12% and ivermectine 7% respectively.

Discussion:

To the best of our knowledge, this is the first online survey in Dhaka city evaluating the pattern and frequency of self-medication practice of prescription-only drugs in COVID-19 outbreak (April to June 2020) in Bangladesh. As the survey population was well-educated adults from different working status, the respondents' participation was expected to represent the sincerest and accurate scenario of the defined survey population.

Almost equal proportion of male and female participated in the study and most of them (50.47%) belonged to the age group of 45-54 years. Highest number of respondents (49.84%) were graduates (Bachelor degree) and lowest with doctoral degree (4.15%). Only 2.06% respondents were from civil service and most of the rest were from non-civil service (36.74%), self-employed or business (29.39%) living in Dhaka city at the time of COVID-19 pandemic.

Among 626 survey population in Dhaka city who had taken medication for COVID-19, only 132 (21.08%) were documented as positive and 78 (12.45%) as negative by RT-PCR test. The rest of 416 (66.45%) had never done the test, but almost 355 (85.33%) had taken medication without doing any test for COVID-19. This finding could be due to having additional distress due to high self-awareness of their health as reported by Roberts T. et al. among people with higher educational status⁹. Having self-medication without detecting COVID-19 among a large number of respondents could also be due to feeling of insecurity influenced by availability of local medical resources, efficiency of public health system, and prevention and control measures taken in pandemic situation¹⁷.

The rate of self-medication of antimicrobial agents like azithromycin (54.15%), doxycycline (40.25%) were found much higher during the outbreak of COVID-19 comparing to 21% and 25% for azithromycin and doxycycline before the pandemic as reported by Chowdhury N et al¹³. Azithromycin was the fifth highest percentage of people with self-medication throughout the previous years, whereas it became most common antibiotic during the present pandemic. On the other hand, ivermectin being the anti-parasitic agent was self-medicated by 77.15% of the respondents. This might be due to the nationwide broadcast of an experience by a team of Bangladeshi physicians and Bangladesh Medical College Hospital (BMCH) claimed as "outstanding results" in 60 patients with COVID-19 patients all of whom recovered in combination of ivermectin and doxycycline¹². This attempt was made on the outcome of an in-vitro study reported as a single treatment by ivermectin is capable of ~5000-fold reduction of viral load at 48 hours in cell culture¹⁴. Self-medication of this drug was found highest (483/626) among the respondents irrespective of test results by RT-PCR. Though the drugs like chloroquine, hydroxychloroquine and azithromycin, was recommended in the treatment protocol of Covid-19 patients in Bangladesh, according to the "National Guidelines on Clinical Management of Coronavirus Disease-2019" published in the health directorate's website¹¹; hydroxychloroquine was used much less (20.44%) than any other antimicrobials without prescription. This could be due to mass publicity and sharing of news in national and international news and social media as the drug can cause hazardous abnormalities in cardiac rhythm in COVID-19 patients, and should be limited only in clinical trials or hospitals with adequate facilities to monitor any cardiac complications, warned by FDA in a safety communication briefing globally¹². The overall the prevalence and dominance of self-

medication of antimicrobials in low and middle income countries were reported around 39% in previous studies before COVID-19 pandemic^{15,16}; but was outrageously higher (88.33%) in Dhaka city during the pandemic.

Considering the sources or advice for medication, only 179 (28.59%) respondents followed or consulted with doctors and rest by media or internet (27.15%), pharmacy or dispensary (24.44%) and friends or family (19.8%). This finding was very much similar to the previous studies that reported the high prevalence of self-medication (including antimicrobials) since people could obtain any drugs from the pharmacies without prescription even in the distant areas of the country¹⁸. Moreover, during pandemic, people struggle to cope with constant news of the spread and effects of COVID-19 on news-media, social-media, internet without having adequate forms of social support and access to doctors as a result of lockdowns and self-isolation^{8,19}. Most of the respondents (37.61%) took antibiotic as self-medication for fever during Covid19 outbreak, followed by throat pain (28.79%), dry cough (14.20%); whereas almost 16.77% respondents had no symptoms whatsoever. Having inappropriate antimicrobials and supplementary medications (zinc, calcium, Vitamin-D) without prescription is associated with the risk of drug interactions, masking symptoms of underlying diseases and most importantly, the development of antimicrobial resistance^{20,21}.

Prices of essential COVID-19 medicines have increased 4% globally since February 2020, as reported by Gustav Ando in Life Science Research and Analysis. Although there is no specific treatment for Covid-19, the drug administration in Bangladesh started working in advance to increase the production of some supportive medicines²². Despite the availability of essential drugs, the increase demand in major cities for self-medication of azithromycin raised almost 37%, hydroxychloroquine 18%, doxycycline 12% and ivermectine 7% respectively. The total amount of buying those medicines in response to COVID-19 situation had unnecessary financial burden to the people in Dhaka. Overall economic hardship in low and middle-income countries have to cope with the added expenditure, that could be avoided by strict regulatory surveillance on self-medication and dispensing without prescription.

Although there is no approved specific medication to prevent or treat COVID-19, this online survey among the educated adults with high socio-economic standings revealed that high prevalence of self-medication of prescription-only drugs was persistent during the outbreak in Dhaka city. Unsolicited news of spread, effects and remedies in media channels, internet; mental stress of lockdown and isolation, insecurity and panic about scarcity of drug and healthcare support might have triggered up the practice self-medication. Psychological distress levels were also influenced by availability of local medical resources, prevention and control measures²³. There is sufficient evidence of increasing resistance to antibiotics in the Bangladesh resulting from irrational and misuse of antibiotics, where sales of antibiotics are not restricted at any level²⁴. Unopposed access to buy antibiotics and its injudicious use might provoke a long-term burden of drug-resistant strains with problems of under and over-dosage, treatment failure and severe adverse effects to vital organs. It also results in delay in care seeking, which results in paradoxical economic loss due to delayed diagnosis and irrational treatment²⁵. Bangladesh is already in the burden of lower ratio of healthcare workers and supply support system, with high rate of community-transmission of COVID-19 and require effective and efficient enforcement of regulation against free display and sales of drugs without prescription and individuals duly authorized.

Conclusion:

The pattern of medication, including self-medication is an important health indicator, which reflect the degree of supply utilization and regulatory enforcement of healthcare services within a community. Immediate acceleration of health education campaigns, strict legislations on dispensing drugs and increasing the quality and access to healthcare are the important interventions that might change the people's health seeking behavior²⁶. Pandemic situation of COVID-19 is likely to sustain for years and will have huge socio-economic and psycho-social impact on people's lifestyle and behavior, as predicted by the WHO and epidemiologists from different regions. Therefore, a vast nationwide survey and surveillance should be done on self-medication of mass population to protect them from the potential risks, overuse shortage and irrational financial burden during the COVID-19 outbreak.

Acknowledgement:

We gratefully acknowledge all the respondents for their spontaneous participation in the study. We also extend our gratitude and thanks to Dr. Farhana Selina for data extraction, logistic and software support.

Funding: No fund or grants to declare.

Conflicts of Interest: None of the author or co-authors has any conflict of interest regarding any financial and personal relationships with other people or organizations that could inappropriately influence (bias) the work.

References:

1. Oyediran O, Ayandrian EO, Olatubi MI et al. Awareness of risk associated with self-medication among patients attending out-patient department of a tertiary hospital in south western Nigeria. *International Journal of Africa Nursing Science* 2019; 10: 110-115.
2. WHO. Report of the WHO Expert Committee on National Drug Policies. Geneva, Ed:1, 1995. Available at: <https://apps.who.int/iris/handle/10665/63068?locale-attribute=en&show=full>
3. Hughes CM, McElnay JC, Fleming GF. Benefits and risks of self-medication. *Drg Saf.* 2001; 24:1027-37.
4. Hussain S, Malik F, Hameed A, Riaz H. Exploring health seeking behavior, medicine use and self-medication in rural and urban Pakistan. *Southern Med Rev.* 2008; 3:32-34.
5. <https://www.who.int/workforcealliance/countries/bgd/en/>
6. Report in Dhaka Tribune, July 21, 2019. Available at: <https://www.dhakatribune.com/health/2019/07/21/patient-doctors-nurses-ratio-bangladesh-lags-far-behind-its-neighbours>
7. Nasir M, Zahan T, Parvin RA et al. Comaparative study on knowledge, attitude and practice of self-medication among the medical and non-medical undergraduate students in Dhaka city. *World Journal of Pharmaceuticals and Life Sciences* 2017; 3(4): 17-20.

8. Kretchy IA, Asiedu-Danso M, Kretchy JP. Medication management and adherence during the COVID-19 pandemic: Perspectives and experiences from low-and middle-income countries. *Research in Social and Administrative Pharmacy*. Available at: <https://www.sciencedirect.com/science/article/pii/S1551741120303326> [IN PRESS]
9. Roberts T, Miguel Esponda G, Krupchanka D, et al. Factors associated with health service utilization for common mental disorders: a systematic review. *BMC Psychiatry* 2018; 18:262.
10. WHO. Available at: https://www.who.int/docs/default-source/searo/bangladesh/covid-19-who-bangladesh-situation-reports/who-covid-19-update-18-20200629.pdf?sfvrsn=6ef8ba08_2
11. National Guidelines on Clinical Management of Coronavirus Disease 2019 (Covid-19), by Disease Control Division, Directorate General of Health Services Ministry of Health & Family Welfare; Government of the People's Republic of Bangladesh. Version-4, 30 March, 2020. Available at: <https://dghs.gov.bd/index.php/en/home/5343-covid-19-update>
12. News Report, available at: <https://thelogicalindian.com/story-feed/awareness/selfmedication-coronavirus-covid-19-pandemic-20327>
13. Chowdhury N, Islam MR, Hasan MM, et al. *International Journal of Pharmacy Teaching & Practices*. 2013;4(1):504-510.
14. LeonCaly, Julian D.Druce, Mike G.Catton, et al. The FDA-approved drug ivermectin inhibits the replication of SARS-CoV-2 in vitro. *Antiviral Research*. 2020; 178: 104787.
15. Ocan M, Obuku EA, Bwanga F, et al. Household antimicrobial self-medication: a systematic review and meta-analysis of the burden, risk factors and outcomes in developing countries. *BMC Public Health*. 2015; 15:742.
16. Morgan DJ, Okeke IN, Laxminarayan R, et al. Non-prescription antimicrobial use worldwide: A systematic review. *Lancet Infect Dis*. 2011; 11: 692–701.
17. Wind TR, Komproe IH. The mechanism that associate community social capital with post-disaster mental health: a multilevel model. *Soc Sci Med* 2012; 75: 1715-20.
18. Islam MS. Self-medications among higher educated population in Bangladesh: an email-based exploratory study. *Internet J Health* 2007; 5: 2.
19. Huibo Li, Siqian Zheng, Fang Liu, et al. Fighting against COVID-19: Innovative strategies for clinical pharmacists. *Res Soc Adm Pharm* 2020; [IN PRESS] Available at: <https://www.sciencedirect.com/science/article/pii/S1551741120303284>
20. Okeke IN, Klugman KP, Bhutta ZA. Antimicrobial resistance in developing countries Part II: strategies for containment. *Lancet Infect Dis*. 2005; 5: 568–80.
21. World Health Organization. Community-Based Surveillance of Antimicrobial use and Resistance in Resource constrained settings. A report on five pilot projects. Geneva, Switzerland: WHO; 2009. Available at: <http://www.apps.who.int/medicinedocs/documents/s16168e.pdf>.
22. DGDA- Medicine supply, prices managed well amid crisis. *The daily Star*; May08, 2020 at <https://www.thedailystar.net/frontpage/news/medicine-supply-prices-managed-well-amid-crisis-1900585> Viewed on: 30 July, 2020.

23. Qiu J, Shen B, Zhao M, et al. A nationwide survey of psychological distress among Chinese people in the COVID-19 epidemic: implications and policy recommendations. *General Psychiatry*. 2020; 33: e100213.
24. Biswas M, Roy MN, Manik IN, et al. Self-medicated antibiotics in Bangladesh: a cross sectional health survey conducted in the Rajshahi city. *BMC Public Health* 2014; 14: 847.
25. Kalyan VS, Sudhakar K, Srinivas P, et al. Evaluation of self-medication practices among undergraduate dental students of tertiary care teaching dental hospital in south India. *Journal of Education and Ethics in Dentistry* 2013; 3(1):21-25.
26. Alghanim, SA. Self-medication practice among patients in a public health care system. *Eastern Mediterranean Health Journal* 2011; 7(5): 409-416.

Figures




Figure 1

Distribution of respondents taken medication according to source of advices




Figure 2

Distribution of respondents taken medication according to symptoms




Figure 3

Raised price of the anti-infective agents during COVID-19 outbreak