Is Outdoor-Resting Behaviour in Malaria Vectors Consistent? Short Report From Northern Ghana.

Majidah Hamid-Adiamoh (✉ majidah.hamid-adiamoh@lshtm.ac.uk)

University of Ghana <https://orcid.org/0000-0002-0101-5445>

Davis Nwakanma

Medical Research Council The Gambia: Medical Research Council Unit The Gambia at the London School of Hygiene and Tropical Medicine

Isaac Sraku

University of Ghana College of Health Sciences

Alfred Amambua-Ngwa

Medical Research Council The Gambia: Medical Research Council Unit The Gambia at the London School of Hygiene and Tropical Medicine

Yaw A Afrane

University of Ghana College of Health Sciences

Short report

Keywords: Mark-release-recapture, resting behaviour, outdoor resting, malaria vectors, northern Ghana

Posted Date: June 7th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-572080/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

Recent reports of a change in the resting behaviour of malaria vectors, from predominantly indoor resting to outdoor resting following blood feeding, have been attributed to selection pressure from use of vector control tools such as indoor residual spraying and long-lasting insecticide-treated nets. Recent studies have observed vectors resting predominantly outdoors in settings where anti-vector tools are extensively deployed. This present study examined if the outdoor resting behaviour in the vector population, is random or indicative of a consistent preference of one resting site over the other.

Methods

Mark-release-recapture (MRR) experiments were conducted with outdoor-resting *Anopheles gambiae* and *An. funestus* mosquitoes collected from pit shelters, animal houses and granaries in two villages in Northern Ghana. Mosquitoes were marked with fluorescent dyes and released indoors. The experiments were controlled with indoor-resting mosquitoes, which were marked and released outdoors. Species of all recaptured mosquitoes were identified and assessed for consistency in their resting behaviour.

Results

A total of 4,460 outdoor-resting mosquitoes comprising *An. gambiae sensu lato (s.l.)* (59%) and *An. funestus* complex (41%) were marked and released. Overall, 31 (0.7%) mosquitoes (25 *An. gambiae s.l.* and 6 *An. funestus* complex) were recaptured mostly from outdoor location. Only 3 of the recaptured mosquitoes were found resting indoors where they were released. The majority of the outdoor-recaptured mosquitoes were *An. arabiensis* (11, 39%), followed by *An. coluzzii* (7, 25%); whereas the indoor-recaptured mosquitoes were mainly (3) *An. coluzzii*. For the control experiment, 324 indoor-resting mosquitoes constituting 313 *An. gambiae s.l.* and 11 *An. funestus* complex were marked and released. However, none of these was recaptured neither indoors nor outdoors.

Conclusions

The mosquitoes demonstrated the tendency to retain their outdoor-resting behaviour. Further investigations are required to ascertain if emerging preference for outdoor resting behaviour in malaria vector populations is consistent or a random occurrence.

Background

Vector control with indoor residual spraying (IRS) and long-lasting insecticidal nets (LLINs) has contributed largely to reduction in malaria incidence and mortality [1]. These tools are both insecticide and indoor-based and target vectors that feed indoors (endophagic) and rest indoors (endophilic) [2–4]. However, recent reports are indicating that vector control tools are inducing behavioural changes in mosquitoes such that some vector species have shifted from indoor-resting behaviour to resting outdoors

(exophilic). This was as documented from studies in Ethiopia [5], Ghana [6, 7], Kenya [8,9], Libreville [10] and Tanzania [11, 12]. Such behavioural diversification in vector population can be detrimental to vector control [13, 14], which mainly target vectors resting indoors. Therefore, efforts to understand emergence of such behaviour are highly essential for elimination agenda.

Mark-release-recapture (MRR) studies are a standard experimental method extensively employed to investigate mosquito biology, ecology, life history and behaviour, measuring parameters including dispersal and flight distance; survival, population size and density, as well as blood feeding, host seeking and reproductive behaviour [15–18]. MRR experiments have been used particularly for field and laboratory studies involving mosquito species such as *Aedes* [19, 20], *Culex* [21, 22] and *Anopheles* [15, 16, 18]; to collect data highly relevant in understanding pathogen transmission, gene flow and development and optimization of vector intervention tools [16, 18, 23].

Outdoor resting behaviour in vector populations may be triggered by heavy use of insecticide-based indoor intervention in settings where control tools are widely implemented [11, 24, 25]. Vectors in these settings avoid contact with insecticides and rest outdoors where no insecticide is used [11, 24–26]. Such behaviour promotes outdoor transmission as vectors can bite unprotected humans outdoors [11, 27].

Northern Ghana has documented outdoor resting in *An. gambiae sensu lato* (s.l.) population [6, 7] but it is not clear whether this behavior only occurs randomly in the vector population. This present study examined if the preference for outdoor resting in the vector population was consistent or not. MRR experiments were conducted using field-collected mosquitoes of unknown age in two villages in northern Ghana, where coverage of IRS and LLINs are high [7, 28, 29].

Materials And Methods

Study sites

The MRR experiments were conducted in two villages in northern Ghana, Kpalsogu (9.33⁰ N, 1.02⁰ W) and Libga (9.35⁰ N, 0.51⁰ W) during the rainy season between July-November 2017 and dry season in December 2017-January 2018. The region has a unimodal rainfall pattern with monthly density between 150–250 mm and a forest vegetation zone with mean daily temperature ranging between 25–30°C; and an average relative humidity between 65–75% [30]. .

This region has previously documented malaria incidence rate of about 40% in under-five children (29,31) and LLINs coverage is > 70% in both villages [7, 29]. Kpasolgu has been under IRS implementation since 2008 [7]. Extensive farming is practiced in both villages, occurring throughout the year as the villages are close to irrigation dams.

Mosquito collection

Live outdoor-resting adult female *Anopheles* mosquitoes were collected from pit shelters, granaries and animal resting shelters. The pit shelters were positioned about 5-10m outside six randomly selected compounds in each village. The mosquitoes were carefully collected using mouth and prokopack electrical aspirators so that they remain alive. Collections were carried out during the early mornings (05.00–07.00).

To compare the trend in behaviour of mosquitoes when resting indoors and also as a control for the experiments, indoor-resting mosquitoes were also collected from sleeping rooms in the compounds where the pit shelters were situated, using mouth and prokopack aspirators at the same time the outdoor mosquitoes were collected.

Mosquito processing, marking and subsequent release.

Captured mosquitoes were immediately transported to the insectary for morphological identification as female *An. gambiae s.l.* and *An. funestus* complex [32]. Following identification, indoor and outdoor mosquitoes were carefully kept in separate cages in the insectary, irrespective of their abdominal status. The mosquitoes were fed with 10% glucose until experiment time at dusk of the same day between 18:00–19:00.

The marking of mosquitoes was carried out 3 hours to release time. Batches of 15–20 mosquitoes from the same compounds were kept in the same paper cup. The mosquitoes were lightly dusted with red fluorescent dye and left in the cup to rest and feed on 10% glucose until when transported to the study sites for release. The indoor (control) specimens were similarly marked with green dye.

At least 200 mosquitoes were released in five sleeping rooms (indoors) of the same compounds where the pit traps were situated. This was done to assess if they would remain indoors to rest or return outdoors where they were originally collected from. The control indoor-collected mosquitoes were released outdoors to ascertain if an alternate trend in the resting behaviour would be demonstrated.

Recapture of released mosquitoes

The mosquitoes were left for 48 hours before recapture was initiated. This offered sufficient time for the mosquitoes to redistribute in their new environment and enabled us to assess if the mosquitoes opted to return to their initial collection location.

To recapture, mosquitoes were sought from sleeping rooms, animal houses, clay pots, granaries and pit traps deploying light traps, exit traps, pyrethrum spray catches as well as mouth and prokopack aspirations. All rooms and compounds in each of the study sites were sampled to recapture the released mosquitoes. Recapturing was done for 7 consecutive days before the next release event. In all, twelve (12) rounds of MRR experiment were conducted within the study period.

Processing and identification of recaptured mosquitoes

All recaptured mosquitoes were morphologically identified as marked and unmarked. Only the marked Anopheles mosquitoes were selected for data collection and assessed for change in resting behaviour. All unmarked mosquitoes were counted and recorded.

The marked recaptured mosquitoes were processed to discriminate the sibling members of *An. gambiae s.l.* and *An. funestus* complex. DNA was extracted from these mosquitoes and molecular species were genotyped using the protocols as previously described [33, 34].

Ethical considerations

Ethical approval was obtained from the Ghana Institutional Review Boards (IRB) of Noguchi Memorial Institute for Medical Research (NMIMR). Village heads were also engaged in discussions where concerns on release of mosquitoes into sleeping rooms were clarified. MRR experiment presents no potential risks as mosquitoes were released in the compounds where they were initially captured and a proportion was to be recaptured.

Results

Recaptured mosquitoes and their species

Overall, a total of 4,460 outdoor-resting mosquitoes were released to indoor space, while 324 indoor-resting mosquitoes were released outdoors as control. The morphological identification of test mosquitoes (outdoor) revealed that 2, 636 (59 %) were *An. gambiae s.l.* while 1, 824 (41%) were *An. funestus* complex (Figure 1). Whereas the control mosquitoes comprised 313 (97%) *An. gambiae s.l.* and 11 (3%) *An. funestus* complex.

Following all the 12 MRR experiments conducted, a total of 3, 950 Anopheles mosquitoes were recaptured, of which 3,919 (99.2%) were unmarked and 31 were marked (0.8%). These unmarked mosquitoes were notably not among the mosquitoes released for the MRR experiments. Subsequent analyses of the outcome of the MRR experiments were focused only on the marked mosquitoes that were recaptured.

Recapture of the marked mosquitoes was achieved from nine (9) out of twelve release experiments. Thirty (31) marked test mosquitoes were recaptured from 4,460 released. This indicates a recapture rate of 0.7%. The breakdown of the recaptured test mosquitoes comprised 25 *An. gambiae s.l.* and 6 *An. funestus* complex (Figure 1), from which twenty eight (28) were resting outdoors, and three (3) indoors. The majority of the mosquitoes recaptured outdoors were *An. arabiensis* (11, 39%), followed by *An. coluzzii* (7, 25%) and *An. funestus s.s.* (6, 21%). *An. gambiae s.s.* constituted the least outdoor-recaptured species (4, 15%) (Table 1). The three mosquitoes recaptured indoors were mainly *An. coluzzii*. None of the control samples that were released outdoors was recaptured from any location, indoor or outdoor.

Table 1: Molecular species of the recaptured test mosquitoes

Mosquito species	# Recaptured	
	Indoor	Outdoor
<i>An. arabiensis</i>	0	11
<i>An. coluzzii</i>	3	7
<i>An. gambiae s.s.</i>	0	4
<i>An. funestus s.s.</i>	0	6

Discussion

Behavioural shift from predominantly indoor to outdoor-resting is becoming widespread in vector populations in settings where anti-vector interventions are extensively deployed [6, 24, 25, 35, 36]. The aim of this study was to examine if outdoor resting behaviour of the malaria vector populations described in the study settings [6, 7] was consistent or it was just a random occurrence. The study observed that the majority of mosquitoes returned outdoors to rest after being released indoors. This pattern of preference for outdoor resting in mosquitoes was consistently demonstrated from nine out of the twelve experiments, despite a low recapture rate. As outdoor resting behaviour in vector population could counteract control efforts [27, 37, 38], research assessing the extent of this behaviour is well-timed and necessary to guide decision making in malaria control programmes.

Switch to outdoor-resting behaviour in the vectors may be as a result of avoidance behaviour from indoor interventions [38, 39], which were heavily used in the study sites. Most sleeping rooms had LLINs and they were also sprayed with pirimiphos methyl, the insecticide that was being used for IRS at the time of the experiments [7]. These mosquitoes might be avoiding contact with the insecticide indoors and chose to rest outdoors, where they may be safe, as there was no outdoor intervention in the villages. This could have also accounted for the predominant outdoor resting preference documented in the vectors in a previous study from this setting [6] and others all over Africa [8, 12, 36, 37], which prompted this investigation.

The highly endophilic vectors: *An. gambiae s.s.*, *An. coluzzii* and *An. funestus s.s.* were recaptured outdoors after being released indoors. This demonstrates a likely switch to outdoor resting behavior because these vector species may be avoiding contact with insecticide indoors and exiting to rest outdoors [40–42]. Change from endophily to exophily in response to control intervention is increasingly being documented in these highly endophilic vectors [8, 12, 25, 35, 44, 45]. This could have negative impact on malaria control as it can promote outdoor and residual transmission in these settings [11, 38, 45].

Anopheles arabiensis was found to predominate the mosquito species found resting outdoors from the recaptured test mosquitoes. This may be because *An. arabiensis* is known to be highly exophilic and zoophilic (preference for animal blood meal) [46–48], where it tends to stay outdoors to rest and feed on animals when there is intervention indoors [4, 49]. This vector species has also been observed to display insecticide avoidance and early-exiting behaviour [12, 27, 42], which make them difficult to control [11].

In this study, the recapture rate was much lower than most MRR studies reported. Several reasons could have accounted for this. One of such reasons is age of mosquitoes [16, 18, 50], which was unknown in this study. Plausibly, if the mosquitoes were old, they may have died within the period of recapture. Indeed, senescence is a factor associated with reduction of mosquito daily survival in the wild [51, 52]. Age was also previously suggested to be responsible for the low recapture rate in a similar experiment [16]. Moreover, predators such as spiders were particularly common in the study areas, and were found inside animal houses that provide favourable environment for the mosquitoes to rest. Other factors including emigration from the study area, climate condition, stress and negative effect of the experimental procedures could have also contributed to this low recapture success as previously documented in other MRR experiments [16, 18, 23].

Conclusions

This study showed that preference for outdoor resting in malaria mosquitoes may be an emerging consistent behaviour. There is a need for further studies to establish this observation in settings where interventions are extensively deployed. Furthermore, a probe into the genetic basis underlying this behavioural change will also be highly essential. This is important as malaria control moves to the elimination phase in sub-Saharan African countries.

Abbreviations

MRR: Mark, release and recapture

DNA: Deoxyribonucleic Acid

IRS: Indoor residual spraying

LLINs: Long-lasting insecticidal nets

SIT : Sterile insect techniques

s.l.: *sensu lato*

s.s.: *sensu stricto*

Declarations

Acknowledgements

We appreciate Messrs Abdul Rahim Mohammed, Dhikrullahi Shittu, Honourable Chris and Osei Kwaku Akuoko for their assistance in the field work.

Funding

This work was supported by funds from a Wellcome Trust DELTAS Africa grant (*DEL-15-007: Awandare*), the National Institute of Health (R01 A1123074 and D43 TW 011513), University of Ghana Research Funds (URF/9/ILG-078/2015-2016), H3Africa PAMGENe project, H3A/18/002 and TWAS Fellowship for Research and Advanced Training programme, 2019 (FR number: 3240307913). Majidah Hamid-Adiamoh was supported by a WACCBIP-Wellcome Trust DELTAS PhD fellowship. The DELTAS Africa Initiative is an independent funding scheme of the African Academy of Sciences (AAS)'s Alliance for Accelerating Excellence in Science in Africa (AESA) and supported by the New Partnership for Africa's Development Planning and Coordinating Agency (NEPAD Agency) with funding from the Wellcome Trust (107755/Z/15/Z: Awandare) and the UK government. The views expressed in this publication are those of the author(s) and not necessarily those of AAS, NEPAD Agency, Wellcome Trust or the UK government.

Authors' contributions

MHA designed, performed the field and laboratory work, analyzed data and drafted the manuscript. YAA conceived and supervised the study, analyzed data and revised the manuscript. AAN and DN supervised study and revised the manuscript. IS, ARM and OKA performed field and laboratory experiments.

Competing Interests

The authors have declared that no competing interests exist.

Ethical Approval

Ethical approval was given by the Institutional Review Board of the Noguchi Memorial Institute for Medical Research (NMIMR), University of Ghana. Verbal consent was also obtained from village heads before all field work.

Consent for publication

Not applicable.

Availability of data and materials

All relevant data are presented within the paper. No supporting information is available.

References

1. WHO. World Malaria Report. World Health Organization. Geneva. 2020.
2. Githeko AK, Service MW, Mbogo CM, Atieli FK. Resting behaviour, ecology and genetics of malaria vectors in large scale agricultural areas of Western Kenya. *Parassitologia*. 1996;38:481–9.
3. Githeko AK, Adungo NI, Karanja DM, Hawley WA, Vulule JM, Seroney IK, et al. Some observations on the biting behavior of *Anopheles gambiae s.s.*, *Anopheles arabiensis*, and *Anopheles funestus* and their implications for malaria control. *Exp Parasitol*. 1996;82:306–15.
4. Coluzzi M, Sabatini A, Petrarca V, Angela Di Deco M. Behavioural divergences between mosquitoes with different inversion karyotypes in polymorphic populations of the *Anopheles gambiae* complex. *Nature*. 1977;266:832–3.
5. Kibret S, Wilson GG. Increased outdoor biting tendency of *Anopheles arabiensis* and its challenge for malaria control in Central Ethiopia. *Public Health*. 2016;141:143–145.
6. Hamid-Adiamoh M, Amambua-Ngwa A, Nwakanma D, D'Alessandro U, Awandare GA, Afrane YA. Insecticide resistance in indoor and outdoor-resting *Anopheles gambiae* in Northern Ghana. *Malar J*. 2020;19:314.
7. Coleman S, Dadzie SK, Seyoum A, Yihdego Y, Mumba P, Dengela D, et al. A reduction in malaria transmission intensity in Northern Ghana after 7 years of indoor residual spraying. *Malar J*. 2017;16:324.
8. Degefa T, Yewhalaw D, Zhou G, Lee MC, Atieli H, Githeko AK, et al. Indoor and outdoor malaria vector surveillance in western Kenya: Implications for better understanding of residual transmission. *Malar J*. 2017;16:443.
9. Machani MG, Ochomo E, Amimo F, Kosgei J, Munga S, Zhou G, et al. Resting behaviour of malaria vectors in highland and lowland sites of western Kenya: Implication on malaria vector control measures. *PLoS One*. 2020;15:e0224718.
10. Mourou J-R, Coffinet T, Jarjaval F, Cotteaux C, Pradines E, Godefroy L, et al. Malaria transmission in Libreville: results of a one year survey. *Malar J*. 2012;11:40.
11. Killeen GF, Govella NJ, Lwetoijera DW, Okumu FO. Most outdoor malaria transmission by behaviourally-resistant *Anopheles arabiensis* is mediated by mosquitoes that have previously been inside houses. *Malar J*. 2016;15:225.
12. Kreppel KS, Viana M, Main BJ, Johnson PCD, Govella NJ, Lee Y, et al. Emergence of behavioural avoidance strategies of malaria vectors in areas of high LLIN coverage in Tanzania. *Sci Rep*. 2020;10:14527.
13. Sougoufara S, Ottih EC, Tripet F. The need for new vector control approaches targeting outdoor biting Anopheline malaria vector communities. *Parasit Vectors*. 2020;13:295.
14. Sougoufara S, Doucouré S, Sembéne PMB, Harry M, Sokhna C. Challenges for malaria vector control in sub-Saharan Africa: Resistance and behavioral adaptations in *Anopheles* populations. *J Vector Borne Dis*. 2017;54:4–15.
15. Saddler A, Kreppel KS, Chitnis N, Smith TA, Denz A, Moore JD, et al. The development and evaluation of a self-marking unit to estimate malaria vector survival and dispersal distance. *Malar J*.

2019;18:441.

16. Epopa PS, Millogo AA, Collins CM, North A, Tripet F, Benedict MQ, et al. The use of sequential mark-release-recapture experiments to estimate population size, survival and dispersal of male mosquitoes of the *Anopheles gambiae* complex in Bana, a west African humid savannah village. *Parasit Vectors*. 2017;10:376.
17. Service MW. Mosquito ecology: Field sampling methods, Second edition. Mosquito ecology: Field sampling methods, Second edition. 1993.
18. Guerra CA, Reiner RC, Perkins TA, Lindsay SW, Midega JT, Brady OJ, et al. A global assembly of adult female mosquito mark-release-recapture data to inform the control of mosquito-borne pathogens. *Parasit Vectors*. 2014;7:276.
19. Vavassori L, Saddler A, Müller P. Active dispersal of *Aedes albopictus*: A mark-release-recapture study using self-marking units. *Parasit Vectors*. 2019;12:583.
20. Russell RC, Webb CE, Williams CR, Ritchie SA. Mark-release-recapture study to measure dispersal of the mosquito *Aedes aegypti* in Cairns, Queensland, Australia. *Med Vet Entomol*. 2005; 9:451–7.
21. Niebylski ML, Meek CL. A self-marking device for emergent adult mosquitoes. *J Am Mosq Control Assoc*. 1989;5:86-90.
22. Ciota AT, Drummond CL, Ruby MA, Drobnack J, Ebel GD, Kramer LD. Dispersal of *Culex* mosquitoes (Diptera: Culicidae) from a wastewater treatment facility. *J Med Entomol*. 2012;49:35-42.
23. Midega JT, Mbogo CM, Mwambi H, Wilson MD, Ojwang G, Mwangangi JM, et al. Estimating dispersal and survival of *Anopheles gambiae* and *Anopheles funestus* along the Kenyan coast by using mark-release-recapture methods. *J Med Entomol*. 2007;44:923-9
24. Gatton ML, Chitnis N, Churcher T, Donnelly MJ, Ghani AC, Godfray HCJ, et al. The importance of mosquito behavioural adaptations to malaria control in Africa. *Evolution*. 2013;67:1218–30.
25. Sokhna C, Ndiath MO, Rogier C. The changes in mosquito vector behaviour and the emerging resistance to insecticides will challenge the decline of malaria. *Clin Microbiol Infect*. 2013.19:902–7.
26. Carrasco D, Lefèvre T, Moiroux N, Pennetier C, Chandre F, Cohuet A. Behavioural adaptations of mosquito vectors to insecticide control. *Curr Opin Insect Sci*. 2019;34:48–54.
27. Killeen GF, Marshall JM, Kiware SS, South AB, Tusting LS, Chaki PP, et al. Measuring, manipulating and exploiting behaviours of adult mosquitoes to optimise malaria vector control impact. *BMJ Glob Heal*. 2017;2:e000212.
28. Monroe A, Asamoah O, Lam Y, Koenker H, Psychas P, Lynch M, et al. Outdoor-sleeping and other night-time activities in northern Ghana: Implications for residual transmission and malaria prevention. *Malar J*. 2015;14:35.
29. Abuaku B, Ahorlu C, Psychas P, Ricks P, Oppong S, Mensah S, et al. Impact of indoor residual spraying on malaria parasitaemia in the Bunkpurugu-Yunyoo District in northern Ghana. *Parasit Vectors*. 2018;11:555.

30. McSweeney C, New M, Lizcano G, Lu X. The UNDP climate change country profiles. Bull Am Meteorol Soc. 2010;91:157–66.
31. Millar J, Psychas P, Abuaku B, Ahorlu C, Amratia P, Koram K, et al. Detecting local risk factors for residual malaria in northern Ghana using Bayesian model averaging. Malar J. 2018;17:343.
32. Gillies MT, Coetzee M. A Supplement to the Anophelinae of the South of the Sahara (Afrotropical Region). Publications of the South African Institute for Medical Research. 1987;55:1–143.
33. Koekemoer LL, Kamau L, Hunt RH, Coetzee M. A cocktail polymerase chain reaction assay to identify members of the *Anopheles funestus* (Diptera: Culicidae) group. Am J Trop Med Hyg. 2002;6:804.
34. Fanello C, Santolamazza F, Della Torre A. Simultaneous identification of species and molecular forms of the *Anopheles gambiae* complex by PCR-RFLP. Med Vet Entomol. 2002;46:1–64.
35. Reddy MR, Overgaard HJ, Abaga S, Reddy VP, Caccone A, Kiszewski AE, et al. Outdoor host seeking behaviour of *Anopheles gambiae* mosquitoes following initiation of malaria vector control on Bioko Island, Equatorial Guinea. Malar J. 2011;10:184.
36. Meyers JI, Pathikonda S, Popkin-Hall ZR, Medeiros MC, Fuseini G, Matias A, et al. Increasing outdoor host-seeking in *Anopheles gambiae* over 6 years of vector control on Bioko Island. Malar J. 2016;15:239.
37. Russell TL, Govella NJ, Azizi S, Drakeley CJ, Kachur SP, Killeen GF. Increased proportions of outdoor feeding among residual malaria vector populations following increased use of insecticide-treated nets in rural Tanzania. Malar J. 2011;10:80.
38. Killeen GF. Characterizing, controlling and eliminating residual malaria transmission. Malar J. 2014;13:330.
39. Coosemans M. Residual transmission of malaria: an old issue for new approaches BT-Anopheles mosquitoes – New insights into malaria vectors. 2013;671–4.
40. Shcherbacheva A, Haario H, Killeen G. Modeling host-seeking behavior of African malaria vector mosquitoes in the presence of long-lasting insecticidal nets. Math Biosci. 2017;295:36–47.
41. Killeen GF, Kiware SS, Okumu FO, Sinka ME, Moyes CL, Massey NC, et al. Going beyond personal protection against mosquito bites to eliminate malaria transmission: population suppression of malaria vectors that exploit both human and animal blood. BMJ Glob Heal. 2017;2:e000198.
42. Govella NJ, Chaki PP, Killeen GF. Entomological surveillance of behavioural resilience and resistance in residual malaria vector populations. Malar J. 2013;12:124.
43. Sougoufara S, Sokhna C, Diagne N, Doucouré S, Sembène PM, Harry M. The implementation of long-lasting insecticidal bed nets has differential effects on the genetic structure of the African malaria vectors in the *Anopheles gambiae* complex in Dielmo, Senegal. Malar J. 2017;16:337.
44. Meyers JI, Pathikonda S, Popkin-Hall ZR, Medeiros MC, Fuseini G, Matias A, et al. Increasing outdoor host-seeking in *Anopheles gambiae* over 6 years of vector control on Bioko Island. Malar J. 2016;15:239.

45. Bamou R, Mbakop LR, Kopya E, Ndo C, Awono-Ambene P, Tchuinkam T, Rono MK, Mwangangi J, Antonio-Nkondjio C. Changes in malaria vector bionomics and transmission patterns in the equatorial forest region of Cameroon between 2000 and 2017. *Parasit Vectors*. 2018;11:464.
46. Massebo F, Balkew M, Gebre-Michael T, Lindtjørn B. Zoophagic behaviour of Anopheline mosquitoes in southwest Ethiopia: opportunity for malaria vector control. *Parasit Vectors*. 2015;8:645.
47. Main BJ, Lee Y, Ferguson HM, Kreppel KS, Kihonda A, Govella NJ, Collier TC, Cornel AJ, Eskin E, Kang EY, Nieman CC, Weakley AM, Lanzaro GC. The Genetic Basis of Host Preference and Resting Behavior in the Major African Malaria Vector, *Anopheles arabiensis*. *PLoS Genet*. 2016;12:e1006303.
48. Bryan JH, Petrarca V, Di Deco MA, Coluzzi M. Adult behaviour of members of the *Anopheles gambiae* complex in the Gambia with special reference to *An. melas* and its chromosomal variants. *Parassitologia*. 1987;29:221–49.
49. Coluzzi M, Sabatini A, Petrarca V, Di Deco MA. Chromosomal differentiation and adaptation to human environments in the *Anopheles gambiae* complex. *Trans R Soc Trop Med Hyg*. 1979;73:483–97.
50. Cho SH, Lee HW, Shin EH, Lee H II, Lee WG, Kim CH, et al. A mark-release-recapture experiment with *Anopheles sinensis* in the northern part of Gyeonggi-do, Korea. *Korean J Parasitol*. 2002;40:139–48.
51. Clements AN, Paterson GD. The Analysis of Mortality and Survival Rates in Wild Populations of Mosquitoes. *J Appl Ecol*. 1981;18:373–99.
52. Styer LM, Carey JR, Wang JL, Scott TW. Mosquitoes do senesce: Departure from the paradigm of constant mortality. *Am J Trop Med Hyg*. 2007;76:111–17.

Figures

Figure 1

Composition of released and recaptured marked test and control mosquitoes

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [GraphicalAbstract.png](#)