

De novo transcriptome in roots of switchgrass (*Panicum virgatum* L.) reveals gene expression dynamic and act network under alkaline salt stress

Pan Zhang

Northeast Agricultural University <https://orcid.org/0000-0002-2628-4852>

Tianqi Duo

Northeast Agricultural University

Fengdan Wang

Northeast Agricultural University

Xunzhong Zhang

Virginia Tech University

Zouzhan Yang

Northeast Agricultural University

Guofu Hu (✉ guofuh2003@163.com)

Northeast Agricultural University <https://orcid.org/0000-0002-3630-4503>

Research article

Keywords: *Panicum virgatum*, alkaline salt stress, ASTTI, transcriptome, WGCNA

Posted Date: December 2nd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-57429/v2>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Genomics on January 28th, 2021. See the published version at <https://doi.org/10.1186/s12864-021-07368-w>.

Abstract

Background: Soil salinization is a major limiting factor for crop cultivation. Switchgrass is a perennial rhizomatous bunchgrass that is considered an ideal plant for marginal lands, including sites with saline soil. Here, we investigated the physiological responses and transcriptome changes in the roots of Alamo (alkali-tolerant genotype) and AM-314/MS-155 (alkali-sensitive genotype) under alkaline salt stress.

Results: Alkaline salt stress significantly affected the membrane, osmotic adjustment and antioxidant systems in switchgrass roots, and the ASTTI values between Alamo and AM-314/MS-155 were divergent at different time points. A total of 108,319 unigenes were obtained after reassembly, including 73,636 unigenes in AM-314/MS-155 and 65,492 unigenes in Alamo. A total of 10,219 DEGs were identified, and the number of upregulated genes in Alamo was much greater than that in AM-314/MS-155 in both the early and late stages of alkaline salt stress. The DEGs in AM-314/MS-155 were mainly concentrated in the early stage, while Alamo showed greater advantages in the late stage. These DEGs were mainly enriched in plant-pathogen interactions, ubiquitin-mediated proteolysis and glycolysis/gluconeogenesis pathways. We characterized 1,480 TF genes into 64 TF families, and the most abundant TF family was the C2H2 family, followed by the bZIP and bHLH families. A total of 1,718 PKs were predicted, including CaMK, CDPK, MAPK and RLK. WGCNA revealed that the DEGs in the blue, brown, dark magenta and light steel blue 1 modules were associated with the physiological changes in roots of switchgrass under alkaline salt stress. The consistency between the qRT-PCR and RNA-Seq results confirmed the reliability of the RNA-seq sequencing data. A molecular regulatory network of the switchgrass response to alkaline salt stress was preliminarily constructed on the basis of transcriptional regulation and functional genes.

Conclusions: The alkaline salt tolerance of switchgrass may be achieved by the regulation of ion homeostasis, transport proteins, detoxification, heat shock proteins, dehydration and sugar metabolism. These findings provide a comprehensive analysis of gene expression dynamic and act network induced by alkaline salt stress in two switchgrass genotypes and contribute to the understanding of the alkaline salt tolerance mechanism of switchgrass and the improvement of switchgrass germplasm.

Background

Soil salinization, a major limiting factor for crop cultivation, seriously restricts global agricultural production and ecological environment construction [1, 2]. Soil salinization may cause neutral salt stress or alkaline salt stress, of which alkaline salt stress is mainly caused by NaHCO_3 and Na_2CO_3 . Alkaline damage, including osmotic, ionic and high pH damage, produces more direct toxic effects on plants than neutral salt stress [3, 4]. Osmotic stress and ionic stress lead to the accumulation of reactive oxygen species (ROS) and trigger enzymatic and nonenzymatic systems to mitigate ROS stress [5, 6]. Excess soluble ions reduce the water potential on the root surface, produce osmotic stress, and lead to water shortage in plants [7, 8]. Additionally the elevated pH suppressed photosynthesis, N metabolism, glycolysis, and growth of maize plants more markedly than NaCl

stress [9]. The seedling emergence and seedling establishment of *Phaseolus vulgaris* were also markedly suppressed under NaHCO₃ stress [10].

When plants are exposed to alkaline salt stress, root is the first tissue affected, and various biochemical and physiological mechanisms are stimulated to deal with the stress [11]. Alkaline stress greatly reduced root growth and root vigour, and induced a marked accumulation of superoxide anions (O₂⁻) and H₂O₂ in rice roots [12]. The increasing pH around the root system also leads to the deposition of metal ions, resulting in the reduction of inorganic anions and the hindrance of plant uptake of mineral nutrients [13]. The roots minimize the distribution of absorbed salt at the tissue and cellular levels to avoid accumulation of toxic concentrations in the cytosol of functional leaves [14]. Wheat roots exhibited greater growth performance in response to alkaline salt stress as a result of increased glutamine synthetase activity and soluble protein contents [15].

Plants respond to alkaline salt stress by regulating the expression of a variety of salt-responsive genes, such as osmoregulatory genes [16], transporters/antiporters[17], transcription factors (TFs) [18], and protein kinases (PKs) [19, 20]. Genes encoding Na⁺ transport proteins are involved in regulating Na⁺ transport under alkaline salt stress [21]. A quantitative trait locus (QTL) detected from NaCl and NaHCO₃ treated rice suggested that the genes controlling the transport of Na⁺, in the form of NaCl and NaHCO₃, may be different or induced in an uncoordinated manner by salt stress [22]. Vacuolar proton pump ATPase (V-H⁺-ATPase) is a multisubunit membrane protein complex that plays a major role in the activation of ion and nutrient transport. Overexpression of *ScVHA-B*, *ScVHA-C* and *ScVHA-H* improves tolerance to alkaline salt stress in transgenic alfalfa [23]. The HD-Zip TF family is one of the largest plant-specific TF superfamilies and plays important roles in the response to abiotic stresses. *Gshdz4*, an HD-Zip gene, showed a high response to alkaline stress in wild soybean treated with 50 mM NaHCO₃ [24]. *MsCBL4* plays an important role in alkaline salt stress tolerance via its influence on the regulation of calcium transport and accumulation [25]. Overexpression of *GsMSRB5a* and *GsCBRLK* in Arabidopsis enhanced alkaline stress tolerance, inhibiting ROS accumulation and modifying the expression of ROS signalling, biosynthesis and scavenging genes. With the continuous development of high-throughput sequencing, some genes and TFs have been identified in switchgrass [26, 27]. These interacting genes form multiple pathways, such as the salt overly sensitive (SOS) pathway, the calcium-dependent protein kinase (CDPK) pathway and the mitogen-activated protein kinase (MAPK) pathway [28].

Switchgrass (*Panicum virgatum* L.), a perennial warm-season C₄ rhizomatous bunchgrass, shows great prospects in terms of its strong adaptability, high water and nitrogen use efficiency, rapid growth and high productivity [29]. As an important ethanol bioenergy plant, switchgrass is considered one of the ideal plants to alleviate soil salinization and can still grow well in alkaline saline soil and marginal soil [30]. Anderson *et al.* showed that three lowland type switchgrass varieties (EG 2101, EG 1101 and EG 1102) had higher emergence rates and biomass yields under moderate to severe salt stress [29]. However, the molecular mechanism of switchgrass tolerance to alkaline salt stress is not well understood.

There may be much potential to be exploited in the responses and molecular mechanisms in switchgrass under alkaline salt stress, but the transcriptional regulation and functional genes play a key role in alkaline salt stress tolerance. Here, the gene expression dynamic and act network were revealed by assaying the physiological and transcriptome changes in roots of two switchgrass genotypes, Alamo (alkali-tolerant genotype) and AM-314/MS-155 (alkali-sensitive genotype), under alkaline salt stress for a period of 24 h. The pattern of association between differentially expressed genes (DEGs) and physiological changes in response to alkaline salt stress were explored by weighted gene coexpression network analysis (WGCNA). According to the expression patterns of signal transduction, kinases and binding proteins, TFs, and physiological responses, a regulatory network model based on functional genes was established to comprehensively illustrate the alkaline salt stress tolerance mechanism of switchgrass.

Results

Changes in alkaline salt tolerance trait index in two switchgrass genotypes

The physiological responses in roots of Alamo (alkali-tolerant genotype) and AM-314/MS-155 (alkali-sensitive genotype) under alkaline salt stress for 0, 3, 6, 12 and 24 h were evaluated by the alkaline salt tolerance trait index (ASTTI). Although the relative water content (RWC) decreased under alkaline salt stress, the ASTTI of RWC showed an upward trend (Fig. 1A). The ASTTI values of the relative electrical conductivity (REC) and malondialdehyde (MDA) content showed a downward trend with stress time (Fig. 1B and C). The REC and MDA contents in Alamo were higher than that in AM-314/MS-15 under all treated time points. The significantly differences of REC between Alamo and AM-314/MS-15 were observed at 6 and 24 h under alkaline stress. The REC in AM-314/MS-15 were 1.16 and 1.18 times higher than that in Alamo, respectively. The root MDA contents in AM-314/MS-155 were 1.88 times higher than that in Alamo at 24 h post alkaline salt stress.

The contents of free proline, soluble protein and soluble sugar were used as indicators of the osmotic adjustment system. The ASTTI value of proline and soluble protein contents first increased with increasing stress duration and then decreased, and the value of ASTTI of soluble sugar gradually declined with the increase of stressed time (Fig. 1 D-F). The proline content in Alamo were significantly higher than that in AM-314/MS-155 at 6 and 24 h post alkaline salt stress. Although the contents of soluble protein only showed significantly difference between Alamo and AM-314/MS-15 at 6 h, the contents of soluble sugar showed significantly differences under all treated time.

The contents of reduced glutathione (GSH), and the activities of superoxide dismutase (SOD), peroxidase (POD) and catalase (CAT) were used as indicators of the antioxidant system. Interestingly, the value of ASTTI of GSH steady raised under alkaline stress, while the value of ASTTI of SOD, POD and CAT increased firstly, and then decreased and re-increased with stress time. These four physiological traits changes significantly between Alamo and AM-314/MS-15 under alkaline salt stress, especially the activities of SOD and POD, they changed significantly at all treated time points.

In addition, we observed that there were significant physiological differences between the two varieties at different time points under alkali stress, especially at 6 and 24 h. Therefore, we chose these two time points for further transcriptome analysis.

De novo assembly and annotation of unigenes

Six cDNA libraries were constructed from total RNA extracted from E5 stage roots of Alamo and AM-314/MS-155 treated with alkaline salt stress for 0, 6 and 24 h. A total of 114.03 Gb of clean data were obtained, with 6.05 Gb of clean data for each sample and a Q30 base percentage of 93.21% or more (Additional file 1: Table S1). Since the whole-genome sequence of switchgrass is not currently publicly available, valid readings from the six libraries were combined for reassembly (Additional file 2: Table S2). A total of 108,319 unigenes were obtained after reassembly, including 73,636 unigenes in AM-314 and 65,492 unigenes in Alamo. The total length, N50 length and mean length were 106,429,710 nt, 1,751 nt, and 982.56 nt, respectively. Functional annotation of assembled sequences was primarily based on BLAST homology searches with the NCBI nonredundant protein (Nr), Evolutionary Genealogy of Genes: Nonsupervised Orthologous Groups (eggNOG), Gene Ontology (GO), Pfam, Swiss-Prot, Eukaryotic Orthologue Groups (KOG), Kyoto Encyclopedia of Genes and Genomes (KEGG), and Clusters of Orthologous Groups (COG) databases, and a total of 66,253 genes were annotated (Additional file 3: Table S3). Among these annotated sequences, 63,663 (96.09%) sequences had significant matches in the Nr database. On the basis of the homology among sequences of different species, 25,805 (40.56%) sequences were found against *Setaria italica*, and 5,124 (8.05%) sequences had clandestine hits with *Sorghum bicolor*, followed by *Zea mays* (4,732, 7.44%), *Oryza sativa* (1,590, 2.50%), *Phaeosphaeria nodorum* (1,564, 2.46%), *Verticillium dahliae* (950, 1.49%), *V. alfalfae* (839, 1.32%), *Phytophthora sojae* (680, 1.07%), *Togninia minima* (627, 0.99%) and *Hordeum vulgare* (584, 0.92%). Only 21,129 (33.21%) annotated sequences were similar to those of other plant species (Additional file 4: Figure S1A). The gene expression levels in response to alkaline salt stress were evaluated by converting the mapped read count for each gene into the expected number of fragments per kilobase of reproduction per million mapped reads (FPKM) to eliminate the effects of transcript length and sequencing differences on computational expression (Additional file 5: Table S4). The gene expression levels were not evenly distributed in the different stress environments in the boxplot diagram of the FPKM values (Additional file 4: Figure S1B). Then, the correlations of each biological sample were evaluated by Pearson correlation coefficients, and a value of r^2 close to 1 indicated a strong correlation between two replicate samples (Additional file 4: Figure S1C). Finally, all the unigenes were used for further identification of DEGs after the exclusion of abnormal samples.

Identification of DEGs associated with alkaline salt stress

To identify the DEGs of the two different genotypes under alkali stress, the expression patterns of DEGs were analysed by comparing 6-h and 24-h libraries with the control library for Alamo and AM-314/MS-155. A total of 10,219 DEGs whose expression was up- or downregulated between samples (fold change ≥ 2 and false discovery rate (FDR) < 0.01) at any pair of alkaline salt-treated points were

identified (Additional file 6: Table S5). In the early stage of alkaline salt stress (0-6 h), 4483 genes (58.02% of the DEGs) were upregulated in AM-314/MS-155, and 867 genes (61.36% of the DEGs) were upregulated in Alamo (Table 1). In the late stage of alkaline salt stress (24 h), 2942 genes (56.17% of the DEGs) were upregulated in AM-314/MS-155, while Alamo had 2732 upregulated genes (61.61% of the DEGs). The proportion of upregulated genes in the tolerant genotype (Alamo) was higher than that in the sensitive genotype (AM-314/MS-155) in both the early and late stages of stress. In addition, the DEGs of AM-314/MS-155 were mainly concentrated in the early stage, while Alamo showed greater advantages in the late stage.

To further analyse the effects of alkaline salt treatment in the two genotypes over time, the expression trends of DEGs were clustered into 16 modules (Fig. 2A). Then, the mainstream gene expression trends (6 groups) were screened over the duration of the stress (Fig. 2B). The first group contained 1,831 enriched genes, which mainly reflected the functional classification of genes expressed abundantly in AM-314/MS-155 but with little or no expression in Alamo (Fig. 2C). A large number of ribosomal proteins and a small number of cytochrome and energy-related genes were enriched in the ribosome and oxidative phosphorylation pathways and gradually downregulated with increasing stress time (Additional file 7: Table S6). The expression level of genes in group 2 (1,492 genes) decreased sharply in the early stage of alkaline salt stress in both switchgrass genotypes. KEGG analysis of the genes in group 2 revealed that most were involved in the ribosome, phenylalanine metabolism, plant hormone signal transduction, and ribosome biogenesis in eukaryotes pathways (Additional file 8: Table S7). The expression of genes in group 3 (1,261 genes) was completely opposite to that in group 1, mainly reflecting the functional classification of genes expressed abundantly in Alamo but with little or no expression in AM-314/MS-155. Genes in this group functioned mostly in the glycolysis/gluconeogenesis, plant-pathogen interaction, amino sugar and nucleotide sugar metabolism, and ubiquitin-mediated proteolysis categories (Additional file 9: Table S8). The expression of genes in group 4 (1,072 genes) increased strongly in the early stage of alkaline salt stress, and most of these genes functioned in starch and sucrose metabolism, valine, leucine and isoleucine degradation, plant hormone signal transduction, galactose metabolism, and fatty acid degradation (Additional file 10: Table S9). Although there was a fast decrease in gene expression levels in both group 5 (718 genes) and group 6 (736 genes), the decrease in the expression of genes in group 5 was mainly in AM-314/MS-155, while the decrease in the expression of genes in group 5 was mainly in Alamo. Genes involved in oxidative phosphorylation, protein processing in the endoplasmic reticulum, glycolysis/gluconeogenesis, and glycerophospholipid metabolism were enriched in group 5 (Additional file 11: Table S10), and genes in group 6 were enriched in ribosome, oxidative phosphorylation, glycolysis/gluconeogenesis, and DNA replication (Additional file 12: Table S11).

Identification of TFs and PKs

In this study, 1,480 TF genes were predicted using iTAK software (<http://itak.feilab.net/cgi-bin/itak/index.cgi>) and were classified into 64 TF families (Additional file 13: Table S12). The most abundant TF families were the C2H2 (153 genes), bZIP (108 genes), bHLH (93 genes), C3H (91 genes), MYB-related (82 genes), NAC (81 genes), WRKY (74 genes), GRAS (57 genes) and AP2/ERF-ERF (52

genes) families. The genes in the bHLH, C2H2, NAC and WRKY families were mainly upregulated under alkali stress, while the genes in the AP2/ERF-ERF, bZIP and MYB-related families showed a relatively balanced number of upregulated and downregulated members (Additional file 14: Figure S2). There were 32 TF genes that showed differential expression in both Alamo and AM-314/MS-155 under alkaline salt stress, including 7 TF genes that showed differential expression only at 6 h, 10 TF genes that showed differential expression only at 24 h, and 15 TF genes that showed differential expression at both 6 and 24 h (Additional file 15: Table S13). Interestingly, the TF genes only differentially expressed at 6 h were all downregulated, and most of them belonged to the MYB-related family. Nine of ten TF genes differentially expressed only at 24 h were unregulated, and most of them belonged to the NAC family. There were 27 upregulated TF genes and 16 downregulated TF genes specifically expressed in Alamo, and many of them were differentially expressed in the late stage of alkali stress (24 h) and belonged to the bZIP, MYB-related and bHLH families (Additional file 16: Table S14).

In addition, a total of 1,718 PKs, such as calcium/calmodulin-dependent protein kinase (CaMK), CDPK, MAPK and receptor-like kinase (RLK), were predicted in AM-314/MS-155 and Alamo under alkaline salt stress for 6 and 24 h (Additional file 17: Table S15). The CBL-interacting protein kinases (CIPKs), which play a role as Ca^{2+} sensors, were upregulated and highly expressed in Alamo under alkaline salt stress for 24 h. RLKs accounted for the largest proportion among many protein kinase families, and the genes belonging to the RLK family maintained a high expression level at 6 h and were then downregulated at 24 h.

Weighted gene coexpression network analysis (WGCNA) of DEGs in response to alkaline salt stress

To further study the patterns of association between the DEGs responding to alkaline salt stress and physiological indicators across the two genotypes, WGCNA was performed to explore the gene modules for synergistic expression. After the filtering of low-expression genes ($\text{FPKM} < 5$), the remaining 7,485 genes were classified into 20 different modules (combined modules) (Fig. 3A).

The blue module showed the highest positive correlation with the REC ($R=0.82$, $P=3e-5$) (Fig. 3B). This module was characterized by a significant upregulation of DEGs in the alkali-sensitive genotype AM-314/MS-155 at 6 and 24 h (Fig. 3C). Genes in this module were significantly enriched in the peroxisome, amino sugar and nucleotide sugar metabolism, starch and sucrose metabolism and plant-pathogen interaction pathways by KEGG (Additional file 18: Figure S3A). The brown module showed a positive correlation with soluble protein ($R=0.64$) and the RWC ($R=0.49$). This module represented the downregulated genes in Alamo under alkaline salt stress. The functions of these genes were mainly concentrated in the plant hormone signal transduction, phenylpropanoid biosynthesis and biosynthesis of amino acid pathways (Additional file 18: Figure S3B). The dark magenta module showed a high positive correlation with MDA, GSH and SOD, with R values of 0.70, 0.68 and 0.63, respectively. This module showed the features of significantly upregulated genes in both Alamo and AM-314/MS-155 under alkaline salt stress for 6 h. Genes in this module were mainly enriched in the amino sugar and nucleotide sugar metabolism, starch and sucrose metabolism, and other functional pathways (Additional file 18:

Figure S3C). The light steel blue 1 module was positively correlated with proline, soluble sugar and CAT, with R values of 0.80, 0.71 and 0.64, respectively (Fig. 4-B). The module mainly showed genes in the alkali-tolerant Alamo that were significantly upregulated under alkaline salt stress for 6 h. These genes were mainly enriched in the functions of phenylalanine metabolism and phenylpropanoid biosynthesis (Additional file 18: Figure S3D).

Verification of RNA-Seq sequencing data by qRT-PCR analysis

The DEGs associated with alkaline salt stress were selected for qRT-PCR assays to verify the accuracy of the RNA-Seq sequencing data. Ten genes were selected randomly from the DEGs that were coexpressed in all sequencing samples. The expression patterns of these ten genes were the same as those of the RNA-Seq assay (Fig. 4A). The expression levels indicated by the qRT-PCR results were strongly correlated with the differential gene expression levels identified by mRNA-seq according to the Pearson correlation coefficients, which were 0.8934 and 0.9144 under alkaline salt stress for 6 and 24 h, respectively (Fig. 4B), and demonstrated the reliability of the RNA-seq data.

Discussion

Comparison of physiological differences between different switchgrass genotypes

Due to the combined effects of osmotic, ionic and secondary stresses, plant growth is severely affected by alkaline salt stress [7]. Alkaline salt stress causes a large amount of soluble salts, such as Na^+ , to flow into plant cells, reduces the water potential on the root surface, and reduces the water absorption of plants [8]. Cell membrane permeability has been used as an effective selection criterion for alkaline salt resistance in plants [31]. With the prolongation of alkali stress, the cell membrane is destroyed, resulting in increased membrane permeability and severe lipid peroxidation, and alkali-tolerant genotypes show an obvious advantage [32]. In our study, the ASTTI value of the root RWC showed an upward trend, while the ASTTI values of the REC and MDA content showed a downward trend. Additionally, the root REC and MDA content in AM-314/MS-155 were significantly higher than those in Alamo at 24 h post alkaline salt stress. These results indicated that Alamo was more alkaline salt tolerant than AM-314/MS-155, and the degree of damage in Alamo was gradually reduced compared with that of AM-314/MS-155.

The accumulation of osmotic adjustment substances, such as proline, soluble protein and soluble sugar, is an important factor in sustaining plant growth under alkaline salt conditions [33]. The relative alkali-tolerant varieties resist alkaline salt stress by enhancing compatible solutes, including proline, thus maintaining cell turgor and water potential to achieve better growth and development [34]. In our study, the accumulation of soluble proteins was relatively lagging, which indicated that proline and soluble sugar could be used as important osmotic regulators in switchgrass under alkali treatment.

Antioxidant enzymes are the most important components of the ROS scavenging system. SOD can catalyse the formation of hydrogen peroxide and oxygen by superoxide radicals, and POD and CAT further scavenge H_2O_2 [35]. In this paper, the SOD activity of the alkali-tolerant Alamo increased rapidly at

3 h. The expression patterns of POD and SOD induced by alkali stress were similar in switchgrass. The nonenzymatic substance GSH in the antioxidant system also has the effect of scavenging ROS, and the glutathione transferase (GST) family is also a very important component of the metabolic detoxification enzyme system [36]. The GSH content of the sensitive genotype AM-314/MS-155 was significantly higher than that of the alkali-tolerant genotype Alamo in the early stage of alkaline salt stress. We found that the glutathione S-transferase genes in sensitive cultivars were significantly upregulated in the early stage of alkaline salt stress, and this may have been controlled by the regulation of the related genes, leading to changes in physiological indicators. This might be due to the consumption of GST activity, which decreased the GSH content of Alamo. These results revealed the physiological responses of switchgrass to complex alkaline salt stress, including the initial injury accumulation phase (0-6 h) and subsequent gradual recovery of infiltration and ion homeostasis phases (6-24 h).

Gene expression in response to alkaline salt stress in two switchgrass genotypes

In this study, a total of 10,219 DEGs were identified in two switchgrass genotypes under alkaline salt stress. The proportion of upregulated genes in Alamo was higher than that in AM-314/MS-155, and the DEGs in AM-314/MS-155 were mainly concentrated in the early stage, suggesting quick adjustments in dealing with the damage caused by alkali-salt stress in the short term. All DEGs clustered into 16 modules, and the mainstream gene trends were grouped into six groups (Fig. 2). Thirteen genes were enriched in the glycolysis/gluconeogenesis pathway, and five of them were hardly expressed in Alamo. Glycolysis/gluconeogenesis was considered to be the key pathway in the process of root development because the component exchanges with other pathways, such as arachidonic acid metabolism, glycan structure degradation, limonene and pinene degradation, and N-glycan degradation, were strongly dependent on its existence [37]. These results suggested that the sensitive genotype was not resistant to alkali stress in the late stage. Some disease resistance proteins were enriched in the plant-pathogen interaction pathway; they all contained the NB-ARC domain, which plays an important role in plant hypersensitivity, and hypersensitivity is a prerequisite for the initiation of defence responses [38, 39].

Identification of alkali-tolerant TFs in two switchgrass genotypes

In this study, bHLH, bZIP, C2H2 and WRKY were significantly enriched at 6 h in AM-314/MS-155, while bZIP, MYB-related and NAC were significantly enriched at 24 h in Alamo. The specific upregulated TFs associated with high alkali tolerance in Alamo were AP2-ERF, bZIP, bHLH and MYB-related. AP2-ERF transcription factors are involved in many physiological processes, including plant growth [40], abiotic stress responses [41], and ethylene and abscisic acid signalling pathways [42]. It was reported that *Gm-ERF3* isolated from soybean could improve the salt tolerance of transgenic tomato [43]. Overexpression of *PvERF001* isolated from the ERF genes of switchgrass could also increase biomass yield and sugar release efficiency in transgenic lines [44]. In this paper, we observed that AP2-ERF family TFs were still significantly upregulated at 24 h and might be used as candidate TFs to improve alkali tolerance in switchgrass. Overexpression of bZIP and bHLH TFs can help plants positively respond to salt stress[45]. [46]. Rong *et al* showed that *OsMYBc* in rice can regulate *OsHKT1* to deal with Na⁺ damage

caused by salt stress and prevent leaf accumulation of Na^+ toxicity [47]. In our study, the bZIP and MYB TFs continuously responded positively to alkaline salt stress at 6 h and 24 h. The bHLH TFs were specifically upregulated at 6 and 24 h in Alamo, indicating the alkaline salt tolerance of Alamo and these TFs could be considered target genes involved in the alkali stress response.

Identification of functional genes and potential regulatory response mechanisms of alkali tolerance in switchgrass under alkaline salt stress

Under alkaline salt stress, the genes encoded transporters responsible for reconstituting osmotic and ion-balanced proteins under alkali stress, such as sodium/hydrogen exchange protein (NHX), alkaline and neutral invertase (CINV2), and heat shock proteins (HSPs) including HSP20, HSP70, Hsp90 and DanJ, were differentially expressed in the two switchgrass genotypes. The ABC transporter and H^+ transport ATPase genes were significantly upregulated at 6 and 24 h in the two genotypes and showed a downward trend in Alamo in the late stage. At the same time, osmotic and ionic stresses can also lead to an increase in the concentration of free Ca^{2+} in the cytosol, which results in downstream gene phosphorylation and protein cascade [48]. The CDPK and MAPK pathways are important signalling pathways in plant responses to salt stress [49]. The MAPK cascade is an important regulator of antioxidant defence [50]. CAMS/CMLS, CIPKs and CDPKs, which function as Ca^{2+} sensors, are involved in many growth, development and stress-induced signal transduction pathways [51, 52]. Overexpression of *OsCIPK1* and *OsCIPK9* not only occurs in response to salt stress but also to ABA, drought and cold stress [53]. In this study, MAPK6 was significantly upregulated at 6 and 24 h in the two genotypes. CIPKs were mainly upregulated at 6 and 24 h in Alamo, while expression in AM-314/MS-155 remained low. These results suggest that the CDPK and MAPK signalling pathways may also mediate the response of switchgrass to alkali stress.

Compared with the two genotypes, many more DEGs were involved in plant hormone signal transduction in AM-314/MS-155 than in Alamo at 6 h, while more DEGs were identified in Alamo at 24 h. Alamo showed differential expression in the ABA signalling pathway at two different time points, while AM-314/MS-155 showed no significantly enriched DEGs. ABA plays a very important role in plant salt-tolerance signal transduction pathways and participates in the physiological and biochemical processes of abiotic stress in plants[54]. In our study, the HVA22 protein, a unique ABA stress-induced protein, was mainly upregulated in Alamo at 24 h and in AM-414/MS-155 at 6 h under alkaline salt stress. The HVA22 protein is induced by environmental stresses such as drought and high salt, which could inhibit GA-mediated programmed cell death of cereal aleurone cells and improve plant resistance to stress [55].

ROS also act as second messengers [56, 57]. Detoxification proteins involved in ROS scavenging were highly expressed at 6 and 24 h. Some DEGs were expressed more in AM/314/MS-155 than in Alamo, but most of the favourable upregulated genes were specifically expressed in the alkali-tolerant genotype Alamo. LEA proteins have high hydrophilicity and can supplement cells with sufficient water during stress and lessen the damage caused by osmotic stress [58]. Dehydrin, a second type of LEA protein, can significantly increase the resistance of plants to high salt and osmotic stress [59]. In this paper, the LEA

protein was downregulated in AM-314/MS-155 but upregulated in Alamo. The dehydrin protein showed specific upregulation only in Alamo, which fully reflected the positive response of Alamo to alkali stress.

The alkaline/neutral sucrase genes were also upregulated at 6 and 24 h under alkaline salt stress, and the number and expression of genes upregulated in Alamo were higher than those in AM-314/MS-155. Alkaline/neutral sucrase is a class of genes unique to plants and photosynthetic bacteria. The total alkaline/neutral invertase activity in wheat leaves increased after osmotic stress or low temperature stress, resulting in increased expression of related genes [60].

These genes are involved in antioxidant systems and transport functions and can aid in plant adaptation to alkaline saline environments by regulating ion homeostasis, transport proteins, detoxification, heat shock proteins, dehydration and sugar metabolism. The interaction of these genes constituted a complex regulatory network (Fig. 5). We hypothesized that switchgrass plants respond to alkaline salt stress through ion and osmotic stress signals. These signals were first sensed by receptors present on the plant cell membrane and then involved in signal transduction through ion channels and carrier proteins. The receptors in the roots may recognize salt signals, causing an instantaneous increase in ROS, leading to the opening of some plant hormone signalling pathways (including those of auxins, abscisic acid, cytokinins, gibberellins, ethylene, jasmonic acid and salicylic acid), biosynthesis of ubiquinone and other terpenoid-quinones, phenylpropanoid metabolism and biosynthetic pathways.

Conclusions

Taken together, we investigated the alkaline salt stress tolerance mechanism of switchgrass by analysing gene expression dynamic and act network in two switchgrass genotypes (AM-314/MS-155 and Alamo). A total of 108,319 unigenes were obtained after reassembly, and 10,219 DEGs were identified in the two switchgrass genotypes under alkaline salt stress. The DEGs in AM-314/MS-155 were mainly concentrated in the early stage, while the DEGs in Alamo showed greater advantages in the late stage. We characterized 1,480 TF genes and 1,718 PKs, and the most abundant TF family was the C2H2 family. The WGCNA results revealed that the DEGs in the blue, brown, dark magenta and light steel blue 1 modules were associated with the physiological changes in roots of switchgrass under alkaline salt stress. The alkaline salt tolerance of switchgrass may be achieved by the regulation of ion homeostasis, transport proteins, detoxification, heat shock proteins, dehydration and sugar metabolism.

Methods

Plant materials and growth conditions

Seeds of two lowland switchgrass (*Panicum virgatum* L.) genotypes (the alkaline salt tolerant genotype 'Alamo' and sensitive genotype 'AM-314/MS-155') were used to conduct this research. These two switchgrass lines were originally obtained from the United States Department of Agriculture Germplasm Center, and the accession number was 'PI 422006' for Alamo and 'PI 421999' for AM-314/MS-155. The

alkaline salt tolerance of these two lines were evaluated by Hu et al [61]. Hand-selected seeds of Alamo and AM-314/MS-155 were surface-sterilized in 5% sodium hypochlorite for 3 h and rinsed 4 to 5 times with sterile distilled water. Then, the seeds were germinated in culture dishes with wet filter papers in a dark growth chamber at 25 °C. Five-day-old seedlings were transported to plastic pots (17.8 cm upper diameter, 10 cm bottom diameter and 23.8 cm deep, with 8 drainage holes at the bottom) with the bottom 1/4 filled with coarse sand and the top 3/4 of the pots filled with fine sand in a greenhouse. Seedlings were kept in the greenhouse for 12 weeks (daytime temperature was 35 ± 1 °C, nighttime temperature was 25 ± 1 °C, 16 h of light and relative humidity of 70-80% RH) and fertilized with half-strength Hoagland's nutrient solution every two days.

Alkaline salt stress treatments

Alkaline salt stress treatments were conducted when the seedlings reached the E5 developmental stage [62]. The alkaline salt solution, which was made of Na₂CO₃ and NaHCO₃ (1:9, v/v), with Na⁺ at 150 mM and pH 9.5, was prepared with half-strength Hoagland's nutrient solution [61]. Roots of the two genotypes were harvested at 0, 3, 6, 12, and 24 h post alkaline salt stress. The experiment was repeated three times with three biological replicates. For each biological replicate, some of the samples were used to determine the relative water content (RWC) and relative electric conductivity (REC) directly, and the rest were frozen in liquid nitrogen immediately and stored at -80 °C for physiological and transcriptome analyses.

ASTTI evaluation by physiological assays

Physiological traits including the RWC, REC, the contents of MDA, free proline, soluble protein, soluble sugar, and reduced GSH, and the activities of SOD, POD and CAT in roots of two switchgrass genotypes at E5 developmental stage were assayed three times with three biological replicates [63-66]. The ASTTI, which was calculated based on the formula: ASTTI = (value of a trait under salt stress condition)/(value of a trait under controlled condition), was used to estimate the alkaline salt tolerance of the two switchgrass genotypes [61, 67].

RNA extraction, library preparation and next-generation sequencing (NGS) analysis

The roots of Alamo and AM-314/MS-155 at E5 developmental stage under alkaline salt stress for 6 and 24 h were chosen for transcriptome analysis. Total RNA was extracted with TRIzol reagent (Invitrogen, USA) according to the manufacturer's protocol, and genomic DNA was removed via digestion with DNase I (TaKaRa, Japan). The experiment was repeated three times with three biological replicates. The RNA sample concentration and quality were determined using a Nanophotometer P330 spectrophotometer (IMPLEN, Germany). A total amount of 3 µg RNA per sample was used as input material for the RNA sample preparations. Sequencing libraries were generated using the NEBNext®Ultra™ RNA Library Prep Kit for Illumina® (NEB, USA) following the manufacturer's recommendations, and index codes were added to attribute sequences to each sample.

The clustering of the index-coded samples was performed on a cBot Cluster Generation System using TruSeq PE Cluster Kit v3-cBot-HS (Illumina) according to the manufacturer's instructions. After cluster generation, the library preparations were sequenced on an Illumina HiSeq 2000 platform, and paired-end reads were generated. Clean reads were obtained by removing reads containing adapters, reads containing poly-N and low-quality reads from the raw data. All downstream analyses were based on clean data with high quality. Transcriptome assembly was accomplished using Trinity Software with min_kmer_cov set to 2 by default and all other parameters set to the default selections [68]. The clean reads of each sample were sequence aligned with the assembled unigene library, and the transcriptome sequencing library was evaluated for quality. Gene function was annotated by using BLAST software based on the NR, Pfam, KOG, COG, eggNOG, Swiss-Prot, KEGG and GO databases [69].

Identification of DEGs, putative TFs and PKs

The paired reads were compared with the unigene library using Bowtie [70] and RNA-seq by expectation maximization (RSEM) [71], and the expression abundance of a unigene was expressed using the FPKM value [72]. Differential expression analysis under the two conditions was performed using the DESeq R package (1.10.1). A false discovery rate (FDR) <0.01 and fold change (FC) ≥ 2 were used as screening criteria, and samples with a strong correlation between samples were selected for subsequent functional expression annotation and enrichment analysis of the DEGs. GO enrichment analysis of the DEGs was implemented by the topGO R package-based Kolmogorov–Smirnov test. KOBAS software was subsequently used to test the statistical enrichment of DEGs in the KEGG pathways [73]. TFs and PKs were predicted using iTAK software (<http://itak.feilab.net/cgi-bin/itak/index.cgi>).

Coexpression network analysis with WGCNA

Coexpression networks were constructed via the WGCNA package in R from all the DEGs [74]. Modules were obtained by the automatic network construction function using blockwise modules with the default settings. The eigengene value was calculated for each module and used to test the association with each physiological index. The total connectivity and intramodular connectivity (function soft connectivity), kME (for modular membership), and kME *p* values were calculated for the DEGs [75].

Validation of NGS data by quantitative real-time PCR

Total RNA was extracted from roots with TRIzol (Invitrogen, USA) from the same group of samples that were used in transcriptome analysis, and cDNA synthesis was performed with SuperMix for qPCR (+gDNA wiper) (Vazyme, China) according to the manufacturer's instructions. Primers used in this study (Additional field 19: Table S16) were designed using primer-BLAST (https://www.ncbi.nlm.nih.gov/tools/primerblast/index.cgi?LINK_LOC=BlastHome). The expression of CYP-5 was used as an internal control [76]. qPCR was performed using ChamQTM Universal SYBR qPCR Master Mix (Vazyme, China) on a LightCycler 480 II58 device (Roche, Switzerland) according to the manufacturers' protocol. The experiment was repeated three times with three biological replicates.

Relative gene expression levels were evaluated according to the $2^{\Delta\Delta CT}$ method [77]. Each gene was tested in triplicate with three biological replicates.

Abbreviations

TF: transcription factor; PK: protein kinases; QTL: quantitative trait locus; V-H+-ATPase: vacuolar proton pump ATPase; ROS: reactive oxygen species; SOS: salt overly sensitive; CDPK: calcium-dependent protein kinase; MAPK: mitogen-activated protein kinase; CaMK: calcium/calmodulin-dependent protein kinase; RLK: receptor-like kinase; DEG: differentially expressed gene; WGCNA: weighted gene coexpression network analysis; ASTTI: alkaline salt tolerance trait index; RWC: relative water content; REC: relative electric conductivity; MDA: malondialdehyde; GSH: reduced glutathione; SOD: superoxide dismutase; POD: peroxidase; CAT: catalase; Nr: NCBI nonredundant protein; eggNOG: Evolutionary Genealogy of Genes: Nonsupervised Orthologous Groups; GO: Gene Ontology; KOG: Eukaryotic Orthologue Groups; KEGG: Kyoto Encyclopedia of Genes and Genomes; COG: Clusters of Orthologous Groups; FPKM: fragmental per kilobase of reproduction per million mapped reads; GST: glutathione transferase; AREB: ABA responsive element binding protein; ABF: ABRE binding factors; HSP: heat shock protein; NGS: next-generation sequencing; RSEM: RNA-seq by expectation maximization.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and material

All relevant supplementary data are provided within this manuscript as Additional files. All the sequencing data generated in this study have been deposited in the Sequence Read Archive database under accession number PRJNA540186. Address is as follows: <http://www.ncbi.nlm.nih.gov/sra>.

Competing interests

The authors declare that they have no competing interests.

Funding

This research was funded by The Natural Science Fund of Heilongjiang Province (LC2016009) and U.S. Department of Energy and U.S. Department of Agriculture's Plant Feedstocks Genomics for Bioenergy Program (DE-SC0008338). Each of the funding bodies granted the funds based on a research

proposal. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Authors' contributions

PZ, GH and XZ designed the research. PZ and TD wrote the paper with contributions and discussion from all of the coauthors. PZ, TD, FW and ZY conducted the research. All authors have read and approved the manuscript.

Acknowledgements

We thank the Biomarker Corporation (Beijing, China) for RNA-Seq sequencing and raw data analysis.

References

1. Deinlein U, Stephan AB, Horie T, Luo W, Xu GH, Schroeder JI: **Plant salt-tolerance mechanisms.** *Trends Plant Sci* 2014, **19**(6):371-379.
2. Zhu J-K: **Plant salt tolerance.** *Trends Plant Sci* 2001, **6**(2):66-71.
3. Shi DC, Wang DL: **Effects of various salt-alkaline mixed stresses on Aneurolepidium chinense (Trin.) Kitag.** *Plant Soil* 2005, **271**(1-2):15-26.
4. Yang CW, Chong JN, Li CY, Kim CM, Shi DC, Wang DL: **Osmotic adjustment and ion balance traits of an alkali resistant halophyte Kochia sieversiana during adaptation to salt and alkali conditions.** *Plant and Soil* 2007, **294**(1-2):263-276.
5. Miller G, Suzuki N, Ciftciyilmaz S, Mittler R: **Reactive oxygen species homeostasis and signalling during drought and salinity stresses.** *Plant Cell & Environment* 2010, **33**(4):453-467.
6. Mittler R, Vanderauwera S, Gollery M, Van BF: **Reactive oxygen gene network of plants.** *Trends in Plant Science* 2004, **9**(10):490-498.
7. Yang YQ, Guo Y: **Elucidating the molecular mechanisms mediating plant salt-stress responses.** *New Phytol* 2018, **217**(2):523-539.
8. Hasegawa PM, Bressan RA, Zhu JK, Bohnert HJ: **Plant cellular and molecular responses to high salinity.** *Annu Rev Plant Physiol Plant Mol Biol* 2000, **51**(51):463-499.
9. Guo R, Shi LX, Yan CR, Zhong XL, Gu FX, Liu Q, Xia X, Li HR: **Ionomic and metabolic responses to neutral salt or alkaline salt stresses in maize (*Zea mays* L.) seedlings.** *BMC Plant Biol* 2017, **17**.
10. Yu S, Yu LH, Hou YL, Zhang YF, Guo W, Xue YW: **Contrasting Effects of NaCl and NaHCO₃ Stresses on Seed Germination, Seedling Growth, Photosynthesis, and Osmoregulators of the Common Bean (*Phaseolus vulgaris* L.).** *Agronomy-Basel* 2019, **9**(8):19.
11. Parida AK, Das AB: **Salt tolerance and salinity effects on plants: a review.** *Ecotoxicol Environ Saf* 2005, **60**(3):324-349.
12. Zhang H, Liu XL, Zhang RX, Yuan HY, Wang MM, Yang HY, Ma HY, Liu D, Jiang CJ, Liang ZW: **Root Damage under Alkaline Stress Is Associated with Reactive Oxygen Species Accumulation in Rice**

(*Oryza sativa* L.). *Front Plant Sci* 2017, **8**.

13. Capula-Rodríguez R, Valdez-Aguilar LA, Cartmill DL, Cartmill AD, Alia-Tejacal I: **Supplementary Calcium and Potassium Improve the Response of Tomato (*Solanum lycopersicum* L.) to Simultaneous Alkalinity, Salinity, and Boron Stress.** *Communications in Soil Science & Plant Analysis* 2016, **47**(4):505-511.
14. Goyal E, Amit SK, Singh RS, Mahato AK, Chand S, Kanika K: **Transcriptome profiling of the salt-stress response in *Triticum aestivum* cv. Kharchia Local.** *Scientific Reports* 2016, **6**:27752.
15. Zhang KH, Tang JR, Wang Y, Kang HY, Zeng J: **The tolerance to saline-alkaline stress was dependent on the roots in wheat.** *Physiol Mol Biol Plants* 2020, **26**(5):947-954.
16. Damien J, Cotsaftis, Olivier, Berthomieu, Pierre, Bohnert, Hans, Davenport, Romola J: **Nomenclature for HKT transporters, key determinants of plant salinity tolerance.** *Trends in Plant Science* 2006, **11**(8):372-374.
17. Huang Y, Guan C, Liu Y, Chen B, Yuan S, Cui X, Zhang Y, Yang F: **Enhanced Growth Performance and Salinity Tolerance in Transgenic Switchgrass via Overexpressing Vacuolar Na⁺ (K⁺)/H⁺ Antiporter Gene (PvNHX1).** *Frontiers in Plant Science* 2017, **8**:458.
18. Hong Y, Zhang H, Huang L, Li D, Song F: **Overexpression of a Stress-Responsive NAC Transcription Factor Gene ONAC022 Improves Drought and Salt Tolerance in Rice.** *Front Plant Sci* 2016, **7**(e0116646):4.
19. Quan R, Lin H, Mendoza I, Zhang Y, Cao W, Yang Y, Shang M, Chen S, Pardo JM, Guo Y: **SCABP8/CBL10, a putative calcium sensor, interacts with the protein kinase SOS2 to protect Arabidopsis shoots from salt stress.** *Plant Cell* 2007, **19**(4):1415-1431.
20. Mehlmer N, Wurzinger B, Stael S, Hofmannrodrigues D, Csaszar E, Pfister B, Bayer R, Teige M: **The Ca²⁺-dependent protein kinase CPK3 is required for MAPK-independent salt-stress acclimation in Arabidopsis.** *Plant Journal* 2010, **63**(3):484-498.
21. Chuamnakthong S, Nampei M, Ueda A: **Characterization of Na⁺ exclusion mechanism in rice under saline-alkaline stress conditions.** *Plant Sci* 2019, **287**.
22. Luo X, Deng HD, Wang P, Zhang XY, Li CJ, Li CJ, Tan JG, Wu GL, Wang YN, Cheng Q *et al*: **Genetic analysis of germinating ability under alkaline and neutral salt stress by a high-density bin genetic map in rice.** *Euphytica* 2020, **216**(7).
23. Wang FW, Wang C, Sun Y, Wang N, Li XW, Dong YY, Yao N, Liu XM, Chen H, Chen XF *et al*: **Overexpression of vacuolar proton pump ATPase (V-H⁺-ATPase) subunits B, C and H confers tolerance to salt and saline-alkali stresses in transgenic alfalfa (*Medicago sativa* L.).** *Journal of Integrative Agriculture* 2016, **15**(10):2279-2289.
24. Cao L, Yu Y, DuanMu HZ, Chen C, Duan XB, Zhu PH, Chen RR, Li Q, Zhu YM, Ding XD: **A novel Glycine soja homeodomain-leucine zipper (HD-Zip) I gene, Gshdz4, positively regulates bicarbonate tolerance and responds to osmotic stress in Arabidopsis.** *BMC Plant Biol* 2016, **16**.
25. An Y, Yang X-X, Zhang L, Zhang J, Du B, Yao L, Li X-t, Guo C: **Alfalfa MsCBL4 enhances calcium metabolism but not sodium transport in transgenic tobacco under salt and saline-alkali stress.** *Plant*

26. Huang Y, Cui X, Cen H, Wang K, Zhang Y: **Transcriptomic analysis reveals vacuolar Na⁺ (K⁺)/H⁺ antiporter gene contributing to growth, development, and defense in switchgrass (*Panicum virgatum* L.).** *Bmc Plant Biology* 2018, **18**(1):57.
27. Shen H, He X, Poovaiah CR, Wuddineh WA, Ma J, Mann DG, Wang H, Jackson L, Tang Y, Jr SC: **Functional characterization of the switchgrass (*Panicum virgatum*) R2R3-MYB transcription factor PvMYB4 for improvement of lignocellulosic feedstocks.** *New Phytol* 2012, **193**(1):121-136.
28. Tuteja N: **Mechanisms of high salinity tolerance in plants.** *Methods Enzymol* 2007, **428**:419-438.
29. Anderson EK, Voigt TB, Kim S, Lee DK: **Determining effects of sodicity and salinity on switchgrass and prairie cordgrass germination and plant growth.** *Industrial Crops and Products* 2015, **64**:79-87.
30. Nageswara-Rao M, Soneji JR, Kwit C, Stewart CN: **Advances in biotechnology and genomics of switchgrass.** *Biotechnol Biofuels* 2013, **6**:15.
31. Farooq S, Azam F: **The use of cell membrane stability (CMS) technique to screen for salt tolerant wheat varieties.** *Journal of Plant Physiology* 2006, **163**(6):629-637.
32. Golldack D, Li C, Mohan H, Probst N: **Tolerance to drought and salt stress in plants: Unraveling the signaling networks.** *Front Plant Sci* 2014, **5**(151):151.
33. Garcia AB, Engler J, Iyer S, Gerats T, Van MM, Caplan AB: **Effects of Osmoprotectants upon NaCl Stress in Rice.** *Plant Physiology* 1997, **115**(1):159-169.
34. Ali Q, Athar HUR, Ashraf M: **Modulation of growth, photosynthetic capacity and water relations in salt stressed wheat plants by exogenously applied 24-epibrassinolide.** *Plant Growth Regulation* 2008, **56**(2):107-116.
35. Neto ADDA, Prisco JT, Enéas-Filho J, Abreu CECD, Gomes-Filho E: **Effect of salt stress on antioxidative enzymes and lipid peroxidation in leaves and roots of salt-tolerant and salt-sensitive maize genotypes.** *Environmental & Experimental Botany* 2006, **56**(1):87-94.
36. Cheng MC, Ko K, Chang WL, Kuo WC, Chen GH, Lin TP: **Increased glutathione contributes to stress tolerance and global translational changes in Arabidopsis.** *Plant Journal* 2015, **83**(5):926-939.
37. Yang J, An D, Zhang P: **Expression Profiling of Cassava Storage Roots Reveals an Active Process of Glycolysis/Gluconeogenesis.** *Journal of Integrative Plant Biology* 2011, **53**(3):193-211.
38. Peyraud R, Dubiella U, Barbacci A, Genin S, Raffaele S, Roby D: **Advances on plant-pathogen interactions from molecular toward systems biology perspectives.** *Plant J* 2017, **90**(4):720-737.
39. Chandra S, Kazmi AZ, Ahmed Z, Roychowdhury G, Kumari V, Kumar M, Mukhopadhyay K: **Genome-wide identification and characterization of NB-ARC resistant genes in wheat (*Triticum aestivum* L.) and their expression during leaf rust infection.** *Plant Cell Reports* 2017, **36**(7):1097-1112.
40. Lee SY, Hwang EY, Seok HY, Tarte VN, Jeong MS, Jang SB, Moon YH: **Arabidopsis AtERF71/HRE2 functions as transcriptional activator via cis-acting GCC box or DRE/CRT element and is involved in root development through regulation of root cell expansion.** *Plant Cell Reports* 2015, **34**(2):223-231.

41. Dong L, Cheng Y, Wu J, Cheng Q, Li W, Fan S, Jiang L, Xu Z, Kong F, Zhang D: **Overexpression of GmERF5, a new member of the soybean EAR motif-containing ERF transcription factor, enhances resistance to Phytophthora sojae in soybean.** *Journal of Experimental Botany* 2015, **66**(9):2635-2647.
42. Kim YH, Jeong JC, Park S, Lee HS, Kwak SS: **Molecular characterization of two ethylene response factor genes in sweetpotato that respond to stress and activate the expression of defense genes in tobacco leaves.** *Journal of Plant Physiology* 2012, **169**(11):1112-1120.
43. Zhang G, Ming C, Li L, Xu Z, Chen X, Guo J, Ma Y: **Overexpression of the soybean GmERF3 gene, an AP2/ERF type transcription factor for increased tolerances to salt, drought, and diseases in transgenic tobacco.** *Journal of Experimental Botany* 2009, **60**(13):3781-3796.
44. Wuddineh WA, Mitra M, Turner GB, Sykes RW, Decker SR, Davis MF, C Neal S: **Identification and Molecular Characterization of the Switchgrass AP2/ERF Transcription Factor Superfamily, and Overexpression of PvERF001 for Improvement of Biomass Characteristics for Biofuel.** *Frontiers in Bioengineering & Biotechnology* 2015, **3**(2):226-227.
45. Wang C, Lu G, Hao Y, Guo H, Guo Y, Zhao J, Cheng H: **ABP9, a maize bZIP transcription factor, enhances tolerance to salt and drought in transgenic cotton.** *Planta* 2017, **246**(6):1-17.
46. Zhou J, Li F, Wang JL, Ma Y, Chong K, Xu YY: **Basic helix-loop-helix transcription factor from wild rice (OrbHLH2) improves tolerance to salt- and osmotic stress in Arabidopsis.** *Journal of Plant Physiology* 2009, **166**(12):1296-1306.
47. Rong W, Wen J, Longyun X, Yakang J, Like S, Wenhua Z: **The Rice High-Affinity Potassium Transporter1;1 Is Involved in Salt Tolerance and Regulated by an MYB-Type Transcription Factor.** *Plant Physiology* 2015, **168**(3):1076-1090.
48. Ruiz MCM, Jung HJ, Webb AAR: **Circadian gating of dark-induced increases in chloroplast- and cytosolic-free calcium in *Arabidopsis*.** *New Phytol* 2020, **225**(5):1993-2005.
49. Xiong L, Schumaker KS, Zhu JK: **Cell Signaling during Cold, Drought, and Salt Stress.** *Plant Cell* 2002, **14**(Suppl):S165.
50. Takáč T, O Š, Vadovič P, Pechan T, Košútová P, Ovečka M, Husičková A, Komis G, J Š: **Proteomic and biochemical analyses show a functional network of proteins involved in antioxidant defense of the *Arabidopsis anp2anp3* double mutant.** *Journal of Proteome Research* 2014, **13**(12):5347-5361.
51. McCormack E, Tsai YC, Braam J: **Handling calcium signaling: *Arabidopsis* CaMs and CMLs.** *Trends in Plant Science* 2005, **10**(8):383-389.
52. Weinl S, Kudla J: **The CBL-CIPK Ca(2+)-decoding signaling network: function and perspectives.** *New Phytol* 2009, **184**(3):517-528.
53. Xiang Y, Huang Y, Xiong L: **Characterization of Stress-Responsive CIPK Genes in Rice for Stress Tolerance Improvement.** *Plant Physiology* 2007, **144**(3):1416-1428.
54. Wang P, Zhao Y, Li Z, Hsu CC, Liu X, Fu L, Hou YJ, Du Y, Xie S, Zhang C *et al*: **Reciprocal Regulation of the TOR Kinase and ABA Receptor Balances Plant Growth and Stress Response.** *Mol Cell* 2018, **69**(1):100-112.

55. Woei-Jiun G, Tuan-Hua H, Thun-Hua DH: **An abscisic acid-induced protein, HVA22, inhibits gibberellin-mediated programmed cell death in cereal aleurone cells.** *Plant Physiology* 2008, **147**(4):1710-1722.
56. Raja V, Majeed U, Kang H, Andrabi KI, John R: **Abiotic stress: Interplay between ROS, hormones and MAPKs.** *Environmental and Experimental Botany* 2017, **137**:142-157.
57. Marcec MJ, Gilroy S, Poovaiah BW, Tanaka K: **Mutual interplay of Ca²⁺ and ROS signaling in plant immune response.** *Plant Sci* 2019, **283**:343-354.
58. Imai R, Chang L, Ohta A, Bray EA, Takagi M: **A lea-class gene of tomato confers salt and freezing tolerance when expressed in *Saccharomyces cerevisiae*.** *Gene* 1996, **170**(2):243-248.
59. Brini F, Hanin M, V, Irar S, Pages M, Masmoudi K: **Functional characterization of DHN-5, a dehydrin showing a differential phosphorylation pattern in two Tunisian durum wheat (*Triticum durum Desf.*) varieties with marked differences in salt and drought tolerance.** *Plant Science* 2007, **172**(1):20-28.
60. Vargas WA, Pontis HG, Salerno GL: **Differential expression of alkaline and neutral invertases in response to environmental stresses: characterization of an alkaline isoform as a stress-response enzyme in wheat leaves.** *Planta* 2007, **226**(6):1535-1545.
61. Hu G, Liu Y, Zhang X, Yao F, Huang Y, Ervin EH, Zhao B: **Physiological Evaluation of Alkali-Salt Tolerance of Thirty Switchgrass (*Panicum virgatum*) Lines.** *PLoS One* 2015, **10**(7):e0125305.
62. Hardin CF, Fu C, Hisano H, Xiao X, Shen H, Stewart CN, Parrott W, Dixon RA, Wang ZY: **Standardization of Switchgrass Sample Collection for Cell Wall and Biomass Trait Analysis.** *BioEnergy Research* 2013, **6**(2):755-762.
63. Dreywood R: **Qualitative test for carbohydrate material.** *Ind Eng Chem, Anal Ed* 1946, **18**(8):499-499.
64. Maehly AC: **The assay of catalases and peroxidases.** In: *Methods Biochem Anal.* John Wiley & Sons, Inc.; 2006: 357-424.
65. Bates LS, Waldren RP, Teare ID: **Rapid determination of free proline for water-stress studies.** *Plant Soil* 1973, **39**(1):205-207.
66. Marcum KB, Anderson SJ, Engelke MC: **Salt Gland Ion Secretion: A Salinity Tolerance Mechanism among Five Zoysiagrass Species.** *Crop Science* 1998, **38**(3):806-810.
67. Tavakkoli E, Fatehi F, Rengasamy P, McDonald GK: **A comparison of hydroponic and soil-based screening methods to identify salt tolerance in the field in barley.** *J Exp Bot* 2012, **63**(10):3853-3867.
68. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L, Raychowdhury R, Zeng Q: **Full-length transcriptome assembly from RNA-Seq data without a reference genome.** *Nature Biotechnology* 2011, **29**(7):644.
69. Altschul SF, Madden TL, Schäffer AA: **Jinghui Zhang, Zheng Zhang, Webb Miller, and David J. Lipman (1997)," Gapped BLAST and PSI-BLAST: a new generation of protein database search.** 1997.
70. Langmead B, Trapnell C, Pop M, Salzberg SL: **Ultrafast and memory-efficient alignment of short DNA sequences to the human genome.** *Genome Biology* 2009, **10**(3):R25.
71. Li B, Dewey CN: **RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome.** *Bmc Bioinformatics* 2011, **12**(1):323-323.

72. Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, Baren MJV, Salzberg SL, Wold BJ, Pachter L: **Transcript assembly and abundance estimation from RNA-Seq reveals thousands of new transcripts and switching among isoforms.** *Nature Biotechnology* 2010, **28**(5):511.
73. Xie C, Mao X, Huang J, Ding Y, Wu J, Dong S, Kong L, Gao G, Li CY, Wei L: **KOBAS 2.0: a web server for annotation and identification of enriched pathways and diseases.** *Nucleic Acids Res* 2011, **39**(Web Server issue):W316-322.
74. Langfelder P, Horvath S: **WGCNA: an R package for weighted correlation network analysis.** *BMC Bioinformatics* 2008, **9**(1):559.
75. Du J, Wang S, He C, Zhou B, Ruan YL, Shou H: **Identification of regulatory networks and hub genes controlling soybean seed set and size using RNA sequencing analysis.** *Journal of Experimental Botany* 2017, **68**(8):1955-1972.
76. Gimeno J, Eattock N, Deynze AV, Blumwald E: **Selection and validation of reference genes for gene expression analysis in switchgrass (*Panicum virgatum*) using quantitative real-time RT-PCR.** *PLoS One* 2014, **9**(3):e91474.
77. Livak KJ, Schmittgen TD: **Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the 2 $\Delta\Delta C_T$ Method.** *METHODS* 2001, **25**:402-408.

Tables

Table 1. Comparison of DEGs between two genotypes at different time points

Items	AM-314/MS-155 (sensitive)			Alamo (tolerant)		
	0h vs 6h	0h vs 24h	6h vs 24h	0h vs 6h	0h vs 24h	6h vs 24h
Total	77,578	83,255	34,658	78,349	77,799	33,172
DEGs	7727	5238	1150	1413	4434	1255
Up-regulated	4483	2942	361	867	2732	675
Down-regulated	3244	2296	789	546	1702	580

Figures




Figure 1

ASTTI evaluation of two lowland switchgrass genotypes (Alamo and AM-314/MS-155) by the physiological changes in roots under alkaline salt stress for 0, 3, 6, 12 and 24 h. A: Relative water content (RWC). B: Relative electric conductivity (REC). C: Malondialdehyde (MDA) content. D: Proline content. E: Soluble protein content. F: Soluble sugar content. G: Glutathione content (GSH). H: Superoxide dismutase (SOD) activity. I: Peroxidase (POD) activity. J: Catalase (CAT) activity. The left Y-axis represented value of physiological traitors, and the right Y-axis represented value of ASTTI. The assays were repeated three times with three biological replicates. The data, shown as means \pm SEs, were subjected to student's t test to determine significant differences. "*" means $p<0.05$, "**" means $p<0.01$.




Figure 1

ASTTI evaluation of two lowland switchgrass genotypes (Alamo and AM-314/MS-155) by the physiological changes in roots under alkaline salt stress for 0, 3, 6, 12 and 24 h. A: Relative water content (RWC). B: Relative electric conductivity (REC). C: Malondialdehyde (MDA) content. D: Proline content. E: Soluble protein content. F: Soluble sugar content. G: Glutathione content (GSH). H: Superoxide dismutase (SOD) activity. I: Peroxidase (POD) activity. J: Catalase (CAT) activity. The left Y-axis represented value of physiological traitors, and the right Y-axis represented value of ASTTI. The assays were repeated three times with three biological replicates. The data, shown as means \pm SEs, were subjected to student's t test to determine significant differences. "*" means $p<0.05$, **" means $p<0.01$.




Figure 2

Clustering model analysis of DEGs. A: the 16 clustering modules, in which S0, S6 and S24 mean the DEGs of alkali-sensitive genotype AM-314/MS-155 under alkali-salt stress for 0, 6 and 24 h, while T0, T6 and T24 mean the DEGs of alkali-tolerant genotype Alamo under alkali-salt stress for 0, 6 and 24 h. B: clustering of the mainstream genes. C: the changing trend of the mainstream genes of two genotypes with stress time.




Figure 2

Clustering model analysis of DEGs. A: the 16 clustering modules, in which S0, S6 and S24 mean the DEGs of alkali-sensitive genotype AM-314/MS-155 under alkali-salt stress for 0, 6 and 24 h, while T0, T6 and T24 mean the DEGs of alkali-tolerant genotype Alamo under alkali-salt stress for 0, 6 and 24 h. B: clustering of the mainstream genes. C: the changing trend of the mainstream genes of two genotypes with stress time.




Figure 3

Correlation analysis between modules and physiological traits by WGCNA (B), and eigengenes expression pattern of 20 modules of WGCNA clustering (B). Each row corresponded a clustering module, each column corresponded a physiological trait, the color of each grid at the row-column intersection indicated the correlation between the module and physiological data, red represented positive correlation, blue represented negative correlation, and the number in each grid represented the correlation coefficient and the P value. Twenty modules were marked with numeric value. 1: antiquewhite4; 2: bisque4; 3: blue; 4: brown; 5: coral2; 6: cyan; 7: darkmagenta; 8: darkorange; 9: darkseagreen4; 10: darkslateblue; 11: honeydew; 12: lightcyan; 13: lightsteelblue; 14: lightsteelblue1; 15: mediumorchid; 16: mediumpurple3; 17: midnightblue; 18: skyblue2; 19: white; 20: yellowgreen. In each module, red represented up-regulation and green represented down-regulation. The following corresponded to the degree of up-and-down adjustment of 18 samples in 6 groups (S0, S6, S24, T0, T6, T24).




Figure 3

Correlation analysis between modules and physiological traits by WGCNA (B), and eigengenes expression pattern of 20 modules of WGCNA clustering (B). Each row corresponded a clustering module, each column corresponded a physiological trait, the color of each grid at the row-column intersection indicated the correlation between the module and physiological data, red represented positive correlation, blue represented negative correlation, and the number in each grid represented the correlation coefficient and the P value. Twenty modules were marked with numeric value. 1: antiquewhite4; 2: bisque4; 3: blue; 4: brown; 5: coral2; 6: cyan; 7: darkmagenta; 8: darkorange; 9: darkseagreen4; 10: darkslateblue; 11: honeydew; 12: lightcyan; 13: lightsteelblue; 14: lightsteelblue1; 15: mediumorchid; 16: mediumpurple3; 17: midnightblue; 18: skyblue2; 19: white; 20: yellowgreen. In each module, red represented up-regulation and green represented down-regulation. The following corresponded to the degree of up-and-down adjustment of 18 samples in 6 groups (S0, S6, S24, T0, T6, T24).




Figure 4

Correlation analysis between modules and physiological traits by WGCNA (B), and eigengenes expression pattern of 20 modules of WGCNA clustering (B). Each row corresponded a clustering module, each column corresponded a physiological trait, the color of each grid at the row-column intersection indicated the correlation between the module and physiological data, red represented positive correlation, blue represented negative correlation, and the number in each grid represented the correlation coefficient and the P value. Twenty modules were marked with numeric value. 1: antiquewhite4; 2: bisque4; 3: blue; 4: brown; 5: coral2; 6: cyan; 7: darkmagenta; 8: darkorange; 9: darkseagreen4; 10: darkslateblue; 11: honeydew; 12: lightcyan; 13: lightsteelblue; 14: lightsteelblue1; 15: mediummorchid; 16: mediumpurple3; 17: midnightblue; 18: skyblue2; 19: white; 20: yellowgreen. In each module, red represented up-regulation and green represented down-regulation. The following corresponded to the degree of up-and-down adjustment of 18 samples in 6 groups (S0, S6, S24, T0, T6, T24).




Figure 4

Correlation analysis between modules and physiological traits by WGCNA (B), and eigengenes expression pattern of 20 modules of WGCNA clustering (B). Each row corresponded a clustering module, each column corresponded a physiological trait, the color of each grid at the row-column intersection indicated the correlation between the module and physiological data, red represented positive correlation, blue represented negative correlation, and the number in each grid represented the correlation coefficient and the P value. Twenty modules were marked with numeric value. 1: antiquewhite4; 2: bisque4; 3: blue; 4: brown; 5: coral2; 6: cyan; 7: darkmagenta; 8: darkorange; 9: darkseagreen4; 10: darkslateblue; 11: honeydew; 12: lightcyan; 13: lightsteelblue; 14: lightsteelblue1; 15: mediummorchid; 16: mediumpurple3; 17: midnightblue; 18: skyblue2; 19: white; 20: yellowgreen. In each module, red represented up-regulation and green represented down-regulation. The following corresponded to the degree of up-and-down adjustment of 18 samples in 6 groups (S0, S6, S24, T0, T6, T24).




Figure 5

Possible molecular mechanisms of switchgrass in response to alkali-salt stress. The red-up arrows mean these genes were up-expressed, the blue-down arrows mean these genes were down-expressed, and the "?" mean the express of these genes were conflicted, or they both up- or down-expressed under alkaline salt stress in roots of two switchgrass genotypes.




Figure 5

Possible molecular mechanisms of switchgrass in response to alkali-salt stress. The red-up arrows mean these genes were up-expressed, the blue-down arrows mean these genes were down-expressed, and the "?" mean the express of these genes were conflicted, or they both up- or down-expressed under alkaline salt stress in roots of two switchgrass genotypes.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [renamedff851.docx](#)
- [renamedff851.docx](#)
- [renamed9c698.tif](#)
- [renamed9c698.tif](#)
- [renameddd9ee6.xlsx](#)
- [renameddd9ee6.xlsx](#)
- [renamed55238.xlsx](#)
- [renamed55238.xlsx](#)
- [renamed3a24b.xlsx](#)
- [renamed3a24b.xlsx](#)
- [renamedf69f3.tif](#)
- [renamedf69f3.tif](#)
- [renamed71f81.xlsx](#)
- [renamed71f81.xlsx](#)
- [renamedac369.xlsx](#)
- [renamedac369.xlsx](#)
- [renamed34cb9.xlsx](#)
- [renamed34cb9.xlsx](#)
- [renameddd8048.xlsx](#)
- [renameddd8048.xlsx](#)
- [renamedf50c8.xlsx](#)
- [renamedf50c8.xlsx](#)
- [renamed23dea.xlsx](#)
- [renamed23dea.xlsx](#)
- [renamed4d0e9.xlsx](#)

- renamed4d0e9.xlsx
- renamedffd1a.xlsx
- renamedffd1a.xlsx
- renamed127e7.xlsx
- renamed127e7.xlsx
- renamede9b97.tif
- renamede9b97.tif
- renamed6989e.docx
- renamed6989e.docx
- renamedfdd13.docx
- renamedfdd13.docx
- renamed1236b.docx
- renamed1236b.docx