

# LncRNA-KCNQ1OT1: A Potential Target in Exosomes Derived From ADSCs for The Treatment of Osteoporosis

**Shanzheng Wang**

The Affiliated Zhongda Hospital of Southeast University

**Jun Jia**

The 904th Hospital, PLA

**Chang-hong Chen** (✉ [changhongchen0917@gmail.com](mailto:changhongchen0917@gmail.com))

Jiangyin Hospital

---

## Research

**Keywords:** Osteoporosis, Adipose derived mesenchymal stem cells, Exosomes, lncRNA-KCNQ1OT1, miR-141-5p

**Posted Date:** June 9th, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-576971/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

## Background

Osteoporosis is a worldwide medical and socioeconomic threat characterized by systemic impairment to bone strength and microstructure. Exosomes derived from adipose-derived stem cells (ADSCs-Exos) have been confirmed to play effective roles in the repair of various tissues and organs. This study aimed to investigate the role of ADSCs-Exos and a novel long non-coding RNA KCNQ10T1 (lnc-KCNQ10T1) played in osteoporosis as well as the underlying mechanism.

## Methods

MC3T3-E1 cells were treated with different doses of TNF- $\alpha$  (0, 1, 2.5, 5, 10 ng/ml) and then co-cultured with ADSCs-Exos or exosomes-derived from lnc-KCNQ10T1-modified ADSCs (KCNQ10T1-Exos). The expression of microRNA (miRNA)-141-5p (miR-141-5p) and lnc-KCNQ10T1 was evaluated by quantitative real-time polymerase chain reaction (qRT-PCR). The protein expression of cleaved-caspase-3, caspase-3 and Bax was determined by Western blot. Cell viability and apoptosis were assessed by cell counting kit 8 (CCK-8) and flow cytometry analysis, respectively. The binding sites between KCNQ10T1 and miR-141-5p were validated by dual-luciferase reporter assay.

## Results

Tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) dose dependently increased miR-141-5p expression, inhibited viability and promoted apoptosis of MC3T3-E1 cells. However, miR-141-5p silencing or co-culture with ADSCs-Exos attenuated these effects. In addition, KCNQ10T1-Exos could more significantly attenuate the induced cytotoxicity and apoptosis compared to ADSCs-Exos. Moreover, miR-141-5p was confirmed as the target of lnc-KCNQ10T1 by luciferase reporter assay.

## Conclusions

ADSCs-Exos attenuated cytotoxicity and apoptosis of TNF- $\alpha$ -induced MC3T3-E1 cells. KCNQ10T1-Exos had a more significant inhibitory effect compared to ADSCs-Exos by the function of sponging miR-141-5p, suggesting that KCNQ10T1-Exos could be promising agents in osteoporosis treatment.

## Background

Osteoporosis is a common skeletal disease characterized by structural disorders of bone mass caused by increased osteoclast activity and reduced osteoblast generation [1]. Clinical treatment strategies are based on the promotion of osteoblast proliferation and osteoblast apoptosis [2, 3]. In osteoporosis pathogenesis, tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) is a pro-inflammatory cytokine which has been revealed to

contribute to osteoporosis by regulating both osteoblasts and osteoclasts [3, 4]. TNF- $\alpha$  can suppress bone formation and inhibit osteoblast differentiation by blocking RUNX2 [5, 6]. Therefore, exploring new pathway to arrest TNF- $\alpha$ -induced cytotoxicity and apoptosis in osteoporosis is necessary.

As a subset of mesenchymal stem cells (MSCs), adipose-derived stem cells (ADSCs) can be easily obtained from adipose tissues and possess the potential of multi-differentiation and self-renewal. ADSCs are found in abundant quantities and can be easily obtained by minimally invasive surgeries [7]. These cells are included in clinical therapeutic strategies that fulfill the general criteria for cell-based regenerative therapies, making them the most adapted cell source for tissue regeneration [8]. Thus, ADSCs might be beneficial MSCs-based strategy for bone formation and structural remodeling in osteoporosis [9, 10].

Exosomes are 40–100 nm bilayer lipid vesicles that are secreted from various cell types into the extracellular space [11]. Exosomes have also been proved to play a significant role in the etiology of bone metabolic diseases, especially osteoporosis [12]. As a previous study shown, exosomes derived from ADSCs (ADSCs-Exos) also contain the same bioactive substances of ADSCs and might play a similar role to ADSCs [13]. Recent researches have demonstrated that ADSCs-Exos had motivating effects in the repair of a variety of tissues and organs [14–16]. Even though, the effects of ADSCs-Exos on cell cytotoxicity and apoptosis in osteoporosis as well as the potential mechanisms remain unclear.

Exosomes are important mediators between cells by transferring molecules, such as long none coding RNAs (lncRNA), microRNAs (miRNAs) and cytokines [17–20]. Evidence indicates that exosomes derived from mmu\_circ\_0000250-modified ADSCs are able to promote wound healing in diabetic mice [21]. Additionally, exosomes derived from miR-188-3p-modified ADSCs can protect Parkinson's disease[22]. Recently, a novel lncRNA, KCNQ1overlapping transcript 1 (KCNQ1OT1) was found to be able to promote the proliferation, metabolism and growth of tumor cells [23]. However, little is known about the role of KCNQ1OT1 in osteoporosis. The aim of this study was to to investigate the role of ADSCs-Exos and lnc-KCNQ1OT1 played in osteoporosis as well as the underlying mechanism.

## Methods

### Isolation and culture of ADSCs

The Institutional Animal Care and Use Committee of Southeast University approved the protocol for the use of animals in this study. Adipose tissues were collected from the inguinal fat pad from 2-3-week-old C57BL/6 mice (SLAC, Shanghai, China), and then rinsed in phosphate-buffered saline (PBS) and cut into 1×1mm pieces. The collected tissues were digested by collagenase type II (Sigma-Aldrich, USA) at 37 °C for 1 h. After digestion, tissues were centrifuged at room temperature (1000rpm, 5min) and the resultant cell pellet was re-suspended in Dulbecco's modified Eagle's medium (DMEM) at the cell density of 5×10<sup>6</sup>/ml. Cells were then cultured at 37°C under a 5% CO<sub>2</sub> atmosphere. The culture medium was replaced every 2-3 days and cells after 3 passages were used in the present study.

## **Induced differentiation potential of isolated ADSCs**

ADSCs were cultured in a 24-well plate at a density of  $4 \times 10^4$  cells/well with basic culture medium (DMEM-LG) containing 10% FBS, 100 U/ml penicillin and 100mg/ml streptomycin. They were subjected to induced differentiation by culturing them in osteogenic (Cyagen Biosciences, Rasmx-90021), adipogenic (Cyagen Biosciences, Rasmx-90031) and chondrogenic (Cyagen Biosciences, Rasmx-90041) medium, respectively. The outcomes were evaluated by Alizarin Red, Oil Red O and Alcian Blue staining, respectively.

## **Isolation and characterization of exosomes**

A Total Exosome Isolation kit (Invitrogen, 4478359, USA) was applied to isolate the total exosomes from the supernatant of ADSCs culture medium according to the manufacturer's protocol. The experimental procedures were conducted as described before [24]. Bicinchoninic acid assay (BCA) protein assay kit was used to measure the concentration of isolated exosomes. The protein levels of CD9, CD63, CD81 and Alix (representative markers of exosomes) were then detected.

## **Exosomes uptake assay**

ADSCs-Exos were labeled with PKH26 (Sigma Aldrich, USA) according to the manufacturer's protocol. Isolated exosomes were resuspended in diluent C (1ml). Then 6  $\mu$ l PKH26 was added into diluent C (1ml). The ADSCs-Exos and PKH26 solutions were mixed for 30 s and then centrifuged ( $120000 \times g$ , 2h, 4°C). Exosomes were resuspended in the complete culture medium and the PKH26-labeled ADSCs-Exos solution was added into MC3T3-E1 cells for incubation. After 24-48h of culture, MC3T3-E1 cells were fixed with 4% formaldehyde for 10 min. DAPI was used to stain the nuclei. Cells were finally observed under a confocal microscope.

## **MC3T3-E1 cells culture**

MC3T3-E1 cells were provided by the American Type Culture Collection (ATCC, USA) and cultured with  $\alpha$ -modified essential medium (Gibco, USA) at 37°C under a 5% CO<sub>2</sub> atmosphere. MC3T3-E1 cells were cultured in 6-well plates at the concentration of  $1 \times 10^6$  cells/well. After 24-48 h of culture, MC3T3-E1 cells were transfected with anti-miR-NC, anti-miR-141-5p, miR-141-5p or miR-NC, which were obtained from GenePharma (Shanghai, China). After transfection, MC3T3-E1 cells were treated with different doses of TNF- $\alpha$  (0, 1, 2.5, 5, 10ng/ml). After 24-48 h of culture, MC3T3-E1 cells were then co-cultured with ADSCs-Exos (25  $\mu$ g/ml, 50  $\mu$ g/ml, 100  $\mu$ g/ml) or KCNQ10T1-Exos (25  $\mu$ g/ml) for 24-48 h.

## **Cell Viability Assay**

The MC3T3-E1 viability was determined by the Cell Counting Kit-8 (CCK-8, Dojindo, Japan) assay. Briefly, cells were seeded into 96-well plates ( $2 \times 10^4$  cell/ml) and incubated for 24h. Then, 10  $\mu$ l CCK-8 solution

was added into each well for incubating 1-2 h (37 °C, 5% CO<sub>2</sub>). Subsequently, the optical density (OD) value was measured at 450 nm using spectrophotometer (Bio-Rad, USA).

### **Cell apoptosis assay**

MC3T3-E1 cell apoptosis was evaluated by Annexin V-FITC/PI Apoptosis Assay Kit (Keygen Biotech, China). Briefly, MC3T3-E1 cells were collected and washed with PBS. A total of 500 µl binding buffer was added to suspend cells. Firstly, 5 µl annexin V-FITC was added, and then 5 µl propidium iodide was added for incubation for 5-15min in the dark at room temperature. Cell apoptosis was analyzed by flow cytometry (Becton-Dickinson, FACS Calibur, USA).

### **Western blot analysis**

The total protein of MC3T3-E1 cell was extracted using RIPA lysis buffer (Beyotime, China) and quantified by BCA assay (Beyotime, China). Equal amounts of proteins (100µg) were separated via BeyoGel™ Plus PAGE (Beyotime, China) and then transferred to a PVDF membrane (Millipore, USA). After transferring, the membranes were blocked with 5% fat-free milk for 1h. The membranes were incubated with primary antibodies (Bax, ab32503, Caspase-3, ab32351, cleaved- Caspase-3, ab32042), which were purchased from Abcam (USA) and GAPDH, 10494-I-AP which was obtained from Proteintech (China) at 4°C overnight. The membranes were incubated with the second antibodies (Goat anti rabbit IgG HRP SE134, Goat anti mouse IgG HRP SE131, Solarbio, China) at 37°C for 1h. The ECL system (CLINX, China) was used for exposing protein bands. The intensity of the bands was analyzed using Image lab (version 3.0, Bio-Rad, USA).

### **Gene expression analysis using quantitative real-time polymerase chain reaction (qRT-PCR)**

Total RNA was extracted from cells using TRIzol reagent (Invitrogen, USA) according to the manufacturer's protocol. For detecting miR-141-5p expression, MicroRNA cDNA Synthesis Kit (Vazyme, China) was applied for reverse transcription of cDNA, followed by qRT-PCR analysis. U6 was employed as the loading control. The primers of miR-141-5p and U6 were purchased from GeneCopoeia (Guangzhou, China). Meanwhile, the expression of lncRNA-KCNQ10T1 was evaluated by Prime Script™ Master Mix (Takara, Japan) and GAPDH was employed as the loading control. The data was calculated using the 2<sup>-DDCt</sup> method. Primers used in this study were shown below:

lncRNA-KCNQ10T1, F: 5'-TTGGTAGGATTTTGTGAGG-3' and

R: 5'-CAACCTTCCCCTACTACC-3';

GAPDH, F: 5-TGGATTTGGACGCATTGGTC-3 and

R: 5-TTTGCACTGGTACGTGTTGAT-3.

### **Dual-luciferase reporter assay**

The wild-type (WT) sequence of lncRNA-KCNQ10T1 containing the miR-141-5p binding sites (KCNQ10T1-WT) and the mutant sequence (KCNQ10T1-MUT) were cloned into pMIR vectors (Promega, USA) respectively. MC3T3-E1 and HEK293 cells were co-transfected with miR-141-5p or miR-NC and KCNQ10T1-WT or KCNQ10T1-MUT and kept for 24 h. The luciferase activity was detected using Dual-Luciferase Reporter Assay System (Promega, USA) in the dark.

## Statistical Analysis

Statistical analysis was performed with SPSS 20.0 software (IBM, Armonk, NY, U.S.A.). All quantitative data were described as mean  $\pm$  SD. One-way ANOVA was used to analyze the statistical differences among three or more groups while unpaired Student's *t* test was applied to analyze the statistical differences between two groups.  $P < 0.05$  was considered statistically significant.

## Results

### **TNF- $\alpha$ increased miR-141-5p expression, suppressed the viability and promoted the apoptosis of MC3T3-E1 cells in a dose-dependent manner**

MC3T3-E1 cells were treated with increased concentrations of TNF- $\alpha$  (0, 1, 2.5, 5, 10 ng/ml). TNF- $\alpha$  dose-dependently increased miR-141-5p expression (Fig. 1a). Meanwhile, MC3T3-E1 cells treated with TNF- $\alpha$  showed reduced cell viability and promoted apoptosis in a dose-dependent manner (Fig. 1b and c). Furthermore, TNF- $\alpha$  increased the protein expression level of Bax and cleaved-Caspase-3 dose-dependently (Fig. 1d).

### **Knockdown of miR-141-5p reversed the effect of TNF- $\alpha$ on MC3T3-E1 cells.**

Since miR-141-5p was up-regulated in TNF- $\alpha$ -treated MC3T3-E1 cells, we knocked down miR-141-5p expression using anti-miR-141-5p to explore its role in TNF- $\alpha$ -induced cytotoxicity and apoptosis. As expected, anti-miR-141-5p reduced the expression of miR-141-5p more significantly in MC3T3-E1 cells compared to anti-miR-NC (Fig. 2a). CCK-8 and Annexin V-FITC/PI assays revealed that the knockdown of miR-141-5p partly reversed the inhibition of cell viability and the promotion of cell apoptosis induced by TNF- $\alpha$  when cells treated with PBS, TNF- $\alpha$ , TNF- $\alpha$ +anti-miR-NC, TNF- $\alpha$ +anti-miR-141-5p (1mg) or TNF- $\alpha$ +anti-miR-141-5p (2mg) respectively (Fig. 2b and c). In line with this result, the increase of Bax and cleaved-Caspase-3 expression induced by TNF- $\alpha$  was inhibited by downregulating miR-141-5p (Fig. 2d).

### **Characteristics of ADSCs and ADSCs-derived exosomes**

ADSCs isolated from C57/BL6 mice had a typical fibroblastic-like morphology (Fig. 3a). Alizarin Red, Oil Red O and Alcian Blue staining was positive after the induced osteogenic, adipogenic and chondrogenic differentiation of ADSCs (Fig. 3b). The immunoblotting showed that the ADSCs-Exos were positive for the exosomes' markers, including CD9, CD81, CD63 and Alix (Fig. 3c). Furthermore, MC3T3-E1 cells were co-cultured with PKH26-labeled ADSCs-Exos and the red fluorescence of PKH26 label was observed in MC3T3-E1 cells 24 h later (Fig.3d), indicating MC3T3-E1 cells could uptake ADSCs-Exos.

## **ADSCs-Exos attenuated the effect of TNF- $\alpha$ on MC3T3-E1 cells.**

To determine the effects of ADSCs-Exos, MC3T3-E1 cells were treated with TNF- $\alpha$  (5 ng/ml) and different doses of ADSCs-Exos (PBS, TNF- $\alpha$ , TNF- $\alpha$ +Exos (25mg), TNF- $\alpha$ +Exos (50mg), TNF- $\alpha$ +Exos (100mg)). ADSCs-Exos attenuated the up-regulation of miR-141-5p induced by TNF- $\alpha$  dose-dependently (Fig. 4a). Similarly, ADSCs-Exos reversed the inhibition of cell viability caused by TNF- $\alpha$  in a dose-dependent manner (Fig. 4b). The elevated protein expression of cleaved caspase-3 and Bax induced by TNF- $\alpha$  was suppressed after co-culture with ADSCs-Exos (Fig. 4c). In line with this, the results of flow cytometry indicated ADSCs-Exos dose-dependently blocked cell apoptosis induced by TNF- $\alpha$  (Fig. 4d).

## **KCNQ10T1-Exos inhibited TNF- $\alpha$ -induced cytotoxicity and apoptosis of MC3T3-E1 cells.**

ADSCs were transfected with LV-KCNQ10T1 or LV-NC for the determination of whether ADSCs transfected LV-KCNQ10T1 into secreted exosomes. At 24 h post LV-KCNQ10T1 transfection, the expression of KCNQ10T1 in ADSCs or exosomes derived from the ADSCs was elevated, but the expression of miR-141-5p was down-regulated compared with that in LV-NC treated group (Fig. 5a and b). In order to confirm whether ADSCs-Exos carrying LV-KCNQ10T1 could deliver KCNQ10T1 into MC3T3-E1 cells, MC3T3-E1 cells were co-cultured with LV-NC-Exos or LV-KCNQ10T1-Exos. KCNQ10T1 expression was up-regulated in MC3T3-E1 cells treated with LV-KCNQ10T1-Exos (Fig. 5c). Next, to explore whether KCNQ10T1-Exos could influence TNF- $\alpha$ -induced cytotoxicity and apoptosis, MC3T3-E1 cells were treated with TNF- $\alpha$  and then co-cultured with medium, ADSCs-Exos, LV-NC-Exos or LV-KCNQ10T1-Exos. The results of CCK-8 showed the co-culture of MC3T3-E1 cells with LV-KCNQ10T1-Exos mitigated the negative effect of TNF- $\alpha$  on cell viability; ADSCs-Exos exerted a weaker stimulative effect on cell viability compared to LV-KCNQ10T1-Exos (Fig. 5d). Flow cytometry analysis indicated that TNF- $\alpha$ -induced cell apoptosis was reversed when MC3T3-E1 cells were co-cultured with LV-KCNQ10T1-Exos; ADSCs-Exos exerted a weaker inhibitory effect on cell apoptosis compared to LV-KCNQ10T1-Exos (Fig. 5e). Consistent with that, the expression of Bax and cleaved caspase-3 in MC3T3-E1 cells was blocked after co-culture of LV-KCNQ10T1-Exos (Fig. 5f).

## **KCNQ10T1 could sponge miR-141-5p**

Given KCNQ10T1 sequence contains potential binding sites for miR-141-5p, we predicted KCNQ10T1 could sponge miR-141-5p (Fig. 6a). The dual-luciferase reporter assay was conducted to confirm the combination between them. As the result showed, overexpressed miR-141-5p weakened the luciferase activity in KCNQ10T1-WT group obviously, but the activity of luciferase reporters containing KCNQ10T1-MUT was not changed significantly in HEK293 and MC3T3-E1 cells (Fig. 6b and c). Compared with ADSC-Exos and LV-NC-Exos, LV-KCNQ10T1-Exos blocked the inhibitory effect of miR-141-5p on the luciferase activity of reporters containing KCNQ10T1-WT (Fig. 6d). Overexpressed KCNQ10T1 decreased the expression of miR-141-5p, yet the down-regulation of KCNQ10T1 increased that (Fig. 6e). Moreover, when MC3T3-E1 cells cultured with LV-KCNQ10T1-Exos, the expression of miR-141-5p was inhibited (Fig. 6f).

## **KCNQ10T1-Exos inhibited the effect of TNF- $\alpha$ in MC3T3-E1 cells by sponging miR-141-5p**

MC3T3-E1 cells were transfected with miR-141-5p or miR-NC to confirm whether KCNQ10T1-Exos attenuate TNF- $\alpha$ -induced cytotoxicity and apoptosis in MC3T3-E1 cells by acting as an miR-146a sponge. After the treatment of TNF- $\alpha$ , MC3T3-E1 cells were transfected with miR-141-5p or miR-NC and co-cultured with ADSCs-Exos, LV-NC-Exos or LV-KCNQ10T1-Exos. The up-regulation of miR-141-5p promoted the inhibitory effect of TNF- $\alpha$  on cell viability and co-culture with ADSCs-Exos or KCNQ10T1-Exos partly reversed this phenomenon; ADSCs-Exos exerted a weaker stimulative effect on cell viability compared to LV-KCNQ10T1-Exos (Fig. 7a). Similarly, the up-regulation of miR-141-5p enhanced cleaved caspase-3 and Bax expression. However, these effects were attenuated following co-culture with ADSC-Exos or LV-KCNQ10T1-Exos (Fig. 7b). In line with this, as flow cytometry showed overexpressed miR-141-5p promoted cell apoptosis but when co-cultured with ADSC-Exos or LV-KCNQ10T1-Exos, the cell apoptosis was inhibited; ADSCs-Exos exerted a weaker inhibitory effect on cell apoptosis compared to LV-KCNQ10T1-Exos (Fig. 7c).

## Discussion

With the deepening understanding of osteobiology, skeletal stem cells and osteoblasts are identified as significant target in the treatment of OP by promoting bone formation and remodeling [25]. Previous studies have proven that decreasing miR-10a-3p and raising CXCL12 can induce the osteogenic differentiation of bone marrow-derived MSCs (BMSCs) [26]. miR-26b could promote BMSCs osteogenesis through directly regulating GSK3 $\beta$  and activating Wnt pathway [27]. MSCs-derived exosomes are involved in multiple physiology and pathology processes, including osteogenesis, bone regeneration and osteoarthritis [28, 29]. BMSC-derived exosomes are able to alleviate radiation-induced bone loss via Wnt/ $\beta$ -catenin signaling pathway. In addition, BMSCs-derived exosomal MALAT1 can promote the progression of osteoblasts [30].

Similar to BMSCs, ADSCs also have the potential of multi-lineage differentiation. In 2001, Zuk et al [31] firstly isolated ADSCs from adipose tissue and found they could differentiate into adipocytes, osteoblasts and chondrocytes under different induction conditions. In addition to the commonness with BMSCs, ADSCs can be easily obtained with abundant sources. Depending on this, ADSCs might be a better choice in the treatment of osteoporosis compared to BMSCs. However, few studies have investigated the roles of ADSCs-Exos, especially in the treatment of osteoporosis. Thus, whether ADSCs-Exos could effectively protect MC3T3-E1 cells from the TNF- $\alpha$ -induced cytotoxicity and apoptosis was studied in the present study.

Our results indicated that TNF- $\alpha$  increased miR-141-5p expression, inhibited cell viability and promoted cell apoptosis in MC3T3-E1 cells. Consistent with this, the expression of cleaved caspase-3 and Bax was also elevated. Currently, ADSCs have been widely used in tissue regeneration and bioengineering. However, with the in-depth investigations, the application of ADSCs has the potential risk of iatrogenic infection, malignant transformation, immune rejection safety issues [32–34]. Compared to ADSCs, ADSCs-Exos have no risk of malignant transformation and can hardly cause the immune rejections [35, 36]. Therefore, ADSCs-Exos can be promising regenerative agents for osteoporosis. To make clear the

effects of ADSCs-Exos in osteoporosis, we attempted to culture TNF- $\alpha$ -treated MC3T3-E1 cells with ADSCs-Exos. Interestingly, ADSCs-Exos promoted cell viability and impeded apoptosis, suggesting that ADSCs-Exos can be promising candidates in the treatment of OP.

Although we've found ADSCs-Exos can be beneficial in the treatment of OP, the underlying mechanism has not been revealed. Exosomes derived from MSCs contain multiple lncRNAs, which can be transported and transferred to other cells to regulate biological functions through targeting downstream genes [37, 38]. In osteoarthritis, exosomal KLF3-AS1 originated from hMSCs boosted cartilage repair and chondrocyte proliferation [39]. KCNQ10T1, a lncRNA which is closely related to cell proliferation, migration and apoptosis, has been reported to be an oncogene in a variety of tumors [23]. Evidence has showed that KCNQ10T1 can promote cell proliferation and migration [40]. KCNQ10T1 was found to enhance glioma cell proliferation and invasion via regulating the miR-375/YAP pathway and thereby accelerated the progression of glioma [41]. Meanwhile, in colorectal cancer, KCNQ10T1 can induce cell apoptosis by sponging miR-329-3p [42]. Herein, KCNQ10T1 was targeted for further investigation in this study. We confirmed the presence of KCNQ10T1 in ADSCs-Exos, and further investigated the role of KCNQ10T1 in the treatment of osteoporosis. KCNQ10T1-Exos had a more significant inhibitory effect on TNF- $\alpha$ -induced cytotoxicity and apoptosis compared to ADSCs-Exos. Therefore, KCNQ10T1-Exos are expected to be superior candidates in osteoporosis treatment.

lncRNAs can function as miRNA sponges by binding miRNAs [43]. As previously reported, lncRNA-HOTAIR could induce the apoptosis of osteoblasts via modulating the expression of miR-138 [44]. In postmenopausal osteoporosis, lncRNA LOXL1-AS1 regulates osteogenic and adipocytic differentiation of BMSCs via sponging miR-196a-5p [45]. For further mechanistic investigations, we predicted KCNQ10T1 sequence contained miR-141-5p binding sites *via* bioinformatics analysis. As reported before, miR-141-5p promoted preeclampsia via regulating MAPK1/ERK2 signaling [46]. In chronic myeloid leukemia, microRNA-141-5p acts as a tumor suppressor by downregulating RAB32 [47]. However, the role of miR-141-5p played in osteoporosis has not been reported. In TNF- $\alpha$ -treated MC3T3-E1 cells, we found increased expression of miR-141-5p. Moreover, the knock-down of miR-141-5p promoted cell viability and inhibited cell apoptosis after the treatment of TNF- $\alpha$ . As dual-luciferase reporter assay showed, miR-141-5p was proved to be the target gene of KCNQ10T1. Furthermore, the rescue experiments revealed that when co-cultured with KCNQ10T1-Exos, the effects induced by miR-141-5p on cell viability and apoptosis in TNF- $\alpha$ -treated MC3T3-E1 cells were partly reversed, suggesting that KCNQ10T1-Exos functioned by sponging miR-141-5p.

## Conclusions

In conclusion, we demonstrated that ADSCs-Exos attenuated the effect of TNF- $\alpha$  on MC3T3-E1 cells. KCNQ10T1-Exos had a more significant inhibitory effect on TNF- $\alpha$ -induced cytotoxicity and apoptosis compared to ADSCs-Exos. KCNQ10T1 exerted its role by sponging miR-141-5p, suggesting that KCNQ10T1-Exos might provide potential precise target for patients with osteoporosis. Further explorations of the pleiotropic effect of KCNQ10T1 and the crosstalk between KCNQ10T1 and miR-141-

5p will provide new insights for developing new treatments to improve the therapeutic efficacy based on ADSCs-Exos.

## Declarations

### Ethics approval

The Institutional Animal Care and Use Committee of Southeast University approved the protocol for the use of animals in this study.

### Consent for publication

Not applicable.

### Availability of data and material

Please contact author for data requests.

### Authors' contributions

SZW performed the experiments, collected data, and drafted the manuscript. JJ participated in the experimental design and experimentation, and helped collect the data. CHC took part in the study design and revised the manuscript. All authors read and approved the final manuscript.

### Competing interests

The authors declare that they have no competing interests.

### Acknowledgments

The funding support from Wuxi Health Committee Research Grants for Top Talent Support Program (2020) and [Nanjing University of Chinese Medicine](#) Research Grant (XZR2020075).

### Funding

Already stated above.

## References

1. Johnston CB and Dagar M. Osteoporosis in Older Adults. *Med Clin North Am.* 2020; 104(5):873-884.
2. Karsenty G. The complexities of skeletal biology. *Nature.* 2003; 423(6937):316-318.
3. Rachner TD, Khosla S and Hofbauer LC. Osteoporosis: now and the future. *Lancet.* 2011; 377(9773):1276-1287.


4. Osta B, Benedetti G and Miossec P. Classical and Paradoxical Effects of TNF-alpha on Bone Homeostasis. *Front Immunol.* 2014; 5:48.
5. Armour KJ, Armour KE, van't Hof RJ, Reid DM, Wei XQ, Liew FY and Ralston SH. Activation of the inducible nitric oxide synthase pathway contributes to inflammation-induced osteoporosis by suppressing bone formation and causing osteoblast apoptosis. *Arthritis Rheum.* 2001; 44(12):2790-2796.
6. Gilbert L, He X, Farmer P, Rubin J, Drissi H, van Wijnen AJ, Lian JB, Stein GS and Nanes MS. Expression of the osteoblast differentiation factor RUNX2 (Cbfa1/AML3/Pebp2alpha A) is inhibited by tumor necrosis factor-alpha. *J Biol Chem.* 2002; 277(4):2695-2701.
7. Fahy N, Alini M and Stoddart MJ. Mechanical stimulation of mesenchymal stem cells: Implications for cartilage tissue engineering. *J Orthop Res.* 2018; 36(1):52-63.
8. Rada T, Reis RL and Gomes ME. Adipose tissue-derived stem cells and their application in bone and cartilage tissue engineering. *Tissue Eng Part B Rev.* 2009; 15(2):113-125.
9. Im GI. Bone marrow-derived stem/stromal cells and adipose tissue-derived stem/stromal cells: Their comparative efficacies and synergistic effects. *J Biomed Mater Res A.* 2017; 105(9):2640-2648.
10. An Y, Zhao J, Nie F, Wu Y, Xia Y and Li D. Parathyroid hormone (PTH) promotes ADSC osteogenesis by regulating SIK2 and Wnt4. *Biochemical and biophysical research communications.* 2019; 516(2):551-557.
11. Raposo G and Stoorvogel W. Extracellular vesicles: exosomes, microvesicles, and friends. *J Cell Biol.* 2013; 200(4):373-383.
12. Murphy C, Withrow J, Hunter M, Liu Y, Tang YL, Fulzele S and Hamrick MW. Emerging role of extracellular vesicles in musculoskeletal diseases. *Mol Aspects Med.* 2018; 60:123-128.
13. Qiu H, Liu S, Wu K, Zhao R, Cao L and Wang H. Prospective application of exosomes derived from adipose-derived stem cells in skin wound healing: A review. *J Cosmet Dermatol.* 2020; 19(3):574-581.
14. Lee M, Ban JJ, Yang S, Im W and Kim M. The exosome of adipose-derived stem cells reduces beta-amyloid pathology and apoptosis of neuronal cells derived from the transgenic mouse model of Alzheimer's disease. *Brain Res.* 2018; 1691:87-93.
15. Cho BS, Kim JO, Ha DH and Yi YW. Exosomes derived from human adipose tissue-derived mesenchymal stem cells alleviate atopic dermatitis. *Stem Cell Res Ther.* 2018; 9(1):187.
16. Fang Y, Zhang Y, Zhou J and Cao K. Adipose-derived mesenchymal stem cell exosomes: a novel pathway for tissues repair. *Cell Tissue Bank.* 2019; 20(2):153-161.
17. Kilchert C, Wittmann S and Vasiljeva L. The regulation and functions of the nuclear RNA exosome complex. *Nature reviews Molecular cell biology.* 2016; 17(4):227-239.
18. Zhang Y, Bi J, Huang J, Tang Y, Du S and Li P. Exosome: A Review of Its Classification, Isolation Techniques, Storage, Diagnostic and Targeted Therapy Applications. *Int J Nanomedicine.* 2020; 15:6917-6934.

19. Sluijter JP, Verhage V, Deddens JC, van den Akker F and Doevendans PA. Microvesicles and exosomes for intracardiac communication. *Cardiovasc Res.* 2014; 102(2):302-311.
20. Colombo M, Raposo G and Thery C. Biogenesis, secretion, and intercellular interactions of exosomes and other extracellular vesicles. *Annual review of cell and developmental biology.* 2014; 30:255-289.
21. Shi R, Jin Y, Hu W, Lian W, Cao C, Han S, Zhao S, Yuan H, Yang X, Shi J and Zhao H. Exosomes derived from mmu\_circ\_0000250-modified adipose-derived mesenchymal stem cells promote wound healing in diabetic mice by inducing miR-128-3p/SIRT1-mediated autophagy. *Am J Physiol Cell Physiol.* 2020; 318(5):C848-C856.
22. Li Q, Wang Z, Xing H, Wang Y and Guo Y. Exosomes derived from miR-188-3p-modified adipose-derived mesenchymal stem cells protect Parkinson's disease. *Mol Ther Nucleic Acids.* 2021; 23:1334-1344.
23. Wang CG, Liao Z, Xiao H, Liu H, Hu YH, Liao QD and Zhong D. LncRNA KCNQ10T1 promoted BMP2 expression to regulate osteogenic differentiation by sponging miRNA-214. *Exp Mol Pathol.* 2019; 107:77-84.
24. Santos JC, Lima NDS, Sarian LO, Matheu A, Ribeiro ML and Derchain SFM. Exosome-mediated breast cancer chemoresistance via miR-155 transfer. *Scientific reports.* 2018; 8(1):829.
25. Marie PJ and Kassem M. Osteoblasts in osteoporosis: past, emerging, and future anabolic targets. *Eur J Endocrinol.* 2011; 165(1):1-10.
26. Liu H, Yi X, Tu S, Cheng C and Luo J. Kaempferol promotes BMSC osteogenic differentiation and improves osteoporosis by downregulating miR-10a-3p and upregulating CXCL12. *Mol Cell Endocrinol.* 2021; 520:111074.
27. Hu H, Zhao C, Zhang P, Liu Y, Jiang Y, Wu E, Xue H, Liu C and Li Z. miR-26b modulates OA induced BMSC osteogenesis through regulating GSK3beta/beta-catenin pathway. *Exp Mol Pathol.* 2019; 107:158-164.
28. Huang CC, Kang M, Lu Y, Shirazi S, Diaz JI, Cooper LF, Gajendrareddy P and Ravindran S. Functionally engineered extracellular vesicles improve bone regeneration. *Acta Biomater.* 2020; 109:182-194.
29. Jin Z, Ren J and Qi S. Exosomal miR-9-5p secreted by bone marrow-derived mesenchymal stem cells alleviates osteoarthritis by inhibiting syndecan-1. *Cell Tissue Res.* 2020; 381(1):99-114.
30. Yang X, Yang J, Lei P and Wen T. LncRNA MALAT1 shuttled by bone marrow-derived mesenchymal stem cells-secreted exosomes alleviates osteoporosis through mediating microRNA-34c/SATB2 axis. *Aging (Albany NY).* 2019; 11(20):8777-8791.
31. Zuk PA, Zhu M, Mizuno H, Huang J, Futrell JW, Katz AJ, Benhaim P, Lorenz HP and Hedrick MH. Multilineage cells from human adipose tissue: implications for cell-based therapies. *Tissue Eng.* 2001; 7(2):211-228.
32. Paula AC, Martins TM, Zonari A, Frade SP, Angelo PC, Gomes DA and Goes AM. Human adipose tissue-derived stem cells cultured in xeno-free culture condition enhance c-MYC expression

- increasing proliferation but bypassing spontaneous cell transformation. *Stem Cell Res Ther.* 2015; 6:76.
33. Tolar J and Neglia JP. Transplacental and other routes of cancer transmission between individuals. *J Pediatr Hematol Oncol.* 2003; 25(6):430-434.
  34. Teng IW, Hou PC, Lee KD, Chu PY, Yeh KT, Jin VX, Tseng MJ, Tsai SJ, Chang YS, Wu CS, Sun HS, Tsai KD, Jeng LB, et al. Targeted methylation of two tumor suppressor genes is sufficient to transform mesenchymal stem cells into cancer stem/initiating cells. *Cancer research.* 2011; 71(13):4653-4663.
  35. Manchon E, Hirt N, Bouaziz JD, Jabrane-Ferrat N and Al-Daccak R. Stem Cells-Derived Extracellular Vesicles: Potential Therapeutics for Wound Healing in Chronic Inflammatory Skin Diseases. *International journal of molecular sciences.* 2021; 22(6).
  36. Jin J, Shi Y, Gong J, Zhao L, Li Y, He Q and Huang H. Exosome secreted from adipose-derived stem cells attenuates diabetic nephropathy by promoting autophagy flux and inhibiting apoptosis in podocyte. *Stem Cell Res Ther.* 2019; 10(1):95.
  37. Liu Y, Lin L, Zou R, Wen C, Wang Z and Lin F. MSC-derived exosomes promote proliferation and inhibit apoptosis of chondrocytes via lncRNA-KLF3-AS1/miR-206/GIT1 axis in osteoarthritis. *Cell Cycle.* 2018; 17(21-22):2411-2422.
  38. Yang Q, Yao Y, Zhao D, Zou H, Lai C, Xiang G, Wang G, Luo L, Shi Y, Li Y, Yang M and Huang X. LncRNA H19 secreted by umbilical cord blood mesenchymal stem cells through microRNA-29a-3p/FOS axis for central sensitization of pain in advanced osteoarthritis. *Am J Transl Res.* 2021; 13(3):1245-1256.
  39. Liu Y, Zou R, Wang Z, Wen C, Zhang F and Lin F. Exosomal KLF3-AS1 from hMSCs promoted cartilage repair and chondrocyte proliferation in osteoarthritis. *Biochem J.* 2018; 475(22):3629-3638.
  40. Sun H, Li Y, Kong H, Dai S and Qian H. Dysregulation of KCNQ10T1 promotes cholangiocarcinoma progression via miR-140-5p/SOX4 axis. *Arch Biochem Biophys.* 2018; 658:7-15.
  41. Ding P, Liang B, Shou J and Wang X. lncRNA KCNQ10T1 promotes proliferation and invasion of glioma cells by targeting the miR375/YAP pathway. *Int J Mol Med.* 2020; 46(6):1983-1992.
  42. Liu X, Zhang Y, Wang Y, Bian C and Wang F. Long non-coding RNA KCNQ10T1 up-regulates CTNND1 by sponging miR-329-3p to induce the proliferation, migration, invasion, and inhibit apoptosis of colorectal cancer cells. *Cancer Cell Int.* 2020; 20:340.
  43. Venkatesh J, Wasson MD, Brown JM, Fernando W and Marcato P. LncRNA-miRNA axes in breast cancer: Novel points of interaction for strategic attack. *Cancer Lett.* 2021; 509:81-88.
  44. Xu SY, Shi P and Zhou RM. Post-menopausal oestrogen deficiency induces osteoblast apoptosis via regulating HOTAIR/miRNA-138 signalling and suppressing TIMP1 expression. *Journal of cellular and molecular medicine.* 2021.
  45. Zhang L, Xie H and Li S. LncRNA LOXL1-AS1 controls osteogenic and adipocytic differentiation of bone marrow mesenchymal stem cells in postmenopausal osteoporosis through regulating the miR-196a-5p/Hmga2 axis. *J Bone Miner Metab.* 2020; 38(6):794-805.

46. Wang Y, Cheng K, Zhou W, Liu H, Yang T, Hou P and Li X. miR-141-5p regulate ATF2 via effecting MAPK1/ERK2 signaling to promote preeclampsia. *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie*. 2019; 115:108953.
47. Bao J, Li X, Li Y, Huang C, Meng X and Li J. MicroRNA-141-5p Acts as a Tumor Suppressor via Targeting RAB32 in Chronic Myeloid Leukemia. *Front Pharmacol*. 2019; 10:1545.


## Figures



**Figure 1**


TNF- $\alpha$  increased miR-141-5p expression, suppressed the viability and promoted the apoptosis of MC3T3-E1 cells in a dose-dependent manner. After MC3T3-E1 cells were treated with increased concentrations of TNF- $\alpha$  (0, 1, 2.5, 5, 10 ng/ml) for 24 h, the expression of miR-141-5p was detected via qRT-PCR (a), cell viability was determined by the CCK-8 assay (b), cell apoptosis was analyzed using flow cytometry analysis (c) and the protein level of caspase-3, cleaved caspase-3 and Bax was detected by

immunoblotting (d). Significant differences ( $P < 0.05$ ) are indicated by asterisks. TNF- $\alpha$ , Tumor necrosis factor- $\alpha$




**Figure 2**

Knockdown of miR-141-5p reversed the effect of TNF- $\alpha$  on MC3T3-E1 cells. anti-miR-141-5p reduced the expression of miR-141-5p more significantly in MC3T3-E1 cells compared to anti-miR-NC (a). CCK-8 and Annexin V-FITC/PI assays revealed that the knockdown of miR-141-5p partly reversed the inhibition of cell viability and the promotion of cell apoptosis induced by TNF- $\alpha$  when cells treated with PBS, TNF- $\alpha$ , TNF- $\alpha$ +anti-miR-NC, TNF- $\alpha$ +anti-miR-141-5p (1 $\mu$ g) or TNF- $\alpha$ +anti-miR-141-5p (2 $\mu$ g) respectively (b, c). In line with this result, the increase of Bax and cleaved-Caspase-3 expression induced by TNF- $\alpha$  was inhibited by downregulating miR-141-5p (d). Significant differences ( $P < 0.05$ ) are indicated by asterisks. TNF- $\alpha$ : Tumor necrosis factor- $\alpha$



**Figure 3**


Characteristics of ADSCs and ADSCs-derived exosomes. The isolated ADSCs exhibited typical fibroblastic-like morphology (a). Alizarin Red, Oil Red O and Alcian Blue staining was positive after the induced osteogenic, adipogenic and chondrogenic differentiation of ADSCs (b). Immunoblotting showed that ADSCs-Exos were positive for Alix, CD63, CD81 and CD9 (c). MC3T3-E1 cells could uptake the PKH26-labeled ADSCs-Exos (d). Red: PKH26-labeled ADSCs-Exos. Blue: nuclei. Exos: exosomes, ADSCs: adipose-derived stem cells



**Figure 4**


ADSCs-Exos attenuated the effect of TNF- $\alpha$  on MC3T3-E1 cells. After MC3T3-E1 cells were treated with TNF- $\alpha$  (5 ng/ml) and different dose of ADSCs-Exos (PBS, TNF- $\alpha$ , TNF- $\alpha$ +Exos (25 $\mu$ g), TNF- $\alpha$ +Exos (50 $\mu$ g), TNF- $\alpha$ +Exos (100 $\mu$ g)), ADSCs-Exos dose-dependently decreased TNF- $\alpha$  induced miR-141-5p expression by qRT-PCR (a). CCK-8 assay revealed that ADSCs-Exos mitigated the inhibition of TNF- $\alpha$  in cell viability in a dose-dependent manner (b). Immunoblotting showed that ADSCs-Exos dose-dependently decreased the

promotion of TNF- $\alpha$  on cleaved caspase-3 and Bax expression (c). Flow cytometry analysis showed that ADSCs-Exos dose-dependently reversed TNF- $\alpha$ -induced cell apoptosis (d). Significant differences ( $P < 0.05$ ) are indicated by asterisks. Exos: exosomes, ADSCs-Exos: exosomes derived from adipose-derived stem cells, TNF- $\alpha$ : tumor necrosis factor- $\alpha$




**Figure 5**

KCNQ10T1-Exos inhibited TNF- $\alpha$ -induced cytotoxicity and apoptosis of MC3T3-E1 cells. The expression of KCNQ10T1 in ADSCs or exosomes derived from the ADSCs was elevated, but the expression of miR-141-5p was down-regulated compared with that in LV-NC treated group after the transfection of LV-KCNQ10T1 (a, b). KCNQ10T1 expression was up-regulated in MC3T3-E1 cells treated with LV-KCNQ10T1-Exos compared to NC-Exos (c). In MC3T3-E1 cells, LV-KCNQ10T1-Exos mitigated the negative effect of TNF- $\alpha$  on cell viability, while ADSCs-Exos exerted a weaker stimulative effect on cell viability compared to LV-KCNQ10T1-Exos (d). When MC3T3-E1 cells were co-cultured with LV-KCNQ10T1-Exos, the TNF- $\alpha$ -induced cell apoptosis was reversed and ADSCs-Exos exerted a weaker inhibitory effect on cell apoptosis compared to LV-KCNQ10T1-Exos (e). The expression of Bax and cleaved caspase-3 in MC3T3-E1 cells was blocked after co-culture of LV-KCNQ10T1-Exos (f). Significant differences ( $P < 0.05$ ) are indicated by asterisks. Exos: exosomes, ADSCs-Exos: exosomes derived from adipose-derived stem cells



**Figure 6**

KCNQ10T1 could sponge miR-141-5p. KCNQ10T1 and miR-141-5p have potential binding sites (a). As dual-luciferase reporter assay showed, miR-141-5p weakened the luciferase activity in KCNQ10T1-WT group obviously, but the activity of luciferase reporters containing KCNQ10T1-MUT was not changed significantly in both HEK293 and MC3T3-E1 cells (b,c). LV-KCNQ10T1-Exos blocked the inhibitory effect of miR-141-5p on the luciferase activity of reporters containing KCNQ10T1-WT compared with ADSC-Exos and LV-NC-Exos (d). KCNQ10T1 suppressed the expression of miR-141-5p, but the sh-KCNQ10T1 increased that (e). The expression of miR-141-5p was down-regulated when MC3T3-E1 cells co-cultured with LV-KCNQ10T1-Exos (f). Significant differences ( $P < 0.05$ ) are indicated by asterisks. Exos: exosomes, ADSCs-Exos: exosomes derived from adipose-derived stem cells



**Figure 7**

KCNQ10T1-Exos inhibited the effect of TNF- $\alpha$  in MC3T3-E1 cells by sponging miR-141-5p. MC3T3-E1 cells were treated with TNF- $\alpha$  and then treated with miR-NC, miR-141-5p, miR-141-5p+ADSCs-Exos, miR-141-5p+LV-NC-Exos or miR-141-5p+LV-KCNQ10T1-Exos respectively. The inhibitory effect of TNF- $\alpha$  on cell viability was blocked by miR-141-5p and the treatment of ADSCs-Exos or KCNQ10T1-Exos partly reversed this phenomenon; ADSCs-Exos exerted a weaker stimulative effect on cell viability compared to LV-KCNQ10T1-Exos (a). MiR-141-5p enhanced the expression of cleaved caspase-3 and Bax but was attenuated following co-culture with ADSC-Exos or LV-KCNQ10T1-Exos (b). As flow cytometry showed, miR-141-5p promoted cell apoptosis while the treatment of ADSC-Exos or LV-KCNQ10T1-Exos inhibited that; ADSCs-Exos exerted a weaker inhibitory effect on cell apoptosis compared to LV-KCNQ10T1-Exos (c). Significant differences ( $P < 0.05$ ) are indicated by asterisks. Exos: exosomes, ADSCs-Exos: exosomes derived from adipose-derived stem cells.

of ADSCs-Exos or KCNQ10T1-Exos partly reversed this phenomenon; ADSCs-Exos exerted a weaker stimulative effect on cell viability compared to LV-KCNQ10T1-Exos (a). MiR-141-5p enhanced the expression of cleaved caspase-3 and Bax but was attenuated following co-culture with ADSC-Exos or LV-KCNQ10T1-Exos (b). As flow cytometry showed, miR-141-5p promoted cell apoptosis while the treatment of ADSC-Exos or LV-KCNQ10T1-Exos inhibited that; ADSCs-Exos exerted a weaker inhibitory effect on cell apoptosis compared to LV-KCNQ10T1-Exos (c) . Significant differences ( $P < 0.05$ ) are indicated by asterisks. Exos: exosomes, ADSCs-Exos: exosomes derived from adipose-derived stem cells