Exploring the Missing Heritability in Subjects With Hearing Loss, Enlarged Vestibular Aqueducts, and A Single or No Pathogenic *SLC26A4* Variant

Jeroen Smits

Radboudumc Department of Human Genetics <https://orcid.org/0000-0002-4241-0900>

Suzanne E. de Bruijn

Radboudumc Department of Human Genetics

Cornelis P. Lanting

Radboudumc

Jaap Oostrik

Radboudumc Department of Human Genetics

Luke O'Gorman

Radboudumc Department of Human Genetics

Tuomo Mantere

Radboudumc Department of Human Genetics

Frans P. M. Cremers

Radboudumc Department of Human Genetics

Susanne Roosing

Radboudumc Department of Human Genetics

Helger G. Yntema

Radboudumc Department of Human Genetics

Erik de Vrieze

Radboudumc Department of Human Genetics

Ronny Derks

Radboudumc Department of Human Genetics

Alexander Hoischen

Radboudumc Department of Human Genetics

Sjoert A.H. Pegge

Radboudumc Afdeling Radiologie en nucleaire beeldvorming

Kornelia Neveling

Radboudumc Department of Human Genetics

Ronald J.E. Pennings

Radboudumc

Hannie Kremer (✉ Hannie.Kremer@radboudumc.nl)

Radboudumc Department of Human Genetics

Research Article

Keywords: Hearing loss, genome sequencing, long-read sequencing, genotype-phenotype correlations, SLC26A4, enlarged vestibular aqueduct

Posted Date: June 4th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-577715/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Human Genetics on August 19th, 2021. See the published version at <https://doi.org/10.1007/s00439-021-02336-6>.

Abstract

Pathogenic variants in *SLC26A4* have been associated with autosomal recessive hearing loss (arHL) and a unilateral or bilateral enlarged vestibular aqueduct (EVA). *SLC26A4* is the second most frequently mutated gene in arHL. Despite the strong genotype-phenotype correlation, a significant part of *SLC26A4* cases remains genetically unresolved. In this study, we investigated a cohort of 28 Dutch index cases diagnosed with HL in combination with an EVA but without (M0) or with a single (M1) pathogenic variant in *SLC26A4*. To explore the missing heritability, short- and long-read whole genome sequencing and optical genome mapping were performed. We found a previously described EVA-associated haplotype (Caucasian EVA (CEVA)) to be significantly enriched in our M1 patient cohort. The haplotype was also present in two M0 cases. Despite extensive genetic analyses, we were not able to prioritize any of the variants present within the haplotype as the likely pathogenic defect, and therefore additional analyses addressing the defect(s) at the RNA, protein, or epigenetic level are required. Whole genome sequencing also revealed splice-altering *SLC26A4* variants in two M1 cases, which are now genetically explained, but no deep-intronic or copy number variants. With these findings, we have provided important insights that will pave the way for elucidating the missing heritability in M0 and M1 *SLC26A4* cases.

Introduction

SLC26A4 encodes the transmembrane anion transporter pendrin and is most abundantly expressed in the inner ear, thyroid gland, kidney, and airways epithelia (Everett et al. 1997; Everett et al. 1999; Pedemonte et al. 2007; Royaux et al. 2000; Royaux et al. 2001). The 780 amino acid protein is part of the solute carrier family 26 and plays a pivotal role in chloride, bicarbonate and iodine transport. In the inner ear, pendrin functions as a $\text{Cl}^-/\text{HCO}_3^-$ exchanger. The protein is expressed in the epithelial cells of the cochlea (outer sulcus and spindle cells), the vestibular labyrinth (transitional cells), and the endolymphatic duct and sac (mitochondrial-rich cells) (Wangemann 2011; Wangemann et al. 2007). Expression of pendrin is essential for the development of the (murine) auditory and vestibular system and for maintaining ion homeostasis in the endolymphatic fluid and the endocochlear potential (Dou et al. 2004; Everett et al. 1999; Royaux et al. 2003; Wangemann 2011).

Defects in *SLC26A4* are among the most frequent causes (up to 10%) of early-onset autosomal recessive hearing loss (arHL); non-syndromic DFNB4 (MIM: 600791) and Pendred syndrome (MIM: 274600) (Sloan-Heggen et al. 2016). Individuals carrying biallelic pathogenic *SLC26A4* variants are affected by variable, often progressive and predominantly sensorineural HL with a congenital or childhood-onset (Lee et al. 2014; Suzuki et al. 2007). In Pendred syndrome, the HL phenotype is accompanied by an iodine organification defect that can lead to thyroid goiter (Fraser 1965). In individuals affected by either syndromic or non-syndromic *SLC26A4*-associated HL, a unilateral or bilateral enlarged vestibular aqueduct (EVA) is observed, which is the most common imaging abnormality in individuals with HL (van Beeck Calkoen et al. 2018; van Beeck Calkoen et al. 2017).

Pathogenic variants in *SLC26A4* have a loss-of-function effect, leading to malfunctioning of the pendrin ion transporter. Besides the antenatal formation of an EVA, this ultimately leads to acidification of the endolymphatic fluids in the inner ear during embryonic development (Griffith and Wangemann 2011; Wangemann 2011). Although the exact molecular pathogenic mechanism remains to be elucidated, the lack of pendrin function ultimately leads to degeneration of the sensory cells in the inner ear (Wangemann 2011).

Despite the strong association between defects of *SLC26A4* and HL combined with an EVA, genetic screening of subjects with this combination of defects often does not reveal biallelic pathogenic variants in *SLC26A4* (coined M2). Cohort studies report that 14–31% of the subjects with an EVA and HL carry a monoallelic pathogenic variant in *SLC26A4* (M1), whereas in 10–65% of the subjects, no potentially pathogenic variant in the coding or splice site regions of the gene can be identified (M0) (Azaiez et al. 2007; Choi et al. 2009; Mey et al. 2019). Segregation analyses performed in family members of M1 subjects, however, do suggest that in 98% of M1 subjects an unidentified or unrecognized variant is present on the *trans SLC26A4* allele (Azaiez et al. 2007; Pryor et al. 2005). In line with this hypothesis, Chattaraj and coworkers reported a ≥ 613 -kb haplotype located upstream of *SLC26A4* that was present in 13 of 16 (81%) of the studied M1 families and that was also enriched in M0 subjects (Chattaraj et al. 2017). The haplotype, referred to as the Caucasian EVA (CEVA) haplotype, is defined by the combination of 12 single nucleotide polymorphisms (SNPs) (allele frequency (AF) 1.9–4.0%) that are located in intergenic or intronic regions. The true genetic defect of the CEVA allele has not been identified yet, but it cannot be excluded that a potential defect was missed due to the technical limitations of short-read sequencing and other standard-of-care tests. The CEVA haplotype was reported to be associated with a less severe HL phenotype as compared to variants in the protein-coding or splice site regions of *SLC26A4* (Chao et al. 2019).

We investigated a Dutch cohort of M1 and M0 subjects with HL and a unilateral or bilateral EVA. All subjects were tested for the presence of the CEVA haplotype, and whole genome sequencing (WGS) was performed to detect potentially missed single nucleotide variants (SNVs), structural variants (SVs), and regulatory or deep-intronic variants. Long-read sequencing and optical genome mapping were performed to reveal a potentially missed SV located on the CEVA haplotype. Variants located within the haplotype were subjected to *in silico* analyses to investigate potential effects on the regulation of *SLC26A4* expression or on splicing. With this study, we provided further insights into *SLC26A4*-associated disease.

Material And Methods

Inclusion criteria and clinical evaluation

Subjects diagnosed with a unilateral or bilateral EVA on CT or MRI and for whom medical genetic testing only revealed a heterozygous (M1) or no pathogenic variant (M0) in *SLC26A4* were eligible to participate in this study. A retrospective cohort of nine subjects with confirmed pathogenic (biallelic) variants in *SLC26A4* was added as a reference cohort (**Online Resource Table S1**).

Medical history was taken from all participants and special attention was paid to non-genetic causes of HL. Results of pure tone, speech, and brainstem evoked response audiometry, performed in a sound-attenuated booth, were collected. Air and bone conduction pure tone thresholds were determined for

frequencies ranging from 0.25 to 8 kHz. Threshold estimates based on brainstem evoked response audiometry were used when pure tone audiometry was not available. Individuals were considered affected when pure tone thresholds for at least three frequencies were above the frequency-specific 95th percentile of age- and sex-specific thresholds (ISO 7029:2017) for the best hearing ear. In the Netherlands, routine newborn hearing screening is carried out by the detection of transient evoked otoacoustic emissions (van der Ploeg et al. 2012). When available, these data were used to determine whether the HL was congenital.

Previously performed CT and MRI scans were retrieved and reassessed by an experienced neuroradiologist (S.A.H.P.). An EVA was defined as a vestibular aqueduct that measured ≥ 2 mm at the operculum and/or ≥ 1 mm at the midpoint (Boston et al. 2007), in accordance with previously published reports on this topic (Chao et al. 2019; Chattaraj et al. 2017). Analyses of pair-wise differences between patient groups were performed with R (R Foundation, Auckland, New Zealand) using multivariate linear regression analysis (using lsmmeans 2.3.0) with a correction for multiple comparisons using the Holm method (Lenth 2016).

Next generation sequencing and variant interpretation

Genomic DNA was isolated from peripheral blood lymphocytes and analyzed by molecular inversion probe (MIP) sequencing, whole exome sequencing (WES) or whole genome sequencing (WGS) (**Online Resource Table S2**). For WES, exome enrichment was performed using the Agilent SureSelect Human All Exome V4 or V5 kits according to the manufacturer's instructions. Subsequently, sequencing was executed on an Illumina HiSeq system by BGI Europe (Copenhagen, Denmark), with a minimal coverage of 20x for 93.77% of the targets and an average coverage of > 100 reads. Read mapping along the hg19 reference genome (GRCh37/hg19) and variant calling was performed using BWA V.0.78 (Li and Durbin 2009) and GATK HaplotypeCaller V.3.3 (McKenna et al. 2010), respectively. An in-house developed pipeline was used for variant annotation and copy number variant (CNV) detection was performed using CoNIFER V.0.2.2.3 (Krumm et al. 2012). WGS was performed by BGI (Hongkong, China) on a BGISEQ500 using a 2x 100 bp paired end module, with a minimal median coverage of 30-fold per genome. Read mapping (GRCh37/hg19) and variant calling was performed as described for WES. Structural variants (SVs) were called using the Manta Structural Variant Caller V.1.1.0 (SV detection based on paired end and split read evidence) (Chen et al. 2016) and CNVs using Control-FREEC (CNV detection based on alterations in read depth) (Boeva et al. 2012). MIP design, sequencing and data analysis were performed as previously described (de Bruijn et al. 2021; Neveling et al. 2017). MIPs were designed to cover exons and exon-intron boundaries of a panel of 120 HL genes (**Online Resource Table S3**). For each targeted region an average coverage of > 500 reads was obtained. A minimal coverage of 20x was reached for 91.78% of the MIPs. CNV detection for *SLC26A4* was performed using a read coverage analysis as previously described (Khan et al. 2020).

Variant prioritization was based on an AF of $\leq 0.5\%$ (gnomAD V2.1.1 (Karczewski et al. 2019) and our in-house exome database ($\sim 15,000$ alleles)), unless specified otherwise. Variant visualization was performed using the IGV software V.2.4 (Broad Institute, Cambridge, MA, USA) (Robinson et al. 2011). Interpretation of missense variants was performed using the *in silico* tools CADD-PHRED (≥ 15) (Kircher et al. 2014), SIFT (≤ 0.05) (Vaser et al. 2015), PolyPhen-2 (≥ 0.450) (Adzhubei et al. 2010) and MutationTaster (deleterious) (Schwarz et al. 2014) to predict potentially deleterious effects. Variants were prioritized if a deleterious effect was predicted by at least two of these tools. Candidate variants were validated by Sanger sequencing and segregation analysis was performed when DNA of family members was available. Primer sequences and PCR conditions are available upon request. Potential effects on splicing of missense, synonymous and intronic variants were assessed using the deep-learning splice prediction algorithm SpliceAI (≥ 0.1) (Jaganathan et al. 2019). The maximum distance between the variant and potential gained or lost splice sites was set to 1000 bp. Predicted splice altering defects were evaluated using an *in vitro* splice assay in HEK293T cells as previously described (Sangermano et al. 2018).

Detection of the CEVA haplotype

After initial SNP-genotyping by Sanger sequencing, genotyping of the CEVA haplotype (Chattaraj et al. 2017) was performed using VNTR marker analysis. DNA segments were amplified by employing touchdown PCR, and subsequent analysis was carried out on an ABI Prism 3730 Genetic Analyzer (Applied Biosystems, Foster City, CA, USA). Genomic positions of the markers were determined using the UCSC genome browser (GRCh37/hg19) (Kent et al. 2002). Alleles were assigned with the GeneMarker software (V.2.6.7, SoftGenetics, State College, PA, USA) according to the manufacturer's instructions. When an individual was suspected of carrying the CEVA haplotype based on VNTR-marker alleles, SNP genotyping by Sanger sequencing was performed to confirm the presence of the twelve SNPs that are located within the haplotype (Chattaraj et al. 2017).

Optical genome mapping

Optical genome mapping (Bionano Genomics, San Diego, CA, USA) was performed as previously described (Mantere et al. 2020; Neveling et al. 2020). Ultra-high molecular weight DNA was isolated from whole peripheral blood (collected in EDTA tubes) using the SP Blood & Cell Culture DNA Isolation Kit (Bionano Genomics, San Diego, CA, USA). CTTAAG labeling was performed using the DLS (Direct Label and Stain) DNA Labeling Kit (Bionano Genomics, San Diego, CA, USA) and the labeled sample was analyzed using a 3x 1,300 Gb Saphyr chip (G2.3) on a Saphyr instrument (Bionano Genomics, San Diego, CA, USA). An effective coverage of 124x was reached, with a label density of 14.63/100 kb and an average N50 of 279 kb. *De novo* assembly (using GRCh37 and GRCh38) and variant annotation were performed using Bionano Solve version 3.4, which includes two separate algorithms for SV and CNV detection. Annotated variants were filtered for rare events as described previously (Mantere et al. 2020). In addition, the genomic region spanning the CEVA haplotype was analyzed visually in Bionano Access version 1.4.3.

PacBio long-read sequencing

Genomic DNA was isolated from peripheral blood according to standard procedures and subjected to long-read genome Hi-Fi sequencing using the SMRT sequencing technology (Pacific Biosciences, Menlo Park, CA, USA). Library preparation was performed using the SMRTbell™ Template Prep Kit 2.0 (Pacific Biosciences, Menlo Park, CA, USA) following manufacturer's instructions. Size selection was performed using a BluePippin DNA size selection system (target fragments ~ 15 – 18 kb). Sequence primer V2 and polymerase 2.0 were used for binding. Subsequently, the SMRTbell library was loaded on an 8M SMRTcell and sequencing was performed on a Sequel II system (Pacific Biosciences, Menlo Park, CA, USA). Circular consensus sequencing (CCS), Hi-Fi reads, were generated using the CCS (v4.2.0) tool and were aligned to the GRCh37/hg19 reference genome with pbmm2 (v.1.3.0). The unique molecular yield was 93.46

Gb and the post-alignment Hi-Fi- coverage was 12x (Mosdepth v0.3.1, (Pedersen and Quinlan 2018)). SV calling was performed using PBSV (v2.4.0) and annotation was applied using an in-house SV annotation pipeline.

Results

Patient inclusion and genetic prescreening

In this study, we included 28 Dutch index cases diagnosed with a unilateral or bilateral EVA and HL. All individuals were prescreened for pathogenic variants in *SLC26A4* (NM_000441.1) in a diagnostic setting and complete coverage of the coding and splice site (+/- 14 nucleotides) regions of *SLC26A4* was confirmed. In 16 individuals, a heterozygous (likely) pathogenic *SLC26A4* variant was reported and these cases were deemed M1. In the remaining 12 individuals, no potentially pathogenic variants were found in the coding or splice site regions of this gene, and these subjects were therefore considered M0. Causative variants in other genes associated with autosomal recessive arHL (Van Camp and Smith) were addressed and excluded by analyzing available sequencing data (WES or MIPs-based) or in WGS data obtained in this study (**Online Resource Table S2**). This revealed no homozygous or compound heterozygous variants that were known or predicted to be pathogenic, except two compound heterozygous variants in *OTOGL* (NM_173591.3) in individual SLC012 (**Online Resource Table S4**). The c.890C > T (p.(Pro297Leu)) variant in *OTOGL* has, however, been reported as (likely) benign in ClinVar (Landrum et al. 2018) and the Deafness Variation Database (Azaiez et al. 2018). The c.1369G > T (p.(Val457Leu)) is considered as a variant of unknown significance in ClinVar. Furthermore, subject SLC012 has progressive high-frequency HL, which differs from the symmetric, moderate, and stable HL associated with *OTOGL* (DFNB84B) (Oonk et al. 2014; Yariz et al. 2012). Therefore, we considered the identified *OTOGL* variants as non-causative.

The CEVA haplotype is enriched in Dutch monoallelic *SLC26A4* cases

In 2017, Chatteraj et al. described the ≥ 613 -kb CEVA haplotype located centromeric of the *SLC26A4* gene to be enriched in M1 *SLC26A4* cases and M0 cases with HL and EVA (Chatteraj et al. 2017). To investigate whether this haplotype is also enriched in the selected Dutch cohort of M0 and M1 *SLC26A4* cases, we screened for the presence of this haplotype using VNTR marker analysis followed by Sanger sequencing of the 12 CEVA-associated SNPs. The CEVA haplotype was detected in 8 out of 16 (50%) M1 individuals and 2 out of 12 (16.7%) M0 subjects (Fig. 1, Table 1). In two M1 individuals (SLC040 and SLC071), only a partial CEVA haplotype was found, harboring 9/12 SNPs. We will refer to this smaller haplotype as the variant 1-CEVA (V1-CEVA) haplotype.

Table 1
Detection of the CEVA haplotype in M1 and M0 individuals

Zeroallelic <i>SLC26A4</i> cases			
	Allele 1		Allele 2
Case	Variant	ClinVar	CEVA
SLC014	c.2059G > T; p.(Asp687Tyr)	UV3	
SLC015	-	-	
SLC017	-	-	
SLC039	-	-	ACACATG-GC-C (CEVA)
SLC043	-	-	
SLC052	-	-	
SLC069	-	-	
SLC070	-	-	
SLC073	-	-	
SLC080	-	-	ACACATG-GC-C (CEVA)
SLC084	-	-	
SLC086	-	-	
Monoallelic <i>SLC26A4</i> cases			
SLC002	c.412G > T; p.(Val138Phe)	UV4	
SLC003	c.131dup; p.(Thr45Aspfs*42)		ACACATG-GC-C (CEVA)
SLC012 ^a	c.707T > C; p.(Leu236Pro)	UV5	ACACATG-GC-C (CEVA)
SLC013	c.1001 + 1G > A; p.(?)		ACACATG-GC-C (CEVA)
SLC018	c.349C > T; p.(Leu117Phe)	UV4	
SLC031	c.1001 + 1G > A; p.(?)		ACACATG-GC-C (CEVA)
SLC032	c.1334T > G; p.(Leu445Trp)	UV5	
SLC036 ^a	c.1246A > C; p.(Thr416Pro)	UV5	ACACATG-GC-C (CEVA)
SLC040 ^a	c.655_656dup; p.(Phe223Alafs*15)		GTTTCATG-GC-C (V1-CEVA)
SLC045	c.1334T > G; p.(Leu445Trp)	UV5	
SLC048	c.706C > G; p.(Leu236Val)	UV5	
SLC056	c.707T > C; p.(Leu236Pro)	UV5	ACACATG-GC-C (CEVA)
SLC071 ^a	c.1334T > G; p.(Leu445Trp)	UV5	GTTTCATG-GC-C (V1-CEVA)
SLC078	c.304G > C; p.(Gly102Arg)	UV5	ACACATG-GC-C (CEVA)
SLC079	c.1001 + 1G > A; p.(?)		ACACATG-GC-C (CEVA)
SLC085	c.706C > G; p.(Leu236Val)	UV5	
Presence of the CEVA haplotype was tested in zeroallelic (M0) and monoallelic (M1) <i>SLC26A4</i> cases with a unilateral or bilateral enlarged vestibular aqueduct. <i>SLC26A4</i> (NM_000441.1) variants reported in ClinVar as (likely) pathogenic (UV4, UV5) were considered causative, whereas variants reported as (likely) benign or of unknown significance were considered non-causative. In ten individuals, the complete CEVA haplotype was detected (ACACATG-GC-C), whereas in two individuals a shorter version of the haplotype was found, consisting of 9/12 CEVA SNPs (GTTTCATG-GC-C; V1). For individuals marked with an ^a , it could be conclusively determined that the (V1-)CEVA haplotype is present on the <i>trans SLC26A4</i> allele. ClinVar, American College of Medical Genetics and Genomics (ACMG) classification of variants as in ClinVar; UV3, variant of unknown significance; UV4, likely pathogenic; UV5, pathogenic.			

The CEVA haplotype has an AF of 2.8% in the 1000G database (28 in 1006 alleles) (Chattaraj et al. 2017; Genomes Project Consortium et al. 2015), and an AF of 3.3% in an in-house control cohort consisting of 322 healthy unrelated individuals (21 in 644 unphased alleles). This implies a significant enrichment of the CEVA haplotype in our M1 cohort (8 in 32 alleles) compared to the 1000G database (p-value 5.419×10^{-6}) and the control cohort (p-value 2.187×10^{-5}) as determined by a two-sided Fisher's exact test). The two M1 cases with the V1-CEVA haplotype were not included in this statistical analysis. Also this V1-CEVA allele is significantly enriched in our M1 cohort as only a single V1-CEVA allele is reported in the 1000G database (1 in 1006 alleles) (Chattaraj et al. 2017) (p-value 0.0027). The CEVA haplotype was not found to be significantly enriched in the M0 cohort (2 in 24 alleles). Although the pathogenicity of the CEVA haplotype is unclear, the significant enrichment of the haplotype within this M1 patient cohort and the patient cohorts (M1 and M0) previously described by Chattaraj and co-workers strongly suggests that a pathogenic defect resides within this haplotype (Chattaraj et al. 2017). Because of this strong association

of the CEVA haplotype with HL and EVA, we considered the M1 individuals carrying the CEVA or the V1-CEVA haplotype as genetically explained (M1/CEVA), and M0 individuals with the CEVA haplotype (M0/CEVA) as monoallelic in further steps of this study. For six M1 individuals, it could not be conclusively determined whether the CEVA haplotype was present *in trans* with the pathogenic *SLC26A4* variant, as the genetic material of family members was not available (Table 1).

Whole genome sequencing reveals potential *SLC26A4* splice and regulatory variants in M1 subjects without the CEVA haplotype

To detect any potentially missed coding or unidentified intronic *SLC26A4* variants or variants located *in cis* regulatory elements of the gene, WGS analysis was performed for all six M1 individuals who could not be genetically explained by the presence of the CEVA haplotype. Additionally, WGS analysis was performed for the two M0/CEVA individuals. In none of these eight cases, SVs overlapping with the *SLC26A4* gene were identified by WGS.

To identify any variants with a potential effect on splicing, the deep-learning algorithm SpliceAI was employed (Jaganathan et al. 2019). In two M1 individuals (SLC048 and SLC085), a rare heterozygous potentially splice altering *SLC26A4* variant was identified (Table 2). For both variants, the predicted splice defect was investigated using an *in vitro* splice assay performed in HEK293T cells. For SLC048, a canonical splice site variant (c.1342-2A > C) was predicted to remove the splice acceptor site. Indeed, the splice assay revealed loss of the acceptor site and usage of an alternative splice acceptor site located thirteen nucleotides downstream (**Online Resource Figure S1A**). This leads to the formation of an out-of-frame exon 12 and premature protein truncation (p.(Ser448Leufs*3)). In SLC085, a synonymous variant (c.471C > T) was identified in exon 5. SpliceAI predicts that this variant strengthens an alternative splice acceptor site (27 nucleotides downstream of the variant). Indeed, an *in vitro* splice assay confirmed that the alternative splice acceptor site is used, which leads to the partial deletion of exon 5 and a truncated protein (p.(Gly139Alafs*6)) (**Online Resource Figure S1B**). The observed splice defect resulting from this synonymous variant underlines the importance of evaluating potential splice effects of all rare variants in coding sequences, using *in silico* prediction splice tools. We considered the two identified splice variants as pathogenic and the HL of the two individuals as genetically explained, thus M2.

Table 2
WGS revealed two heterozygous splice variants in *SLC26A4*

Case	Class	Genome	cDNA	Protein	In-house AF (%)	gnomAD AF (%)	CADD_PPHRED	SIFT	PPH2	MutationTaster	SpliceAI	CI
SLC048	M1	Chr7:107335064A > C	c.1342-2A > C	p.Ser448Leufs*3	0.00	-	21.7	NA	NA	NA	0.99 (AS loss)	U
SLC085	M1	Chr7:107314664C > T	c.471C > T	p.Gly139Alafs*6,=	-	0.00	0.725	NA	NA	NA	0.59 (AS gain)	-

Whole genome sequencing (WGS) revealed two potentially splice altering variants in *SLC26A4*. Variants are selected based on an allele frequency of $\leq 0.5\%$ in gnomAD and the in-house database. Scores that meet the thresholds for pathogenicity as described in the methods section are indicated in red. The predicted effect on splicing was confirmed in an *in vitro* splice assay that was performed in HEK293T cells (**Figure S1**). Genome; Genomic position according to GRCh37/hg19; In-house AF, allele frequency (%) in an in-house database (~ 7,500 exomes); GnomAD AF, allele frequency (%) in gnomAD database V2.1.1; CADD_PPHRED, Combined Annotation Dependent Depletion PHRED score; SIFT, Scale-Invariant Feature Transform; PPH2, PolyPhen-2 score; MutationTaster (p MutationTaster score with probability (0–1); spliceAI, splicing prediction score; AS, acceptor site; ClinVar, American College of Medical Genetics and Genomic (ACMG) classification of variants as in ClinVar;UV4, likely pathogenic; NA, not applicable.

To explore variants that are potentially located within a *cis* regulatory element of *SLC26A4*, we extracted all (predicted) human enhancer and promoter elements that are associated with the *SLC26A4* gene from the GeneHancer (Fishilevich et al. 2017) and EnhancerAtlas (Gao and Qian 2020) databases (**Online Resource Table S5**). GeneHancer V5 is a collection of both predicted and experimentally validated enhancer-to-gene and promoter-to-gene interactions, based on information integrated from multiple resources: ENCODE (Dunham et al. 2012), Ensembl (Zerbino et al. 2015), FANTOM5 (Andersson et al. 2014), VISTA (Visel et al. 2007), dbSuper (Khan and Zhang 2016), EPDnew (Dreos et al. 2013), UCNEbase (Dimitrieva and Bucher 2013) and CraniofacialAtlas (Wilderman et al. 2018). For each regulatory element, a gene interaction score (> 7) and element confidence score (> 0.7) are provided. The EnhancerAtlas V2 is a database providing enhancer annotations in different species based on experimental datasets determined in several tissues and cell types.

WGS data were analyzed for variants located within these elements and two rare potentially regulatory variants were identified in two M1 individuals (SLC002 and SLC045) (**Online Resource Table S6**). Both variants are located in a predicted enhancer element of *SLC26A4* according to GeneHancer. The variants have a low nucleotide conservation score (PhyloP, ≤ 1.0) and are not located in a conserved region (UCSC genome browser (Kent et al. 2002), Conservation comparative genomics track). Additionally, we investigated the possibility of the loss of a transcription factor binding site using the JASPAR database (Fornes et al. 2020), which was not the case. Since we did not find any strong indication of a functional effect for the two variants, we considered them non-pathogenic.. Nevertheless, only a reporter assay can completely exclude a potential regulatory effect of the variants on *SLC26A4* expression.

A FOXI1 missense variant is revealed in three unrelated index cases

Several studies have suggested a potential digenic inheritance for *SLC26A4* variants and variants in *KCNJ10* and *FOXI1* (Pique et al. 2014; Yang et al. 2009; Yang et al. 2007). Additionally, a more recent study suggested digenic inheritance with pathogenic variants in *EPHA2* (Li et al. 2020). We screened all remaining genetically unexplained individuals (M1, M0/CEVA and M0) for variants in these genes with an AF $\leq 5\%$ (gnomAD V2.1.1). In cases for which only MIP sequencing data was available, coding regions and exon-intron boundaries of *FOXI1* and the regions harboring the reported pathogenic variants in *EPHA2* (c.1063G > A; p.(G355R), c.1532C > T; (p.T511M), NM004431.4) were analyzed using Sanger sequencing. In three individuals (SLC039; M0/CEVA, SLC052; M0 and SLC069; M0) a c.677C > T (p.(Thr226Ile)) *FOXI1* (NM_012188.4) missense variant was identified (Table 3). The variant was not identified in any of the

M1/CEVA or the two M2 cases. *FOXI1* encodes the Forkhead transcription factor FOXI1, a key transcriptional regulator of *SLC26A4* (Yang et al. 2007). Segregation analysis has confirmed that the *FOXI1* variant is not co-inherited with the CEVA allele in individual SLC039, which is in line with digenic inheritance. The Thr226 residue is located outside of the conserved forkhead DNA-binding domain of FOXI1 (amino acids 94–211) (Yang et al. 2007) and none of the *in silico* tools used for analysis predicted a deleterious effect of the c.677C > T variant. Nevertheless, the variant seems to be enriched in individuals diagnosed with HL and EVA, and we consider the c.677C > T *FOXI1* variant as an interesting candidate for functional validation.

Table 3
Rare variants identified in *EPHA2*, *FOXI1* and *KCNJ10*

Case	Class	Gene	Transcript	cDNA	Protein	In-house AF (%)	gnomAD AF (%)	CADD_PPHRED	SIFT	PPH2	Mutation Taster	SpliceAI	ClinVar
SLC017	M0	<i>EPHA2</i>	NM_004431.4	c.2627G > A	p.(Arg876His)	2.36	1.70	32	0	0.769	NA	0.03	UV
SLC039	M0/CEVA	<i>FOXI1</i>	NM_012188.4	c.677C > T	p.(Thr226Ile)	0.56	0.37	11	0.14	0.109	P	0.03	UV
SLC052	M0	<i>EPHA2</i>	NM_004431.4	c.1941G > T	p.(Thr647=)	1.09	0.55	7.309	NA	NA	NA	0.05	UV
SLC052	M0	<i>EPHA2</i>	NM_004431.4	c.1896G > A	p.(Leu632=)	0.76	0.05	3.197	NA	NA	NA	0.05	UV
SLC052	M0	<i>FOXI1</i>	NM_012188.4	c.677C > T	p.(Thr226Ile)	0.56	0.37	11	0.14	0.109	P	0.03	UV
SLC069	M0	<i>FOXI1</i>	NM_012188.4	c.677C > T	p.(Thr226Ile)	0.56	0.37	11	0.14	0.109	P	0.03	UV

Available sequencing datasets of monoallelic (M1, M0/CEVA) and zeroallelic (M0) individuals were screened for variants in *EPHA2*, *FOXI1* and *KCNJ10* with an allele frequency of $\leq 5\%$ in gnomAD (V2.1.1). Scores that meet the thresholds for pathogenicity as described in the methods section are indicated in red. In-house AF, allele frequency (%) in in-house database (~7,500 exomes); GnomAD AF, allele frequency (%) in gnomAD database V2.1.1; CADD_PPHRED, Combine Annotation Dependent Depletion PHRED score; SIFT, Scale-Invariant Feature Transform; PPH2, PolyPhen-2 score; MutationTaster (prob), MutationTaster score with probability (0–1); spliceAI, splicing prediction score; ClinVar, American College of Medical Genetics and Genomics (ACMG) classification of variants as in ClinVar; UV1, benign; UV2, likely benign; NA, not available; P, polymorphism.

In case SLC017, a heterozygous missense variant in *EPHA2* was detected (c.2627G > A (p.(Arg876His))). Although the variant is predicted to be pathogenic by *in silico* predictors, it has a relatively high AF of 1.70% (gnomAD) and 2.36% (in-house database) and is reported as benign (UV1) in ClinVar. Because the variant was only found in an M0 *SLC26A4* case, a potential digenic inheritance of pathogenic *SLC26A4* variants and the newly identified *EPHA2* variant could not be addressed.

To summarize, the CEVA haplotype or a short CEVA haplotype (V1-CEVA) was detected in 12 of the 28 index cases (16 M1, 12 M0) that were included in our study (Fig. 1). In two individuals (M1), an *SLC26A4* splice variant was identified using WGS. After performing these genetic analyses by which the enrichment of the (V1-)CEVA haplotype in M1 cases was demonstrated, we consider the HL in 12 individuals to be associated with *SLC26A4* defects and these subjects to be genetically explained (2 M2, 10 M1/CEVA), six individuals are considered M1 (4 M1, 2 M0/CEVA), and ten individuals are still considered M0. Additionally, in three individuals (1 M0/CEVA, 2 M0) a potentially pathogenic variant in *FOXI1* was found.

Determination of boundaries of the CEVA haplotype

To identify the true pathogenic defect located on the CEVA haplotype, an in-depth analysis of this genomic region was performed. Firstly, the exact boundaries of the genomic region shared by CEVA haplotype carriers were determined using VNTR marker analysis. For two individuals with the complete CEVA haplotype and the two subjects with the V1-CEVA haplotype, DNA samples of family members were available, allowing reliable determination of the marker alleles located within the haplotype. A shared haplotype of 0.89 Mb delimited by markers D7S501 and D7S2459 was identified (Fig. 2 and **Online Resource Figure S2**). Although the V1-CEVA haplotype shares the marker alleles with the complete CEVA haplotype, the absence of SNPs 1–3 potentially delimits the shared haplotype even more (0.57 Mb, CEVA SNP 3-D7S2459). The remaining eight individuals with the complete CEVA haplotype share identical marker alleles in the 0.89 Mb-sized region, although they could not be conclusively assigned to the haplotype as no segregation analysis could be performed. For individual SLC003, a deviating repeat length was identified for marker D7S2420. As we cannot exclude a rare event to be responsible for the change in allele length, this marker was still considered part of the shared CEVA haplotype.

Short-read WGS did not reveal a pathogenic defect on the CEVA haplotype

To identify the pathogenic defect on the CEVA haplotype, short-read WGS was performed in two individuals (SLC012 & SLC036) carrying the CEVA haplotype *in trans* with a pathogenic variant in *SLC26A4* (M1/CEVA). All heterozygous variants with an AF $\leq 5\%$ in gnomAD that were shared between the two individuals and located within the determined boundaries of the CEVA haplotype were analyzed (**Online Resource Table S7**). In total, 20 shared variants remained and included the 12 original SNPs that previously defined the CEVA haplotype (Chattaraj et al. 2017). Sixteen of the shared variants are located in intronic regions, but for none of them, a significant effect (score ≥ 0.1) on splicing is predicted by SpliceAI. Two variants are located within a *cis* regulatory element of *SLC26A4* according to the GeneHancer database, however, these also show overlap with a long interspersed nuclear element (LINE) repeat element. One variant (CEVA SNP9) has a high nucleotide evolutionary conservation score (PhyloP, 2.769 [range – 14, 3]). No overlapping SVs or CNVs were detected within or overlapping with the CEVA haplotype and shared by the two individuals.

Optical genome mapping & long-read sequencing

To investigate the possibility that SVs were missed using short-read sequencing, optical genome mapping (Bionano Genomics) was performed using ultra-high molecular weight DNA isolated from peripheral blood cells of individual SLC012. Optical genome mapping identified a total of 6,565 SVs, of which none were within the CEVA region (D7S501-D7S2459; chr7:106,440,266 – 107,360,254). Two SV calls (both calling the same 2,196 bp insertion between chr7:107,367,549 and 107,373,585) were located just outside this region (**Online Resource Figure S3A**). This insertion was also called in 100% of our current optical genome mapping control cohort (Levy-Sakin et al. 2019), strongly suggesting that this reflects a reference problem rather than a real SV. Additionally, there were 22 CNV calls, of which none were within the CEVA region.

Subsequently, PacBio long-read sequencing was performed on genomic DNA isolated from individual SLC079. SV analysis of the long-read sequencing data revealed a total of 55,205 SVs, of which 12 within the CEVA region. After careful interrogation of the SVs, all of them were considered false positives based on SV length, and presence of the SVs in an in-house control dataset. The CEVA haplotype region was also manually inspected in the IGV software, which did not reveal any indications for potential SVs (**Online Resource Figure S3B**). Interestingly, the insertion event that was detected with optical genome mapping and located just outside the CEVA region was also present in the long-read sequencing data (chr7:107,370,573, 1,612 bp insertion). This insertion was also present in available in-house control sequencing data, supporting the hypothesis the variant concerns a reference problem and is not a true SV.

A comparable severity of hearing loss in the M1/CEVA and M2 cohorts

As the CEVA haplotype was reported to be associated with a less severe HL phenotype than variants in the protein-coding or splice site regions of *SLC26A4* (Chao et al. 2019), we addressed genotype-phenotype correlations in our cohort. We were able to retrieve pure tone audiometry of all subjects except for SLC071; for this subject, complete audiometry from only one ear was available (**Online Resource Figure S2**). The original CT or MRI scans of subjects SLC018 and SLC032 could not be retrieved. However, written reports of the imaging were available. Data on thyroid gland function were not consistently available and were therefore not included in this study. We applied the methods of Chao et al. 2019 to compare the severity of HL between four subject groups (M0, M1, M1/CEVA, and M2 Fig. 3, Table 4) (Chao et al. 2019). The M1/CEVA group includes the M1/V1-CEVA subjects. Bilateral EVA was present in 7 of 10 (70%) M1/CEVA subjects, in all 4 M1 subjects without the CEVA haplotype, and 7 of 10 (70%) M0 subjects without the CEVA haplotype. All 11 M2 subjects (reference cohort, SLC048 and SLC085) had bilateral EVA. The median pure tone audiometry in the M2 group (85 dB HL, n = 20) was not significantly different from that of the M1/CEVA group (84 dB HL, n = 16) and the M1 group (79.5 dB HL, n = 8) (p-values 0.8300 and 0.7142, respectively, all adjusted for age). Also, no difference was observed between the M1/CEVA group and the M1 group (p = 0.8782). In contrast, when we compare the M2 and M1/CEVA groups with the M0 group, we observed significant differences in the severity of HL (p = 0.0015 and p = 0.0135, respectively). When compared to Chao et al, subjects in our study displayed a similar degree of median HL in the M2 group (86.3 and 85 dB in Chao et al. 2019 and the present study, respectively), more severe HL in the M1/CEVA group (47.5 and 84 dB, respectively) and less severe HL in the M0 group (54.4 and 42 dB HL, respectively). Slight age differences were seen between the groups presented in Chao et al. 2019 and those in the current study (**Online Resource Table S8**). Chao and co-workers did not report audiometric data for the M1 group without the CEVA haplotype *in trans*, presumably due to the small sample size. Overall, in contrast to the study by Chao et al. (2019), the present study showed that subjects with biallelic pathogenic variants in the coding regions and splice sites of *SLC26A4* have a degree of HL that is similar to that of subjects with a monoallelic *SLC26A4* variant and the CEVA haplotype. Due to the small sample size, we could not test the hypothesis that the CEVA haplotype acts as a modifier in M0 subjects as reported previously (Chao et al. 2019).

Table 4
Clinical evaluation of affected individuals

Class	Case	Gender	Age of onset (yr)	Otoscopy examination	Newborn hearing screening	Motor development	Imaging		Audiometric evaluation		
							CT	MRI	Subject age (yr)	PTA (0.5-4kHz)	
M0	SLC014	M	PC	N	3rd time pass	Delayed	bil EVA		4	> 120	55
	SLC015	F	PS	myringosclerosis	N	NR		uni EVA R	25	29	66
	SLC017	M	PC	N	NA	NR	bil EVA		5	10	59
	SLC043	F	PC	N	N	NR	uni EVA L		17	10	95
	SLC052	F	33	N	NA	NR	bil EVA		33	20	> 120
	SLC069	F	15	N	NA	NR	bil EVA		19	43	42
	SLC070	M	PC	N	NA	N		bil EVA	19	43	31
	SLC073	M	U	L: atelectatic middle ear, retracted eardrum	NA	NR		uni EVA R	12	11	18
	SLC084	F	PC	N	3rd time pass	Delayed	bil EVA		6	29	38
	SLC086	M	PC	N	N	Delayed	bil EVA		6	54	41
M0/CEVA	SLC039	F	2-4	N	NA	NR		bil EVA	24	46	70
	SLC080	F	5	N	N	N	bil EVA		6	40	53
M1	SLC002	F	U	N	NA	NR	bil EVA		18	83	76
	SLC018	M	PC	N	NA	N	bil EVA ^a		16	75	101
	SLC032	F	PC	N	NA	NR	bil EVA ^a		59	119	> 120
	SLC045	F	PC	N	N	N	bil EVA		7	20	69

Table 4
(continued): Clinical evaluation of affected individuals

Class	Case	Gender	Age of onset (yr)	Otoscopy examination	Newborn hearing screening	Motor development	Imaging		Audiometric evaluation		
									Subject age (yr)	PTA (0.5-4kHz)	
							CT	MRI		R	L
M1/CEVA	SLC003	M	15	N	NA	NR	uni EVA R		17	83	-1
	SLC012	M	PC	N	NA	NR	bil EVA		17	> 120	53
	SLC013	F	U	N	NA	NR	bil EVA		22	36	9
	SLC031	M	PS	N	2nd time pass	N	uni EVA L		16	0	68
	SLC036	F	PC	N	NA	NR	bil EVA		20	73	68
	SLC040	M	U	N	U	N		uni EVA L	7	5	78
	SLC056	M	PC	N	NA	NR	bil EVA		14	108	93
	SLC071	M	PC	N	N	N	bil EVA		3	85	NT
	SLC078	F	PC	N	NA	U	bil EVA		10	114	93
	SLC079	F	C	N	R	N	bil EVA		2	110	108
M2	SLC048	M	PC	N	NA	NR	bil EVA		8	105	71
	SLC085	M	C	N	R	N	bil EVA		2	23	85
	SLC087	F	C	N	R	Delayed		bil EVA	5	65	63
	SLC088	F	4	N	NA	N		bil EVA	17	85	105
	SLC089	M	U	N	U	U		bil EVA	10	58	98
	SLC090	F	3	N	NA	U		bil EVA	41	116	110
	SLC091	F	2–4	N	N	Delayed	bil EVA		12	63	69
	SLC092	F	C	N	R	N	bil EVA		4	79	48
	SLC093	M	PC	N	U	Delayed		bil EVA	8	71	98
	SLC094	M	PC	sclerotic eardrum L	NA	U	bil EVA		37	101	113
	SLC095	F	C	N	R	N	bil EVA		1	NT	89

Age of onset (AoO), age of onset in years as reported by the subjects. Subject age, the age at which the audiometric data of the last two columns were obtained, in general the last audiogram. Newborn hearing screening was introduced in the Netherlands in 2006. ^aonly written report available. Y, years; PTA, pure tone average, mean of 0.5, 1, 2 and 4 kHz air conduction thresholds; M, male; F, female; R, right; L, left, PC, age of onset of HL is presumably congenital, based on anamnesis; C, age of onset of HL is congenital, based on newborn hearing screening; PS, subject reported onset of HL during primary school, exact age unknown; NR, not reported; NT, not tested; N, no abnormalities; R, refer in newborn hearing screening, failed in test; U, unknown, CT, computed tomography; MRI, magnetic resonance imaging; uni EVA L/R, unilateral enlarged vestibular aqueduct in left or right ear; bil EVA, bilateral enlarged vestibular aqueduct.

Discussion

In this study we investigated 28 genetically unexplained Dutch index cases with HL and a unilateral or bilateral EVA. To elucidate the missing heritability in monoallelic *SLC26A4* cases, who represent 14–31% of subjects with HL and EVA (Azaiez et al. 2007; Mey et al. 2019), extensive genomic analyses as well as

phenotyping were performed. Important findings in this study were (1) the enrichment of a shared (V1-)CEVA haplotype in M1 *SLC26A4* cases, (2) two novel *SLC26A4* splice variants and (3) the identification of a *FOXI1* variant in three subjects suggesting a contribution of this variant to the etiology of HL and EVA. Furthermore, the genotype-phenotype analyses revealed that the severity of the HL associated with biallelic variants (M2) in *SLC26A4* is comparable to the HL associated with a monoallelic variant in *SLC26A4* with or without the CEVA haplotype (M1 and M1/CEVA).

The V1-CEVA haplotype was also previously reported in a single M1 case and suggested to refine the CEVA haplotype or to harbor a different genetic defect (Chattaraj et al. 2017). The shared VNTR marker alleles of the V1-CEVA and the CEVA haplotype suggest that V1-CEVA refines the boundaries of the shared genomic region to 0.57 Mb.

We anticipated that a pathogenic variant co-segregates with the CEVA haplotype. Therefore, we subjected the shared genomic region to extensive genomic analyses that included WES, short- and long-read WGS, and optical genome mapping, to reveal any potential variants missed or misinterpreted in earlier studies. None of the applied sequencing or imaging techniques revealed rare SVs that overlap or are present within the CEVA haplotype. In the light of the proven accuracy and efficacy of especially optical genome mapping and long-read sequencing in SV detection (Chaisson et al. 2019), we deem it unlikely that any SVs within the CEVA region escaped detection. Additionally, we evaluated all SNVs with an AF $\leq 5\%$ (gnomAD) present within the region for predicted regulatory or splice altering effects but for none of the 20 SNVs a potential effect was predicted by SpliceAI. Two SNVs overlap with a potential regulatory element of *SLC26A4* (GeneHancer, EnhancerAtlas), and one variant is present within the intronic regions of this gene. However, all three variants are located within a highly repetitive element (LINE). Although little is known about the effects of genetic variation within LINE elements, a potential effect on the methylation landscape and consequently gene expression levels has been suggested (Xie et al. 2009) and such an effect can therefore not be excluded for the three indicated variants. For the remaining SNVs, no potential effects on transcript splicing or gene regulation were predicted. Nevertheless, we cannot rule out combinatory effects of the SNVs, since they are all located in *cis*. A thorough experimental (multi-omic) analysis is required to optimally assess the effects of the identified variants. RNA studies can be performed to detect quantitative or qualitative changes affecting the *SLC26A4* transcripts. A defect observed on the RNA level could provide valuable insights that may point towards the true pathogenic defect, and prioritize one, or a combination, of the variants on the CEVA allele. However, *SLC26A4* is not or at extremely low levels expressed in readily accessible patient cell types (e.g., fibroblasts and blood cells). The same holds true for induced pluripotent stem cells or otic progenitor cells (Hosoya et al. 2017). However, Hosoya and co-workers have successfully developed a protocol that allows the differentiation of otic progenitor cells into outer sulcus-like cells that express *SLC26A4* at high levels. This protocol could potentially be a powerful tool to evaluate the consequences of CEVA haplotype at the RNA level.

SLC26A4 is not the only gene present within the CEVA haplotype, which also spans *BCAP29*, *COG5*, *DUS4L*, *HBP1*, *PIK3CG*, and *PRKAR2B*. For none of these genes, pathogenic variants associated with (syndromic) HL have been reported, nor has a function in the inner ear been described. The majority of the CEVA-associated SNVs (16/20) are located within an intronic region of these genes, however, for none of these variants a splice altering effect is predicted by SpliceAI.

Since the genetic defect on the CEVA haplotype could not be pinpointed by the genetic analyses, we could not determine whether the AF of the defect is lower than that of the CEVA haplotype and more in line with the expected frequency based on the prevalence of HL (1:1,000 newborns (Morton and Nance 2006)) and the genetic heterogeneity of the condition. Alternatively, the CEVA haplotype could be considered a hypomorphic allele, of which the penetrance depends on the contribution of other co-existing (common) variants.

Not all M0 or M1 *SLC26A4* cases could be genetically explained by the presence of the CEVA haplotype. Therefore, digenic inheritance with variants in *EPHA2*, *FOXI1*, and *KCNJ10* was also explored as a potential explanation for the missing heritability. Digenic inheritance of defects in *SLC26A4* and *EPHA2* has recently been reported in two Japanese Pendred syndrome cases (Li et al. 2020). A c.1063G > A (p.(Gly355Arg)) and a c.1532C > T (p.(Thr511Met)) variant in *EPHA2* were each found 'in trans' with a reported pathogenic variant in *SLC26A4* (Deafness Variation Database (Azaiez et al. 2018)). *EPHA2* was identified as a binding partner of pendrin, with a crucial role in regulating pendrin localization (Li et al. 2020). The identified variants in *EPHA2* were predicted to be pathogenic by several *in silico* predictions tools. However, the c.1532C > T variant has a relatively high allele frequency of 3.03% in the East Asian population, including 11 homozygotes (gnomAD). Yet, in the present study, we did not obtain indications for digenic inheritance of variants in *SLC26A4* and *EPHA2* in subjects with HL and EVA. Besides for *EPHA2*, a digenic mechanism has also been reported and debated for variants in *SLC26A4* and *KCNJ10* or *FOXI1*, with currently no consensus (Jonard et al. 2010; Landa et al. 2013; Pique et al. 2014; Yang et al. 2009; Yang et al. 2007). *FOXI1* is a transcriptional regulator of *SLC26A4* (Yang et al. 2007). We identified a c.677C > T (p.(Thr226Ile)) *FOXI1* variant in three subjects (2 M0/*FOXI1* and 1 M0/CEVA/*FOXI1*). This variant was previously detected in an individual diagnosed with Pendred syndrome and a monoallelic pathogenic *SLC26A4* variant (Pique et al. 2014). The variant has an allele frequency of 0.71% in non-Finnish Europeans (gnomAD) and affects an amino acid residue located outside the DNA-binding domain but close to the nuclear localization signal (NucPred (Brameier et al. 2007)). Previously reported pathogenic *FOXI1* variants have been shown to affect the DNA-binding properties of the protein (Enerbäck et al. 2018). We speculate that a variant affecting the localization motif of the protein could potentially have a loss of function effect as well. Although the variant is classified as benign in the Deafness Variation Database, we identified the variant three times in our cohort of genetically unexplained *SLC26A4* cases and combined with the fact that it has been reported in a previous study (Pique et al. 2014), this suggests that the variant might actually contribute to the etiology of HL and EVA. This is in line with the expansion of the endolymphatic compartment and an audio-vestibular phenotype observed in *Foxi1*^{-/-} mice (Hulander et al. 2003). *In situ* hybridization of the endolymphatic duct and sac of these mice revealed complete absence of *Slc26a4* mRNA expression. Functional studies, among which cellular localizations assays, are warranted to evaluate the effect of the c.677C > T *FOXI1* variant. We did not identify likely pathogenic variants in *KCNJ10* (AF $\leq 5\%$) in our cohort.

WGS did not reveal strong candidate regulatory variants based on data derived from enhancer databases and transcription factor binding site predictions. Nevertheless, interpretation of regulatory variants is still considered complex and is limited by the lack of available epigenetic datasets for the inner ear. Also, no SVs overlapping with *SLC26A4* were detected using WGS, suggesting a limited contribution of SVs to the mutational landscape of *SLC26A4*. This is in line with earlier observations described in literature (Liu et al. 2021; Pique et al. 2014). For the monoallelic cases (M1, M0/CEVA), no long-read sequencing or

optical genome mapping was performed. As it is generally accepted that most SVs could not be accurately detected using short-read sequencing approaches only (Chaisson et al. 2019), it cannot be excluded that causative SVs are present but missed due to technical limitations.

The present study did not confirm that the CEVA allele is associated with a milder HL compared to *SLC26A4* variants affecting the protein-coding sequences, as indicated by Chao et al. (Chao et al. 2019). They discerned a significantly milder HL in their cohort of M1/CEVA subjects (n = 20 ears, median 47.5 dB HL) than we have seen in our cohort of M1/CEVA subjects (n = 16 ears, median 84 dB HL). A possible explanation for this discrepancy could be the progression of HL combined with a ~ 5-year difference in average age between the cohorts (7.5 and 12.8 years, respectively). Progression of HL is seen in up to 39.6% of EVA-ears (Alemi and Chan 2015), with progression rates of ~ 3.5 – ~5.5 dB/y (Govaerts et al. 1999; Jackler and de la Cruz 1989). On the other hand, the older subjects in our M1/CEVA cohort show less severe HL than the younger subjects, which is questioning the relationship with age. Furthermore, there is also an average age difference of 5 years between the M2 groups in both studies (13.2 years and 18.4 years, respectively), while the severity of HL is comparable (85 and 86.3 dB HL, respectively).

The reported variability of the auditory phenotype associated with EVAs (Arjmand and Webber 2004; Gopen et al. 2011; Griffith and Wangemann 2011) may be another explanation for the observed differences in severity of HL in both studies. In literature, many prognostic factors such as genotype, EVA size and morphology, age, head trauma, and gender are reported as underlying explanations for this variability, although some of these studies draw contradicting conclusions (Alemi and Chan 2015; Archibald et al. 2019; Ascha et al. 2017; Gopen et al. 2011; Miyagawa et al. 2014; Rah et al. 2015; Saeed et al. 2021). In the same line, Song et al reported intrafamilial differences in the severity of hearing loss in siblings with the same biallelic variants in *SLC26A4* (Song et al. 2014). Larger sample sizes are needed to confirm or reject the hypothesis that the CEVA haplotype is associated with a milder HL phenotype.

The significant difference in HL severity between the M2 and M1/CEVA groups versus the M0 group suggests that *SLC26A4* defects have a prognostic value which can be strengthened in the future by the identification of the underlying genetic defects in subjects of the M0 group.

In conclusion, the HL and EVA in 12 of the 28 studied subjects could be associated with *SLC26A4*. In addition, we have identified genetic factors that might (partially) explain the phenotype in four additional subjects. However, we could not pinpoint the genetic defect that is present on the CEVA haplotype. The arrival of third-generation sequencing techniques, the expansion of epigenetic and transcriptomic datasets and the increasing understanding of non-coding, structural, and regulatory variants will aid in solving the missing heritability in *SLC26A4* in the coming years. This is of great importance for counseling patients about the underlying cause and expected prognosis of their HL. Furthermore, as variants in *SLC26A4* are a frequent cause of HL (Sloan-Heggen et al. 2016), it is an interesting target for the development of a genetic therapy (Kim et al. 2019). Although the involved molecular defect of the CEVA haplotype is still not resolved, the high prevalence of the CEVA haplotype suggests that a significant portion of monoallelic *SLC26A4* cases can be associated with *SLC26A4* defects by testing for the presence of this haplotype.

Declarations

Acknowledgements

The authors thank Lieke Lamers, Jolinda Put and Evelien Verwoerd for their assistance in genetic analyses and Ronald van Beek, Ellen Kater-Baats, Marcel Nelen and Michiel Oorsprong for technical support. We would like to thank Mieke Wesdorp for including subjects in the study and Galuh D.N. Astuti and Christian Gilissen for their contribution to statistical and bioinformatic analyses. We thank the Radboud Genomics Technology Center, Radboud University Medical Center Nijmegen, for their technical assistance.

Funding

This study was financially supported by a DCMN Radboudumc grant (to H.K. and F.P.M.C.) and by a grant of the Heinsius-Houbolt foundation (to R.J.E.P. and H.K). TM was supported by the Sigrid Jusélius Foundation. This research was part of the Netherlands X-omics Initiative and partially funded by NWO (The Netherlands Organization for Scientific Research; project 184.034.019).

Conflict of interest

The authors have no conflict of interest to declare.

Ethics approval

This study was approved by the medical ethics committee of the Radboud University Medical Center (registration number: NL33648.091.10) and was carried out according to the Declaration of Helsinki.

Consent to participate

Written informed consent was obtained from all participants or their legal representatives.

Consent for publication

Written informed consent was obtained from all participants or their legal representatives.

Availability of data and material

Not applicable

Code availability

Not applicable

Author's contributions

JJS and SEdB co-designed the study and wrote the manuscript. JJS included subjects in the study and collected samples, performed subject evaluation and analyzed audiometric data, SEdB and JO conducted experiments and analyzed genetic sequencing data. CPL and SAHP analyzed audiometric and imaging data and revised the manuscript. LO, TM, RD, AH and KN were involved in long-read sequencing and optical genome mapping, LO, AH and KN revised the manuscript. EdV, FPMC and SR discussed experimental design and critically read and revised the manuscript. HGY analyzed diagnostic WES data and revised the manuscript. RJEP co-designed and supervised the clinical part of the study, and revised the manuscript. HK co-designed the study, supervised the project and wrote the manuscript.

Collaborators

The DOOFNL consortium is a Dutch nationwide collaboration on hereditary hearing loss and consists of M.F. van Dooren, S.G. Kant, H.H.W. de Gier, E.H. Hoefsloot, M.P. van der Schroeff (ErasmusMC, Rotterdam, the Netherlands), L.J.C. Rotteveel, F.G. Ropers (LUMC, Leiden, the Netherlands), J.C.C. Widdershoven, J.R. Hof, E.K. Vanhoutte (MUMC+, Maastricht, the Netherlands), I. Feenstra, H. Kremer, C.P. Lanting, R.J.E. Pennings, H.G. Yntema (Radboudumc, Nijmegen, the Netherlands), R.H. Free and J.S. Klein Wassink-Ruiter (UMCG, Groningen, the Netherlands), R.J. Stokroos, A.L. Smit, M.J. van den Boogaard (UMC, Utrecht, the Netherlands) and F.A. Ebbens, S.M. Maas, A. Plomp, T.P.M. Goderie, P. Merkus and J. van de Kamp (Amsterdam UMC, Amsterdam, the Netherlands).

References

1. Adzhubei IA, Schmidt S, Peshkin L, Ramensky VE, Gerasimova A, Bork P, Kondrashov AS, Sunyaev SR (2010) A method and server for predicting damaging missense mutations. *Nat Methods* 7: 248-9. doi: 10.1038/nmeth0410-248
2. Alemi AS, Chan DK (2015) Progressive Hearing Loss and Head Trauma in Enlarged Vestibular Aqueduct: A Systematic Review and Meta-analysis. *Otolaryngology–Head and Neck Surgery* 153: 512-517. doi: 10.1177/0194599815596343
3. Andersson R, Gebhard C, Miguel-Escalada I, Hoof I, Bornholdt J, Boyd M, Chen Y, Zhao X, Schmidl C, Suzuki T, Ntini E, Arner E, Valen E, Li K, Schwarzfischer L, Glatz D, Raithel J, Lilje B, Rapin N, Bagger FO, Jørgensen M, Andersen PR, Bertin N, Rackham O, Burroughs AM, Baillie JK, Ishizu Y, Shimizu Y, Furuhashi E, Maeda S, Negishi Y, Mungall CJ, Meehan TF, Lassmann T, Itoh M, Kawaji H, Kondo N, Kawai J, Lennartsson A, Daub CO, Heutink P, Hume DA, Jensen TH, Suzuki H, Hayashizaki Y, Müller F, Forrest ARR, Carninci P, Rehli M, Sandelin A, The FC (2014) An atlas of active enhancers across human cell types and tissues. *Nature* 507: 455-461. doi: 10.1038/nature12787
4. Archibald HD, Ascha M, Gupta A, Megerian C, Otteson T (2019) Hearing loss in unilateral and bilateral enlarged vestibular aqueduct syndrome. *International Journal of Pediatric Otorhinolaryngology* 118: 147-151. doi: 10.1016/j.ijporl.2018.12.023
5. Arjmand EM, Webber A (2004) Audiometric findings in children with a large vestibular aqueduct. *Arch Otolaryngol Head Neck Surg* 130: 1169-74. doi: 10.1001/archotol.130.10.1169
6. Ascha MS, Manzoor N, Gupta A, Semaan M, Megerian C, Otteson TD (2017) Vestibular Aqueduct Midpoint Width and Hearing Loss in Patients With an Enlarged Vestibular Aqueduct. *JAMA otolaryngology–head & neck surgery* 143: 601-608. doi: 10.1001/jamaoto.2016.4522
7. Azaiez H, Booth KT, Ephraim SS, Crone B, Black-Ziegelbein EA, Marini RJ, Shearer AE, Sloan-Heggen CM, Kolbe D, Casavant T, Schnieders MJ, Nishimura C, Braun T, Smith RJH (2018) Genomic Landscape and Mutational Signatures of Deafness-Associated Genes. *Am J Hum Genet* 103: 484-497. doi: 10.1016/j.ajhg.2018.08.006
8. Azaiez H, Yang T, Prasad S, Sorensen JL, Nishimura CJ, Kimberling WJ, Smith RJ (2007) Genotype-phenotype correlations for SLC26A4-related deafness. *Hum Genet* 122: 451-7. doi: 10.1007/s00439-007-0415-2
9. Boeva V, Popova T, Bleakley K, Chiche P, Cappel J, Schleiermacher G, Janoueix-Lerosey I, Delattre O, Barillot E (2012) Control-FREEC: a tool for assessing copy number and allelic content using next-generation sequencing data. *Bioinformatics* 28: 423-425. doi: 10.1093/bioinformatics/btr670
10. Boston M, Halsted M, Meinzen-Derr J, Bean J, Vijayasekaran S, Arjmand E, Choo D, Benton C, Greinwald J (2007) The large vestibular aqueduct: a new definition based on audiologic and computed tomography correlation. *Otolaryngol Head Neck Surg* 136: 972-7. doi: 10.1016/j.otohns.2006.12.011
11. Brameier M, Krings A, MacCallum RM (2007) NucPred—Predicting nuclear localization of proteins. *Bioinformatics* 23: 1159-1160. doi: 10.1093/bioinformatics/btm066
12. Chaisson MJP, Sanders AD, Zhao X, Malhotra A, Porubsky D, Rausch T, Gardner EJ, Rodriguez OL, Guo L, Collins RL, Fan X, Wen J, Handsaker RE, Fairley S, Kronenberg ZN, Kong X, Hormozdiari F, Lee D, Wenger AM, Hastie AR, Antaki D, Anantharaman T, Audano PA, Brand H, Cantsilieris S, Cao H, Cerveira E, Chen C, Chen X, Chin C-S, Chong Z, Chuang NT, Lambert CC, Church DM, Clarke L, Farrell A, Flores J, Galeev T, Gorkin DU, Gujral M, Guryev V, Heaton WH, Korlach J, Kumar S, Kwon JY, Lam ET, Lee JE, Lee J, Lee W-P, Lee SP, Li S, Marks P, Viaud-Martinez K, Meiers S, Munson KM, Navarro FCP, Nelson BJ, Nodzak C, Noor A, Kyriazopoulou-Panagiotopoulou S, Pang AWC, Qiu Y, Rosanio G, Ryan M, Stütz A, Spierings DCJ, Ward A, Welch AE, Xiao M, Xu W, Zhang C, Zhu Q, Zheng-Bradley X, Lowy E, Yakneen S, McCarroll S, Jun G, Ding L, Koh CL, Ren B, Flicek P, Chen K, Gerstein MB, Kwok P-Y, Lansdorp PM, Marth GT, Sebat J, Shi X, Bashir A, Ye K, Devine SE, Talkowski ME, Mills RE, Marschall T, Korbel JO, Eichler EE, Lee C (2019) Multi-platform discovery of haplotype-resolved structural variation in human genomes. *Nature Communications* 10: 1784. doi: 10.1038/s41467-018-08148-z
13. Chao JR, Chattaraj P, Munjal T, Honda K, King KA, Zalewski CK, Chien WW, Brewer CC, Griffith AJ (2019) SLC26A4-linked CEVA haplotype correlates with phenotype in patients with enlargement of the vestibular aqueduct. *BMC Medical Genetics* 20: 118. doi: 10.1186/s12881-019-0853-4

14. Chattaraj P, Munjal T, Honda K, Rendtorff ND, Ratay JS, Muskett JA, Risso DS, Roux I, Gertz EM, Schaffer AA, Friedman TB, Morell RJ, Tranebjaerg L, Griffith AJ (2017) A common SLC26A4-linked haplotype underlying non-syndromic hearing loss with enlargement of the vestibular aqueduct. *J Med Genet* 54: 665-673. doi: 10.1136/jmedgenet-2017-104721
15. Chen X, Schulz-Trieglaff O, Shaw R, Barnes B, Schlesinger F, Källberg M, Cox AJ, Kruglyak S, Saunders CT (2016) Manta: rapid detection of structural variants and indels for germline and cancer sequencing applications. *Bioinformatics* 32: 1220-1222. doi: 10.1093/bioinformatics/btv710
16. Choi BY, Madeo AC, King KA, Zalewski CK, Pryor SP, Muskett JA, Nance WE, Butman JA, Brewer CC, Griffith AJ (2009) Segregation of enlarged vestibular aqueducts in families with non-diagnostic SLC26A4 genotypes. *J Med Genet* 46: 856-61. doi: 10.1136/jmg.2009.067892
17. de Bruijn SE, Smits JJ, Liu C, Lanting CP, Beynon AJ, Blankevoort J, Oostrik J, Koole W, de Vrieze E, Cremers CWRJ, Cremers FPM, Roosing S, Yntema HG, Kunst HPM, Zhao B, Pennings RJE, Kremer H (2021) A RIPOR2 in-frame deletion is a frequent and highly penetrant cause of adult-onset hearing loss. *Journal of Medical Genetics* 58: 96. doi: 10.1136/jmedgenet-2020-106863
18. Dimitrova S, Bucher P (2013) UCNEbase—a database of ultraconserved non-coding elements and genomic regulatory blocks. *Nucleic Acids Research* 41: D101-D109. doi: 10.1093/nar/gks1092
19. Dou H, Xu J, Wang Z, Smith AN, Soleimani M, Karet FE, Greinwald JH, Jr., Choo D (2004) Co-expression of pendrin, vacuolar H⁺-ATPase alpha4-subunit and carbonic anhydrase II in epithelial cells of the murine endolymphatic sac. *J Histochem Cytochem* 52: 1377-84. doi: 10.1177/002215540405201014
20. Dreos R, Ambrosini G, Cavin Périer R, Bucher P (2013) EPD and EPDnew, high-quality promoter resources in the next-generation sequencing era. *Nucleic Acids Research* 41: D157-D164. doi: 10.1093/nar/gks1233
21. Dunham I, Kundaje A, Aldred SF, Collins PJ, Davis CA, Doyle F, Epstein CB, Frietze S, Harrow J, Kaul R, Khatun J, Lajoie BR, Landt SG, Lee B-K, Pauli F, Rosenbloom KR, Sabo P, Safi A, Sanyal A, Shoresh N, Simon JM, Song L, Trinklein ND, Altschuler RC, Birney E, Brown JB, Cheng C, Djebali S, Dong X, Dunham I, Ernst J, Furey TS, Gerstein M, Giardine B, Greven M, Hardison RC, Harris RS, Herrero J, Hoffman MM, Iyer S, Kellis M, Khatun J, Kheradpour P, Kundaje A, Lassmann T, Li Q, Lin X, Marinov GK, Merkel A, Mortazavi A, Parker SCJ, Reddy TE, Rozowsky J, Schlesinger F, Thurman RE, Wang J, Ward LD, Whitfield TW, Wilder SP, Wu W, Xi HS, Yip KY, Zhuang J, Bernstein BE, Birney E, Dunham I, Green ED, Gunter C, Snyder M, Pazin MJ, Lowdon RF, Dillon LAL, Adams LB, Kelly CJ, Zhang J, Wexler JR, Green ED, Good PJ, Feingold EA, Bernstein BE, Birney E, Crawford GE, Dekker J, Eltnitski L, Farnham PJ, Gerstein M, Giddings MC, Gingeras TR, Green ED, Guigó R, Hardison RC, Hubbard TJ, Kellis M, Kent WJ, Lieb JD, Margulies EH, Myers RM, Snyder M, Stamatoyannopoulos JA, Tenenbaum SA, et al. (2012) An integrated encyclopedia of DNA elements in the human genome. *Nature* 489: 57-74. doi: 10.1038/nature11247
22. Enerbäck S, Nilsson D, Edwards N, Heglin M, Alkanderi S, Ashton E, Deeb A, Kokash FEB, Bakhsh ARA, Van't Hoff W, Walsh SB, D'Arco F, Daryadel A, Bourgeois S, Wagner CA, Kleta R, Bockenhauer D, Sayer JA (2018) Acidosis and Deafness in Patients with Recessive Mutations in FOXI1. *Journal of the American Society of Nephrology : JASN* 29: 1041-1048. doi: 10.1681/ASN.2017080840
23. Everett LA, Glaser B, Beck JC, Idol JR, Buchs A, Heyman M, Adawi F, Hazani E, Nassir E, Baxeavanis AD, Sheffield VC, Green ED (1997) Pendred syndrome is caused by mutations in a putative sulphate transporter gene (PDS). *Nat Genet* 17: 411-22. doi: 10.1038/ng1297-411
24. Everett LA, Morsli H, Wu DK, Green ED (1999) Expression pattern of the mouse ortholog of the Pendred's syndrome gene (Pds) suggests a key role for pendrin in the inner ear. *Proceedings of the National Academy of Sciences of the United States of America* 96: 9727-9732. doi: 10.1073/pnas.96.17.9727
25. Fishilevich S, Nudel R, Rappaport N, Hadar R, Plaschkes I, Iny Stein T, Rosen N, Kohn A, Twik M, Safran M, Lancet D, Cohen D (2017) GeneHancer: genome-wide integration of enhancers and target genes in GeneCards. *Database (Oxford)* 2017: bax028. doi: 10.1093/database/bax028
26. Fornes O, Castro-Mondragon JA, Khan A, van der Lee R, Zhang X, Richmond PA, Modi BP, Correard S, Gheorghe M, Baranašić D, Santana-Garcia W, Tan G, Chèneby J, Ballester B, Parcy F, Sandelin A, Lenhard B, Wasserman WW, Mathelier A (2020) JASPAR 2020: update of the open-access database of transcription factor binding profiles. *Nucleic Acids Research* 48: d87-d92. doi: 10.1093/nar/gkz1001
27. Fraser GR (1965) Association of Congenital Deafness with Goitre (Pendred's Syndrome): A Study of 207 Families. *Ann Hum Genet* 28: 201-49. doi: 10.1111/j.1469-1809.1964.tb00479.x
28. Gao T, Qian J (2020) EnhancerAtlas 2.0: an updated resource with enhancer annotation in 586 tissue/cell types across nine species. *Nucleic Acids Research* 48: d58-d64. doi: 10.1093/nar/gkz980
29. Genomes Project Consortium, Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM, Korbel JO, Marchini JL, McCarthy S, McVean GA, Abecasis GR (2015) A global reference for human genetic variation. *Nature* 526: 68-74. doi: 10.1038/nature15393
30. Gopen Q, Zhou G, Whittemore K, Kenna M (2011) Enlarged vestibular aqueduct: Review of controversial aspects. *The Laryngoscope* 121: 1971-1978. doi: <https://doi.org/10.1002/lary.22083>
31. Govaerts PJ, Casselman J, Daemers K, De Ceulaer G, Somers T, Offeciers FE (1999) Audiological findings in large vestibular aqueduct syndrome. *Int J Pediatr Otorhinolaryngol* 51: 157-64. doi: 10.1016/s0165-5876(99)00268-2
32. Griffith AJ, Wangemann P (2011) Hearing loss associated with enlargement of the vestibular aqueduct: mechanistic insights from clinical phenotypes, genotypes, and mouse models. *Hearing research* 281: 11-17. doi: 10.1016/j.heares.2011.05.009
33. Hosoya M, Fujioka M, Sone T, Okamoto S, Akamatsu W, Ukai H, Ueda HR, Ogawa K, Matsunaga T, Okano H (2017) Cochlear Cell Modeling Using Disease-Specific iPSCs Unveils a Degenerative Phenotype and Suggests Treatments for Congenital Progressive Hearing Loss. *Cell Rep* 18: 68-81. doi: 10.1016/j.celrep.2016.12.020
34. Hulander M, Kiernan AE, Blomqvist SR, Carlsson P, Samuelsson EJ, Johansson BR, Steel KP, Enerbäck S (2003) Lack of pendrin expression leads to deafness and expansion of the endolymphatic compartment in inner ears of Foxi1 null mutant mice. *Development* 130: 2013-25. doi: 10.1242/dev.00376
35. Jackler RK, de la Cruz A (1989) The large vestibular aqueduct syndrome. *The Laryngoscope* 99: 1238-1243. doi: 10.1288/00005537-198912000-00006

36. Jaganathan K, Kyriazopoulou Panagiotopoulou S, McRae JF, Darbandi SF, Knowles D, Li YI, Kosmicki JA, Arbelaez J, Cui W, Schwartz GB, Chow ED, Kanterakis E, Gao H, Kia A, Batzoglu S, Sanders SJ, Farh KK-H (2019) Predicting Splicing from Primary Sequence with Deep Learning. *Cell* 176: 535-548.e24. doi: 10.1016/j.cell.2018.12.015
37. Jonard L, Niasme-Grare M, Bonnet C, Feldmann D, Rouillon I, Loundon N, Calais C, Catros H, David A, Dollfus H, Drouin-Garraud V, Duriez F, Eliot MM, Fellmann F, Francannet C, Gilbert-Dussardier B, Gohler C, Goizet C, Journal H, Mom T, Thuillier-Obstoy M-F, Couderc R, Garabédian EN, Denoyelle F, Marlin S (2010) Screening of SLC26A4, FOXI1 and KCNJ10 genes in unilateral hearing impairment with ipsilateral enlarged vestibular aqueduct. *International Journal of Pediatric Otorhinolaryngology* 74: 1049-1053. doi: <https://doi.org/10.1016/j.ijporl.2010.06.002>
38. Karczewski KJ, Francioli LC, Tiao G, Cummings BB, Alfoldi J, Wang Q, Collins RL, Laricchia KM, Ganna A, Birnbaum DP, Gauthier LD, Brand H, Solomonson M, Watts NA, Rhodes D, Singer-Berk M, England EM, Seaby EG, Kosmicki JA, Walters RK, Tashman K, Farjoun Y, Banks E, Poterba T, Wang A, Seed C, Whiffin N, Chong JX, Samocha KE, Pierce-Hoffman E, Zappala Z, O'Donnell-Luria AH, Minikel EV, Weisburd B, Lek M, Ware JS, Vittal C, Armean IM, Bergelson L, Cibulskis K, Connolly KM, Covarrubias M, Donnelly S, Ferreira S, Gabriel S, Gentry J, Gupta N, Jeandet T, Kaplan D, Llanwarne C, Munshi R, Novod S, Petrillo N, Roazen D, Ruano-Rubio V, Saltzman A, Schleicher M, Soto J, Tibbetts K, Tolonen C, Wade G, Talkowski ME, Neale BM, Daly MJ, MacArthur DG (2019) Variation across 141,456 human exomes and genomes reveals the spectrum of loss-of-function intolerance across human protein-coding genes. *bioRxiv*: 531210. doi: 10.1101/531210
39. Kent WJ, Sugnet CW, Furey TS, Roskin KM, Pringle TH, Zahler AM, Haussler D (2002) The human genome browser at UCSC. *Genome Res* 12: 996-1006. doi: 10.1101/gr.229102
40. Khan A, Zhang X (2016) dbSUPER: a database of super-enhancers in mouse and human genome. *Nucleic Acids Research* 44: D164-D171. doi: 10.1093/nar/gkv1002
41. Khan M, Cornelis SS, Pozo-Valero MD, Whelan L, Runhart EH, Mishra K, Bults F, AlSwaiti Y, AlTalibishi A, De Baere E, Banfi S, Banin E, Bauwens M, Ben-Yosef T, Boon CJF, van den Born LI, Defoort S, Devos A, Dockery A, Dudakova L, Fakin A, Farrar GJ, Sallum JMF, Fujinami K, Gilissen C, Glavač D, Gorin MB, Greenberg J, Hayashi T, Hettinga YM, Hoischen A, Hoyng CB, Hufendiek K, Jäggle H, Kamakari S, Karali M, Kellner U, Klaver CCW, Kousal B, Lamey TM, MacDonald IM, Matynia A, McLaren TL, Mena MD, Meunier I, Miller R, Newman H, Ntozini B, Oldak M, Pieterse M, Podhajcer OL, Puech B, Ramesar R, Rütter K, Salameh M, Salles MV, Sharon D, Simonelli F, Spital G, Steehouwer M, Szaflik JP, Thompson JA, Thuillier C, Tracewska AM, van Zweeden M, Vincent AL, Zanlonghi X, Liskova P, Stöhr H, Roach JND, Ayuso C, Roberts L, Weber BHF, Dhaenens CM, Cremers FPM (2020) Resolving the dark matter of ABCA4 for 1054 Stargardt disease probands through integrated genomics and transcriptomics. *Genetics in Medicine* 22: 1235-1246. doi: 10.1038/s41436-020-0787-4
42. Kim M-A, Kim SH, Ryu N, Ma J-H, Kim Y-R, Jung J, Hsu C-J, Choi JY, Lee K-Y, Wangemann P, Bok J, Kim U-K (2019) Gene therapy for hereditary hearing loss by SLC26A4 mutations in mice reveals distinct functional roles of pendrin in normal hearing. *Theranostics* 9: 7184-7199. doi: 10.7150/thno.38032
43. Kircher M, Witten DM, Jain P, O'Roak BJ, Cooper GM, Shendure J (2014) A general framework for estimating the relative pathogenicity of human genetic variants. *Nature genetics* 46: 310-315. doi: 10.1038/ng.2892
44. Krumm N, Sudmant PH, Ko A, O'Roak BJ, Malig M, Coe BP, Project NES, Quinlan AR, Nickerson DA, Eichler EE (2012) Copy number variation detection and genotyping from exome sequence data. *Genome Res* 22: 1525-32. doi: 10.1101/gr.138115.112
45. Landa P, Differ A-M, Rajput K, Jenkins L, Bitner-Glindzic M (2013) Lack of significant association between mutations of KCNJ10 or FOXI1 and SLC26A4 mutations in pendred syndrome/enlarged vestibular aqueducts. *BMC Medical Genetics* 14: 85. doi: 10.1186/1471-2350-14-85
46. Landrum MJ, Lee JM, Benson M, Brown GR, Chao C, Chitipiralla S, Gu B, Hart J, Hoffman D, Jang W, Karapetyan K, Katz K, Liu C, Maddipati Z, Malheiro A, McDaniel K, Ovetsky M, Riley G, Zhou G, Holmes JB, Kattman BL, Maglott DR (2018) ClinVar: improving access to variant interpretations and supporting evidence. *Nucleic Acids Res* 46: D1062-D1067. doi: 10.1093/nar/gkx1153
47. Lee HJ, Jung J, Shin JW, Song MH, Kim SH, Lee JH, Lee KA, Shin S, Kim UK, Bok J, Lee KY, Choi JY, Park HJ (2014) Correlation between genotype and phenotype in patients with bi-allelic SLC26A4 mutations. *Clin Genet* 86: 270-5. doi: 10.1111/cge.12273
48. Lenth RV (2016) Least-Squares Means: The R Package lsmeans. *Journal of Statistical Software* 69: 1-33. doi: 10.18637/jss.v069.i01
49. Levy-Sakin M, Pastor S, Mostovoy Y, Li L, Leung AKY, McCaffrey J, Young E, Lam ET, Hastie AR, Wong KHY, Chung CYL, Ma W, Sibert J, Rajagopalan R, Jin N, Chow EYC, Chu C, Poon A, Lin C, Naguib A, Wang W-P, Cao H, Chan T-F, Yip KY, Xiao M, Kwok P-Y (2019) Genome maps across 26 human populations reveal population-specific patterns of structural variation. *Nature Communications* 10: 1025. doi: 10.1038/s41467-019-08992-7
50. Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25: 1754-60. doi: 10.1093/bioinformatics/btp324
51. Li M, Nishio S-y, Naruse C, Riddell M, Sapski S, Katsuno T, Hikita T, Mizapourshafiyi F, Smith FM, Cooper LT, Lee MG, Asano M, Boettger T, Krueger M, Wietelmann A, Graumann J, Day BW, Boyd AW, Offermanns S, Kitajiri S-i, Usami S-i, Nakayama M (2020) Digenic inheritance of mutations in EPHA2 and SLC26A4 in Pendred syndrome. *Nature Communications* 11: 1343. doi: 10.1038/s41467-020-15198-9
52. Liu YL, Wang LL, Wen J, Mei LY, He CF, Jiang L, Feng Y (2021) Application value of high-throughput gene copy number variation detection in the diagnosis of enlarged vestibular aqueduct. *Zhonghua Yi Xue Za Zhi* 101: 103-107. doi: 10.3760/cma.j.cn112137-20201010-02794
53. Mantere T, Neveling K, Pebrel-Richard C, Benoist M, van der Zande G, Kater-Baats E, Baatout I, van Beek R, Yammine T, Oorsprong M, Olde-Weghuis D, Majdali W, Vermeulen S, Pauper M, Lebbar A, Stevens-Kroef M, Sanlaville D, Smeets D, Dupont JM, Hoischen A, Schluth-Boland C, El Khattabi L (2020) Next generation cytogenetics: genome-imaging enables comprehensive structural variant detection for 100 constitutional chromosomal aberrations in 85 samples. *bioRxiv*: 2020.07.15.205245. doi: 10.1101/2020.07.15.205245
54. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernysky A, Garimella K, Altshuler D, Gabriel S, Daly M, DePristo MA (2010) The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome research* 20: 1297-1303. doi: 10.1101/gr.107524.110

55. Mey K, Muhamad AA, Tranebjaerg L, Rendtorff ND, Rasmussen SH, Bille M, Cayé-Thomasen P (2019) Association of SLC26A4 mutations, morphology, and hearing in pendred syndrome and NSEVA. *Laryngoscope* 129: 2574-2579. doi: 10.1002/lary.27319
56. Miyagawa M, Nishio S-y, Usami S-i, The Deafness Gene Study C (2014) Mutation spectrum and genotype–phenotype correlation of hearing loss patients caused by SLC26A4 mutations in the Japanese: a large cohort study. *Journal of Human Genetics* 59: 262-268. doi: 10.1038/jhg.2014.12
57. Morton CC, Nance WE (2006) Newborn hearing screening—a silent revolution. *N Engl J Med* 354: 2151-64. doi: 10.1056/NEJMra050700
58. Neveling K, Mantere T, Vermeulen S, Oorsprong M, van Beek R, Kater-Baats E, Pauper M, van der Zande G, Smeets D, Weghuis DO, Stevens-Kroef MJ, Hoischen A (2020) Next generation cytogenetics: comprehensive assessment of 48 leukemia genomes by genome imaging. *bioRxiv*: 2020.02.06.935742. doi: 10.1101/2020.02.06.935742
59. Neveling K, Mensenkamp AR, Derks R, Kwint M, Ouchene H, Steehouwer M, van Lier B, Bosgoed E, Rikken A, Tychon M, Zafeiropoulou D, Castelein S, Hehir-Kwa J, Tjwan Thung D, Hofste T, Lelieveld SH, Bertens SM, Adan IB, Eijkelenboom A, Tops BB, Yntema H, Stokowy T, Knappskog PM, Høberg-Vetti H, Steen VM, Boyle E, Martin B, Ligtenberg MJ, Shendure J, Nelen MR, Hoischen A (2017) BRCA Testing by Single-Molecule Molecular Inversion Probes. *Clin Chem* 63: 503-512. doi: 10.1373/clinchem.2016.263897
60. Onk AM, Leijendeckers JM, Huygen PL, Schraders M, del Campo M, del Castillo I, Tekin M, Feenstra I, Beynon AJ, Kunst HP, Snik AF, Kremer H, Admiraal RJ, Pennings RJ (2014) Similar phenotypes caused by mutations in OTOG and OTOGL. *Ear Hear* 35: e84-91. doi: 10.1097/aud.0000000000000008
61. Pedemonte N, Caci E, Sondo E, Caputo A, Rhoden K, Pfeffer U, Di Candia M, Bandettini R, Ravazzolo R, Zegar-Moran O, Galletta LJ (2007) Thiocyanate transport in resting and IL-4-stimulated human bronchial epithelial cells: role of pendrin and anion channels. *J Immunol* 178: 5144-53. doi: 10.4049/jimmunol.178.8.5144
62. Pedersen BS, Quinlan AR (2018) Mosdepth: quick coverage calculation for genomes and exomes. *Bioinformatics* 34: 867-868. doi: 10.1093/bioinformatics/btx699
63. Pique LM, Brennan ML, Davidson CJ, Schaefer F, Greinwald J, Jr., Schrijver I (2014) Mutation analysis of the SLC26A4, FOXI1 and KCNJ10 genes in individuals with congenital hearing loss. *PeerJ* 2: e384. doi: 10.7717/peerj.384
64. Pryor SP, Madero AC, Reynolds JC, Sarlis NJ, Arnos KS, Nance WE, Yang Y, Zalewski CK, Brewer CC, Butman JA, Griffith AJ (2005) SLC26A4/PDS genotype-phenotype correlation in hearing loss with enlargement of the vestibular aqueduct (EVA): evidence that Pendred syndrome and non-syndromic EVA are distinct clinical and genetic entities. *Journal of Medical Genetics* 42: 159-165. doi: 10.1136/jmg.2004.024208
65. Rah YC, Kim AR, Koo J-W, Lee JH, Oh S-h, Choi BY (2015) Audiologic presentation of enlargement of the vestibular aqueduct according to the SLC26A4 genotypes. *The Laryngoscope* 125: E216-E222. doi: 10.1002/lary.25079
66. Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, Mesirov JP (2011) Integrative genomics viewer. *Nature biotechnology* 29: 24-26. doi: 10.1038/nbt.1754
67. Royaux IE, Belyantseva IA, Wu T, Kachar B, Everett LA, Marcus DC, Green ED (2003) Localization and functional studies of pendrin in the mouse inner ear provide insight about the etiology of deafness in pendred syndrome. *J Assoc Res Otolaryngol* 4: 394-404. doi: 10.1007/s10162-002-3052-4
68. Royaux IE, Suzuki K, Mori A, Katoh R, Everett LA, Kohn LD, Green ED (2000) Pendrin, the protein encoded by the Pendred syndrome gene (PDS), is an apical porter of iodide in the thyroid and is regulated by thyroglobulin in FRTL-5 cells. *Endocrinology* 141: 839-45. doi: 10.1210/endo.141.2.7303
69. Royaux IE, Wall SM, Karniski LP, Everett LA, Suzuki K, Knepper MA, Green ED (2001) Pendrin, encoded by the Pendred syndrome gene, resides in the apical region of renal intercalated cells and mediates bicarbonate secretion. *Proc Natl Acad Sci U S A* 98: 4221-6. doi: 10.1073/pnas.071516798
70. Saeed HS, Kenth J, Black G, Saeed SR, Stivaros S, Bruce IA (2021) Hearing Loss in Enlarged Vestibular Aqueduct: A Prognostic Factor Systematic Review of the Literature. *Otology & Neurotology* 42. doi: 10.1097/MAO.0000000000002843
71. Sangermano R, Khan M, Cornelis SS, Richelle V, Albert S, Garanto A, Elmelik D, Qamar R, Lugtenberg D, van den Born LI, Collin RWJ, Cremers FPM (2018) ABCA4 midgenes reveal the full splice spectrum of all reported noncanonical splice site variants in Stargardt disease. *Genome Res* 28: 100-110. doi: 10.1101/gr.226621.117
72. Schwarz JM, Cooper DN, Schuelke M, Seelow D (2014) MutationTaster2: mutation prediction for the deep-sequencing age. *Nature Methods* 11: 361. doi: 10.1038/nmeth.2890
73. Sloan-Heggen CM, Bierer AO, Shearer AE, Kolbe DL, Nishimura CJ, Frees KL, Ephraim SS, Shibata SB, Booth KT, Campbell CA, Ranum PT, Weaver AE, Black-Ziegelbein EA, Wang D, Azaiez H, Smith RJH (2016) Comprehensive genetic testing in the clinical evaluation of 1119 patients with hearing loss. *Hum Genet* 135: 441-450. doi: 10.1007/s00439-016-1648-8
74. Song MH, Shin J-W, Park H-J, Lee K-A, Kim Y, Kim U-K, Jeon JH, Choi JY (2014) Intrafamilial phenotypic variability in families with biallelic SLC26A4 mutations. *The Laryngoscope* 124: E194-E202. doi: 10.1002/lary.24504
75. Suzuki H, Oshima A, Tsukamoto K, Abe S, Kumakawa K, Nagai K, Satoh H, Kanda Y, Iwasaki S, Usami S (2007) Clinical characteristics and genotype-phenotype correlation of hearing loss patients with SLC26A4 mutations. *Acta Otolaryngol* 127: 1292-7. doi: 10.1080/00016480701258739
76. van Beeck Calkoen EA, Merkus P, Goverts ST, van de Kamp JM, Mulder MF, Sanchez Aliaga E, Hensen EF (2018) Evaluation of the outcome of CT and MR imaging in pediatric patients with bilateral sensorineural hearing loss. *Int J Pediatr Otorhinolaryngol* 108: 180-185. doi: 10.1016/j.ijporl.2018.02.022
77. van Beeck Calkoen EA, Sanchez Aliaga E, Merkus P, Smit CF, van de Kamp JM, Mulder MF, Goverts ST, Hensen EF (2017) High prevalence of abnormalities on CT and MR imaging in children with unilateral sensorineural hearing loss irrespective of age or degree of hearing loss. *Int J Pediatr Otorhinolaryngol* 97: 185-191. doi: 10.1016/j.ijporl.2017.04.002
78. Van Camp G, Smith R Hereditary Hearing Loss Homepage. Available from: <https://hereditaryhearingloss.org>.
79. van der Ploeg CP, Uilenburg NN, Kauffman-de Boer MA, Oudesluys-Murphy AM, Verkerk PH (2012) Newborn hearing screening in youth health care in the Netherlands: National results of implementation and follow-up. *Int J Audiol* 51: 584-90. doi: 10.3109/14992027.2012.684402

80. Vaser R, Adusumalli S, Leng SN, Sikic M, Ng PC (2015) SIFT missense predictions for genomes. *Nature Protocols* 11: 1. doi: 10.1038/nprot.2015.123
81. Visel A, Minovitsky S, Dubchak I, Pennacchio LA (2007) VISTA Enhancer Browser—a database of tissue-specific human enhancers. *Nucleic Acids Research* 35: D88-D92. doi: 10.1093/nar/gkl822
82. Wangemann P (2011) The role of pendrin in the development of the murine inner ear. *Cell Physiol Biochem* 28: 527-34. doi: 10.1159/000335113
83. Wangemann P, Nakaya K, Wu T, Maganti RJ, Itza EM, Sanneman JD, Harbidge DG, Billings S, Marcus DC (2007) Loss of cochlear HCO₃⁻ secretion causes deafness via endolymphatic acidification and inhibition of Ca²⁺ reabsorption in a Pendred syndrome mouse model. *Am J Physiol Renal Physiol* 292: F1345-53. doi: 10.1152/ajprenal.00487.2006
84. Wilderman A, VanOudenhove J, Kron J, Noonan JP, Cotney J (2018) High-Resolution Epigenomic Atlas of Human Embryonic Craniofacial Development. *Cell Reports* 23: 1581-1597. doi: 10.1016/j.celrep.2018.03.129
85. Xie H, Wang M, Bischof J, Bonaldo MdF, Soares MB (2009) SNP-based prediction of the human germ cell methylation landscape. *Genomics* 93: 434-440. doi: 10.1016/j.ygeno.2009.01.005
86. Yang T, Gurrola JG, Wu H, Chiu SM, Wangemann P, Snyder PM, Smith RJH (2009) Mutations of KCNJ10 Together with Mutations of SLC26A4 Cause Digenic Nonsyndromic Hearing Loss Associated with Enlarged Vestibular Aqueduct Syndrome. *The American Journal of Human Genetics* 84: 651-657. doi: 10.1016/j.ajhg.2009.04.014
87. Yang T, Vidarsson H, Rodrigo-Blomqvist S, Rosengren SS, Enerback S, Smith RJ (2007) Transcriptional control of SLC26A4 is involved in Pendred syndrome and nonsyndromic enlargement of vestibular aqueduct (DFNB4). *Am J Hum Genet* 80: 1055-63. doi: 10.1086/518314
88. Yazar KO, Duman D, Zazo Seco C, Dallman J, Huang M, Peters TA, Sirmaci A, Lu N, Schraders M, Skromne I, Oostrik J, Diaz-Horta O, Young JI, Tokgoz-Yilmaz S, Konukseven O, Shahin H, Hettterschijs L, Kanaan M, Oonk AM, Edwards YJ, Li H, Atalay S, Blanton S, Desmidt AA, Liu XZ, Pennings RJ, Lu Z, Chen ZY, Kremer H, Tekin M (2012) Mutations in OTOGL, encoding the inner ear protein otogelin-like, cause moderate sensorineural hearing loss. *Am J Hum Genet* 91: 872-82. doi: 10.1016/j.ajhg.2012.09.011
89. Zerbino DR, Wilder SP, Johnson N, Juettemann T, Flicek PR (2015) The Ensembl Regulatory Build. *Genome Biology* 16: 56. doi: 10.1186/s13059-015-0621-5

Figures

Figure 1

Overview of genetic analyses performed in zeroallelic and monoallelic SLC26A4 cases. (a-b) To explain the missing heritability in zeroallelic (M0, n=12) and monoallelic (M1, n=16) SLC26A4 cases, different genetic analyses were performed. Firstly, individuals were screened for the presence of the CEVA haplotype (M0/CEVA (M0C), n=2; M1/CEVA (M1C), n=10). Secondly, whole genome sequencing (WGS) was performed in all monoallelic cases (M0/CEVA, M1) to identify potential structural, splice (M2, n=2) or regulatory variants. Lastly, sequencing data were screened for potentially pathogenic variants in the EPHA2, FOXI1 and KCNJ10 genes. Digenic inheritance has been previously suggested for variants in these genes and the SLC26A4 gene. In three cases (M0/FOXI1 (M0F), n=2, CEVA/FOXI1 (M1F), n=1), a potentially pathogenic variant in FOXI1 (NM_012188.4, c.677C>T) was identified

Figure 2

Determination of the boundaries of the shared CEVA haplotype. (a) The CEVA haplotype was detected in 10 individuals, in an additional 2 individuals (SLC040 and SLC071, indicated with *), a smaller haplotype was found, termed V1-CEVA. To determine the boundaries of the CEVA haplotype, VNTR marker analysis was performed. The shared haplotype (0.89 Mb, CEVA; 0.57 Mb V1-CEVA is marked in orange. For marker D7S2420 (light-orange) a deviating CA-repeat length was determined in SLC003. Nevertheless, the marker is still considered to be potentially part of the shared haplotype as a change or repeat length cannot be excluded. Genomic positions (Mb) are according to the UCSC Genome Browser (GRCh37/hg19). (b) A schematic overview of the identified shared CEVA haplotype (D7S501-D7S2459). Positions of the CEVA-associated SNPs and the genes located within the haplotype region (CEVA, D7S501-D7S2459; V1-CEVA, SNP3-D7S2459) have been indicated. All SNPs are located within intronic or intergenic regions. Genomic positions of the CEVA-associated SNPs are provided in Table S7. SLC26A4 (NM_000441.1) is only partially included (exons 10/21) in the shared haplotype

Figure 3

Results of audiometric evaluation in affected individuals. PTA0.5-4 kHz for ears with an EVA. Each dot represents the hearing level of an ear with an enlarged vestibular aqueduct, allocated to genotype class (M2, M1/CEVA, M1, M0/CEVA and M0). The M1/CEVA group also includes subjects with an M1/V1-CEVA genotype. For an objective comparison, the same methods as used by Chao et al. (2019) were applied

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SmitsSLC26A4OnlineResources.pdf](#)