Big Data Palaeoecology reveals significant variation in Black Death mortality in Europe

Adam Izdebski

Max Planck Institute for the Science of Human History

Piotr Guzowski

University of Białystok <https://orcid.org/0000-0002-6494-4217>

Radosław Poniak

Faculty of History and International Relations, University of Białystok

Lucrezia Masci

Department of Earth Science, Sapienza University of Rome

Jordan Palli

Dendrologylab, Department of Agriculture and Forest Sciences (Dafne), University of Tuscia

Cristiano Vignola

Department of Environmental Biology, Sapienza University of Rome

Martin Bauch

Leibniz Institute for the History and Culture of Eastern Europe (GWZO)

Carlo Coccozza

Max Planck Institute for the Science of Human History

Ricardo Fernandes

Max Planck Institute for the Science of Human History

Fredrik Ljungqvist

Stockholm University <https://orcid.org/0000-0003-0220-3947>

Timothy Newfield

Georgetown University <https://orcid.org/0000-0003-1451-5024>

Andrea Seim

Chair of Forest Growth, Institute of Forest Sciences, Albert Ludwig University of Freiburg

Daniel Abel-Schaad

Department of Botany, Faculty of Sciences, Universidad de Granada <https://orcid.org/0000-0003-3915-8342>

Francisca Alba-Sánchez

Department of Botany, Faculty of Sciences, Universidad de Granada <https://orcid.org/0000-0003-0387-1533>

Leif Björkman

Viscum Pollenanalys & Miljöhistoria

Achim Brauer

Helmholtz-Zentrum Potsdam Deutsches GeoForschungsZentrum GFZ

Alex Brown

Wessex Archaeology, Portway House

Sambor Czerwiński

Climate Change Ecology Research Unit, Adam Mickiewicz University

Ana Ejarque

CNRS, Université Clermont Auvergne, GEOLAB

Magdalena Fiłoc

Department of Palaeobiology, Faculty of Biology, University of Białystok <https://orcid.org/0000-0003-1226-2676>

Assunta Florenzano

Laboratory of Palynology and Palaeobotany, Department of Life Sciences, University of Modena and Reggio Emilia

Erik Fredh

Museum of Archaeology, University of Stavanger

Ralph Fyfe

University of Plymouth

Nauris Jasiunas

Department of Geography, University of Latvia

Piotr Kołaczek

Climate Change Ecology Research Unit, Adam Mickiewicz University

Katerina Kouli

National and Kapodistrian University of Athens <https://orcid.org/0000-0003-1656-1091>

Radka Kozáková

Institute of Archeology, Academy of Sciences of the Czech Republic

Mirosława Kupryjanowicz

Department of Palaeobiology, Faculty of Biology, University of Białystok

Per Lagerås

The Archaeologists, National Historical Museums

Mariusz Lamentowicz

Climate Change Ecology Research Unit, Adam Mickiewicz University

Matts Lindblad

Southern Swedish Forest Research Centre, Swedish University of Agricultural Sciences

Jose Antonio López-Sáez

CSIC

Reyes Luelmo-Lautenschlaeger

Environmental Archaeology Research Group, Institute of History, CSIC

Katarzyna Marcisz

Climate Change Ecology Research Unit, Adam Mickiewicz University <https://orcid.org/0000-0003-2655-9729>

Florence Mazier

Department of Environmental Geography, GEODE UMR 5602, Jean Jaurès University

Scott Mensing

Department of Geography

Anna Maria Mercuri

Università di Modena e Reggio Emilia <https://orcid.org/0000-0001-6138-4165>

Krystyna Milecka

Anthropocene Research Unit, Faculty of Geographical and Geological Sciences, Adam Mickiewicz University

Yannick Miras

CNRS, HNHP UMR 7194, Muséum National d'Histoire Naturelle, Institut de Paléontologie Humain

Agnieszka Noryśkiewicz

Institute of Archaeology, Faculty of History, Nicolaus Copernicus University

Elena Novenko

Faculty of Geography, Lomonosov Moscow State University

Milena Obremska

Institute of Geological Sciences, Polish Academy of Sciences

Anna Pędziszewska

Laboratory of Palaeoecology and Archaeobotany, Department of Plant Ecology, Faculty of Biology, University of Gdańsk

Sebastián Pérez-Díaz

Department of Geography, Urban and Regional Planning, Universidad de Cantabria

Gianluca Piovesan

Università della Toscana <https://orcid.org/0000-0002-3214-0839>

Aleks Pluskowski

University of Reading <https://orcid.org/0000-0002-4494-7664>

Petr Pokorný

Charles University in Prague

Anneli Poska

Department of Geology, Tallinn University of Technology

Triin Reitalu

Department of Geology, Tallinn University of Technology

Manfred Rösch

Department of Prae- and Early History and West Asian Archaeology, University of Heidelberg

Laura Sadori

Sapienza Università di Roma <https://orcid.org/0000-0002-2774-6705>

Carla Sá Ferreira

School of Natural and Built Environment, Queen's University Belfast

David Sebag

University of Rouen <https://orcid.org/0000-0002-6446-6921>

Michał Słowiński

Institute of Geography and Spatial Organization Polish Academy of Sciences <https://orcid.org/0000-0002-3011-2682>

Migle Stančikaitė

Institute of Baltic Region History and Archaeology, University of Klaipėda

Normunds Stivrins

Department of Geography, University of Latvia

Irene Tunno

Center for Accelerator Mass Spectrometry (CAMS), Lawrence Livermore National Laboratory

Siim Veski

Department of Geology, Tallinn University of Technology

Agnieszka Wacnik

W. Szafer Institute of Botany, Polish Academy of Sciences

Alessia Masi (✉ alessia.masi@uniroma1.it)

Sapienza University of Rome <https://orcid.org/0000-0001-9822-9767>

Article

Keywords: Pandemic, Ancient DNA Research, Bacterium Yersinia Pestis, Landscape Change, Interregional Differences

Posted Date: July 2nd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-581474/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Version of Record: A version of this preprint was published at Nature Ecology & Evolution on February 10th, 2022. See the published version at <https://doi.org/10.1038/s41559-021-01652-4>.

Abstract

The Black Death is the most reknown pandemic in human history, believed by many to have killed half of Europe's population. However, despite the advances in ancient DNA research that allowed for the successful identification of the pandemic's causative agent (bacterium *Yersinia pestis*), our knowledge of the Black Death is still limited, based primarily on medieval texts available for single areas of Western Europe. In our study we remedy this situation and we focus in particular on the scale of the Black Death mortality. We collected data on landscape change from 261 coring sites (lakes and wetlands) located in 19 European countries. We used two independent methods of analysis to evaluate whether the changes we see in the landscape at the time of the Black Death agree with the hypothesis that half of the population died within a single year in each of the 21 regions we studied. We discovered that while the Black Death had devastating impact in some regions, it had negligible or no impact in others. The inter-regional differences in the Black Death mortality across Europe demonstrate the significance of cultural, ecological, economic and climatic factors that mediate the dissemination and impact of the disease. The complex interplay of these factors, along with the identification of the pathogen that caused disease outbreaks, should be the focus of future research on historical pandemics.

Summary Paragraph

Few doubt the mid-14th-century Afro-Eurasian plague pandemic, the Black Death, killed tens of millions of people. In Western Asia and Europe, where its spread and mortality are best understood, upwards of 50% of the population is thought to have died within ~ 5 years.^{1–4} Whole-genome sequencing confirms the pandemic as a novel introduction of the zoonotic bacterium *Yersinia pestis*.^{5,6} Yet, despite advances in paleogenetics and generations of written-source-based research on the cultural and economic transformations plague is credited with accelerating, from the 'Great Divergence' to the Renaissance^{7,8}, much about the Black Death's demographic impact remains poorly understood. The regionality of the plague's mortality is particularly underexplored, owing to the availability of written sources and the limits of traditional historical methods. Here we pioneer a new approach, Big Data Palaeoecology (BDP), that leverages the field of palynology to evaluate the demographic impact of the Black Death on a regional scale across Europe, independent of written sources. Our analysis of 1,634 pollen samples from 261 sites, reflecting landscape change and agricultural activities, demonstrates Black Death mortality was far more spatially heterogenous than previously recognised. Strikingly, BDP provides independent confirmation of the devastating toll historical sources evidence of the Black Death in some European regions, while establishing conclusively for the first time that the Black Death did not affect all regions equally. We attribute this mortality variation to climatic, cultural, ecological, economic and societal factors, which influenced *Y. pestis* dissemination and prevalence, generating regionally unique outcomes.

Estimating Black Death Mortality

Multidisciplinary studies are redefining the Black Death. In recent years, paleogeneticists have confirmed the pandemic's *Y. pestis* identity and established that the outbreak seeded novel plague reservoirs in

Europe^{5,6}. Archeologists and historians meanwhile have begun to put Sub-Saharan Africa on the Black Death map⁹, to fill in lacunae in our understanding of the pandemic's Mediterranean and European spread^{10,11}, and to explore the pandemic's origins in Central Asia and dissemination in East Asia, drawing on evolutionary biology and paleogenetics¹². But while multiple disciplines have reassessed the pandemic's spatiotemporality, the mortality of the outbreak, estimates of which have drawn attention to the Black Death for centuries, remains underexplored and limited to unidisciplinary, written-source-based approaches.

Medieval mortality data are scarce and highly fragmentary. 19th -c. historians of the Black Death based their assertion that the pandemic claimed 25% or more of European lives on assessments of qualitative narrative sources¹³. Since the mid-20th c., historians have painstakingly built-up multiple more instructive case studies of Black Death mortality for regions comparatively well-endowed with administrative sources allowing for statistical analysis (e.g., regions of England, France, Italy, the Netherlands)^{14–16}. Some of these cases studies have argued for a death toll in the range of 50% or more. These studies have been increasingly considered representative of the Black Death's broader mortality, and employed as proxies for Black Death mortality in European regions where direct evidence for the pandemic is nonexistent (e.g. Bohemia, Finland), slight (single sentences or vague passages: e.g. Moravia, Hungary and Scotland), or available but neither detailed nor quantitative (most regions, including areas of England, France, Italy, the Netherlands)¹.

By treating case studies of relatively well-documented regions as predictive of the pandemic's death toll in all regions the pandemic touched, histories of the Black Death implicitly rest on the untested assumption that plague mortality was uniform across regions regardless of local cultural, ecological, economic, and societal contexts, and therefore that the prevalence of *Y. pestis*, an ecologically and epidemiologically complex disease¹⁷, was comparable across Europe. This methodology has lent itself to estimates of an aggregate European death toll of ~ 50 % (~ 50 million deaths)^{1,4}, though studies accounting for regional source scarcities have estimated mortality to have fallen below that mark^{8,18,19}. Problematically, most of the quantitative sources drawn upon to build cases studies of Black Death mortality relate to urban contexts, which owing to their crowding and generally poor sanitation may have suffered higher plague mortality than rural areas^{20,21}, and in the mid-14th -c. upwards of 75–95% of the population of all European regions was rural²². Although every pandemic, plague or not, is distinct, a Black Death mortality of ~ 50% far exceeds the demographic losses sustained during the late-19th -c. third plague pandemic, including China and India which were then severely affected^{23,24}.

Big-data Palaeoecology (B D P) And Black Death Mortality

Here we pioneer an alternative approach (discussed in detail in **Methods**). BDP allows us to evaluate Black Death mortality across Europe using quantitative paleoenvironmental datasets which can be employed for spatial statistical comparisons. Our dataset consists of fossil pollen counts from 261 radiocarbon-dated sediment cores from 19 present-day European countries (Fig. 1; Supplementary Data

1). Pollen data can be used to assess past demographic dynamics as human pressure on the landscape in the preindustrial period was directly dependent on the availability of rural labour. We focus on the period between 1250 and 1450 CE for which 1,634 pollen-analysed sediment samples are available (Supplementary Data 2). In order to compare trends between different European bioclimatic zones, we assembled pollen types (taxa) representing different plants from species to family level into four standardised summary indicators, based on their subsistence value for human economy or their ecological needs as reflected in two major ecological indices, the Ellenberg light indicator and Niinemets and Valladares shade tolerance scale: (1) Cereals; (2) Herding (pasturelands); (3) Fast forest secondary succession (pioneer shrubs and trees growing on former fields/pastures within 5–10 years of abandonment); (4) Slow forest secondary succession (mature-late successional woodland on abandoned fields/pastures). We discuss the results of the Ellenberg-based indicators below, but the Niinemets-based indicators yielded the same results in 19 out of 21 regions and the differences do not bear on our discussion (Extended Data Figs. 1 and 2).

Validity Test: Sweden And Poland

We validated our BDP approach by examining two well-studied, but contrasting, regional case studies of Black Death mortality, in Sweden and Poland. An earlier multi-disciplinary analysis discovered significant contraction in cereal cultivation, as well as a more general economic and demographic decline, in Southern Sweden following the Black Death²⁵. In contrast, historians have long demonstrated that Central Europe, Poland particularly, experienced economic growth over the 14th c.²⁶ Our BDP approach independently corroborates these trajectories. We validate the ability of our pollen datasets to reflect the extent of Black Death mortality by comparing them to historical data of national tax payments made to the pope (Peter's Pence)^{27–29} (Extended Data Fig. 3).

Landscape And Demographic Change: Four Major Scenarios

Having validated our approach, we analysed the pollen dataset for all of Europe. We focused on contrasting the four summary pollen indicator values on a regional scale for sub-periods of 100 years before and after the Black Death. We discuss the results of this analysis here (Fig. 2); we also present our complimentary 50-year (1301–1350 to 1351–1400) and 25-year (1325–1350 to 1351–1375) analyses (Extended Data Figs. 4 and 5). For cereals, they returned the same results for 20 out of 21 regions for 50-year and 15 out of 19 regions for 25-year analyses, confirming the robustness of our conclusions.

BDP employs bootstrapping for evaluating statistical significance of the differences between the pre- and post-Black Death sub-periods on a regional level. In this way, we identified four scenarios of post-Black Death agricultural change based on the cereal pollen indicator (Fig. 2): (1) substantial and statistically significant increase, reflecting arable expansion and limited Black Death mortality (top panel); (2) modest and statistically insignificant increase, suggesting stability or slow-paced agrarian growth (upper middle panel); (3) modest and statistically insignificant decrease, suggesting stagnation or some contraction of agrarian activities, possibly stemming from more limited demographic losses or economic disruption

caused by the plague (lower middle panel); (4) substantial and statistically significant decrease, reflecting arable contraction and pronounced Black Death mortality (bottom panel). Scenarios 1 and 2 falsify the theory that Black Death mortality was everywhere significant. Scenario 3 and 4 demonstrate Black Death mortality was devastating in some regions, but that the extent of the mortality varied regionally.

Changes in the herding indicator reveal trajectories similar to those of cereals in most cases or no change. We discovered only one region (SW Germany) where there occurred a statistically significant decline in cultivation in parallel with a statistically significant increase in herding (some indication of this is also apparent in Greece), suggesting a shift to livestock production related to agrarian labour shortages³⁰. The fast forest succession indicator increases in a statistically significant way in Central Italy and Central France, confirming field abandonment. In Central Italy, this is accompanied by discernible reforestation (statistically significant increase in the slow forest succession indicator), attesting to significant regional Black Death mortality and slow demographic and economic recovery.

To confirm the robustness of our results, BDP combines statistical with independent spatial approaches (Fig. 3). The latter yielded results identical for all four BDP indicators to our statistical approach in our 100-year period analysis and for our 50-year (1301–1350 to 1351–1400) and 25-year (1325–1350 to 1351–1375) period analyses (Extended Data Fig. 7–10).

Black Death: A Diverse And Entangled Phenomenon

Figure 4 visualizes the spatial distribution of the four trajectories of post-Black Death landscape change from Fig. 2, demonstrating that Black Death mortality varied significantly between European regions. The pandemic was immensely destructive in some areas, but in others it had a far lighter touch. Strikingly, BDP identifies a sharp agricultural decline in several regions of Europe, independently corroborating analyses of historical sources that suggest high mortality in regions of Scandinavia, France, Western Germany, Greece, and Central Italy¹ and lending further validation to our approach. At the same time, there is much evidence for continuity and uninterrupted agricultural growth in Central and Eastern Europe, and several regions of Western Europe, particularly in Ireland and Iberia. In this way, BDP complicates histories of the Black Death that assume *Y. pestis* was uniformly prevalent across Europe and that the pandemic everywhere had a devastating demographic impact.

The significant variability of Black Death mortality that our BDP approach identifies remains to be explained, but local cultural, demographic, economic, environmental and societal contexts likely influenced *Y. pestis* prevalence, morbidity and mortality. The spread of the pandemic depended on numerous factors, which would have generated compound effects, feedback loops, and regionally unique outcomes. Plague is an ecologically and epidemiologically complex zoonotic disease, maintained by sylvatic rodents and their fleas, and transmittable to and between people via multiple pathways, including commensal and sylvatic rodent flea bites, respiratory secretions, direct contact with infected animals, human ectoparasites (fleas and lice), and fomites¹⁷. The behaviour of *Y. pestis* hosts and vectors, and their capacity to efficiently transmit the pathogen, is partly constrained by complex interactions with

seasonal climate variability and local ecological conditions, both in anthropogenic and rural environments (cities, villages and fields vs. mountains, forests or wetlands³¹). Regional variation in population density and distribution, ectoparasite burdens, living conditions, and commensal rodent populations and their fleas, undoubtedly mattered. Local climatic contexts were shown to have strongly determined third plague pandemic dynamics in Asia^{32,33}. In Europe, where the Black Death spread over several years (1347–1352 for the regions considered here), different seasons and annually variable climatic conditions may have influenced *Y. pestis* prevalence and the pandemic's mortality. Further, while hypothesized links between early-14th-c. famines and the Black Death remain to be substantiated, and while *Y. pestis* is often lethal¹⁷, it has been shown Black Death mortality was selective and the immunological and nutritional heterogeneity of the populations the pathogen interacted with would have ensured uneven toll^{34,35}.

That climatic, cultural, ecological and economic factors shaped regional Black Death outcomes is well-illustrated by the initial phases of the pandemic in Europe. Cereal trade is thought to have been instrumental for the introduction of the Black Death to Mediterranean Europe, and along established conduits of commerce and communication, ecological factors, associated contingency effects and historical path dependency mattered from the outset. Before the pandemic arrived in the Crimea, the volume of cereal trade between Italy and the Black Sea was sizeable, yet blocked by embargo^{11,36,37}. High demand in Southern Europe for Black Sea cereals from 1345 onwards was associated with a period of excessive precipitation and cooling^{38,39} negatively affecting cereal supplies in Italy and beyond, 1345–47^{8,36,40,41}. Cereal imports from Black Sea coasts resumed once the situation improved in 1347, and by early 1348 Venetian merchants had filled many Italian granaries with Black Sea produce^{11,36,37,42} and introduced plague to Europe. Plague outbreaks disseminated from major cereal ports in southern and north-western Italy from January 1348^{1,43}. To the contrary, plague hardly spread in north-western Italy, which was independent from overseas cereal imports^{36,42,44}. Local circumstances were shaping the outcome of the pandemic on regional scales from the outset.

To conclude, our BDP approach shows Black Death mortality was far more spatially heterogeneous than previously thought. This significant variation in Black Death mortality may be explained by the pathogen's entanglement with a dynamic nexus of climatic, cultural, demographic, ecological and societal factors that determined its prevalence and the pandemic's mortality in any given region. That the pandemic was immensely destructive in some regions, but not all, falsifies the practice, common in Black Death studies, of predicting one region's experience on the basis of another's. Regional mortality outcomes must be reconstructed using local sources, including BDP. As a few well-documented case studies of destructive Black Death mortality in Europe have informed estimates of Black Death mortality not only in other European regions, but also in several regions of Africa and Asia, our findings have significant implications both for the wider history of the Black Death and for how historical disease outbreaks are reconstructed.

Declarations

Acknowledgements

The authors acknowledge the following funding sources: Estonian Research Council # PRG323, PUT1173 (AP, TR, NS SV); European Research Council # FP7 263735 (AB, AP), # MSC 655659 (AE); Georgetown Environmental Initiative (TN); Latvian Council of Science # LZP-2020/2-0060 (NS, NJ); LLNL-JRNL-820941 (IT); NSF award # GSS-1228126 (SM); Polish-Swiss Research Programme # 013/2010 CLIMPEAT (ML), # 086/2010 CLIMPOL (AW); Polish Ministry of Science and Higher Education # N N306 275635 (MK); Polish National Science Centre # 2019/03/X/ST10/00849 (ML), # 2015/17/B/ST10/01656 (ML), # 2015/17/B/ST10/03430 (MS), # 2018/31/B/ST10/02498 (MS), # N N304 319636 (AW); SCIEX # 12.286 (KM); Spanish Ministry of Economy and Competitiveness # REDISCO-HAR2017-88035-P (JALS); Spanish Ministry of Education, Culture and Sports # FPU16/00676 (RLL); Swedish Research Council # 421-2010-1570 (PL), # 2018-01272 (FCL, AS); Volkswagen Foundation Freigeist Fellowship Dantean Anomaly (MB).

Author contributions

AI, AM and PG designed the study. AI drafted the paper with contributions from MB, RF, AM, TN, PG, RP, and CV. AM, LM, JP and CV created the pollen database, with the support of JALS (Spain), PL (Sweden), and TR (Estonia). AI, RP, RF and CC carried out the analyses. All other authors provided palynological data and contributed to the text and the interpretation.

Competing interests

The authors declare no competing interests.

Materials & Correspondence

Please address your inquiries to Alessia Masi, alessia.masi@uniroma1.it, and Adam Izdebski, izdebski@shh.mpg.de

References

1.
Benedictow, O. L. *The Black Death 1346-1353: the complete history*. (Boydell, 2004).
2.
Mohammed, M. *Peste, contagion et martyre: Histoire du fléau en Occident médiéval*. (Publisud, 2005).

3.

Varlik, N. *Plague and empire in the early modern Mediterranean world: the Ottoman experience, 1347-1600*. (Cambridge University Press, 2015).

4.

Aberth, J. *The Black Death: a new history of the great mortality in Europe, 1347-1500*. (Oxford University Press, 2021).

5.

Bos, K. I. *et al.* A draft genome of *Yersinia pestis* from victims of the Black Death. *Nature* **478**, 506–510 (2011).

6.

Spyrou, M. A. *et al.* Phylogeography of the second plague pandemic revealed through analysis of historical *Yersinia pestis* genomes. *Nat Commun* **10**, 1–13 (2019).

7.

Herlihy, D. Climate and Documentary Sources: A Comment. *The Journal of Interdisciplinary History* **10**, 713–717 (1980).

8.

Campbell, B. *The Great Transition: Climate, Disease and Society in the Late-Medieval World*. (Cambridge University Press, 2016).

9.

Chouin, G. L. Reflections on plague in African history (14th–19th c.). *Afriques* **9**, 2228 (2018).

10.

Roosen, J. & Curtis, D. R. Dangers of Noncritical Use of Historical Plague Data. *Emerging Infectious Diseases* **24**, 103–110 (2018).

11.

Barker, H. Laying the Corpses to Rest: Grain, Embargoes, and *Yersinia pestis* in the Black Sea, 1346–48. *Speculum* **96**, 97–126 (2020).

12.

Green, M. The Four Black Deaths. *American Historical Review* **125**, 1601–1631 (2020).

13.

Hecker, J. *Der schwarze Tod*. (1832).

14.

Hatcher, J. *Plague, population, and the English economy, 1348-1530*. (Macmillan, 1977).

15.

Ormrod, W. M. & Lindley, P. G. *The Black Death in England, 1348-1500*. (Paul Watkins, 1996).

16.

Roosen, J. & Curtis, D. R. The 'light touch' of the Black Death in the Southern Netherlands: an urban trick? *The Economic History Review* **72**, 32–56 (2019).

17.

Dubyanskiy, V. M. & Yeszhanov, A. B. Ecology of *Yersinia pestis* and the Epidemiology of Plague. in *Yersinia pestis: Retrospective and Perspective* (eds. Yang, R. & Anisimov, A.) 101–170 (Springer Netherlands, 2016). doi:10.1007/978-94-024-0890-4_5.

18.

Gottfried, R. S. *The Black Death: natural and human disaster in medieval Europe*. (Robert Hale, 1983).

19. Christakos, G. *Interdisciplinary public health reasoning and epidemic modelling: the case of Black Death*. (Springer, 2005).

20.

Olea, R. A. & Christakos, G. Duration of urban mortality for the 14th-century Black Death epidemic. *Hum Biol* **77**, 291–303 (2005).

21.

Alfani, G. Plague in seventeenth-century Europe and the decline of Italy: an epidemiological hypothesis. *European Review of Economic History* **17**, 408–430 (2013).

22.

Neithard, B. L'essor (Xe - XIVe siècle). in *Histoire des populations de l'Europe* (eds. Dupâquier, J. & Bardet, J.-P.) vol. 1: Des origines aux prémices de la révolution démographique 168–184 (Fayard, 1997).

23.

Echenberg, M. *Plague Ports: The Global Urban Impact of Bubonic Plague, 1894-1901*. (NYU Press, 2007).

24.

Tumbe, C. *Pandemics and Historical Mortality in India*.

<https://web.iima.ac.in/assets/snippets/workingpaperpdf/17719931472020-12-03.pdf> (2020).

25.

Lagerås, P. *et al.* Abandonment, agricultural change and ecology. in *Environment, society and the Black Death: an interdisciplinary approach to the late-medieval crisis in Sweden* (ed. Lagerås, P.) 30–68 (Oxbow Books, 2016).

26.

Małowist, M. *Western Europe, Eastern Europe and world development, 13th-18th centuries: collection of essays of Marian Małowist*. (Brill, 2010).

27.

Guzowski, P., Kuklo, C. & Poniak, R. O metodach pomiaru natężenia epidemii i zaraz w preindustrialnej Europie w demografii historycznej. in *Epidemie w dziejach Europy: konsekwencje społeczne, gospodarcze i kulturowe* (eds. Polek, K. & Sroka, Ł. T.) 119–144 (Wydawnictwo Naukowe Uniwersytetu Pedagogicznego, 2016).

28.

Myrdal, J. The forgotten plague: the Black Death in Sweden. in *When disease makes history: epidemics and great historical turning points* (ed. Hämmäläinen, P.) 141–186 (Helsinki University Press, 2006).

29.

Myrdal, J. The Black Death in the North. in *Living with the Black Death* (eds. Bisgaard, L. & Søndergaard, L.) 63–84 (University Press of Southern Denmark; Gazelle [distributor, 2011]).

30.

Poos, L. *Rural Society after the Black Death*. (Cambridge University Press, 2011).

31.

Slavin, P. Out of the West: formation of a permanent plague reservoir in South-Central Germany (1349-1356) and its implications. *Past & Present* gtaa028 (2021) doi:10.1093/pastj/gtaa028.

32.

Xu, L. *et al.* Nonlinear effect of climate on plague during the third pandemic in China. *PNAS* **108**, 10214–10219 (2011).

33.

Xu, L. *et al.* Wet climate and transportation routes accelerate spread of human plague. *Proc Biol Sci* **281**, 20133159 (2014).

34.

DeWitte, S., Stature and frailty during the Black Death: the effect of stature on risks of epidemic mortality in London, A.D. 1348–1350. *J Arch Sci* **39**, 1412-1419 (2012).

35.

Crespo, F. & Lawrenz, M.B., Heterogeneous Immunological Landscapes and Medieval Plague: An Invitation to a New Dialogue between Historians and Immunologists. *The Medieval Globe* 1, 229-258 (2014).

36.

Pinto, G. Firenze e la carestia del 1346-47: Aspetti e problemi delle crisi annonarie alla metà del '300. *Archivio Storico Italiano* **130**, 3–84 (1972).

37.

Faugeron, F. *Nourrir la ville: ravitaillement, marchés et métiers de l'alimentation à Venise dans les derniers siècles du Moyen Âge.* (2014).

38.

Luterbacher, J. *et al.* European summer temperatures since Roman times. *Environ. Res. Lett.* **11**, 024001 (2016).

39.

Esper, J. *et al.* Eastern Mediterranean summer temperatures since 730 CE from Mt. Smolikas tree-ring densities. *Clim Dyn* **54**, 1367–1382 (2020).

40.

Campbell, B. The European Mortality Crises of 1346-52 and Advent of the Little Ice Age. in *Famines During the 'Little Ice Age' (1300-1800) Socionatural Entanglements in Premodern Societies* 19–41 (Springer, 2017).

41.

Bauch, M. & Engel, A. Die 1340er Jahre als Schlüsseljahrzehnt der ‚Great Transition‘. Eine klimahistorische Perspektive auf den Vorabend des Schwarzen Todes. in *Pest! Sonderausstellung des LWL-Museums für Archäologie - Westfälisches Landesmuseum Herne* (eds. Berner, A., Leenen, S. & Maus, S.) 76–82 (WBG, 2019).

42.

Hübner, H.-J. *Quia bonum sit anticipare tempus. Die kommunale Versorgung Venedigs mit Brot und Getreide vom späten 12. bis ins 15. Jahrhundert.* (Lang, 1998).

43.

Del Panta, L. *Le epidemie nella storia demografica italiana. (secoli XIV-XIX).* (Loescher, 1980).

44.

Santoro, C. *Gli uffici del Comune di Milano e del Dominio visconteo-sforzesco <1216-1515>.* (Giuffrè, 1968).

Methods

Big Data Palaeoecology (BDP) approach: pollen-inferred landscape change and preindustrial demography

Recently, paleoenvironmental data derived from tree rings or ice cores have been employed to approximate changes in human economic activity related to past epidemics, as well as to warfare and climatic variability^{45,46}. However, none of these proxies is directly related to human demography or provides a basis to estimate variation in Black Death mortality on a regional scale across Europe.

On the contrary, in preindustrial economies, rural labour availability and the spatial scale of cereal cultivation were directly related. An increase in the extent and intensity of cereal cultivation – as reflected in pollen data – would have required not only a predilection and demand for cereals, but greater availability of labour and thus population growth or significant immigration. The maintenance of existing agricultural activity, in turn, would have required relatively stable population levels.^{47–49} The uniform ~50% mortality postulated for the Black Death across Europe should have resulted in a large and significant decline of cereal cultivation and parallel forest regrowth across Europe, as previously demonstrated for mid-14th c. Sweden²⁵ and singular sites in some regions of Western Europe⁵⁰. This result agrees with the fact that Black Death mortality could be high among people at productive age, as illustrated for England^{51,52}. Moreover, even in the case of England, a comparatively commercialised and adaptive rural economy in mid-14th-c. Europe, the loss of 50% of population led to a significant decline in the total area under cultivation (as documented by heterogeneous written sources)⁵³. In Italy, another

well-developed economy at that time, the expansion of large estates following the Black Death also did not compensate for the general loss of cereal productivity⁵⁴. This effect, high mortality driving arable contraction, must have been yet more pronounced in more subsistence-oriented and less adaptive economies, with limited surplus production, such as in regions of the Iberian Peninsula, Germany, Sweden, and particularly East-Central Europe. Importantly, palaeoecological evidence for arable contraction may be indicative, to some extent, of not only rural population decline but also urban population decline in the region, as there is evidence in some areas, following the pandemic, of rural-to-urban migration, of country-dwellers repopulating urban centres¹⁰. Possibly less common was intraregional rural migration, as marginal lands were abandoned for better quality soils, which were more likely to remain under cultivation^{25,55}.

BDP data collection

Existing online palynological databases (the European Pollen Database (EPD)⁵⁶ www.europeanpollendatabase.net, and the Czech Quaternary Palynological Database (PALYCZ)⁵⁷, <https://botany.natur.cuni.cz/palycz/>), as well as personal contacts of the study authors and a systematic publication search were employed to identify palynological sites in Europe reaching the required chronological and resolution quality for the study of the last millennium. In order to enable statistical analysis, we included only sites clustered in well-defined historical-geographical regions, excluding isolated sites even if the quality of a site's data was very good. Data of sufficient quality and amount from regions for which the Black Death is well-studied, notably Central and Northern England and the Low Countries, is not presently available; to the best of our knowledge, for each of these regions there currently is not more than a single isolated site⁵⁰, which does not allow for the application of statistical approaches.

In total, 261 pollen records with the average temporal resolution of 58 years and ¹⁴C-age control (or varve chronology), have been collected. The age-depth models of the sequences have been provided by authors in original publications, by the EPD or developed through the Clam package (version 2.3.4) of R software for the purpose of this study. The analytical protocol for pollen extraction and identification is reported in the original publications. The Pollen Sum includes all the terrestrial taxa with some exceptions based on the selection done in the original publications. The full list of sequences, exclusions from the Pollen Sum, age-depth models and full references are reported in Supplementary Data 1.

The taxa list has been normalized by applying the EPD nomenclature. In this respect, the general name Cichorioideae includes Asteraceae subf. Cichorioideae of the EPD and PALYCZ nomenclatures, which primarily refers to the fenestrate pollen of the Cichorieae tribe⁵⁸. Ericaceae groups *Arbutus unedo*, *Calluna vulgaris*, *Vaccinium* and different *Erica* pollen types, whereas deciduous *Quercus* comprehends both *Q. robur* and *Q. cerris* pollen types⁵⁹. Rosaceae refers to both tree and herb species of the family. Finally, *Rumex* includes *R. acetosa* type, *R. acetosella*, *R. crispus* type, *Rumex/Oxyria* and *Urtica* groups *U. dioica* type and *U. pilulifera*.

BDP summary pollen indicators

In order to connect changes visible in the pollen data to human demographic trajectories, we assembled four summary pollen indicators that describe specific landscapes related to human activity. They reflect different degrees of demographic pressure on the landscape (cereal cultivation, pastoral activities, which are less-labour intensive than cereal cultivation, abandonment and rewilding) as well as different durations of land abandonment that might have occurred post-Black Death. Our indicators account for the fact that Europe is a continent rich in natural heritage, with a wide range of landscapes and habitats and a remarkable wealth of flora and fauna, shaped by climate, geomorphology and human activity. In order to ensure uniform interpretation of the indicators, we relied on criteria that can be applied to all European landscapes regardless of their local specificity. Cereals and herding are directly related to human activities and are barely influenced by spatial differences. More complex is the succession of natural plants with their ecological behaviour and inter-species competition. For this reason, we relied on existing quantitative indicators of plant ecology.

The Ellenberg L – light availability indicator⁶⁰ provides a measure of sunlight availability in woodlands and consequently of tree-canopy thickness, reflecting the scale of the natural regeneration of woodland vegetation after cultivation or pasture activities⁶¹. Nonetheless, ecological studies have suggested that geographic and climatic variability between different European regions can influence the Ellenberg indicator system^{62–65}. The original indicators were primarily designed for Central Europe⁵⁸, but several studies developed Ellenberg indicators for other regions, reflecting the specific ecology of the selected taxa (British Isles⁶⁶; Czech Republic⁶⁷; Greece⁶⁸; Italy⁶⁹; Sweden⁷⁰). Plants with a L values between 5 and 8 are listed in fast succession indicator, the ones with L values ranging from 1 to 4 are included in slow succession indicator. The result is the following list:

1) **Cereals**: only cultivated cereals have been included: *Avena/Triticum* type, Cerealia type, *Hordeum* type, *Secale*. 2) **Herding** includes pastoral indicators linked to the redistribution of human pressure: *Artemisia*, Cichorioideae, *Plantago lanceolata* type, *Plantago major/media* type, *Polygonum aviculare* type, *Rumex*, *Trifolium* type, *Urtica*, *Vicia* type. 3) **Fast Succession** comprises indicators of relatively recent reforestation of cultivated land after abandonment: *Alnus*, *Betula*, *Corylus*, Ericaceae, *Fraxinus ornus*, *Juniperus*, *Picea*, *Pinus*, *Populus*, deciduous *Quercus*, Rosaceae. 4) **Slow Succession** includes indicators of secondary succession established after several decades of abandonment: *Abies*, *Carpinus betulus*, *Fagus*, *Fraxinus*, *Ostrya/Carpinus orientalis*, *Quercus ilex* type.

In order to validate the indicators overcoming the regional limits of Ellenberg values, a different subdivision has been provided following the Niinemets and Valladares shade tolerance scale for woody species of the Northern Hemisphere⁷¹. The subdivision of taxa in the Fast and Slow succession indicators remains the same with only three changes: *Fraxinus ornus* and *Picea* move from Fast to Slow succession and *Fraxinus* from Slow to Fast succession. Extended Data Fig. 1 and 2 show that the two groupings yield the same results, which confirms the reliability of our indicators. There is only one clear exception (Russia), with one more region where smaller-scale diversion occurs for only one indicator,

Slow Succession (Norway). The different indicator behaviour results from the different attribution of *Picea* in our two sets of succession indicators: at high latitude, *Picea* characterizes the final stage of the ecological succession and hence its different attribution results in different summary indicator values in Russia for the two stages of ecological succession, fast and slow.

Please note our summary indicators are not designed to reflect the entirety of the landscape and reconstruct all of its different components. Rather, they are a means of approximating changes in the landscape related to the types of human activities, and their intensity, as much as they relate to demographic changes in human populations using and inhabiting these landscapes.

BDP analytical methods: statistical and spatial

To control for local specificity, pollen percentages of every taxon from each pollen site were standardised. From the taxa percent in a given year the arithmetic mean calculated for the observations from the period 1250–1450 was subtracted and the result divided by the standard deviation for the 1250–1450 period. Standardized taxa results were assembled for each site into four BDP summary indicators. Since each indicator has different numbers of taxa, the sum of standardized taxa values calculated for a given year and site was divided by the number of taxa in the indicator. For the purposes of replication, this standardised pollen dataset, comprising the four indicators for each sample from each site, is available as Supplementary Data 1.

This dataset has been analysed in two ways, statistically and spatially.

For the statistical approach, standardized regional indices of landscape transformation were created for each region by calculating the average value for all sites within the region, for each of the subperiods analysed in the study (1250–1350 and 1351–1450; 1301–1350 and 1351–1400; 1325–1350 and 1351–1375). Differences between means for each subperiod were measured by the use of the bootstrapping based on 10,000 resamples. The 90% and 95% confidence intervals were estimated with the bias-corrected and accelerated method (BCa)⁷². These results are visualized in Fig. 3 for the comparison of the sub-periods of 1250–1350 vs 1351–1450, and in Extended Data Fig. 7 and 9 for the comparison of the sub-periods of 1300–1350 vs 1351–1400 and 1325–1350 vs 1351–1375, respectively.

For the spatial approach, we employed the Bayesian model AverageR developed within the Pandora & IsoMemo initiatives (<https://pandoraapp.earth/>) to map the distribution of pollen indices across Europe. AverageR is a generalized additive model that has been previously described⁷³. It relies on a thin plate regression spline⁷⁵ to predict new, unseen data using the following model:

$$Y_i = g(\text{longitude}, \text{latitude}) + \epsilon_i$$

where

Y_i : independent variable for site i

$g(\text{longitude, latitude})$: spline smoother

$\epsilon_i \sim N(0, \sigma\epsilon)$: error term

The spline smoother can be written as $X\beta$ where X is a fixed design matrix and β is the parameter vector. Surface smoothing is controlled by employing a Bayesian smoothing parameter estimated from the data and trades-off bias against variance to make the optimal prediction⁷⁵. This parameter β is assumed to follow a normal distribution: $\beta \sim N(0, 1/\delta\lambda^*P)$, where P is a so-called penalty matrix of the thin plate regression spline, which penalizes second derivatives⁷⁵. The δ parameter is by default set to 1 but this can be adjusted to suit smoothing needs for each application. In our study δ was set at 0.9 to match the preferred spatial scale of analysis for our dataset (c. 250 to 500km).

AverageR was employed to generate smoothed surfaces for three sets of temporal bins (1250–1350 vs 1351–1450, as well as 1300–1350 vs 1351–1400 and 1325–1350 vs 1351–1375) and for the four BDP indicators (Extended Data Fig. 6, 8 and 10). For the same indicator the difference between the two temporal bins was plotted (Fig. 3; Extended Data Fig. 7 and 9).

Data Availability: All data generated or analysed during this study are included in this published article (as the Supplementary Data).

References – methods

45.

Ljungqvist, F. C. *et al.* Linking European building activity with plague history. *Journal of Archaeological Science* **98**, 81–92 (2018).

46.

McConnell, J. R. *et al.* Pervasive Arctic lead pollution suggests substantial growth in medieval silver production modulated by plague, climate, and conflict. *PNAS* **116**, 14910–14915 (2019).

47.

Izdebski, A. *A rural economy in transition: Asia Minor from Late Antiquity into the Early Middle Ages*. (Taubenschlag Foundation, 2013).

48.

Roberts, N. *et al.* Europe's lost forests: a pollen-based synthesis for the last 11,000 years. *Scientific Reports* **8**, 716 (2018).

49.

Bevan, A. *et al.* The changing face of the Mediterranean – Land cover, demography and environmental change: Introduction and overview. *The Holocene* 095968361982668 (2019)
doi:10.1177/0959683619826688.

50.

Yeloff, D. & Van Geel, B. Abandonment of farmland and vegetation succession following the Eurasian plague pandemic of ad 1347–52. *Journal of Biogeography* **34**, 575–582 (2007).

51.

Razi, Z. *Life, marriage, and death in a medieval parish: economy, society, and demography in Halesowen, 1270-1400*. (University Press, 1980).

52.

DeWitte, S. N. Age Patterns of Mortality During the Black Death in London, A.D. 1349–1350. *J Archaeol Sci* **37**, 3394–3400 (2010).

53.

Broadberry, S. N. *British economic growth, 1270-1870*. (University Press, 2015).

54.

Cortonesi, A. L'agricoltura italiana fra XIII e XIV secolo: vecchi e nuovi paesaggi. in *Medioevo delle campagne: rapporti di lavoro, politica agraria, protesta contadina*. (eds. Cortonesi, A. & Piccinni, G.) 15–56 (Viella, 2011).

55.

Abel, W. *Agrarkrisen und Agrarkonjunktur. Eine Geschichte der Land- und Ernährungswirtschaft Mitteleuropas seit dem hohen Mittelalter*. (Parey, 1978).

56.

Fyfe, R. *et al.* The European Pollen Database: past efforts and current activities. *Vegetation History and Archaeobotany* **18**, 417–424 (2009).

57.

Kuneš, P., Abraham, V., Kovářík, O., Kopecký, M., & PALYCZ contributors. Czech Quaternary Palynological Database – PALYCZ: review and basic statistics of the data. *Preslia* **81**, 209–238 (2009).

58.

Florenzano, A., Marignani, M., Rosati, L., Fascetti, S. & Mercuri, A. M. Are Cichorieae an indicator of open habitats and pastoralism in current and past vegetation studies? *Plant Biosystems - An International Journal Dealing with all Aspects of Plant Biology* **149**, 154–165 (2015).

59.

Smit, A. A scanning electron microscopical study of the pollen morphology in the genus *Quercus*. *Acta Botanica Neerlandica* **22**, 655–665 (1973).

60.

Ellenberg, H. *et al.* Zeigerwerte von Pflanzen in Mitteleuropa. *Scripta Geobotanica* **18**, 1–258 (1992).

61.

Dzwonko, Z. Assessment of light and soil conditions in ancient and recent woodlands by Ellenberg indicator values. *Journal of Applied Ecology* **38**, 942–951 (2001).

62.

Diekmann, M. & Lawesson, J. E. Shifts in ecological behaviour of herbaceous forest species along a transect from northern central to North Europe. *Folia Geobotanica* **34**, 127–141 (1999).

63.

Gégout, J.-C. & Krizova, E. Comparison of indicator values of forest understory plant species in Western Carpathians (Slovakia) and Vosges Mountains (France). *Forest Ecology and Management* **182**, 1–11 (2003).

64.

Hájková, P., Hájek, M., Apostolova, I., Zelený, D. & Dítě, D. Shifts in the ecological behaviour of plant species between two distant regions: evidence from the base richness gradient in mires. *Journal of Biogeography* 071103055558002-??? (2007) doi:10.1111/j.1365-2699.2007.01793.x.

65.

Wasof, S. *et al.* Ecological niche shifts of understorey plants along a latitudinal gradient of temperate forests in north-western Europe. *Global Ecology and Biogeography* **22**, 1130–1140 (2013).

66.

Hill, M., Mountford, J., Roy, D. & Bunce, R. *Ellenberg's indicator values for British plants. ECOFACT Volume 2 Technical Annex*. (Institute of Terrestrial Ecology, 1999).

67.

Chytrý, M., Tichý, L., Dřevojan, P., Sádlo, J. & Zelený, D. Ellenberg-type indicator values for the Czech flora. *Preslia* **90**, 83–103 (2018).

68.

Böhling, N., Greuter, W. & Raus, T. Zeigerwerte der Gefäßpflanzen der Südägäis (Griechenland). Indicator values of the vascular plants in the Southern Aegean (Greece). *Braun-Blanquetia* **32**, 1–106 (2002).

69.

Pignatti, S., Menegoni, P. & Pietrosanti, S. Biondificazione attraverso le piante vascolari. Valori di indicazione secondo Ellenberg (Zeigerwerte) per le specie della Flora d'Italia. *Braun-Blanquetia* **39**, 1–97 (2005).

70.

Diekmann, M. Use and improvement of Ellenberg's indicator values in deciduous forests of the Boreo-nemoral zone in Sweden. *Ecography* **18**, 178–189 (1995).

71.

Niinemets, Ü. & Valladares, F. Tolerance to Shade, Drought, and Waterlogging of Temperate Northern Hemisphere Trees and Shrubs. *Ecological Monographs* **76**, 521–547 (2006).

72.

Hall, P. Theoretical Comparison of Bootstrap Confidence Intervals. *The Annals of Statistics* **16**, 927–953 (1988).

73.

Cubas, M. *et al.* Latitudinal gradient in dairy production with the introduction of farming in Atlantic Europe. *Nature Communications* **11**, 2036 (2020).

74.

Wood, S. N. Thin plate regression splines. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* **65**, 95–114 (2003).

75.

Groß, M. Modeling body height in prehistory using a spatio-temporal Bayesian errors-in variables model. *AStA Adv Stat Anal* **100**, 289–311 (2016).

Figures

Figure 1

Location of pollen-analysed sediment cores used in this study. Circle size reflects the number of samples per site for the period of 1250-1450 CE, different colours reflect division of sites into regional clusters for the purpose of the analysis presented in Figure 3.

Figure 2

Post-Black Death regional-scale changes in four BDP pollen indicators. Difference between the means of 100-yr periods of 1250-1350 and 1351-1450 CE, with the standard deviation. Statistical significance is based on bootstrap estimates. The indicators are presented in four rows from statistically significant increases in standardised mean percentages of cereal pollen (top) to statistically significant declines in standardised mean percentages of cereal pollen (bottom).

Figure 3

Spatial extrapolation showing 1250-1350 CE versus 1351-1450 CE temporal variation in the BDP pollen indicators a) Cereal b) Herding c) Fast Succession d) Slow Succession. Based on results from Extended Data Fig. 6.

Figure 4

BDP-determined regional scenarios of Black Death demographic impact. Colours reflect centennial-scale changes in the cereal pollen indicators from Fig. 2. Background map with political borders of 14th-c. Europe (©Hans Sell, Michelle O'Reilly, Adam Izdebski, MPI SHH).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SIGuide.docx](#)
- [SupplementaryData1.xlsx](#)
- [SupplementaryData2.pdf](#)
- [ExtendedDataFigures.docx](#)