Risk Factors Associated With the Presence of *Mycobacterium Bovis* in Macroscopic Lesions Suspected as Being Caused by Bovine Tuberculosis Detected in Slaughterhouses

Flávio Gomes Barcelos

UFMT: Universidade Federal de Mato Grosso

Tais Ramalho dos Anjos

Universidade de Cuiabá - Campus Beira Rio

Leila Auxiliadora de Arruda Alencar

UFMT: Universidade Federal de Mato Grosso

Maria Júlia Sudária de Freitas

Universidade de Cuiabá: Universidade de Cuiabá

Mariano Martinez Espinosa

UFMT: Universidade Federal de Mato Grosso

Ricardo Cesar Tavares Carvalho

Universidade de Cuiabá: Universidade de Cuiabá

Eduardo Eustáquio de Souza Figueiredo (✉ figueiredoeduardo@ufmt.br)

Universidade Federal de Mato Grosso <https://orcid.org/0000-0002-2608-4324>

Research Article

Keywords: Mycobacterium bovis, risk factors, nested q-PCR, slaughter.

Posted Date: June 9th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-581494/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Semina: Ciências Agrárias on February 25th, 2022. See the published version at <https://doi.org/10.5433/1679-0359.2022v43n2p713>.

Abstract

Mycobacterium bovis is a bacterium belonging to the *Mycobacterium tuberculosis* complex that causes tuberculosis in cattle and in other domestic and wild animals, as well as in humans. Disease control measures are carried out by slaughtering animals tested positive in the intradermal tuberculinization test and sanitation of their original living spaces, in addition to epidemiological surveillance carried out through the sanitary inspection of bovine carcasses in slaughterhouses. In the latter, official inspection services collect samples from macroscopic lesions suspected of bovine tuberculosis, which are then sent for laboratory analysis. Knowledge concerning the variables associated with the occurrence of *M. bovis* can aid in decision-making regarding control and disease eradication efforts. In this context, the aim of this study was to identify the risk factors for a positive *M. bovis* diagnosis in suspected bovine tuberculosis lesions obtained during epidemiological surveillance activities in the state of Mato Grosso, Brazil. A total of 105 suspicious lesions were analyzed using the Nested Polymerase Chain Reaction (nested q-PCR) method, of which 14 (13.33%) tested positive for *M. bovis*. Univariate and bivariate statistical analyses indicated that the variable “animal slaughter” was the only risk factor presenting statistical significance associated with the diagnosis of *M. bovis* ($p < 0.05$), demonstrating that macroscopic lesions suspected as being caused by bovine tuberculosis from animals with an *in vivo* diagnosis were 2.82 - fold more likely to result in a positive *M. bovis* diagnosis by molecular tests.

Introduction

Bovine tuberculosis (bTB) is a low prevalence disease. However, when diagnosed, it causes economic losses by reducing the zootechnical value of infected animals and herds, as well as consequent meat and milk production losses (Azami and Zinsstag 2018).

The etiologic agent of bovine tuberculosis is *Mycobacterium bovis* (*M. bovis*), a bacterium belonging to the *Mycobacterium tuberculosis* Complex (CMT), distributed worldwide and affecting cattle and several mammal species, including humans. Humans acquire the infection through contact with aerosols expelled by infected animals or mainly, through the consumption of milk and other foods from contaminated animals. It is, therefore, a serious public health problem, as it is a zoonosis (Medeiros et al. 2010; Zarden et al. 2013; Carvalho et al. 2016).

Epidemiological surveillance in slaughterhouses is applied in bTB monitoring, where suspicious lesions are identified and sent for laboratory diagnosis by bacteriological and molecular tests (Araújo et al. 2014a), allowing for the tracking of remaining disease *foci*. In the state of Mato Grosso, Brazil, the Joint Ordinance Sedraf - INDEA/MT 009 of November 14, 2014, established mandatory collection of granulomatous lesions displaying bTB characteristics from cattle slaughtered in slaughterhouses, followed by their shipment to a laboratory indicated by the health authority and notification of the injuris to the Agricultural Defense Service (Mato Grosso, 2014).

Characterization of bTB risk factors is important to determine situations that favor the persistence of this disease in a certain locality and to define strategies for locating remaining disease *foci*, thus aiding local health authorities in identifying regions or properties with animals that present a greater risk of contracting or spreading the disease.

In this context, the aim of this study was to identify the risk factors concerning the occurrence of *Mycobacterium bovis* in macroscopic lesions suspected as being caused by bovine tuberculosis found in bovine carcasses in the state of Mato Grosso, Brazil.

Material And Methods

Study area

This study was carried out in the state of Mato Grosso, Midwestern Brazil, which is categorized into four livestock circuits as described by Negreiros et al. (2009), called CP 01 - Pantanal; CP 02 - Milk; CP 03 - Fattening; CP 04 - Breeding; according to the predominant animal production characteristic, local ecosystem and animal transit network.

Sampling and zootechnical data

A total of 105 suspected bTB lesion samples were collected between September 2017 and May 2018, stored in sterile packaging and frozen during the post-mortem inspection procedure carried out in slaughterhouses in the state of Mato Grosso by Federal (SIF) and state (SISE) inspection authorities. Zootechnical information was also obtained during sampling, including cattle breed, sex, age group and geographic location of the animals' property of origin, for later data analysis. Simultaneously, sanitary inspection and collection of suspicious lesions were carried out following normal routine in all slaughterhouses in the state.

DNA extraction and quantification

Sample preparation and DNA extraction from bovine tissue fragments containing suspicious lesions were performed according to Figueiredo et al. (2012), using the commercial Dneasy Blood and Tissue kit (Qiagen®). Following extraction, DNA was quantified by the fluorometry method using the QUBIT™ 2.0 Kit (Invitrogen®).

Nested q-PCR

Nested q-PCR was performed using the TaqMan PCR Master Mix (Applied Biosystems®), primers and probe for the Rv2807 gene (Applied Biosystems®), specific for species belonging to the *Mycobacterium tuberculosis* complex (CMT) and primers and probe for the TbD1 gene (Applied Biosystems®), specific for *M. bovis*, based on the method described by Araújo et al. (2014a) and Araújo et al. (2014b) and modified by Carvalho et al. (2015).

The Nested q-PCR reaction occurred in two stages. During the first stage, a conventional PCR was performed to amplify the Rv2807 and TbD1 regions using external primers, *Forward Outer* GGCGGTGGCGGAGTTGAAGGCGATGAG, *Reverse Outer* CATTGCTGCGTAATTCGATCA; *Forward Outer* TTATGGCGGCCACACCCACCCAAAACAG, *Reverse Outer* GACCTTGGGCGCCTCAT, respectively and template DNA at 75 ng. This reaction was prepared in a total volume of 25 µL and was carried out according to the following amplification conditions: 95°C for 4 minutes for initial denaturation, 35 cycles at 95°C for 90 seconds for denaturation, 60°C for 30 seconds for ringing, 72°C for 45 seconds for extension and a final extension step carried out at 72 ° C for 3 minutes.

During the second stage of the reaction, a nested q-PCR was performed to amplify the Rv2807 and TbD1 regions using internal primers, TaqMan probe, *Forward Internal* CATTGCTGCGTAATTCGATCA, *Reverse Internal* GACCTTGGGCGCCTCAT, *Probe* 6FAM CATCCACACCTGTTCG MGBNFQ; *Forward Internal* GCGGTCTTCGCCAATGTT, *Reverse Internal* GCAGCCGATGGAATTGCT, *Probe* 6FAM TCGCGCAAGGCGA MGBNFQ, respectively and 3 µL of the amplicon obtained during the first stage. This reaction was prepared in a total volume of 12.5 µL and carried out according to the following amplification conditions: 95°C for 10 minutes for initial denaturation and 35 cycles at 95 ° C for 15 seconds for denaturation and 60°C for 30 seconds for annealing/extension.

Data analysis

For the statistical and descriptive analyses, the information contained in the lesion receiving forms was considered, comprising breed, sex, age group, geographic location and sanitary slaughter. The breed variable comprised two categories, Zebu and other breeds, the age group variable was formed by the categories 12–24 months, 24–36 months and > 36 months, variable geographic location comprised the livestock circuit categories 01 to 04, and the variable slaughter was composed of categories YES (positive animals in the intradermal tuberculinization test) and NO (animals without an *in vivo* diagnosis).

A descriptive analysis of the sample variables and a bivariate analysis for the molecular *M. bovis* diagnosis using Fisher's exact test, chi-square tests and the crude prevalence ratio, with a 95% confidence interval and a significance level of less than 5% ($p < 0.05$), were performed. The variables associated by the chi-square test were selected to adjust a Poisson regression model. The SPSS® Software (v. 20) was used for all statistical analyses.

Results

Univariate analysis

A univariate analysis (Table 1) was applied to indicate the probability of 105 suspected bTB lesions harvested from cattle in the state of Mato Grosso to be associated with one of the established variables, *i.e.*, sex, breed, age group, livestock circuit, slaughter and a molecular diagnosis of *M. bovis*.

Table 1

Frequency and distribution analysis of 105 suspected bovine tuberculosis lesions collected from cattle in the state of Mato Grosso, Brazil. Frequency, percentage and 95% confidence intervals for the variables sex, breed, age group, livestock circuit, slaughter, and a molecular diagnosis of *Mycobacterium bovis*, from September 2017 to May 2018, are presented.

Variable	Frequency (n = 105)	Percentage (%)	95% CI
Sex			
Female	36	34.29	(25.30–44.19)
Male	69	65.71	(55.81–74.70)
Bovine species			
Zebu	86	81.90	(73.19–88.74)
Others	19	18.11	(11.26–26.81)
Age range			
12–24 months	20	19.05	(12.04–27.87)
24–36 months	48	45.71	(35.96–55.72)
> 36 months	37	35.24	(26.16–45.17)
Livestock circuit			
CP 01 (Pantanal)	12	11.43	(6.05–19.11)
CP 02 (milk)	25	23.81	(16.04–33.11)
CP 03 (Fattening)	1	1.0	(0.02–5.19)
CP 04 (Breeding)	67	63.8	(53.86–72.96)
Sanitary slaughter			
No	98	93.33	(86.74–97.28)
Yes	7	6.67	(2.72–13.25)
Molecular <i>M. bovis</i> diagnosis			
Negative	91	86.67	(78.64–92.51)
Positive	14	13.33	(7.49–21.36)
*95% CI: Confidence Interval 95%.; CP: Livestock Circuit.			

From the total of 105 suspected bTB lesions collected, 86.67% (91/105) [IC95%: 78.64–92.51] tested negative for *M. bovis* and 13.33% (14/105) [IC95%: 7.49–21.36] tested positive.

Of the total of suspected injuries, 65.71% [IC95%: 55.81–74.70] were from male cattle and 34.29% [IC95%: 25.30–44.19] from females, indicating a statistically significant difference concerning sex, based

on the confidence interval. Regarding breed, 81.9% [IC95%: 73.19–88.74] of the suspected bTB lesions were obtained from zebu cattle, while 18.11% [IC95%: 11.26–26.81] were from other categories. With regard to age group, and 19.05% [IC95%: 12.0–27.87] of the suspected bTB lesions were obtained from cattle aged 12 to 24 months, 45.71% [IC95%: 35.96–55.72], from individuals aged 24 to 36 months and 35.24% [IC95%: 26.16–45.17] from animals over 36 months (Table 1).

The stratification concerning livestock circuit origin indicated that 63.8% [IC95%: 53.86–72.96] of the suspected bTB lesions originated from the livestock breeding circuit (CP 04), followed by the milk breeding circuit (CP 02), at 23.81% [IC95%: 16.04–33.11], Pantanal livestock circuit (CP 01), 11.43% [IC95%: 6.05–19.11] and fattening livestock circuit (CP 03), 1% [IC95%: 0.02–5.19]. The health slaughter variable indicated that only 6.67% [IC95%: 2.72–13.25] of the suspected bTB lesions were obtained from animals sent for slaughter, while 93.33% [IC95%: 86.74–97.28] of the harvested lesions came from commercial slaughter.

Bivariate Analysis

The associations between the molecular *M. bovis* diagnosis results and the other independent variables (Table 2), identified by a bivariate analysis, allowed for the determination that slaughter ($p = 0.048$) was the only variable presenting statistical significance ($p < 0.05$).

Table 2

Frequency distribution concerning the variables gender, breed, age group, livestock circuit, slaughter and molecular *Mycobacterium bovis* diagnosis in suspicious lesions obtained during the epidemiological surveillance of bovine tuberculosis in the state of Mato Grosso, Brazil, from September 2017 to May 2018.

Variable	Category	N° of injuries	Molecular diagnosis Positive for <i>M. bovis</i>		p value
			N°	%	
Sex	Female	36	6	16.67	0.549
	Male	69	8	11.60	
Breed	Others	19	2	10.53	1.00
	Zebu	86	12	13.95	
Age range	12–24 months	20	3	15	0.713
	24–36 months	48	5	10.42	
	> 36 months	37	6	16.22	
Livestock circuit	CP 01 (Pantanal)	12	0	0	0.206
	CP 02 (milk)	25	5	20	
	CP 03 (Fattening)	1	0	0	
	CP 04 (Breeding)	67	9	13.43	
Sanitary slaughter	No	98	11	11.22	0.048*
	Yes	7	3	42.86	
* significant at 95% CI; N: number of injuries; CP: Livestock Circuit.					

The factors associated with the molecular *M. bovis* diagnosis and the other assessed variables, after grouping the most relevant categories, again demonstrated that only slaughter ($p = 0.048$) exhibited a statistically significant difference (Table 3).

Table 3

Frequency distribution and prevalence ratio concerning the occurrence of *Mycobacterium bovis* by variable, in injuries obtained from the epidemiological surveillance regarding bovine tuberculosis in the state of Mato Grosso, from September 2017 to May 2018.

Variable	Category	<i>M. bovis</i> results				RP _b	95% CI	p < 0.05
		Nº Pos	%	Nº Neg	%			
Sex	Female	6	16.7	30	83.3	1.43	(0.54–3.83)	0.549*
	Male	8	11.6	61	88.4	1.00		
Breed	Zebu	12	13.9	74	86.1	1.32	(0.32–5.44)	1.0*
	Others	2	10.5	17	89.5	1.0		
Age Range	> 36 months	6	16.2	31	83.8	1.38	(0.52–3.67)	0.557*
	≤ 36 months	8	11.8	60	88.2	1.00		
Livestock circuit	CP 02 (milk)	5	20.0	9	80.0	1.49	(0.55–4.02)	0.515*
	CP 04 (Breeding)	9	13.4	58	86.6	1.00		
Sanitary slaughter	Yes	3	42.9	4	57.1	3.82	(1.38–10.60)	0.048
	No	11	11.2	87	86.7	1.00		
* Fisher's exact test; ** chi-square test for 2 x 2 type table; RP _b : Prevalence Ratio or Risk; Pos .: Positive; Neg .: Negative; p: p-value (< 0.005); 95% CI: 95% confidence interval; CP: Livestock Circuit.								

An analysis of the regression model of the equation indicated that the only risk factor associated with *M. bovis* diagnosis was slaughter (p < 0.05), due to a statistically significant difference. The data is presented in Table 4, indicating the positivity of suspicious bTB lesions from bovines tested positive to the intradermal tuberculin skin test.

Table 4
Risk factors associated with the occurrence of *Mycobacterium bovis* in macroscopic post-mortem lesions suspected of being caused by bovine tuberculosis obtained during the epidemiological surveillance for bovine tuberculosis in the state of Mato Grosso, Brazil, from September 2017 to May 2018.

Variable	RP _a	IC95%	Valor de p
Sanitary slaughter			
Yes	3.82	(1.38–10.60)	0.01*
No	1.00		
RPa: adjusted Prevalence Risk; CI: Confidence Interval			

Discussion

The present study considered the risk factors associated with slaughtered cattle variables and the stratification of the property of origin of animals presenting suspicious lesions for bTB in the state of Mato Grosso. The univariate analysis concerning sex indicated a statistically significant difference. However, when performing the bivariate analysis, this variable became non-statistically different, demonstrating that samples with suspicious lesions collected by the bTB surveillance service do not display an associated positive diagnostic relationship to sex. This corroborates the results described by several authors (Biffa et al. 2011; Clegg, Good and More 2016; Okeke et al. 2014; Pascual-Linaza et al. 2017).

Regarding the univariate analysis, lesions obtained from zebu cattle were statistically different compared to other breeds. However, the bivariate analysis indicated no significant association between zebu breeds and other breeds, most likely due to the fact that meat production in the state of Mato Grosso is mainly of zebu cattle, which is more representative in the sampling due to of the amount of cattle sent for slaughter during the study period. Similarly, studies carried out in Nigeria indicated no statistically significant differences concerning positive *M. bovis* diagnoses associated with breed, thus corroborating the results obtained herein (Akinseye et al. 2018; Okeke et al. 2014).

A statistically significant difference in the univariate analysis concerning suspicious lesions associated with age group was observed only between 12–24 months and 24–36 months, and no statistically significant difference between these two age groups were observed in relation to > 36 months. However, the bivariate analysis demonstrated that the age group variable was not significantly associated to lesion positivity for *M. bovis*, similar to that reported by Okeke et al. (2014) in Nigeria. Other authors described statistically significant difference, pointing to a higher risk of positive bTB lesions associated with older animals (Akinseye et al. 2018; Biffa et al. 2011; Boukary et al. 2012; Pascual-Linaza et al. 2017). Other reports associated young animals to a higher frequency of severe bTB lesions (Biffa et al. 2012),

demonstrating that lesions obtained from older animals are less likely to test positive for *M. bovis* (Shittu et al. 2013).

The univariate analysis regarding livestock circuits indicated no statistically significant difference in the frequency of injuries obtained from CP 01 and CP 02, and a statistically significant difference between the CP 03 and CP 04 circuits. In the bivariate analysis, however, no significant difference was observed among the livestock circuits, despite the constant frequency of bTB outbreaks in CP 02 (Leite) and CP 04 (Breeding). The bTB outbreaks were associated with dairy properties, where the health problem was attributed to greater animal density and long production cycles, corroborating Néspoli et al. (2016).

The health slaughter variable was statistically different concerning injuries obtained from animals positive for bTB (07/105) and animals commercially sent for slaughter (98/105) in the univariate analysis. In the bivariate analysis, slaughter was the only statistically different ($p < 0.05$) variable, indicating that animals with a positive diagnosis for bTB (positive for *in vivo* diagnosis) were 2.82 - fold more likely to present suggestive lesions confirmed as being caused by bTB by the nested q-PCR test.

Regarding the descriptive statistical analysis, 42.86% (07/03) of the injuries originating from slaughtered animals (07/105), were confirmed as positive for the molecular diagnosis of *M. bovis*. Cazola et al. (2015) reported that, from a total of 13 cattle positive in the intradermal test, 84.6% (11/13) were confirmed for *M. bovis* by the bacterial culture method (Cazola et al. 2015), demonstrating a high rate of disease confirmation. Fráguas et al. (2008) indicated that, from a total of 97 bovine animals reactive to the intradermal test, only 11.34% (11/97) were confirmed as *M. bovis* in bacterial cultures and postulated that this low bacterial recovery rate may be related to the difficulty of carrying out *M. bovis* cultures, which requires a minimum amount of viable bacilli, in addition to a decontamination technique applied during the process that can render this microorganism unfeasible.

Zarden et al. (2013) followed the slaughter of 24 cattle reactive to the tuberculin skin test, with 29.16% ($n = 7$) presenting suspicious lesions for bTB. When submitting samples without lesions for laboratory analysis, it was found that 73.16% ($n = 19$) of the animals tested positive when using bacterial culture, while 75.16% ($n = 18$) tested positive by the molecular method. The authors concluded that the combination of these methods increased sensitivity and specificity, with 95.8% ($n = 23$) of the animals testing positive for *M. bovis*, reinforcing that inconclusive tuberculinization results presented greater diagnosis chances through suspicious lesions. Figueiredo et al. (2009) performed a pool of samples of individual macroscopic lesions from 34 animals reactive to the tuberculin skin test and obtained a 50% ($n = 17$) positive diagnosis in the bacterial culture method, with 88.24% ($n = 15$) of the samples confirmed through multiplex-PCR.

Araújo et al. (2014a) collected 80 lesion samples from animals reactive to the intradermal tuberculin skin test, and observed presence of *M. bovis* DNA in 61.25% ($n = 49$) by the nested-PCR technique, with a greater number of animals testing positive for the pathogenic agent when compared to bacterial culture in the group of animals without information or without tuberculin tests ($p < 0.05$), that is, animals that were commercially sent to slaughter and observed as presenting lesions in the sanitary inspection line.

Risk factors may vary between surveys, due to regional differences, such as bTB prevalence, climatic variation, breed characteristics, management, production cycle, degree of property technification and strains. Each region may present intrinsic risk factors that must be considered when applying risk-based surveillance strategies, providing better resource application and promoting improvements in local agricultural defense management.

Taking into account that sex, breed, age group and livestock circuit were not significantly different or associated with factors that influence the diagnosis of bTB in the state of Mato Grosso and that slaughter sanitary conditions were statistically different ($p < 0.05$) in confirming the diagnosis of bovine tuberculosis, a greater investment in the *in vivo* diagnosis of the disease is suggested, as a way of increasing the chances of identifying and tracking new bovine tuberculosis *foci* in the state of Mato Grosso.

Given the above, the combination of intradermal and molecular tuberculin diagnostic tests potentiate the identification of *M. bovis* and the location of remaining disease *foci*, suggesting that the adopted strategy is appropriate for maintaining the health profile of the state of Mato Grosso.

Declarations

Acknowledgements

The authors are grateful to INDEA, Mato Grosso, Brazil, for the partnership in providing bovine samples suspected of having bovine tuberculosis

Funding

The authors also acknowledge financial support from the Fundação de Amparo a Pesquisa do Estado de Mato Grosso (Foundation for Research Support in Mato Grosso State) FAPEMAT Grant No. 366047/2017.

Conflict of interest

The authors declare that they have no conflicts of interest.

Ethic approval

The project was approved by the Committee on Ethics in Animal Use of Federal University of Mato Grosso (CEUA protocol 23108.226903/2017-72).

Consent to participate

not applicable.

Availability of data and material

not applicable

Code availability

not applicable

Authors' contributions

FGB: Investigation, Methodology, Writing – Original Draft, Writing – Review and Editing, Visualization, Project administration. **TRA:** Formal Analysis, Data Curation, Writing – Review and Editing, Visualization. **LAAA:** Methodology. **MJSF:** Methodology. **MME:** Statistical methodology. **RCTC:** Writing – Review and Editing. **EESF:** Conceptualization, Resources, Writing – Review and Editing, Visualization, Supervision.

References

1. Akinseye, V. O., Adebayo, M. D., Genesis, O. O., Adelakun, O. D., Cadmus, S. I., 2018. Prevalence and risk factors of mycobacterial infections in farm and trade cattle in southwestern Nigeria. *Tropical animal health and production*, 50(4), 761-772.
2. Araújo, C. P., Osorio, A. L. A., Jorge, K. S., Ramos, C. A. N., S Filho, A. F., Vidal, C. E. S., Roxo, E., Nishibe, C., Almeida, N. F., Júnior, A. A. F., Silva, M. R., Neto, J. D. B., Cerqueira, V. D., Zumárraga, M. J., Araújo, F. R., 2014a. Detection of *Mycobacterium bovis* in bovine and bubaline tissues using nested-PCR for TbD1. *PLoS One*, 9(3).
3. Araújo, C. P., Osório, A. L. A., Jorge, K. S., Ramos, C. A., Souza Filho, A. F., Vidal, C. E., Vargas, A. P. C., Roxo, E., Rocha, A. S., Suffys, P. N., Fonseca Júnior, A. A., Silva, M. R., Barbosa Neto, J. D., Cerqueira, V. D., Araújo, F. R., 2014b. Direct detection of *Mycobacterium tuberculosis* complex in bovine and bubaline tissues through nested-PCR. *Brazilian Journal of Microbiology*, 45(2), 633-640.
4. Azami, H. Y., Zinsstag, J. Economics of Bovine Tuberculosis: A One Health Issue. In: Chambers, M., Gordon, S., Olea-Popelka, F., Barrow, P. (Eds.). *Bovine Tuberculosis*. London, UK: British Library, 2018. cap. 3, p. 31-42.
5. Biffa, D., Inangolet, F., Bogale, A., Oloya, J., Djønne, B., Skjerve, E., 2011. Risk factors associated with prevalence of tuberculosis-like lesions and associated mycobacteria in cattle slaughtered at public and export abattoirs in Ethiopia. *Tropical animal health and production*, 43(2), 529-538.
6. Biffa, D., Bogale, A., Godfroid, J., Skjerve, E., 2012. Factors associated with severity of bovine tuberculosis in Ethiopian cattle. *Tropical animal health and production*, 44(5), 991-998.
7. Boukary, A. R., Thys, E., Rigouts, L., Matthys, F., Berkvens, D., Mahamadou, I., Yenikoye, A., Saegerman, C., 2012. Risk factors associated with bovine tuberculosis and molecular characterization of *Mycobacterium bovis* strains in urban settings in Niger. *Transboundary and emerging diseases*, 59(6), 490-502.
8. Carvalho, R. C. T., Furlanetto, L. V., Maruyama, F. H., de Araújo, C. P., Barros, S. L. B., do Nascimento Ramos, C. A., Dutra, V., Araújo, F. R., Paschoalin, V. M. F., Nakazato, L., de Souza Figueiredo, E. E.,

2015. Evaluation of the efficiency of nested q-PCR in the detection of *Mycobacterium tuberculosis* complex directly from tuberculosis-suspected lesions in post-mortem macroscopic inspections of bovine carcasses slaughtered in the state of Mato Grosso, Brazil. *Meat science*, 106, 11-15.
9. Carvalho, R. C. T., Vasconcellos, S. E. G., de Azevedo Issa, M., Soares Filho, P. M., Mota, P. M. P. C., de Araújo, F. R., Carvalho, A. C. S., Gomes, H. M., Suffys, P. N., de Souza Figueiredo, E. E., Paschoalin, V. M. F., 2016. Molecular typing of *Mycobacterium bovis* from cattle reared in Midwest Brazil. *PloS one*, 11(9).
10. Cazola, D. D. O., Jorge, K. D. S., Zumárraga, M. J., Souza-Filho, A. F., Araújo, F. R., Osório, A. L. A., 2015. Identificação e genotipagem de *Mycobacterium bovis* em bovinos positivos no teste intradérmico para tuberculose em Mato Grosso do Sul. *Pesquisa Veterinária Brasileira*, 35(2), 141-147.
11. Clegg, T. A., Good, M., More, S. J., 2016. Risk factors for cattle presenting with a confirmed bTB lesion at slaughter, from herds with no evidence of within-herd transmission. *Preventive veterinary medicine*, 126, 111-120.
12. Figueiredo, E. E. D. S., Silvestre, F. G., Campos, W. N., Furlanetto, L. V., Medeiros, L., Lilenbaum, W., Fonseca, L. S., Silva, J. T., Paschoalin, V. M. F., 2009. Identification of *Mycobacterium bovis* isolates by a multiplex PCR. *Brazilian Journal of Microbiology*, 40(2), 231-233.
13. Figueiredo, E. E. S., Júnior, C. A. C., Furlanetto, L. V., Silva, F. G. S., Duarte, R. S., Silva, J. T., Lilenbaum, W., Paschoalin, V. M. F., 2012. Molecular techniques for identification of species of the *Mycobacterium tuberculosis* complex: The use of multiplex PCR and an adapted HPLC method for identification of *Mycobacterium bovis* and diagnosis of bovine tuberculosis. In: CARDONA, P.-J. (Ed.) *Understanding tuberculosis: global experiences and innovative approaches to the diagnosis*, 411.
14. Fráguas, S. A., Cunha-Abreu, M. S., dos Reis Ferreira, A. M., Marassi, C. D., Oelemann, W., de Souza Fonseca, L., Ferreira, R., Lilenbaum, W., 2008. Estudo comparativo de métodos complementares para o diagnóstico da tuberculose bovina em animais reagentes à tuberculinização. *Revista Brasileira de Ciência Veterinária*, 15(3), 117–121.
15. Mato Grosso, 2014. Portaria Conjunta SEDRAF-INDEA/MT n. 009, 14 de novembro de 2014. Institui o regulamento técnico do plano de vigilância para a Erradicação da Tuberculose bovina no Estado de Mato Grosso e cria o Programa Estadual de Submissão de Granuloma do Estado de Mato Grosso. *Diário Oficial do Estado de Mato Grosso*. Cuiabá, MT, 13 nov. 2014.
16. Medeiros, L. D. S., Marassi, C. D., Figueiredo, E. E. S., Lilenbaum, W., 2010. Potential application of new diagnostic methods for controlling bovine tuberculosis in Brazil. *Brazilian Journal of Microbiology*, 41(3), 531-541.
17. Negreiros, R. L., Dias, R. A., Ferreira, F., Ferreira Neto, J. S., Gonçalves, V. S. P., Silva, M. D. C. P., Figueiredo, V. C. F., Lôbo, J. R., Freitas, J. Amaku, M., 2009. Situação epidemiológica da brucelose bovina no Estado de Mato Grosso. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, 61, 56-65.
18. Néspoli, J. M. B., Negreiros, R. L., Amaku, M., Dias, R. A., Ferreira, F., Telles, E. O., Heinemann, M. B., Grisi-Filho, J. H. H., Gonçalves, V. S. P., Neto, J. S. F., 2016. Epidemiological situation of bovine

- tuberculosis in the state of Mato Grosso, Brazil. *Semina: Ciências Agrárias*, 37(5), 3589-3599.
19. Okeke, L. A., Cadmus, S., Okeke, I. O., Muhammad, M., Awoloh, O., Dairo, D., Waziri, E. N., Olayinka, A., Nguku, P. M., Fawole, O., 2014. Prevalence and risk factors of *Mycobacterium tuberculosis* complex infection in slaughtered cattle at Jos South Abattoir, Plateau State, Nigeria. *The Pan African Medical Journal*, 18(Suppl 1).
 20. Pascual-Linaza, A. V., Gordon, A. W., Stringer, L. A., & Menzies, F. D., 2017. Efficiency of slaughterhouse surveillance for the detection of bovine tuberculosis in cattle in Northern Ireland. *Epidemiology & Infection*, 145(5), 995-1005.
 21. Shittu, A., Clifton-Hadley, R. S., Ely, E. R., Upton, P. U., Downs, S. H., 2013. Factors associated with bovine tuberculosis confirmation rates in suspect lesions found in cattle at routine slaughter in Great Britain, 2003–2008. *Preventive Veterinary Medicine*, 110(3-4), 395-404.
 22. Zarden, C. F. O., Marassi, C. D., Carvalho, A. C., Figueiredo, E. E. S., Lilenbaum, W., 2013. Bacteriological and molecular detection of *Mycobacterium bovis* in cattle with inconclusive results to intradermal tuberculin tests. *Epidemiology & Infection*, 141(7), 1390-1393.