

Coblation Versus Diode Laser Tonsillectomy: A Comparative Study

Ameer A. Alaqeedy

University Of Anbar, College of Medicine, Department of Surgery/Otolaryngology

Rasheed Ali Rashid

Tikrit University, College of Medicine, Department of Surgery/ Otolaryngology

Raid M. Al-Ani (✉ med.raed.alani2003@uoanbar.edu.iq)

University Of Anbar, College of Medicine, Department of Surgery/ Otolaryngology

<https://orcid.org/0000-0003-4263-9630>

Research Article

Keywords: Coblation, Diode Laser, Tonsillectomy, Comparative Study

Posted Date: June 4th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-582705/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Introduction: The study aimed to compare the 2 procedures of tonsillectomy (coblation and diode laser) concerning the operative time, the amount of blood loss, postoperative pain, and other complications.

Materials and Methods: A comparative study was conducted at many Private Hospitals, Baghdad, Iraq from February 2019- February 2020. Coblation and laser tonsillectomy were performed on the same patient (one for each side) which was blinded to the patients or their caregivers and the surgeon who did the tonsillectomy. Data concerning the age, gender, indications for tonsillectomy, operative time, the amount of the blood loss, post-tonsillectomy pain by VAS scores, and other complications were recorded for each patient.

Results: Out of 62 participants, there were 34 (54.8%) females. The majority were from the age group ≤ 18 years ($n=54$, 87.1%). The operative time and amount of the intraoperative blood loss were significantly lower in the coblation than a diode laser tonsillectomy. However, the postoperative pain scores in the diode laser were a statistically significant difference less than the coblation group at the periods 3 hours, 8 hours, 24 hours, and 7 days ($P\text{-value}<0.05$). While there was no statistically significant difference between the two groups at 2 weeks postoperative period ($P\text{-value}=0.392$). No primary or secondary postoperative bleeding in all patients.

Conclusion: coblation was better than diode laser because of shorter operative time and less intraoperative blood loss. However, the diode laser technique had lesser postoperative pain at the postoperative periods 3 hours, 8 hours, 24 hours, and 7 days than coblation.

Introduction

The Palatine tonsils are a pair of lymphoid tissue located on the lateral wall of the oropharynx between the palatoglossus (anterior pillar) and palatopharyngeus (posterior pillar). One of the most common operations in the field of otolaryngology is a tonsillectomy, which is most commonly indicated in cases of recurrent tonsillitis or obstructive upper airway due to tonsillar enlargement (1)(2). Conventional dissection tonsillectomy is considered a gold standard procedure for a long period (3). Tonsillectomy is performed on both children and adult populations. Although, the procedure is common but still carries significant complications particularly postoperative pain and bleeding. There are many surgical methods of tonsillectomy used in daily practice, without consensus among otolaryngologists on the best procedure (4).

The drawback of the conventional dissection tonsillectomy is an open postoperative wound that heals by secondary intention with its sequel of postoperative pain and bleeding. Therefore, otolaryngologists use many other procedures for removal of the tonsils such as laser, electrocautery, coblation, radio-frequency, monopolar or bipolar diathermy, and harmonic scalpel (5).

The term coblation is derived from controlled ablation. It was first discovered by Hira V. Thapliyal and Philip E. Eggers (6). It depends on radiofrequency energy. It passes through an isotonic sodium chloride and creates a plasma field. As a result, free sodium ions are dissociated, and able to destroy the intercellular bonds and hence dissociation of the tissues. This process is achieved at a lower temperature (60–70°C) than the electrocautery (400–600°C). The cooling effect of the irrigating isotonic saline protects the surrounding tissues from the high heat (7)(8).

There are various types of lasers have been used in tonsillectomy procedure like CO₂, KTP, NDYAG, and diode (9). The advantages of the diode laser are good thermal effect on the perifocal tissues with shallow depth of penetration, thus carries few side effects to the deep tissues. Therefore, it is very useful in the tonsillectomy technique because of low postoperative complications with excellent outcomes (10).

There are various investigations compare the various techniques of tonsillectomy (3)(5)(8)(11)(12)(13). However, few studies investigate the comparison between the coblation and diode laser tonsillectomy in children (14)(15). Hence, we aimed to compare the coblation and diode laser tonsillectomy in pediatrics and adult populations regarding the operative time, intraoperative blood loss, and postoperative pain and complications.

Materials And Methods

This double-blinded randomized controlled trial was conducted at many Private Hospitals, Baghdad city, Iraq. The study covered the period of one year (February 2019 to February 2020). Patients of any age and both sexes with an indication of tonsillectomy (recurrent tonsillitis or tonsillar enlargement) were enrolled in the study. The exclusion criteria included patients with tonsillectomy and other procedures like adenoidectomy, and grommet insertion, previous attack of quinsy, patients who didn't wish to participate in the study, patients with bleeding disorders, and those who lost of follow-up.

The patients were prepared for surgery by a diode laser on one side and coblation cautery on the other side. Both the patient and his attendant had not told which method was used for which side, by this, the patient will be as a control for symptoms while the signs of healing were looked for which are different from patient to patient. Besides the otolaryngologist who did the operation was not participating in the statistical analysis of the data. Informed consent was taken from the patients or their parents. The current study was approved by the Ethical Approval Committee of the University of Anbar (reference number 33, 8-4-2021).

We also recorded the time of surgery and amount of blood loss as the size, vascularity, state of inflammation, and adhesion would be more closely correlated in the same person than between two different patients.

Surgical Method

All cases had general anesthesia in a supine position with a draping of the whole body exposing only the mouth, then a Boyle –Davis mouth gag was inserted with Draffin suspensors applied. The tonsil was held by tonsil holding forceps and pulled medially, by using either coblation or diode laser handle, the anterior pillar was incised till reaching the tonsil capsule, then the tonsil dissected till reaching the inferior pole starting from the superior one.

In the coblation tonsillectomy group, we used a Coblator® II system (ENTec, Sunnyvale, California, USA) and the Evac T&A Plasma Wand (ENTec) was used at an intensity setting of 7 for dissection of the tonsils at the subcapsular plane. Bleeding points were secured using an intensity setting of 4 on coagulation mode.

In the diode laser tonsillectomy group, a diode laser (980-nm wavelength with a fiber-optic delivery system; Ceralas™ D25, 980 ± 30 nm, 25 W, model 003; CeramOptec, Bonn, Germany) was used. The machine was set to 5 W continuous beams for subcapsular dissection. Besides, bleeding points were secured by diode laser. The laser beam was delivered via a 0.6 mm Endostat fiber. Routine precautions for the safety of theatre personnel were followed during the use of the laser. The amount of blood loss was estimated by using standard pieces of gauze which were soaked in normal saline then squeezed then weighed by an electric scale before and after it is used to dry blood oozing from the incision or dissection.

The time taken to remove each tonsil was measured by minutes and any complications that occurred during surgery were recorded.

After surgery, the patients were kept in the ward, monitored for vital signs and signs of bleeding regularly (six-hourly for 24 hours) then they were discharged home.

The two sides were compared for pain in the throat and healing signs at 3 hours, 8 hours, 24 hours, 7 days, and 2 weeks post-operatively. Intravenous and oral paracetamol per body weight was the analgesic used according to the standard pain control regime in all cases given 3 times per day for the first day. Thereafter, the analgesic is used according to the need of the patients. Additionally, prophylactic amoxicillin for the first 10 post-tonsillectomy periods was given for all patients. All patients were seen at 7, 14, and 28 postoperative days in the outpatient clinic.

Pain assessment depends on a 0–10 visual analog scale (VAS) score of pain, the mean ± SD was measured for each subject. The VAS was valid and used in the past for the assessment of postoperative pain in children and adults after tonsillectomy (16)(17).

The data were entered and analyzed using IBM SPSS version 25 for windows. The categorical variables were presented in a simple table or figure of the frequencies and percentages. Chi-Square test used for the comparison of the categorical variables. While independent sample T-Test was used to compare the means of the variables. P-value at < 0.05 considered a statistically significant difference.

Results

Sixty-two patients fulfilled the inclusion criteria and enrolled in the study. The age ranged from 8 to 31 years with a mean age of 10.84 ± 7.418 . The most involved age group was ≤ 18 years ($n=54, 87.1\%$). Females ($n = 34, 54.8\%$) out the number of the males ($n = 28, 45.2\%$), with a male/female ratio of 1/1.21 Table 1.

Table 1
The distribution of the 62 patients according to the age group and gender.

Gender	Age groups		Total
	≤ 18 years	> 18 years	
	N/%	N/%	N/%
Males	27 (96.4)	1 (3.6)	28 (45.2)
Females	27 (79.4)	7 (20.6)	34 (54.8)
Total	54 (87.1)	8 (12.9)	62 (100)

Recurrent infection comprises 64.5% ($n = 40$), of the tonsillectomy indications and tonsillar enlargement 35.5% ($n = 22$) Fig. 1.

The mean operative time for coblation and diode laser tonsillectomy was 10.39 minutes ± 2.059 and 11.61 minutes ± 2.228 respectively with a statistically significant difference between the two groups (P -value = 0.002). Besides, there was a statistically significant difference between the two groups (P -value = 0.003) regarding the amount of the mean blood loss (21.61 ml ± 3.973 for coblation and 23.87 ml ± 4.287 diode laser tonsillectomy). The mean VAS of pain at the 3, 8 hours, 24 hours, 7 days, and 2 weeks for the coblation group were 8.71 ± 0.857 , 8.45 ± 0.953 , 7.24 ± 0.824 , 0.31 ± 0.841 , and 0.10 ± 0.564 respectively, while in the diode laser group 8.10 ± 0.918 , 7.97 ± 0.923 , 6.65 ± 0.832 , 0.03 ± 0.178 , and 0.03 ± 0.178 respectively. Besides, the mean VAS of pain was statistically significantly less in the diode laser than the coblation group at the first 4 postoperative periods (P -value < 0.05). While there was no statistically significant difference between the 2 groups at the 2 weeks postoperative period (P -value = 0.392) Table 2. There were no postoperative primary or secondary bleeding or other complications in all patients. At 4 weeks follow-up visit, the tonsillar fossae of all patients were completely healed without sloughing and necrotic tissues.

Table 2

The relationship between the type of operation and the mean operative time, mean blood loss, and VAS scores post-operative pain at different periods in 62 patients.

Variable	Tonsillectomy		P-value
	Coblation	Diode laser	
Mean operative time	10.39 ± 2.059	11.61 ± 2.228	0.002
Mean blood loss	21.61 ± 3.973	23.87 ± 4.287	0.003
Post-operative pain			
3 hours	8.71 ± 0.857	8.10 ± 0.918	0.000
8 hours	8.45 ± 0.953	7.97 ± 0.923	0.005
24 hours	7.24 ± 0.824	6.65 ± 0.832	0.000
7 days	0.31 ± 0.841	0.03 ± 0.178	0.013
2 weeks	0.10 ± 0.564	0.03 ± 0.178	0.392

Discussion

Despite tonsillectomy is one of the most common surgical procedures performed by an otolaryngologist through various techniques, postoperative bleeding and pain remain complications of this operation. The otolaryngologists try to practice and compare different procedures of tonsillectomy to achieve the best way regarding the shorter operative time with less blood loss, avoidance of intra and postoperative complications including mainly post-tonsillectomy hemorrhage and pain. We aimed to compare the diode laser and coblation tonsillectomy both in children and adults. The strength of the study was the use of both procedures in the same patient (coblation on one side and diode laser on the other side) to lessen the bias. Similar studies used 2 methods of tonsillectomy in the same patient (11)(18). The study revealed that the coblation was better than the diode laser method regarding the shorter operative time and less blood loss during the procedure. However, the diode laser was superior to the coblation tonsillectomy concerning postoperative pain.

Although coblation and laser tonsillectomies are relatively new techniques, if one compares them with the cold steel dissection method, they have gained popularity in the otolaryngology practice owing to the advancement in the technology of their machines.

In Iraq, few randomized controlled trials studied the comparison between the cold steel dissection method with CO₂ laser (19), coblation and bipolar diathermy (12), and coblation versus dissection method (20). This study was the first study in Iraq compared the diode laser and coblation tonsillectomy in children and adults.

Powell et al. in 1997 studied the effect of radiofrequency on the tissue of the tongue in animal models, both in vitro and in vivo. They concluded that radiofrequency can reduce the volume of the tongue in a perfect and controlled way (21). Since that time, many otolaryngologists start the use of coblation tonsillectomy in their daily clinical practice [5](12)(22)(18). The outcomes were good in studies involving a small size of patients (22)(18). Moreover, many randomized controlled studies comparing the coblation with other techniques of tonsillectomy revealed the superiority of this kind over other procedures regarding the operative time, amount of intraoperative blood loss, and postoperative pain (14)(15). Basu et al. study reported that harmonic scalpel tonsillectomy is better than coblation in the following parameters operative time, intraoperative blood loss volume, and postoperative bleeding (5). Accordingly, the study is considered this method as an advanced way of tonsillectomy. Other studies reported equivocal results with other procedures.

In a study from China by Zhou et al.(13), they compare retrospectively the coblation (300 patients) with coblation and tie tonsillectomy (215 patients) in the adult population regarding the operative time, blood loss during the procedure, post-tonsillectomy pain and complications, and full recovery and return to normal activities. The study concluded that the coblation tonsillectomy alone was statistically significantly superior to coblation with tie regarding all the above-mentioned parameters except post-tonsillectomy bleeding, coblation with a tie was significantly less post-operative bleeding than coblation tonsillectomy alone.

A recent systematic review study by Ahmed and Arya (23) reviewed 15 randomized controlled trial articles that studied tonsillectomy by laser or other procedures. They reported that the majority of laser tonsillectomy was performed by CO2 laser (n = 665, 60.3%), followed by potassium-titanyl-phosphate (KTP) laser (n = 238, 21.6%), and the least diode laser (n = 199, 18.1%). While, tonsillectomies performed by other techniques were 792 (64.7%) by cold dissection, 238 (19.4%) by diathermy, and 194 (15.8%) by coblation method. They concluded that there is an overall improvement in laser tonsillectomy in comparison with other techniques and this may be due to increased familiarity with laser surgeries and increment in the number of laser centers across the globe. However, our study revealed that the coblation tonsillectomy was better than the diode laser regarding the operative time and blood loss volume during surgery. The better results of coblation in comparison with diode laser in operative time and intraoperative blood loss may be attributed to the larger probe size of the coblation handle with a wider surface area of effect concerning the diode laser handle. Similar findings were observed in previous studies (14)(15). But the results of our study (significantly lower VAS scores of postoperative pain in diode laser in comparison with coblation technique) in contrast to the above-mentioned studies concerning postoperative pain. The significantly less postoperative pain at least in the first postoperative week with diode laser may be due to the precise cut with less necrotic tissues and less damage to surrounding structures.

A recent study by Kumar et al. (24) reported that the operative time and bleeding during surgery were significantly low for CO2 laser in comparison with the cold steel dissection tonsillectomy group. Pain score was comparable in early post-operative days but was high towards the end of the first week. Our

results revealed that the pain scores were significantly less in the laser group in the first 7 days post-tonsillectomy in comparison with coblation (P-value < 0.05). While at 2 weeks, there was no statistically significant difference (P-value = 0.392) between the two groups. This may be attributed to two causes, firstly, the laser has a desensitizing effect on the cut ends of the nerve fibers in the few postoperative days, thereafter, the pain sensation is increased. Secondly, the thermal effect of the laser needs more time to heal, in addition to the formation of a thick slough in the tonsillar area with possible late-onset of contraction and pain (25). Of note, secondary infection may occur beneath the thick slough layer with the increased chances of the development of secondary hemorrhage. Therefore, it is suggested to use a prophylactic antibiotic in the post-tonsillectomy period by many researchers to avoid such complications.

Post-tonsillectomy bleeding (primary or secondary) is a major complication of tonsillectomy with variable incidences (0.95%-7.8%) among previous investigations (2)(8)(5)(14)(26). The difference in these incidences may be attributed to the following; type of tonsillectomy, primary or secondary bleeding, children or adult patients, and whether an antibiotic is used as prophylactic or not. Many investigations have observed different incidences of delayed post-tonsillectomy bleeding between children and adults, with more incidence in adults than children (2) (8). Fortunately, our study didn't report a case of post-tonsillectomy bleeding. This may be an incidental finding.

The parameters in defining the improvement of tonsillectomy operation are shorter operative time, less intraoperative blood loss, low incidence of primary or secondary post-tonsillectomy bleeding, less postoperative pain, and early recovery and return to usual activities. Despite, many researchers studied these parameters among various tonsillectomy methods (12)(13)(14)(18), there is still no consensus about the best method of tonsillectomy. Therefore, the following factors should be taken into consideration when choosing the tonsillectomy procedure; availability of devices and instrumentations, the experience of the surgeon, preference of the patients or parents, and the cost of the procedure.

Although the current study tried to avoid result bias by comparing the two sides of the same patient still it did not bypass the intrapersonal differences in size, fibrosis, and blood supply between the two tonsils of the same patient. This actually cannot be overcome and may have some effect on the results but the arbitrary selection of the sides may minimize this effect. Even though the time in preparing the 2 devices didn't take into consideration when calculating the operative time in the current study, it had a drawback on the patients as it prolongs the total operative time with its risk of anesthesia. Therefore, we don't advise using 2 techniques of tonsillectomy in the same patient in daily clinical practice as here we used them for research purposes.

Conclusion

The study revealed that the coblation tonsillectomy was better than the diode laser concerning the operative time and amount of blood loss. However, the diode laser had significant low postoperative pain VAS scores in the first 7 days postoperatively.

Declaration

The authors declare no competing interests.

References

1. Windfuhr JP, Toepfner N, Steffen G, Waldfahrer F, Berner R. Clinical practice guideline: tonsillitis II. Surgical management. *Eur Arch Oto-Rhino-Laryngology*. 2016;273(4):989–1009.
2. Al-Ani RM. Post-tonsillectomy bleeding: Incidence and risk factors running title: PTB: Incidence and risk factors. *IRAQI JOURNAL OF COMMUNITY Med*. 2012;25(3).
3. Pinder DK, Wilson H, Hilton MP. Dissection versus diathermy for tonsillectomy. *Cochrane Database Syst Rev*. 2011;(3).
4. Burton MJ, Doree C. Coblation versus other surgical techniques for tonsillectomy. *Cochrane database Syst Rev*. 2007;(3).
5. Basu S, Sengupta A, Dubey AB, Sengupta A. Harmonic scalpel versus coblation tonsillectomy a comparative study. *Indian J Otolaryngol Head Neck Surg*. 2019;71(4):498–503.
6. Thiagarajan B. Coblation Tonsillectomy our experience. *Otolaryngol online J [Online]*. 2014;4(1.5).
7. Derkay CS, Maddern BR. Innovative techniques for adenotonsillar surgery in children: introduction and commentary. *Laryngoscope*. 2002;112(S100 S100):2.
8. Belloso A, Chidambaram A, Morar P, Timms MS. Coblation tonsillectomy versus dissection tonsillectomy: postoperative hemorrhage. *Laryngoscope*. 2003;113(11):2010–3.
9. Sayin I, Cingi C. Recent medical devices for tonsillectomy. *Hippokratia*. 2012;16(1):11.
10. Sedlmaier B, Bohlmann P, Jakob O, Reinhardt A. Outpatient diode laser tonsillotomy in children with tonsillar hyperplasia. Clinical results. *HNO*. 2010;58(3):244–54.
11. Kassab AN, Ahmed MR, Saber M, Mekawy S. Comparative intraindividual ablative tissue effects of diode laser 980 nm versus radiofrequency in tonsillar hypertrophy management. *ACTA Otorhinolaryngol Ital*. 2019;39(3):150.
12. Saeed HI, Emin A, Yaseen MA, Saleh AM. A comparative study between coblation and bipolar electrocautery tonsillectomy in children. *Zanco J Med Sci (Zanco J Med Sci)*. 2019;23(1):81–9.
13. Zhou X, Xu A, Zhen X, Gao K, Cui Z, Yue Z, et al. Coblation tonsillectomy versus coblation tonsillectomy with ties in adults. *J Int Med Res*. 2019;47(10):4734–42.
14. Elbadawey MR, Hegazy HM, Eltahan AE, Powell J. A randomised controlled trial of coblation, diode laser and cold dissection in paediatric tonsillectomy. *J Laryngol Otol*. 2015;129(11):1058–63.
15. Matin MA, Sabur MA, Azad MAK, Islam MN. Coblation versus Diode Laser Tonsillectomy in Children: A Randomized Controlled Trial. *Bangladesh J Otorhinolaryngol*. 2017;23(2):140–5.
16. Warnock FF, Lander J. Pain progression, intensity and outcomes following tonsillectomy. *Pain*. 1998;75(1):37–45.

17. Arbin L, Enlund M, Knutsson J. Post-tonsillectomy pain after using bipolar diathermy scissors or the harmonic scalpel: a randomised blinded study. *Eur Arch Oto-Rhino-Laryngology*. 2017;274(5):2281–5.
18. Timms MS, Temple RH. Coblation tonsillectomy: a double blind randomized controlled study. *J Laryngol Otol*. 2002;116(6):450.
19. Al-Rubeai TM, Bader HM, Ali SF. CO2 Laser Tonsillectomy: A Comparison with the Standard Technique. *Med J Babylon*. 2009;6(2).
20. Tahyr SE, Yaseen MA-R. Coblation versus dissection tonsillectomy in children. *Zanco J Med Sci (Zanco J Med Sci)*. 2018;22(1):25–31.
21. Powell NB, Riley RW, Troell RJ, Blumen MB, Guilleminault C. Radiofrequency volumetric reduction of the tongue: a porcine pilot study for the treatment of obstructive sleep apnea syndrome. *Chest*. 1997;111(5):1348–55.
22. Temple RH, Timms MS. Paediatric coblation tonsillectomy. *Int J Pediatr Otorhinolaryngol*. 2001;61(3):195–8.
23. Ahmed J, Arya A. Lasers in Tonsillectomy: Revisited With Systematic Review. *Ear, Nose Throat J*. 2021;100(1_suppl):14S-18S.
24. Kumar A, Kumar S, Krishnan A. A Comparative Analysis of Outcomes of Conventional Cold Dissection Versus Laser Tonsillectomy in Pediatric Cases in a Tertiary Care Hospital in Haryana. *Indian J Otolaryngol Head Neck Surg [Internet]*. 2021; Available from: <https://doi.org/10.1007/s12070-020-02301-1>
25. Auf I, Osborne JE, Sparkes C, Khalil H. Is the KTP laser effective in tonsillectomy? *Clin Otolaryngol Allied Sci*. 1997;22(2):145–6.
26. Sheikh NA, Nadeem K. Efficacy of prophylactic antibiotics in reducing post-tonsillectomy haemorrhage. *J Ayub Med Coll Abbottabad*. 2020;32(1):94–8.

Figures




Figure 1

The indication for tonsillectomy of the 62 patients.