A Systematic Review of the Literature Regarding Physical, Cognitive and Emotional Outcomes in Older Adults Practitioners of Physical Exercise

Miguel Ángel Araque-Martínez

Universidad de Almería <https://orcid.org/0000-0003-4375-186X>

Eva María Artés-Rodríguez

Universidad de Almería

Antonio Jesús Casimiro-Andújar (✉ casimiro@ual.es)

University of Almería <https://orcid.org/0000-0001-8864-0309>

Pedro Jesús Ruiz-Montero

Universidad de Granada

Research article

Keywords: Physical activity, ageing, review process, cognition, emotion

Posted Date: August 28th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-58739/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: This systematic review provides an overview about those papers with a specific physical exercise intervention aimed to improve physical, cognitive and/or emotional outcomes in healthy older people aged 60 or over. Studies with no intervention whose sample were active healthy older adults were also included.

Methods: An exhaustive literature search was done through two databases, including studies from January 2000 to July 2020.

Results: Of the 2006 identified records, 66 met the inclusion criteria and were selected. Results from the review showed a wide variety of older adults and lengths of the interventions, from five weeks (the shortest one) to five years. Aerobic exercise, either isolated or combined with strength performance, was also the most common type of exercise recognized in this systematic review.

Conclusions: Findings also suggest that the number of studies with all, physical, cognitive and emotional outcomes have been increasing during recent years in healthy older adults.

Background

The number of people aged 65 or over in 2050 will be doubled in comparison to the current one, from 703 million people at the moment to a future figure of 1.5 billion [1]. In fact, life expectancy is also increasing, from 72.6 years in 2019 to a probable figure of 77.1 years in 2050 [2].

According to the American College of Sport Medicine (ACSM) [3], ageing is associated with some physiologic changes that modify physical and mental health among other factors. In addition, becoming older has some effects on individuals, such as an acceleration in the decline of aerobic capacity and strength [4], memory deterioration [5], decline of motivation and increase of negative emotions [6] or an increment on the prevalence of cognitive impairment and dementia [7, 8].

The role of physical activity (PA) is gaining attention in this group of people because of the evidence that shows the positive effects of it on healthy older adults, such as the prevention of some diseases including type 2 diabetes, strokes, hypertension and some types of cancer [9]. Furthermore, there are also beneficial effects of PA on unhealthy elders with dementia [10], depression [11], mild cognitive impairment (MCI) [12, 13], and Alzheimer's [14] or Parkinson's disease [15].

Physiologically, regular physical exercise increases maximal oxygen consumption ($VO_2\text{max}$) [16], physical function [17], strength and cardiorespiratory fitness [18]. Because of that, physical performance is improved in healthy elderly practitioners of PA [19].

In order to determine what type of activity and how often it should be done, the ACSM and the American Heart Association (AHA) proposed a guidelines recommendation [20] describing frequency and type of

activities to perform, summarizing them in a minimum of 150 min per week for achieving substantial health benefits and including strength, flexibility and cardiorespiratory capacity exercises.

In relation to some of the components of quality of life (i.e., sensorial functioning, social participation, perceptions of death and dying, or intimacy), there seems to be an improvement in them [21] after getting regular PA. Personal growth and well-being also get better with regular PA [22]. A study with older adults ranging from 65 to 96 years old concluded that a weekly activity, such as PA outside the home, was associated with an improvement in both emotional state and cognitive function [23].

It is necessary to highlight brain health as an important factor related to regular PA [24] due to its effects on cognition, such as an enhancement of executive function and information processing speed [25] or beneficial effect on peripheral blood concentrations of brain-derived neurotrophic factor (BDNF) [13, 42]. Regular PA increases the function of neurotransmitters, such as serotonin, noradrenaline or dopamine; therefore, the motivation, happiness and satisfaction are going to be increased in practitioners [26, 27].

Consequently, there is research focused on the relation between PA and cognitive function, which confirms that PA in healthy older adults enhances existing cognitive function and prevents or delays progression of cognitive diseases, such as Alzheimer's dementia [28], in both older adults with and without cognitive impairment [29]. In fact, combined physical and cognitive interventions have been proven as an effective tool for improving cognition in healthy older adults [30].

Overall, considering how important and useful it is to be updated about different physical exercise interventions in an older adult population, the present systematic review pretends to highlight them, taking the recent literature into consideration.

We have performed a systematic review to summarize those supervised physical exercise interventions aimed to improve physical, emotional and/or cognitive performance in healthy older adults aged 60 or over. Studies without interventions in non-sedentary older adults with an active lifestyle have also been included [31–44].

Broadly, this review is organized following different aspects, including original source of information, sample characteristics, study design, objective, duration and characteristics of the intervention in case it exists, and the main outcomes under investigation.

Methods

This systematic review has followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines [45]. A PRISMA flow diagram (Fig. 1) summarizes the information on the phases of the systematic review process.

Search Strategies

A systematic search for relevant studies was conducted on two of the most respected databases in the field of social sciences and education, ISI Web of Knowledge and Scopus-Elsevier. From these, we contemplated all peer-reviewed publications from the current century, between January of 2000 and July of 2020. We consider two decades as an appropriate period of time, which includes literature from recent years and allows us to observe the gradual process of change and development in this area. In order to be as accurate as possible, we decided which terms should be taken into account. Consequently, all the publications had to contain at least one of the following terms: 'older adults' (older adults, elderly, seniors, elders); 'cognition' (cognition, cognitive function, cognitive flexible, memory); 'physical activity' (physical activity, physical exercise, exercise); 'mood' (mood, state of mood); and 'emotions' (depression, sadness, anxiety, happiness, self-esteem, optimism, anger, hostility).

The search included those articles where the terms were found in titles and abstracts.

Apart from that, all the references from the retrieved articles were reviewed looking for possible additional publications related to physical exercise interventions in healthy older adults, as well as those where samples were active older adults.

Selection Criteria

Studies were included according to the following criteria: (1) complete original study; (2) written in English; (3) involving healthy or active people aged 60 years and over; and (4) related to PA and the physical exercise field.

According to these criteria, healthy people were considered as those who did not suffer any kind of mental (e.g., Alzheimer's or Parkinson's disease), emotional (e.g., depression) or physical (e.g., arthritis) diagnosed impairment, while active older adults were those who exercised regularly (e.g., marathon runners) [33, 40].

Studies were excluded if they were (1) not written in English; (2) a review, theoretical or protocol article; (3) book or chapter in a book; (4) related to other areas, such as medicine [46] or genetics [47]; or (5) involving unhealthy older adults 60 years old and over.

Two independent expert reviewers screened the titles and abstracts; secondly, they evaluated full-text articles independently. In case of disagreement, a third reviewer was required to solve it. Those reviewers, who are members of different universities with much experience working with older adults and research in physical exercise in the area of health and sport sciences, followed the considerations previously described.

Quality Assessment

A total of 25 out of the 66 selected papers in this systematic review were indexed on PEDro, the physiotherapy evidence database (<https://www.pedro.org.au/>). Their scores are shown in Table 1. From the indexed papers, there were 4 articles with a score lower than four points, whereas the number of

studies with a score equal to or higher than four points was 21. In addition, the criteria from PEDro for 2/10, 8/10, 9/10 and 10/10 were not included in Table 1, because there were not in any study.

There were also 41 selected studies that were not rated by using the criteria from PEDro. The quality of these studies was independently assessed by two reviewers. In case of disagreement, a third reviewer's opinion was taken into account.

Results

Study selection

A total of 2006 studies were identified using the databases as described in the Search Strategies section. After the removal of 45 duplicated papers, 1749 papers were also excluded based on titles and abstracts. The remaining 212 articles were screened, and from these, 146 were excluded after a full-text evaluation due to the exclusion criteria previously described, leaving 66 studies that were eligible for this review (Fig. 1). These 66 papers were classified according to different aspects, including original source of information, sample characteristics, study design, objective, duration and characteristics of the intervention in case it exists, and the main outcomes under investigation.

Frequency of published studies and geographical distribution

As we can see in Fig. 2, the 66 selected articles have been mostly ($n = 57$; 86%) published during the last 10 years (2011–2020), and from these, 38 (67%) were published 2016–2020, so it seems to be an identified pattern in the number of publications with the course of time. In recent years, 2019 was the most prolific year with 12 papers of the total (18%) contained in this systematic review.

On the other hand, there were 6 years (from 2000 to 2004 and 2008) with no selected publications.

In relation to the geographical distribution of the sample, Brazil (with seven papers (11%)) and the United States (with five papers (8%)) are the countries with the most published articles in this area that have been accepted in our review. On this point, it is necessary to clarify that the geographical distribution of the samples in some studies have not been specified. Specifically, it was not indicated in 16 articles (24%). The remaining papers, including those from Brazil and the United States, have been developed with participants from different countries in Europe (22), where Spain, Greece and Norway are the countries with the most publications, as well as Japan or Thailand in Asia (11), while Brazil and the United States, as previously described, had most of the publications in the Americas (14). From all the included papers, there was an unusual sample of participants from two articles [65, 75] composed of people from different countries and/or continents.

Participant characteristics

The sample size varied from eight participants [62] to 1193 [34] participants. The ages of the participants ranged from 60, which was the minimum accepted in this paper, to 99 years old [79].

In terms of gender, most of the papers (73%) had mixed samples, although there was a minority with men exclusively (6%) and a bigger percentage of studies (21%) comprised mainly of women. Three papers included in this systematic review [36, 66, 95] did not distinguish between men or women in their samples.

According to a PA profile, we found more articles (32%) with a sample composed of sedentary older adults in comparison with those with an active sample (24%). Those active practitioners took part in different studies doing regular activities, such as yoga [32], athletics [33] and dancing [43].

However, there were also studies (41%) that did not specify the PA profile of their samples, as well as those whose samples were comprised of both active and sedentary people [42, 44].

In those papers in which there was no physical exercise intervention, we only included the physically active sample.

Type of intervention

One of the peculiarities of this systematic review is the inclusion of those papers in which there was no physical exercise intervention, but the articles were related to physical exercise in older adult practitioners. In this way, there were 14 studies (21%) out of the 66 included where there was not a physical exercise intervention. The rest of studies included physical exercise interventions with differences in the length of those, from five weeks (the shortest one) [61] to five years [65]. In relation to the weekly frequency of physical exercise sessions, most of the interventions had two (29%) or three (52%) sessions per week in comparison to those with four [71] or five [69] weekly sessions. Moreover, we found papers where the number of weekly sessions varied depending on the intervention group [63, 77, 79].

According to the type of physical exercise during the intervention, aerobic training was the most common one whether isolated (44%) or combined with strength training (17%). The combination of different types of activities, such as cognitive and physical training [58]; outdoor leisure activities and callisthenic exercises [64]; or aerobic, strength and balance exercises [88] was also interesting. These types of exercise combinations have been defined in Table 2 of this study as concurrent interventions.

Research Methodology

In terms of methodology, almost all studies (97%) followed a quantitative methodology, whereas only two [72, 75] used a mixed model. Therefore, none of the articles included in this systematic review followed a qualitative methodology.

However, there was less of a difference between the design of studies. Half of the number of articles ($n = 33$; 50%) had a quasi-experimental study design in contrast to the other half with a randomized controlled trial study design.

Main Outcomes Investigated

This systematic review is focused on summarizing different types of studies where the main outcomes investigated are related to physical, cognitive and/or emotional performance. According to this, physical performance included some outcomes such as muscle strength [86], heart rate variability [36], stability [57], VO₂peak [81], functional balance [95] or gait speed [87]. The studies on executive function [37], attention [41], memory [48], processing speed [93] or concentration [85] made allusion to cognitive performance, while studies of depression [49], anxiety [71], stress [52], health-related quality of life [80], mood [84] or happiness [96] were related to emotional performance. Those outcomes investigated from studies included in this systematic review that were not related to physical, cognitive and/or emotional performance were not included in Table 2.

Discussion

This systematic review aimed to summarize those studies of healthy older adults aged 60 and over with or without a physical exercise intervention programme where the main outcomes investigated have been related to physical, cognitive and/or emotional performance. We found 66 suitable studies fitting the pre-established inclusion criteria.

These outcomes were, mainly, related to physical performance [53, 55], although emotional and cognitive outcomes were also studied, isolated [54, 78] or combined [66] with the physical ones. This combination of physical, cognitive and emotional outcomes was observed in 14 (21%) out of the 66 chosen studies, where six of those 14 papers were published in 2019 [33, 50, 59, 71, 83, 87]. According to this, it is conceivable that there is a current tendency of studying these combinations of physical, cognitive and emotional outcomes together with older adults. It is also interesting how technology is used for improving those outcomes through exergames [51, 74].

Hogan et al. (2013) [100] suggested important benefits in older adults for both affective and cognitive performance after a single bout of moderate exercise, whereas Tarazona-Santabalbina et al. (2016) [101] published the physical, cognitive, emotional and social effects of a multicomponent exercise intervention in frail older adults. In reference to reviews and meta-analysis publications, there is considerable published research focused on the effects of physical exercise on cognition in healthy (Carvalho et al., 2014 [28]; Gomes-Osman et al., 2018 [29]; Zhu et al., 2016 [30]; Kelly et al., 2014 [102]) and unhealthy (Karssemeijer et al., 2017 [13]; Öhman et al., 2014 [103]) older adults.

Investigating cognitive or emotional outcomes, apart from physical outcomes, during physical exercise interventions may contribute to establish more connections between body, brain and emotions to the benefit of science and particularly to older adults. According to the 52 accepted papers in this systematic review in which there was a physical exercise intervention, in 24 of them (46%), the number of physical exercise min per week was 150 or more. If we compare these results and current guidelines of physical exercise for older adults [3, 20, 24], we could suggest that a considerable number of selected papers did

not contribute to the minimum of weekly min, which these guidelines established to achieve benefits in physical health: 150 min of moderate intensity exercise or 75 min of vigorous exercise. According to this, an intervention that does not follow these recommendations could not be successful for practitioners. Thus, this result may insinuate that, in the future, physical exercise interventions should follow current recommendations for weekly min in order to obtain greater benefits to health. In the overall analysis, the results found in this systematic review may prove that physical exercise interventions in healthy older adults have been increasing during recent years. An update on these interventions could provide useful information, such as the length, type of exercise or the effects produced by itself, to those professionals who are in daily contact with older adults.

We considered it appropriate to focus this systematic review in healthy older adults in order to establish differences with the unhealthy older adults. However, there are also published reviews [104, 105], which include both healthy and unhealthy older adults in the same paper. In that case, data or results from those papers should be carefully observed, taking into account possible differences and conclusions to come to.

The sample of studies included in this systematic review was located in two databases, ISI Web of Knowledge and Scopus-Elsevier. We acknowledge that by limiting the search to a couple of databases, we could not capture all possible information. It would be interesting to expand the search to other databases with the purpose of giving access to more information to interested populations. Moreover, of all the included papers, none followed a qualitative methodology, which could also contribute to providing more information in this area. In relation to the quality of the included studies, a remarkable number of them were not indexed on PEDro's criteria. This result suggests some methodologic limitations.

Conclusions

The results of the present paper suggest there were a remarkable number of studies about this topic, which has been increasing during recent years. This increment shows how current and important it is to investigate the way physical exercise is being practiced in healthy older adults and the connection it has with cognition and emotions. These studies show a large methodological heterogeneity in intervention characteristics (e.g., number of participants, length of the intervention or type of exercise). The current systematic review illustrates how physical, cognitive and emotional outcomes could be, or not be, simultaneously investigated following a specific physical exercise intervention or under no intervention but being focused on active older adults who regularly exercise. However, the results of these investigations are not included in this paper. Future research should also focus on results in order to provide useful information about the effects of these kinds of studies.

Abbreviations

ACSM: American College of Sport Medicine; PA: Physical activity; MCI: Mild Cognitive Impairment; VO₂max: Maximal oxygen consumption; AHA: American Heart Association; BDNF: Brain-derived

neurotrophic factor; PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-Analyses; MPO: Myeloperoxidase; TEE: Total energy expenditure; AEE: Activity-related energy expenditure; IL-6: Serum interleukin-6; DMT: Dance movement training.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

The datasets analysed during the current study are available in the ISI-Web of Knowledge (<https://www.webofknowledge.com>) and Scopus-Elsevier (<https://www.scopus.com/home.uri>) repositories.

Competing interests

The authors declare that they have no competing interests

Funding

Not applicable

Author's contribution

MAA-M, PJR-M, EVAR and ACA were involved in the conception and design of the systematic review. MAA-M generated and conducted the search. EVAR and PJR-M were involved in the data collection, extraction and synthesis. ACA revised the manuscript critically for important intellectual content. All authors contributed to the writing of the paper, read and approved the final manuscript.

Acknowledgements

Not applicable

References

1. World Population Ageing 2019. Highlights. Geneva: World Health Organization; 2019.
2. World Population Prospects 2019. Highlights. Geneva: World Health Organization; 2019.

3. Chodzko-Zajko WJ, Proctor DN, Fiatarone Singh MA, Minson CT, Nigg CR, Salem GJ, et al. Exercise and physical activity for older adults [Internet]. Vol. 41, *Medicine and Science in Sports and Exercise*. Med Sci Sports Exerc; 2009 [cited 2020 Jul 28]. p. 1510–30. Available from: <https://pubmed.ncbi.nlm.nih.gov/19516148/>
4. Fleg JL, Morrell CH, Bos AG, Brant LJ, Talbot LA, Wright JG, et al. Accelerated longitudinal decline of aerobic capacity in healthy older adults. *Circulation*. 2005;112(5):674–82.
5. Peter R. Ageing and the brain. *Postgrad Med J*. 2006;82:84-88.
6. Carstensen LL, Mikels JA. At the intersection of emotion and cognition - Aging and the positivity effect. *Curr Dir Psychol Sci*. 2005;14(3):117–21.
7. Bishop NA, Lu T, Yankner BA. Neural mechanisms of ageing and cognitive decline. *Nature*. 2010; 464(7288):529–35.
8. Yuan J, Zhang Z, Wen H, Hong X, Hong Z, Qu Q, et al. Incidence of dementia and subtypes: A cohort study in four regions in China. *Alzheimers Dement*. 2016;12(3):262–71.
9. Vogel T, Brechat P-H, Lepretre P-M, Kaltenbach G, Berthel M, Lonsdorfer J. Health benefits of physical activity in older patients: a review. *Int J Clin Pract*. 2009;63(2):303–20.
10. Blankevoort CG, Van Heuvelen MJG, Boersma F, Luning H, De Jong J, Scherder EJA. Review of effects of physical activity on strength, balance, mobility and ADL performance in elderly subjects with dementia. *Dement Geriatr Cogn Disord*. 2010;30(5):392–402.
11. Lavretsky H, Alstein LL, Olmstead RE, Ercoli LM, Riparetti-Brown M, Cyr NS, et al. Complementary use of Tai Chi Chih augments escitalopram treatment of geriatric depression: A randomized controlled trial. *Am J Geriatr Psychiatry*. 2011;19(10):839–50.
12. Naidu AS, Vasudev A, Burhan AM, Ionson E, Montero-Odasso M. Does Dual-Task Gait Differ in those with Late-Life Depression versus Mild Cognitive Impairment? *Am J Geriatr Psychiatry*. 2019;27(1):62–72.
13. Karssemeijer EGAE, Aaronson JAJ, Bossers WJW, Smits TT, Olde Rikkert MGMM, Kessels RPCR. Positive effects of combined cognitive and physical exercise training on cognitive function in older adults with mild cognitive impairment or dementia: A meta-analysis. *Ageing Res Rev*. 2017;40:75–83.
14. Salazar MCR, Báez ALM, Gallego EAQ, Granada LMR. Hatha yoga effects on Alzheimer patients (AP). *Acta Colomb Psicol*. 2017;20(1):139–53.
15. Rios Romenets S, Anang J, Fereshtehnejad S-M, Pelletier A, Postuma R. Tango for treatment of motor and non-motor manifestations in Parkinson's disease: A randomized control study. *Complement Ther Med*. 2015;23(2):175–84.
16. Malbut KE, Dinan S, Young A. Aerobic training in the 'oldest old': the effect of 24 weeks of training. *Age Ageing*. 2002;31(4):255–60.
17. Geirsdottir OG, Arnarson A, Ramel A, Briem K, Jonson P V, Thorsdottir I. Muscular strength and physical function in elderly adults 6-18 months after a 12-week resistance exercise program. *Scand J Public Health*. 2015;43(1):76–82.

18. Ruiz-Montero PJ, Ramiro, MT, Ramiro-Sánchez T, García-Marmol E. Pilates-Aerobic exercise program's effects in physical fitness level and quality of life related to physical and mental health in elderly women. *Psychol Soc Educ.* 2020;12(2):91-105.
19. Vaughan S, Wallis M, Polit D, Steele M, Shum D, Morris N. The effects of multimodal exercise on cognitive and physical functioning and brain-derived neurotrophic factor in older women: a randomised controlled trial. *Age Ageing.* 2014;43(5):623–9.
20. Nelson ME, Rejeski WJ, Blair SN, Duncan PW, Judge JO, King AC, et al. Physical Activity and Public Health in Older Adults: Recommendation from the American College of Sports Medicine and the American Heart Association. *Med Sci Sports Exerc.* 2007;39(8):1435–45.
21. Figueira HA, Figueira AA, Cader SA, Guimaraes AC, De Oliveira RJ, Figueira JA, et al. Effects of a physical activity governmental health programme on the quality of life of elderly people. *Scand J Public Health.* 2012;40(5):418–22.
22. Hogan CL, Catalino LI, Mata J, Fredrickson BL. Beyond emotional benefits: Physical activity and sedentary behaviour affect psychosocial resources through emotions. *Psychol Health.* 2015;30(3):354–69.
23. Petersen J, Austin D, Mattek N, Kaye J. Time Out-of-Home and Cognitive, Physical, and Emotional Wellbeing of Older Adults: A Longitudinal Mixed Effects Model. *PLoS One.* 2015;10:10.
24. Powell KE, King AC, Buchner DM, Campbell WW, DiPietro L, Erickson KI, et al. The Scientific Foundation for the Physical Activity Guidelines for Americans, 2nd Edition. *J Phys Act Health.* 2019;16(1):1–11.
25. Coetsee C, Terblanche E. The effect of three different exercise training modalities on cognitive and physical function in a healthy older population. *Eur Rev Aging Phys Act.* 2017;14.
26. Davranche K, Audiffren M. Facilitating effects of exercise on information processing. *J Sports Sci.* 2004;22(5):419–28.
27. Secher NH, Seifert T, Van Lieshout JJ. Cerebral blood flow and metabolism during exercise: Implications for fatigue. *J Appl Physiol.* 2008;104(1):306–14.
28. Carvalho A, Rea IM, Parimon T, Cusack BJ. Physical activity and cognitive function in individuals over 60 years of age: A systematic review. *Clin Interv Aging.* 2014;9:661–82.
29. Gomes-Osman J, Cabral DF, Morris TP, McInerney K, Cahalin LP, Rundek T, et al. Exercise for cognitive brain health in aging: A systematic review for an evaluation of dose. *Neurol Clin Pract.* 2018;8(3):257–65.
30. Zhu X, Yin S, Lang M, He R, Li J. The more the better? A meta-analysis on effects of combined cognitive and physical intervention on cognition in healthy older adults. *Ageing Res Rev.* 2016;31:67–79.
31. Haslacher H, Michlmayr M, Batmyagmar D, Perkmann T, Ponocny-Seliger E, Scheichenberger V, et al. Physical Exercise Counteracts Genetic Susceptibility to Depression. *Neuropsychobiology.* 2015;71(3):168–75.

32. Afonso RF, Balardin JB, Lazar S, Sato JR, Igarashi N, Santaella DF, et al. Greater cortical thickness in elderly female yoga practitioners-A cross-sectional study. *Front Aging Neurosci.* 2017;9.
33. Batmyagmar D, Kundi M, Ponocny-Seliger E, Lukas I, Lehrner J, Haslacher H, et al. High intensity endurance training is associated with better quality of life, but not with improved cognitive functions in elderly marathon runners. *Sci Rep.* 2019;9:1.
34. Brady S, D'Ambrosio LA, Felts A, Rula EY, Kell KP, Coughlin JF. Reducing Isolation and Loneliness Through Membership in a Fitness Program for Older Adults: Implications for Health. *J Appl Gerontol.* 2020;39(3):301–10.
35. de Oliveira LSSCB, Souza EC, Rodrigues RAS, Fett CA, Piva AB. The effects of physical activity on anxiety, depression, and quality of life in elderly people living in the community. *Trends Psychiatry Psychother.* 2019;41(1):36–42.
36. Dupuy O, Bosquet L, Fraser SA, Labelle V, Bherer L. Higher cardiovascular fitness level is associated to better cognitive dual-task performance in Master Athletes: Mediation by cardiac autonomic control. *Brain Cogn.* 2018;125:127–34.
37. Haslacher H, Perkmann T, Lukas I, Barth A, Ponocny-Seliger E, Michlmayr M, et al. Myeloperoxidase levels predict executive function. *Int J Sports Med.* 2012;33(12):1034–8.
38. Haslacher H, Michlmayr M, Batmyagmar D, Perkmann T, Ponocny-Seliger E, Scheichenberger V, et al. rs6295 [C]-allele protects against depressive mood in elderly endurance athletes. *J Sport Exerc Psychol.* 2015;37(6):637–45.
39. Haslacher H, Nistler S, Batmyagmar D, Ponocny-Seliger E, Perkmann T, Scherzer TM, et al. Low vitamin D levels do not predict hyperglycemia in elderly endurance athletes (but in controls). *PLoS One.* 2016;11:6.
40. Haslacher H, Ratzinger F, Perkmann T, Batmyagmar D, Nistler S, Scherzer TM, et al. A combination of routine blood analytes predicts fitness decrement in elderly endurance athletes. *PLoS One.* 2017;12:5.
41. Liu S, Li L, Liu Z, Guo X. Long-term Tai Chi experience promotes emotional stability and slows gray matter atrophy for elders. *Front Psychol.* 2019;10.
42. Rosanti S, da Silva GE, dos Santos FH. Longitudinal effects of physical activity on self-efficacy and cognitive processing of active and sedentary elderly. *Dement e Neuropsychol.* 2014;8(2):187–93.
43. Serra MM, Alonso AC, Peterson M, Mochizuki L, Greve JMD, Garcez-Leme LE. Balance and muscle strength in elderly women who dance samba. *PLoS One.* 2016;11:12.
44. Stone RC, Rakhamilova Z, Gage WH, Baker J. Curling for confidence: Psychophysical benefits of curling for older adults. *J Aging Phys Act.* 2018;26(2):267–75.
45. Urrútia G, Bonfill X. PRISMA declaration: A proposal to improve the publication of systematic reviews and meta-analyses. *Med Clin (Barc).* 2010;135(11):507–11.
46. Ates Bulut E, Soysal P, Yavuz I, Kocyigit SE, Isik AT. Effect of Vitamin D on Cognitive Functions in Older Adults: 24-Week Follow-Up Study. *Am J Alzheimers Dis Other Dement.* 2019;34(2):112–7.

47. Szentes N, Tékus V, Mohos V, Borbély É, Helyes Z. Exploratory and locomotor activity, learning and memory functions in somatostatin receptor subtype 4 gene-deficient mice in relation to aging and sex. *GeroScience*. 2019;41(5):631–41.
48. Antunes HKM, Santos-Galduroz RF, De Aquino Lemos V, Bueno OFA, Rzezak P, de Santana MG, et al. The influence of physical exercise and leisure activity on neuropsychological functioning in older adults. *Age (Omaha)*. 2015;37:4.
49. Azizan A, Justine M. Effects of a Behavioral and Exercise Program on Depression and Quality of Life in Community-Dwelling Older Adults. *J Gerontol Nurs*. 2016;42(2):45–54.
50. Bouaziz W, Schmitt E, Vogel T, Lefebvre F, Leprêtre P-M, Kaltenbach G, et al. Effects of a short-term Interval Aerobic Training Programme with active Recovery bouts (IATP-R) on cognitive and mental health, functional performance and quality of life: A randomised controlled trial in sedentary seniors. *Int J Clin Pract*. 2019;73:1.
51. Carrasco M, Ortiz-Maqués N, Martínez-Rodríguez S. Playing with Nintendo Wii Sports: Impact on Physical Activity, Perceived Health and Cognitive Functioning of a Group of Community-Dwelling Older Adults. *Act Adapt Aging*. 2020;44(2):119–31.
52. Čekanauskaitė A, Skurvydas A, Žlibinaitė L, Mickevičienė D, Kilikevičienė S, Solianik R. A 10-week yoga practice has no effect on cognition, but improves balance and motor learning by attenuating brain-derived neurotrophic factor levels in older adults. *Exp Gerontol*. 2020;138.
53. Chutimakul L, Sukonthasab S, Kritpet T, Vannalee C. Effect of modified Khon dance performance on functional fitness in older Thai persons. *J Heal Res*. 2018;32(6):432–9.
54. Coubard OA, Duretz S, Lefebvre V, Lapalus P, Ferrufino L. Practice of contemporary dance improves cognitive flexibility in aging. *Front Aging Neurosci*. 2011;3:1–12.
55. Douka S, Zilidou VI, Lilou O, Manou V. Traditional dance improves the physical fitness and well-being of the elderly. *Front Aging Neurosci*. 2019;11.
56. Ericson H, Skoog T, Johansson M, Wåhlin-Larsson B. Resistance training is linked to heightened positive motivational state and lower negative affect among healthy women aged 65–70. *J Women Aging*. 2018;30(5):366–81.
57. Filar-Mierzwa K, Długosz M, Marchewka A, Dąbrowski Z, Poznańska A. The effect of dance therapy on the balance of women over 60 years of age: The influence of dance therapy for the elderly. *J Women Aging*. 2017;29(4):348–55.
58. González-Palau F, Franco M, Bamidis P, Losada R, Parra E, Papageorgiou SG, et al. The effects of a computer-based cognitive and physical training program in a healthy and mildly cognitive impaired aging sample. *Aging Ment Heal*. 2014;18(7):838–46.
59. Gothe NP, Ehlers DK, Salerno EA, Fanning J, Kramer AF, McAuley E. Physical Activity, Sleep and Quality of Life in Older Adults: Influence of Physical, Mental and Social Well-being. *Behav Sleep Med*. 2019; doi: 10.1080/15402002.2019.1690493.
60. Haraldstad K, Rohde G, Stea TH, Lohne-Seiler H, Hetlelid K, Paulsen G, et al. Changes in health-related quality of life in elderly men after 12 weeks of strength training. *Eur Rev Aging Phys Act*. 2017;14:1.

61. Lee P-L, Yang Y-C, Huang C-K, Hsiao C-H, Liu T-Y, Wang C-Y. Effect of exercise on depressive symptoms and body balance in the elderly. *Educ Gerontol*. 2017;43(1):33–44.
62. Lindahl E, Tilton K, Eickholt N, Ferguson-Stegall L. Yoga reduces perceived stress and exhaustion levels in healthy elderly individuals. *Complement Ther Clin Pract*. 2016;24:50–6.
63. Matsouka O, Kabitsis C, Harahousou Y, Trigonis I. Mood alterations following an indoor and outdoor exercise program in healthy elderly women. *Percept Mot Skills*. 2005;100(3):707–15.
64. Matsouka O, Bebetos E, Trigonis I, Simakis S. The effects of an outdoor exercise program on mood states among the elderly. *World Leis J*. 2010;52(1):34–40.
65. Pandya SP. Yoga Education Program for Improving Memory in Older Adults: A Multicity 5-Year Follow-Up Study. *J Appl Gerontol*. 2020;39(6):576–87.
66. Pereira JR, Gobbi S, Teixeira CVL, Nascimento CMC, Corazza DI, Vital TM, et al. Effects of Square-Stepping Exercise on balance and depressive symptoms in older adults. *Motriz Rev Educ Fis*. 2014;20(4):454–60.
67. Solberg PA, Hopkins WG, Ommundsen Y, Halvari H. Effects of three training types on vitality among older adults: A self-determination theory perspective. *Psychol Sport Exerc*. 2012;13(4):407–17.
68. Solberg PA, Halvari H, Ommundsen Y. Linking exercise and causality orientations to change in well-being among older adults: Does change in motivational variables play a role? *J Appl Soc Psychol*. 2013;43(6):1259–72.
69. Starkweather AR. The effects of exercise on perceived stress and IL-6 levels among older adults. *Biol Res Nurs*. 2007;8(3):186–94.
70. Tada A. Psychological effects of exercise on community-dwelling older adults. *Clin Interv Aging*. 2018;13:271–6.
71. Vaccaro MG, Izzo G, Ilacqua A, Migliaccio S, Baldari C, Guidetti L, et al. Characterization of the Effects of a Six-Month Dancing as Approach for Successful Aging. *Int J Endocrinol*. 2019; doi: [1155/2019/2048391](https://doi.org/10.1155/2019/2048391).
72. Zafar M, Bozzorg A, Hackney ME. Adapted Tango improves aspects of participation in older adults versus individuals with Parkinson's disease. *Disabil Rehabil*. 2017;39(22):2294–301.
73. Zanuso S, Sieverdes JC, Smith N, Carraro A, Bergamin M. The effect of a strength training program on affect, mood, anxiety, and strength performance in older individuals. *Int J Sport Psychol*. 2012;43(1):53–66.
74. Zheng H, Li J, Salmon CT, Theng Y-L. The effects of exergames on emotional well-being of older adults. *Comput Human Behav*. 2020;110.
75. Fox KR, Stathi A, McKenna J, Davis MG. Physical activity and mental well-being in older people participating in the Better Ageing Project. *Eur J Appl Physiol*. 2007;100(5):591–602.
76. Alghadir AH, Gabr SA, Al-Eisa ES. Effects of Moderate Aerobic Exercise on Cognitive Abilities and Redox State Biomarkers in Older Adults. *Oxid Med Cell Longev*. 2016; doi: [1155/2016/2545168](https://doi.org/10.1155/2016/2545168).

77. Hunter GR, Bickel CS, Fisher G, Neumeier WH, McCarthy JP. Combined aerobic and strength training and energy expenditure in older women. *Med Sci Sports Exerc.* 2013;45(7):1386–93.
78. Kim Y-S, O’Sullivan DM, Shin S-K. Can 24 weeks strength training reduce feelings of depression and increase neurotransmitter in elderly females? *Exp Gerontol.* 2019;115:62–8.
79. Anderson-Hanley C, Snyder AL, Nimon JP, Arciero PJ. Social facilitation in virtual reality-enhanced exercise: Competitiveness moderates exercise effort of older adults. *Clin Interv Aging.* 2011;6(1):275–80.
80. Flegal KE, Kishiyama S, Zajdel D, Haas M, Oken BS. Adherence to yoga and exercise interventions in a 6-month clinical trial. *BMC Complement Altern Med.* 2007;7.
81. Flodin P, Jonasson LS, Riklund K, Nyberg L, Boraxbekk CJ. Does aerobic exercise influence intrinsic brain activity? An aerobic exercise intervention among healthy old adults. *Front Aging Neurosci.* 2017;9.
82. Kang D-W, Bressel E, Kim D-Y. Effects of aquatic exercise on insulin-like growth factor-1, brain-derived neurotrophic factor, vascular endothelial growth factor, and cognitive function in elderly women. *Exp Gerontol.* 2020;132.
83. Vranceanu T, Esmail A, Berryman N, Predovan D, Vu TTM, Villalpando JM, et al. Dance your stress away: comparing the effect of dance/movement training to aerobic exercise training on the cortisol awakening response in healthy older adults. *Stress.* 2019;22(6):687–95.
84. Brown AK, Liu-Ambrose T, Tate R, Lord SR. The effect of group-based exercise on cognitive performance and mood in seniors residing in intermediate care and self-care retirement facilities: a randomised controlled trial. *Br J Sports Med.* 2009;43(8):608–14.
85. Cassilhas RC, Viana VAR, Grassmann V, Santos RT, Santos RF, Tufik S, et al. The impact of resistance exercise on the cognitive function of the elderly. *Med Sci Sports Exerc.* 2007;39(8):1401–7.
86. Cassilhas RC, Antunes HKM, Tufik S, de Mello MT. Mood, anxiety, and serum IGF-1 in elderly men given 24 weeks of high resistance exercise. *Percept Mot Skills.* 2010;110(1):265–76.
87. Imaoka M, Nakao H, Nakamura M, Tazaki F, Maebuchi M, Ibuki M, et al. Effect of multicomponent exercise and nutrition support on the cognitive function of older adults: A randomized controlled trial. *Clin Interv Aging.* 2019;14:2145–53.
88. Laredo-Aguilera JA, Carmona-Torres JM, García-Pinillos F, Latorre-Román PÁ. Effects of a 10-week functional training programme on pain, mood state, depression, and sleep in healthy older adults. *Psychogeriatrics.* 2018;18(4):292–8.
89. Martins RA, Coelho E Silva MJ, Pindus DM, Cumming SP, Teixeira AM, Veríssimo MT. Effects of strength and aerobic-based training on functional fitness, mood and the relationship between fatness and mood in older adults. *J Sports Med Phys Fitness.* 2011;51(3):489–96.
90. Noopud P, Suputtitada A, Khongprasert S, Kanungsukkasem V. Effects of Thai traditional dance on balance performance in daily life among older women. *Aging Clin Exp Res.* 2019;31(7):961–7.
91. Tsai C-L, Wang C-H, Pan C-Y, Chen F-C. The effects of long-term resistance exercise on the relationship between neurocognitive performance and GH, IGF-1, and homocysteine levels in the

- elderly. *Front Behav Neurosci*. 2015;9.
92. Cruz-Ferreira A, Marmeleira J, Formigo A, Gomes D, Fernandes J. Creative Dance Improves Physical Fitness and Life Satisfaction in Older Women. *Res Aging*. 2015;37(8):837–55.
93. Eckardt N, Braun C, Kibele A. Instability Resistance Training improves Working Memory, Processing Speed and Response Inhibition in Healthy Older Adults: A Double-Blinded Randomised Controlled Trial. *Sci Rep*. 2020;10:1.
94. Hars M, Herrmann FR, Gold G, Rizzoli R, Trombetti A. Effect of music-based multitask training on cognition and mood in older adults. *Age Ageing*. 2014;43(2):196–200.
95. Joung HJ, Lee Y. Effect of Creative Dance on Fitness, Functional Balance, and Mobility Control in the Elderly. *Gerontology*. 2019;65(5):537–46.
96. Khazaee-pool M, Sadeghi R, Majlessi F, Rahimi Froushani A. Effects of physical exercise programme on happiness among older people. *J Psychiatr Ment Health Nurs*. 2015;22(1):47–57.
97. Machacova K, Vankova H, Volicer L, Veleta P, Holmerova I. Dance as Prevention of Late Life Functional Decline Among Nursing Home Residents. *J Appl Gerontol*. 2017;36(12):1453–70.
98. Oken BS, Zajdel D, Kishiyama S, Flegal K, Dehen C, Haas M, et al. Randomized, controlled, six-month trial of yoga in healthy seniors: Effects on cognition and quality of life. *Altern Ther Health Med*. 2006;12(1):40–7.
99. Solberg PA, Kvamme NH, Raastad T, Ommundsen Y, Tomten SE, Halvari H, et al. Effects of different types of exercise on muscle mass, strength, function and well-being in elderly. *Eur J Sport Sci*. 2013;13(1):112–25.
100. Hogan CL, Mata J, Carstensen LL. Exercise Holds Immediate Benefits for Affect and Cognition in Younger and Older Adults. *Psychol Aging*. 2013;28(2):587–94.
101. Tarazona-Santabalbina FJ, Gómez-Cabrera MC, Pérez-Ros P, Martínez-Arnau FM, Cabo H, Tsaparas K, et al. A Multicomponent Exercise Intervention that Reverses Frailty and Improves Cognition, Emotion, and Social Networking in the Community-Dwelling Frail Elderly: A Randomized Clinical Trial. *J Am Med Dir Assoc*. 2016;17(5):426–33.
102. Kelly ME, Loughrey D, Lawlor BA, Robertson IH, Walsh C, Brennan S. The impact of exercise on the cognitive functioning of healthy older adults: A systematic review and meta-analysis. *Ageing Res Rev*. 2014;16(1):12–31.
103. Öhman H, Savikko N, Strandberg TE, Pitkälä KH. Effect of physical exercise on cognitive performance in older adults with mild cognitive impairment or dementia: A systematic review. *Dement Geriatr Cogn Disord*. 2014;38:347–65.
104. van Uffelen JGZ, Paw MJMCA, Hopman-Rock M, van Mechelen W. The Effects of Exercise on Cognition in Older Adults With and Without Cognitive Decline: A Systematic Review. *Clin J Sport Med*. 2008;18(6):486–500.
105. Coelho FGDM, Gobbi S, Andreatto CAA, Corazza DI, Pedroso RV, Santos-Galduróz RF. Physical exercise modulates peripheral levels of brain-derived neurotrophic factor (BDNF): A systematic review of experimental studies in the elderly. *Arch Gerontol Geriatr*. 2013;56(1):10–5.

Tables

Due to technical limitations, table PDF is only available as a download in the Supplemental Files section.

Figures

Fig. 1

Figure 1

Flowchart of the study selection process using the PRISMA framework

Fig. 2

Figure 2

Number of articles by year of publication

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Prismachecklist.pdf](#)
- [Tables.pdf](#)