Injectisome T3SS Subunits As Potential Chaperones In The Extracellular Export of *Pectobacterium Carotovorum* Subsp. *Carotovorum* Bacteriocins Carocin S1 And Carocin S3 Secreted Via Flagellar T3SS

Hang-Cheng Chen

China Medical University Hospital

Reymund C. Derilo

National Chung Hsing University

Han-Ling Chen

National Chung Hsing University

Tzu-Rung Li

National Chung Hsing University

Ruchi Briam James S. Lagitnay

National Chung Hsing University

Yung-Chieh Chan

National Chung Hsing University

Yutin Chuang

National Chung Hsing University

Duen-Yau Chuang (✉ chuang@dragon.nchu.edu.tw)

National Chung Hsing University

Research Article

Keywords: Carocin, pathogens, injectisome, hypothesize

Posted Date: June 15th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-590298/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Pectobacterium carotovorum subsp. *carotovorum* (*Pcc*) causes soft-rot disease in a wide variety of plants resulting in economic losses worldwide. It produces various types of bacteriocin to compete against related plant pathogens. Studies on how bacteriocins are extracellularly secreted are conducted to understand the mechanism of interbacterial competition. In this study, the secretion of the low-molecular-weight bacteriocins (LMWB) Carocin S1 and Carocin S3 produced by a multiple-bacteriocin producing strain of *Pcc*, 89-H-4, was investigated. Tn5 insertional mutagenesis was used to generate a mutant, TH22-6, incapable of LMWBs secretion. Sequence and homology analyses of the gene disrupted by transposon Tn5 insertion revealed that the gene *sctT*, an essential component of the injectisome type III secretion machinery (T3aSS), is required for the secretion of the bacteriocins. This result raised a question regarding the nature of the secretion mechanism of *Pcc* bacteriocins which was previously discovered to be secreted via T3bSS, a system that utilizes the bacterial flagellum for extracellular secretions. Our previous report has shown that bacteriocin Carocin S1 cannot be secreted by mutants that are defective of T3bSS-related genes such as *flhA*, *flhC*, *flhD* and *fliC*. We knocked out several genes making up the significant structural components of both T3aSS and T3bSS. The findings led us to hypothesize the potential roles of the T3aSS-related proteins, SctT, SctU and SctV, as flagellar T3SS chaperones in the secretion of *Pcc* bacteriocins. This current discovery and the findings of our previous study helped us to conceptualize a unique Type III secretion system for bacteriocin extracellular export which is a hybrid of the injectisome and flagellar secretion systems.

Introduction

Many bacteria often produce extracellular proteins to inhibit the growth of other competing bacterial species (1, 2). Such extracellular proteins produced are commonly called bacteriocins. Bacteriocins are usually found in Gram-positive or Gram-negative bacteria (3, 4, 5), and the functions in different bacterial species are similar. Most of the structural genes of bacteriocins (*e. g. E. coli* and *P. aeruginosa* bacteriocins) are located on the plasmid (6). For *Pectobacterium carotovorum* subsp. *carotovorum* (*Pcc*), the genes responsible for bacteriocins production is in the chromosomes (7, 8, 9, 10). When the bacterial growth environment changes and may cause harm, the structural genes will be activated to produce bacteriocin proteins to attack other related bacterial species. Because bacteriocin has the function of inhibiting the growth of other competing species, we can often see in some related literature that the bacteriocin protein produced by the isolated harmless bacterial species is transferred to other bacteria. Bacteriocins always inhibit the growth of closely related harmful bacteria (11), which shows that bacteriocins have a high application value and economic significance.

The classification of most bacteriocins is based on the producing bacteria, molecular weight, chemical structure, and functional mechanism (3, 12, 13). Gram-positive bacteria are often divided into four categories based on the presence of disulfide bonds or single sulfide bonds on the peptide chain of bacteriocins (14). For the Gram-negative bacteria, however, there is no complete classification basis, so the bacteriocins are roughly classified into two, which is based on their molecular weights, i.e. high-

molecular-weight bacteriocins (HMWB) and low-molecular-weight bacteriocins (LMWB). The strain *Pcc*, which is known to produce LMWBs Carocin S1 (7), Carocin D (8), Carocin S2 (9), and Carocin S3 (10), was selected in this study. These studies revealed that *Pcc* produces bacteriocins of different characteristics (7, 9). The bacteriocins contain two proteins, one responsible for the antimicrobial activity (killing protein), and the other for inhibition of the microbial activity of the killing protein (immunity protein). The killing proteins are organized in functional domains with receptor binding, translocation, and DNase or RNase activity (8).

When bacteria produce bacteriocins, how to deliver the killer protein outside of the cell through various secretion systems is also a topic of interest to many scholars. Several studies have shown that many gram-negative bacteria utilize the Type III secretion system (T3SS). During the course of an infection, several gram-negative pathogens, including *Shigella* (15), *Salmonella* (16), enteropathogenic *Escherichia coli* (17), and *Yersinia* species (18), use T3SS as injection devices to deliver multiple virulence proteins, referred to as effectors, directly into the cytosol of infected cells (19). Moreover, findings from our previous study have shown that the secretion of the LMWB Carocin S1 from *Pcc* is dependent on the flagellar Type III secretion system which also controls the bacterium's cell motility and cell size (20).

The T3SSs are intricate bacterial machinery that gives gram-negative bacteria a unique and distinctive mechanism for virulence, enabling them to deliver effector proteins into their host's cell cytoplasm bypassing the extracellular environment (21). These protein secretion mechanisms include the injectisome (T3aSS) and flagellar (T3bSS) systems (22).

The T3bSS, being the most ancestral type of T3SS, builds the bacterial flagellum which are commonly found in both gram-positive and gram-negative bacteria (22, 23). This extracellular structure enables bacterial motility in both surface and liquid environments (22, 24, 25). An evolutionary related nanomachine of the T3SS is the T3aSS, a needle-like nanomachine that has undergone an evolution, not only for the assembly of the injectisome, but also for energizing effector proteins translocation during invasion and colonization (26, 27, 28). The T3aSS and T3bSS share various homologies in terms of their structures. Both entail around 10 cytoplasmic and inner membrane proteins with the high resemblance in sequence and membrane topology (25, 26, 27). These machineries have three basic structural parts: the basal body, the MS ring, and the rod that passes through the periplasmic space (26, 29, 30, 31, 32). These two secretion systems, however, differ in their external structure. The T3aSS features a needle with a translocon pore located at its end, while the T3bSS contains a hook and a filament (26, 31, 32). How these two nanomachines are assembled is highly organized. They are regulated and constructed in highly explicit order. The assembly of T3aSS injectisome depends on varying affinities and kinetics of protein-protein interactions. On the other hand, assembly of the flagellum for T3bSS is based on a transcriptional hierarchy of three promoter classes (33, 34, 35).

Our previous study on the secretion mechanism of *Pcc*'s LMWB has shown that Carocin S1 is secreted via T3bSS (20). The findings revealed that *flhA*, *flhC*, *flhD* and *fliC*, genes involved in the flagellar virulence machinery, are required for the delivery of the bacteriocin outside of the cell. However, in this current

study, we found that *sctT*, a T3aSS-related gene, functions similarly in the secretion of Carocin S1 and Carocin S3 in the multiple bacteriocin-producing strain of *Pcc*. Thus, we hypothesized that the bacterium is capable of secreting bacteriocins through both the flagellar and injectosome T3SS, similar to *Burkholderia*, a bacterium capable of encoding multiple T3SSs (36, 37). Further deletions of various genes involved in the assembly of the flagellar and injectosome T3SS, however, raised an interesting question about the intricacy in the export mechanism of these *Pcc* bacteriocins. This led us to hypothesize a unique interaction of various T3SS proteins in the secretion of *Pcc* bacteriocins.

Results

Isolation and Detection of Tn5 insertion mutants

Pcc could undergo mutation and produce progeny strains with rifampicin resistance genes. These mutants still produce bacteriocins even after mutations. In this paper, we chose H-rif-8-6, which was from the rifampicin resistant strain 89-H-4, as the recipient of bacterial conjugation, and *E. coli* (1830) with the kanamycin resistance gene transposon Tn5 as the donor of conjugative reproduction. After mating, it is expected that the pJB4JI in *E. coli* could be transmitted into the *Pcc*. To screen the conjugated *Pcc*, we used modified Drigalski's medium with antibiotics kanamycin and rifamycin (data not shown). Subsequently, the bacteriocin assays were performed to find out the strains whose low-molecular-weight bacteriocin-related genes have been successfully disrupted (Fig. 1).

The mutant strain TH22-6 has shown an indication of the Tn5 insertion. The results have shown that Transposon Tn5 has successfully disrupted the low-molecular-weight bacteriocin production-related genes in the strain (38). Hence, we selected the mutant strain TH22-6 for further investigation. To further confirm the insertion, P3 and P4 primers were designed to amplify the specific *npfII* resistance gene on Tn5 (about 500 bp in length). In this technique, the PCR method can be used to confirm whether Transposon Tn5 has been inserted into the DNA of the wild strain (data not shown).

Amplification of the DNA at the Tn5 insertion junction site

Two sets of primers in different directions are designed to perform Thermal Asymmetric Interlaced PCR (TAIL-PCR). This was done to produce two DNA fragments extending outward from the Transposon Tn5 gene. Since the two sets of primers are both outside the Transposon Tn5 gene, the two DNA fragments will also contain genes that may have been disrupted in addition to the Tn5 gene. The ABI PRISM DNA Sequencing System 373S (Perkin-Elmer Corporation) was used to decipher the DNA sequence.

We amplified the Tn5 insertion junction DNA from TH22-6 genomic DNA by TAIL-PCR, and one or more PCR products were obtained. The genomic DNA of *Pcc*'s insertional mutant TH22-6 was used as the template DNA. The isolated chromosomal DNA was treated with *EcoRI* and confirmed with 0.7% agarose electrophoresis gel. Using specific primers PF1 ~ 3 and PR1 ~ 3 and random primer N2 (Table 2), and the above-mentioned TH22-6 chromosomal DNA as template DNA, we performed TAIL-PCR. Supplementary Fig. 1 shows the results of the TAIL-PCR.

Table 2
Primers used in this study

Primer	Sequence (5'→3')
PR-1	5'- GCCGAAGAGAACACAGATTTAGCCCA
PR-2	5'- CCGCACGATGAAGAGCAGAAGTT
PR-3	5'- CAGATCTCTGGAAAACGGGAAAGG
PF-1	5'- AGAGAACACAGATTTAGCCCAGTCGG
PF-2	5'- CCGCACGATGAAGAGCAGAAGTTAT
PF-3	5'- GATCCTGGAAAACGGGAAAGGTTC
N-1	5'- NGTCGA(G/C)(A/T)GANA(A/T)GAA
N-2	5'- GTNCGA(C/G)(A/T)CANA(A/T)GTT
N-3	5'- (A/T)GTGNAG(A/T)ANCANAGA
P-3	5'- CTCGACGTTGTCACTGAAGCGGGAAG
P-4	5'- AAAGCACGAGGAAGCGGTCAGCCCAT
PJ1	5'-GTTTTTTCAGCCATTGTCGC
PB1	5'-TCTGGCTTTCTGAACTTTGC
sctT-sen	5'-TGAAGCTTATGAGCCCAGTG
sctT-anti	5'-TAATAAGCTTTGGTGCAGCC
flhA-sen	5'-TCACTCAACGTTGCATCTAC
flhA-anti	5'-CAAGATGTTGGCCAACAGATG
sctV-sen	5'-TCCGATATAGGTGTTGAGGC
sctV-anti	5'-CCAGCCAGTTAATGATGTGC
sctU-sen	5'-CAATCGTCCTGAACTGTTGG
sctU-anti	5'-GTACTGACTGCACCATGCTC
sctD-sen	5'-GAAGCACCGTGGTGTGGAAG
sctD-anti	5'-GCCTTCTGGATAGCGTTGATG
sctQ-sen	5'-CGTGGCAGCCACTCGAGTGTGAAC
sctQ-anti	5'-GCAGACAAGGATCCTCAGCGGAATC
flgH-sen	5'-ATGGCGAATAAATGGCGTTG
flgH-anti	5'-GGAAGAAACGCTGTAACCAC

Primer	Sequence (5'→3')
flgF-sen	5'-GAATCGCAGCCTGTCACAAC
flgF-anti	5'-TGTGCCAGATCCTCTGCAAG
flhB-sen	5'-AAACAGAAGCTTCCACTCCC
flhB-anti	5'-ACGCCGCAGTCGACGCTTC
fliR-sen	5'-TGATACCAGCCAACTCAGTC
fliR-anti	5'-CCAGCCGATCAAAGAATTTCG
fliG-sen	5'-ATGACCCTGACAGGAACAG
fliG-anti	5'-TTAGACATAAGCATCCTCGC
fliE-sen	5'-CAGTCGCTATGTCCCCTAAC
fliE-anti	5'-TACACCTGCATGCTCATCAC
flgG-sen	5'-TACCCGTGACGGTTCATTTC
flgG-anti	5'-TTGTAGCATCTGATCATACG
^a All primers were purchased from MDE Bio Inc., Taipei, Taiwan	

For TAIL-PCR products sequencing, an automated DNA sequencer 373S (ABI) was used, and a nucleotide sequence of 2249 bp was obtained. The nucleotide sequence and deduced amino acid of T3SS protein were compared by the BLAST programs of National Center for Biotechnology Information server (National Library of Medicine, USA). Sequence data were compiled with DNASIS-Mac software (Hitachi, Japan). The results show that at least two complete open reading frames (ORF1 and ORF2) were present, and Tn5 was in the ORF2, between 2248th and 2249th base pair (data not shown). Based on the nucleotide sequence of TAIL-PCR product, the DNA probe (431 Probe) which was designed near the Tn5 insertion region, was hybridized with the digested wild-type strain genomic DNA (Supplementary Fig. 2).

Thereafter, we sliced the DNA agarose electrophoresis gel, then purified and cloned it into pBR322 for genome library. Finally, we acquired the construction which bears the nucleotide of the interrupted gene (Supplementary Fig. 3). The nucleotide was sequenced and its length is 3031 base pairs (Fig. 2).

Sequence Analysis and Homology

Sequence Walking was performed to identify the two DNA sequences of TAIL-PCR products. It was found that the two single-band PCR products amplified by primers PF-A3 and N2, and the single band PCR products amplified by PL3 and N1, has a total length of 2249 bp. The TAIL-PCR products amplified by specific primers PF1 ~ 3 and PR1 ~ 3 with the TAIL-PCR product amplified by specific primers PF-A1 ~ 3 and PL1 ~ 3 were also compared. Both have inserted mutations by Tn5 and the TAIL-PCR products extend outward from PCR amplification.

We performed sequence analysis using (BLAST) by the National Institutes of Health Medical Library (NCBI) database (Fig. 3). The result showed that ORF2 shows high homology with the YsaT protein of T3SS in *Sodalis glossinidius* (61% homology) and *Yersinia enterocolitica* (45% homology). Another high homology cassette (ORF1) in the nucleotide has shown 74% homology with YsaR, also an assembly of T3SS.

After compiling the sequence with DNASIS-Mac software, it indicates that ORF2 encodes 762 base pairs and a protein of 254 amino acids, which shows high homology with the YsaT protein of the T3SS in *Sodalis glossinidius* and *Yersinia enterocolitica*. Moreover, the hydrophobicity of YsaT-like amino acid was arranged to and fro (Supplementary Fig. 4). The YsaT protein of these two strains is one of the essential proteins in their bacteriocin secretory systems. Hence, we speculate that the gene mutated by the insertion of Transposon Tn5 may be a gene that secretes bacteriocin. The gene was later designated as *sctT*, in accordance with the unified secretion and cellular translocation (Sct) nomenclature.

Complementation and analysis of *flhA* and *sctT* genes

Wild-type H-rif-8-6 was used as a control and transformed with plasmids containing the *flhA* (*flhA*-KO) and *sctT* (TH22-6) genes. The mutations in the *flhA* gene (*flhA*-KO) and *sctT* (TH22-6) gene were complemented by the introduction of the *flhA* and *sctT* genes, and the effects of these respective genes on bacteriocin production were evaluated.

Neither the TH22-6 (*sctT*-KO, *sctT*::Tn5) nor *flhA*-KO (*flhA*::Kan) mutant strains could secrete bacteriocins Carocin S1 and Carocin S3. However, the ability of these strains to secrete bacteriocins was recovered after complementation by transformation with the *flhA* and *sctT* genes, respectively, which have shown inhibition zone diameters comparable in size to that of wild type (Fig. 4A). After transformation, all deletion strains harboring their respective complementing plasmids secreted LMWB (Fig. 4B). It can be observed that knocking down *sctT* and *flhA* disables the extracellular export of bacteriocins Carocin S1 and Carocin S3 (Fig. 4A, No. 2 and 3). However, it can be observed that the bacteriocins are still expressed as displayed in the RT-PCR experiment (Fig. 4B No. 2 and No. 3).

Deletion of the T3aSS and T3bSS-Related Genes and Bacteriocin Activity Assay

We knocked out several genes making up the flagellar and injectisome T3SS to find out how the deletion affects the bacteriocin secretion. The results have shown that deletion of *sctU*, *sctV*, *flgG*, *fliE* and *fliR* genes disabled the secretion of bacteriocin. This suggests that these genes are essential and are required in the secretion of Carocin.

Results from our previous research have shown the significance of *flhA*, *flhC*, *flhD*, and *fliC* genes in regulating the synthesis of bacterial flagella in *Pcc* and that Carocin S1 utilizes this secretion machinery (T3bSS) (20). The results of the present study have shown that further deletion of T3bSS-related genes

such as *flgG*, *fliE*, *fliR*, and *flhA* similarly disrupts the extracellular export of Carocin S1 and Carocin S3 (Fig. 5). This result further supports the previous findings that the T3bSS is used in the secretion of *Pcc* bacteriocin. However, it can also be observed that deletion of various T3aSS-related genes like the *sctT*, *sctU*, and *sctV* likewise affects the Carocin S1 and Carocin S3 secretion. Interestingly, deletion of the T3bSS gene *flhB* did not restrain bacteriocin secretion to the outside of the cell but knocking out its homologous gene in the T3aSS, *sctU*, inhibited the expression of the bacteriocins to the extracellular environment.

The current findings of this study have shown an interesting interaction between the T3aSS and T3bSS. From these findings, we hypothesize that the genes *sctT*, *sctU*, and *sctV* play important roles in the synthesis of chaperones for the delivery of bacteriocins Carocin S1 and Carocin S3.

Discussion

This experiment uses Transposon Tn5 to disrupt the genes related to LMWB production, and TAIL-PCR method was used to amplify the PCR product (2249bp DNA fragment) extending from both ends of Tn5. The 2249 bp DNA was found after the NCBI Genbank alignment sequence. The part disrupted by Transposon Tn5 has a complete gene (*sctT*). The amino acid sequence of the disrupted gene is 45% homologous with the YsaT protein of *Yersinia enterocolitica* and 61% homologous with *Sodalis glossinidius*. Literature (39, 40) indicates that these proteins are known to be involved in the construction of T3SS. When the *sctT* gene was disrupted, *Pcc* LMWBs no longer express (high molecular weight bacteriocins still exist). Hence, we speculate that the gene disrupted by Tn5 is likely to be a protein component involved in the expression of LMWBs.

We compared the DNA fragment obtained from the chromosomal DNA of H-rif-8-6 (3031 bp DNA fragment), NCBI Genbank protein sequence alignment, and TAIL-PCR product (2249 bp DNA fragment), and the results were roughly the same. The *sctT* gene is located at the end of the 3031 bp DNA fragment. There are about two to three ORFs in the front and rear regions of the *sctT* gene. Most of these ORFs are related to the constituent proteins of the T3SS.

Much of our understanding of the T3SS is derived from the study of pathogenic bacteria and their interactions with plant and animal hosts (41). T3SSs of bacteria are specific export machineries for virulence that allow their translocation to eukaryotic cells. The SctT protein is one of the injectisome T3SS apparatus. It is part of the inner membrane machinery of the T3aSS (42). It belongs to the five highly conserved inner membrane proteins (SctRSTUV) which are necessary for the function of the pathogenic T3SS, though, their individual functions are unclear. It is worth noting that these proteins show a high degree of sequence homology to components of the evolutionarily related flagellar secretion system (T3bSS) and may represent a functional core, serving critical chemical roles in initiating or powering protein secretion. SctRSTUV are important in the organized, stepwise assembly of the T3aSS basal body.

The role of SctT is not well understood, but it has been identified as a T3aSS protein based on sequence homology among multiple bacterial species (43). This study would be the first to report the function of SctT in the secretion of an effector protein. By the complementary experiment, the result showed the *sctT* gene expression was a requirement for the LMWBs, Carocin S1 and Carocin S3, secretion. The current result on the role of *sctT*, a gene that encodes a protein in the T3aSS assembly, have led us to further investigate some possible interactions between the T3aSS and T3bSS genes in the secretion of *Pcc* bacteriocins.

Previous findings (20) have shown that Carocin S1 is secreted via T3bSS. This study, however, has shown that its secretion is also affected by deletion of a T3aSS-related gene. According to Sun et al. (36) and Denise et al. (37), some bacteria encode multiple T3SSs (e. g. *Burkholderia*) to interact with multiple types of eukaryotic hosts. Also, according to Matsuda et al. (45), some effectors can evolve to be recognized by multiple systems (e.g. a toxin from *Vibrio*). In this case, it might be possible that *Pcc* can also encode multiple secretion systems, and the bacteriocins Carocin S1 and Carocin S3 are recognized by both secretion machineries.

To test this assumption, we performed series of deletions of multiple genes related and significant to both the T3aSS and T3bSS. We found that deletion of T3aSS-related genes (*sctT*, *sctU* and *sctV*) and T3bSS-related genes (*flgG*, *fliE*, *fliR* and *flhA*) did impede the secretion of bacteriocin. Interestingly, deletion of any of the T3bSS-related gene (*flgG*, *fliE*, *fliR* and *flhA*) totally halts the secretion of bacteriocin. Similarly, omission of T3aSS-related genes (*sctT*, *sctU*, and *sctV*) impedes the secretion of bacteriocin. These results did not support our initial assumption. If *Pcc* is capable of encoding both types of T3SS like what Sun et al. (36) and Denise et al. (37) explained about *Burkholderia*, disabling one secretion system (e. g. T3aSS) could possibly not disrupt the delivery of Carocin outside the cell because the presence of the other machinery (T3bSS) serves as an alternative system for secretion. The results of the experiment, and that of the previous study (20), nevertheless, reveals that the *Pcc* bacteriocin could be secreted by a single secretion machinery that utilizes the genes related to both T3aSS and T3bSS. Thus, this led to the rejection of the assumption that *Pcc* can encode both types of T3SS.

We therefore hypothesized that the inability of *Pcc* to secrete Carocin S1 and Carocin S3 resulted from the absence of T3aSS genes *sctT*, *sctU* and *sctV*, which encode chaperone proteins for the bacteriocin secretion via T3bSS. Figure 6 shows the hypothesized flagellar secretion system, with T3aSS proteins acting as chaperones.

SctT, SctU and SctV are known to be part of the conserved components of the T3aSS (23, 45, 46). The inner membrane ring of the T3SS is associated with the export apparatus, which is assembled by members of five families of transmembrane proteins (SctRSTUV) and presumably forms a transport channel for T3SS substrates (46, 47, 48, 49, 50). This study would be the first to report the potential role of these T3aSS proteins in the delivery of an effector protein via T3bSS. As inferred from the results of this study, these proteins serve as potential chaperones in the secretion of bacteriocins Carocin S1 and Carocin S3 in *Pcc*.

Secretion chaperones are of great significance to bacteria because intact chaperones are needed for the export of substrates (51, 52) or for the regulation of virulence gene expression (53). T3SS chaperones are either required for or regulate effector translocation into host cells. Hence, identifying the chaperones involved in the export of substrates is essential to better understand the secretion systems of bacteria. Literatures stipulate that secretion of most effectors was reduced but not abolished in the absence of their chaperones (54, 49, 55, 56, 57, 58). The result of the present study, however, shows that the absence of chaperones halts the secretion of the effector proteins in *Pcc*. Recent reviews on flagellar chaperones, on the other hand, have stated that flagellar chaperones might be required mainly to prevent premature interactions between subunits of the flagellum by masking their interaction domains (59). For example, the flagellar system uses cytoplasmic chaperones (60) such as the FlgN for FlgL and FlgK (hook-related proteins), FliT for FlgL and FlgK (filament-capping proteins), and FliS for FliC (flagellin protein) (61, 62, 63).

It was proposed that flagellar chaperones mediate interactions between the various subunits of the flagellum by binding to the carboxy-terminal region of the substrate (61, 62, 63, 64). These C-terminal domains are thought to be the glue by which individual filament subunits polymerize together during assembly of a growing flagellum (66). Unless these domains are stabilized by a bound chaperone, the monomeric forms are swiftly oligomerized. This has been proven experimentally. Flagellar chaperones such as the FliS (62, 63, 64) and FlgN (66) prevent premature polymerization.

In this report, Carocin S1 and Carocin S3 are detected in the RT-PCR in small amounts (see Fig. 5). However, the bacteriocin assay revealed that these bacteriocins were not extracellularly secreted. Similarly, previous reports (62, 66) have shown that substrates were detected in lower amounts and were not secreted in the absence of their chaperones. Parsot et al. (67) explains that premature associations of flagellar proteins in the cytoplasm result in their instability. However, by masking the interacting domains in the components of the flagellum, chaperones prevent the premature associations among proteins. Thus, in the absence of chaperones, these processes would possibly be altered. In our experiment, deletion of genes associated with the chaperone proteins have prevented the extracellular secretion of Carocin S1 and Carocin S3. The bacteriocins have been produced but not extracellularly secreted perhaps due to non-assembly of the flagellar components essential for their secretion which could have resulted from the possible instability and premature associations of the various flagellar proteins.

Arguably, the principal role of flagella-specific T3SS chaperones is to prevent early polymerization of cognate flagella substrates. Moreover, assembly of filament subunits into the growing flagellum is strictly sequential. This implies a strict secretion order, which may result if each chaperone-substrate complex interacts with a different affinity to a component of the flagella-specific T3SS (65).

The roles of the T3aSS-related proteins, SctT, SctU and SctV, in the secretion of *Pcc* bacteriocins could be any of the common roles of chaperones in the flagellar T3SS. Whether they act to protect substrates from premature associations with self or other proteins, to guide substrates to the correct T3SS, to establish a secretion order among the substrates, or to couple gene transcription to the process of apparatus

assembly, their specific function in the secretion of Carocin S1 and Carocin S3 remains an unanswered question and needs further investigation and confirmation.

Conclusion

As shown herein, the extracellular export of *Pcc* LMWBs Carocin S1 and Carocin S3 are dependent on secretion machinery involving genes related to both the injectisome (T3aSS) and flagellar (T3bSS) secretion systems. This study is the first to report such unique interaction between genes and proteins of both types of T3SS. Results of gene deletions revealed that *Pcc* bacteriocins are possibly secreted by the T3bSS which utilizes T3aSS-related proteins SctT, SctU, and SctV as secretion chaperones. Moreover, it was found that these chaperone proteins do not act individually, rather, they work in concert in the bacteriocin secretion. However, how they interact with the bacteriocins, and what are their specific role as flagellar T3SS chaperones need further investigations.

Methods

Bacterial strains, plasmids, media, and growth conditions

The bacterial strains and plasmids used in the study are shown in Table 1. Propagation of the *Pectobacterium carotovorum* subsp. *carotovorum* strains was done in 1.4% nutrient agar (NA) at 28°C or in LB medium with 5 g/L of NaCl with shaking. On the other hand, *E. coli* strains were propagated at 37°C in a modified LB medium with shaking. Rifampicin, kanamycin, and ampicillin (50 mg/L) were added to the media when required.

Table 1
Bacteria and plasmids used in this study

Bacterium or plasmid	Relevant characteristics	Source
<i>E.coli</i>		
1830	<i>pro⁻ met^r Km^r Nm^r</i> , containing transposon	Gantotti et al. [38]
	Tn5 on the "sucidal" plasmid pJB4JI	
DH5α	<i>supE44hsdR17recA1endA1gyrA1thi-1relA1</i>	Hanahan; Reusch et al [72, 74]
<i>Pcc</i>		
89-H-4	putative biocontrol agent	Laboratory stock
H-rif-8-6	89-H-4, <i>Rif^r</i>	this work
Ea1068	wild type	Laboratory stock
SP33	wild type	Laboratory stock
TH22-6	H-rif-8-6, <i>sctT::Tn5, Rif^r, Kan^r</i>	this work
FlhA-KO	H-rif-8-6, <i>flhA::Kan, Rif^r, Kan^r</i>	Chan et al. [9]
SctV-KO	H-rif-8-6, <i>sctV::Kan, Rif^r, Kan^r</i>	this work
SctU-KO	H-rif-8-6, <i>sctU::Kan, Rif^r, Kan^r</i>	this work
SctD-KO	H-rif-8-6, <i>sctD::Kan, Rif^r, Kan^r</i>	this work
SctQ-KO	H-rif-8-6, <i>sctQ::Kan, Rif^r, Kan^r</i>	this work
FlgH-KO	H-rif-8-6, <i>flgH::Kan, Rif^r, Kan^r</i>	this work
FlgF-KO	H-rif-8-6, <i>flgF::Kan, Rif^r, Kan^r</i>	this work
FlhB-KO	H-rif-8-6, <i>flhB::Kan, Rif^r, Kan^r</i>	this work
FliR-KO	H-rif-8-6, <i>fliR::Kan, Rif^r, Kan^r</i>	this work
FliG-KO	H-rif-8-6, <i>fliG::Kan, Rif^r, Kan^r</i>	this work
FliE-KO	H-rif-8-6, <i>fliE::Kan, Rif^r, Kan^r</i>	this work
FlgG-KO	H-rif-8-6, <i>flgG::Kan, Rif^r, Kan^r</i>	this work
plasmid		

Bacterium or plasmid	Relevant characteristics	Source
pACYC177	<i>Amp^r</i> , <i>Kan^r</i> , low copy number	Chang et al. [75]
pBR322	<i>Amp^r</i> , <i>Kan^r</i>	Bolivar et al. [68]
pB47	<i>Amp^r</i> , <i>Kan^r</i> , 3.0kb DNA fragment contain <i>sctT</i>	this work
pBSCTT	<i>Amp^r</i> , <i>sctT</i>	this work
pBFA	<i>Amp^r</i> , <i>flhA</i>	this work
<i>Amp^r</i> indicates ampicillin resistance, <i>Rif^r</i> indicates rifampicin resistance, and <i>Kan^r</i> indicates Kanamycin resistance.		

Bacterial mating

For bacterial mating, the membrane-filter method was carried out as described by Gantotti et al. (38). Overnight cultures of *Pcc* (recipient) H-rif-8-6 and *E. coli* (donor) 1830 were evenly spread onto 0.22 µm pore size membrane filters (Millipore, Inc. Bedford, MA) which were placed on LB agar media and incubated overnight at 28°C. After conjugation, suspensions of the progeny were appropriately diluted and were grown at 28°C for 24 to 48 h on modified Drigalski's agar plates (with rifampicin and kanamycin, 100 µg ml⁻¹). Colonies were isolated for the bacteriocin production test.

Bacteriocin assays

Bacteriocin production of the isolates was examined using the double-layer method as previously described (68). The isolates were placed on hard IFO-802 (with 1.4% agar) and soft IFO-802 (with 0.65% agar) medium. The cells were first incubated for 12 hours for colonies to form. The colonies were then exposed to ultraviolet irradiation before incubating for another 12 hours. Thereafter, the cells were treated with chloroform and finally covered with soft agar containing the indicator cells. An inhibition zone of indicator-cell (SP33 or Ea1068) growth around the colony indicates bacteriocin production.

Preparation of genomic DNA, plasmid DNA and RNA

The procedures of plasmid preparation, genomic DNA isolation, and DNA manipulation were performed according to Sambrook (69). For RNA preparation, exponentially growing *E. coli* DH5α cells were harvested (OD₅₉₅ of about 6.0). Trizol reagent (Invitrogen, USA) was used in isolating the RNA which was resuspended in diethylpyrocarbonate (DEPC)-treated water. RNA concentration was determined at OD₂₆₀ absorption and was analyzed by electrophoresis on 1.5% formaldehyde-morpholinepropanesulfonic-agarose gel.

TAIL-PCR and Restriction DNA library screening

Previously detailed protocols were utilized for the general polymerase chain reaction (PCR) (69) and thermal asymmetric interlaced PCR (TAIL-PCR) (70).

Specific primers at both ends of Tn5, namely PF1, PF2, PF3, PR1, PR2, and PR3 were designed for TAIL-PCR. Using the above primers for amplification, two PCR products with unknown DNA sequences at both ends of Tn5 were obtained. However, after the sequence analysis, it was found that the two segments were the same PCR products. Hence, two sets of two PCR products were designed using this known DNA sequence of about 300 bp.

ABI PRISM Dye Terminator Cycle Sequencing Ready Reaction kit (Applied Biosystems, Foster City, CA) was used in the sequence analysis of the TAIL-PCR products. Cycle sequencing was carried out in a GeneAmp System 9600 thermocycler (Applied Biosystems). The sequencing was carried out using an ABI 373S automated DNA sequencer 373S (Applied Biosystems) based on the manufacturer's protocol.

Hybridization and Southern blots

Southern blots were performed according to the DIG Application Manual (Roche, USA). A 543-bp DNA fragment (431 probe) was amplified with pJI and pBI primers, subcloned into pGEM-T Easy vector (Promega Inc., USA), and labeled using a Random Primed DNA Labeling Kit (Roche Diagnostics, USA). The genomic DNA of the wild-type strain H-rif-8-6 was digested with various restriction endonucleases, with sites located outside the putative open reading frame. Samples were electrophoresed and analyzed with Southern blotting. After detection using the 431 probe, the DNA from positive gel slices was purified and cloned into pBR322 to give the carocin-producing plasmid pB47. The pB47 construct was isolated and detected as above with the 431 probe.

Southern and colony hybridizations, probe labeling, and detection were performed by using a DIG DNA Labeling and Detection kit (Boehringer Mannheim GmbH, Mannheim, Germany). Hybridization was performed overnight, and the membrane was washed according to the recommendations of the manufacturer. DNA electrophoresis, restriction digestion, ligation, and transformation for *E. coli* were carried out as described by Sambrook et al. (69). Plasmid DNA transformation for *Pcc* was performed by the methods of Hinton et al. (71) and Hanahan (72).

Subcloning of *sctT* gene from H-rif-8-6

The DNA fragment of SctT was amplified by PCR from H-rif-8-6 using oligonucleotide primers SctT-sen and SctT-anti. The PCR product was subcloned into pGEM-T Easy vector by TA cloning (Promega Inc., USA). The *sctT* gene containing product was digested with restriction enzymes *Hind*III and subcloned into plasmid pBR322. The new plasmid was designated pBSctT. One hundred transformed colonies were isolated using selective LB agar containing 100 µg/ml of ampicillin after the transfer of pBSctT into *E. coli* DH5α. The presence of the *sctT* gene was detected by colony hybridization using the 431 probe and electrophoresis after digestion with *Hind*III to yield the expected 10-Kb DNA fragment bearing SctT protein. The pSctT plasmid was isolated from DH5α/pSctT and transferred into the insertion mutants of

Pcc TH22-6. One hundred colonies were isolated by selection on modified Drigalski's medium containing 50 µg/ml of kanamycin, rifampicin, and ampicillin. The *sctT* gene was detected as previously described.

Construction of the null alleles of *flhB*, *flgH*, *flgF*, *fliR*, *fliE*, *flgG*, *fliG*, *sctD*, *sctQ*, *sctV*, and *sctU* genes

The various genes were isolated from pBSctT by digesting with different restriction enzymes which cleave at two sites in pBSctT and thereby conveniently delete the target genes from the operon. The resulting plasmid was designated and labeled accordingly.

A kanamycin-resistant gene from pACYC177 was isolated, made blunt-ended using a DNA-blunting kit (Takara Co., Tokyo, Japan), and inserted in the unique restriction site of the target genes. The resulting plasmid was designated consequently (see Table 1). The gene-Kan was re-isolated and linearized after restriction enzyme digestion, which deleted the ampicillin resistance gene and replication site of the plasmid.

The linearized construct was transferred into H-rif-8-6, resulting in the homologous replacement of the native genes and generating null alleles. The DNA fragments of the target gene were amplified by PCR from H-rif-8-6. After PCR amplification using pairs of oligonucleotide primers (see Table 2), the partial DNA fragments were isolated and subcloned into plasmid pBR322 to generate the needed plasmid.

RNA preparation and RT-PCR

Bacteriocin synthesis medium (BSM; 0.5% sucrose, 0.1% NH₄Cl, 0.2% KH₂PO₄, and 0.02% MgSO₄·7H₂O (pH = 7.5) was used to produce bacteriocins Carocin S1 and Carocin S3. Total RNA was extracted from cells (*Pcc* harboring constructs) that were grown without antibiotics at 28°C. To determine the stability of the strains, culture samples (8 ml each; with rifampicin [0.2 mg/ml] added when cell density was ~ 150 Klett units to block bacterial contamination) were withdrawn at various time points and transferred to tubes containing 5 mL of ice-cold water. Total RNA was extracted using Trizol (Invitrogen, Carlsbad, CA) according to the manufacturer's protocol.

On the other hand, the Reverse Transcription-PCR (RT-PCR) using AMV Reverse Transcriptase (Promega, USA) was carried out based on the instructions provided by the manufacturer. One microgram (1 µg) of RNA was subjected to RT-PCR. Reverse primer (see Table 2) was used in first strand cDNA synthesis. The RT mixtures were diluted and used as templates in a PCR reaction with the pairs of primers (Table 2).

Computer analysis of sequence data

The nucleotide sequence and the deduced amino acid sequence of *sctT* were compared using the BLAST and FASTA programs of the National Center for Biotechnology Information server (National Library of Medicine, USA). Sequence data were compiled by DNASIS-Mac software (Hitachi, Tokyo, Japan).

References

1. Cotter PD, Hill C, Ross RP. Bacteriocins: developing innate immunity for food. *Nat Rev Microbiol.* 2005;3:777–788. <https://doi.org/10.1038/nrmicro1273>
2. Hibbing ME, Fuqua C, Parsek MR, Peterson SB. Bacterial competition: surviving and thriving in the microbial jungle. *Nat Rev Microbiol.* 2010;8(1):15–25. <https://doi.org/10.1038/nrmicro2259>
3. Tagg JR, Dajani AS, Wannamaker LW. Bacteriocins of gram-positive bacteria. *Bacteriol Rev.* 1976;40(3):722.
4. Kotelnikova EA, Gelfand MS. Bacteriocin production by Gram-positive bacteria and the mechanisms of transcriptional regulation. *Russ J Genet.* 2002;38(6):628–41. <https://doi.org/10.1023/A:1016035700012>
5. Chi H, Holo H. Synergistic antimicrobial activity between the broad spectrum bacteriocin garvicin KS and nisin, farnesol and polymyxin B against gram-positive and gram-negative bacteria. *Curr Microbiol.* 2018;75(3):272–7. <https://doi.org/10.1007/s00284-017-1375-y>
6. Eklund MW, Poysky FT, Mseitif LM, Strom MS. Evidence for plasmid-mediated toxin and bacteriocin production in *Clostridium botulinum* type G. *Appl Environ Microbiol.* 1988;54(6):1405–8.
7. Chuang DY, Chien YC, Wu HP. Cloning and expression of the *Erwinia carotovora* subsp. *carotovora* gene encoding the low-molecular-weight bacteriocin Carocin S1. *J Bacteriol.* 2007;189(2):620–6. <https://doi.org/10.1128/JB.01090-06>
8. Roh E, Park TH, Kim MI, Lee S, Ryu S, Oh CS, Rhee S, Kim DH, Park BS, Heu S. Characterization of a new bacteriocin, Carocin D, from *Pectobacterium carotovorum* subsp. *carotovorum* Pcc21. *Appl Environ Microbiol.* 2010;76(22):7541–9. <https://doi.org/10.1128/AEM.03103-09>
9. Chan YC, Wu JL, Wu HP, Tzeng KC, Chuang DY. Cloning, purification, and functional characterization of Carocin S2, a ribonuclease bacteriocin produced by *Pectobacterium carotovorum*. *BMC Microbiol.* 2011;11(1):99. <https://doi.org/10.1186/1471-2180-11-99>
10. Wang JW, Derilo RC, Lagitnay RB, Wu HP, Chen KI, Chuang DY. Identification and characterization of the bacteriocin Carocin S3 from the multiple bacteriocin producing strain of *Pectobacterium carotovorum* subsp. *carotovorum*. *BMC Microbiol.* 2020;20(1):1–3. <https://doi.org/10.1186/s12866-020-01955-9>
11. Mishra C, Lambert J. Production of anti-microbial substances by probiotics. *Asia Pac J Clin Nutr.* 1996;5:20–4.
12. Kalmokoff ML, Banerjee SK, Cyr T, Hefford MA, Gleeson T. Identification of a new plasmid-encoded sec-dependent bacteriocin produced by *Listeria innocua*. *Appl Environ Microbiol.* 2001;67(9):4041–7. <https://doi.org/10.1128/AEM.67.9.4041-4047.2001>
13. Baysse C, Meyer JM, Plesiat P, Geoffroy V, Michel-Briand Y, Cornelis P. Uptake of pyocin S3 occurs through the outer membrane ferripyoverdine type II receptor of *Pseudomonas aeruginosa*. *J Bacteriol.* 1999;181(12):3849–51. <https://doi.org/10.1128/JB.181.12.3849-3851.1999>

14. Dale C, Young SA, Haydon DT, Welburn SC. The insect -endosymbiont *Sodalis glossinidius* utilizes a type III secretion system for cell invasion. Proc Natl Acad Sci USA. 2001 Feb 13;98(4):1883-8. <https://doi.org/10.1073/pnas.98.4.1883>
15. Espina, M., A. J. Olive, R. Kenjale, D. S. Moore, S. F. Ausar, R. W. Kaminski, E. V. Oaks, C. R. Middaugh, W. D. Picking, and W. Picking. 2006. IpaD localizes to the tip of the type III secretion system needle of *Shigella flexneri*. Infect Immun. 74:4391–4400. <https://doi.org/10.1128/IAI.00440-06>
16. Park D, Lara-Tejero M, Waxham MN, Li W, Hu B, Galán JE, Liu J. Visualization of the type III secretion mediated *Salmonella*-host cell interface using cryo-electron tomography. Elife. 2018;7:e39514. <https://doi.org/10.7554/eLife.39514.001>
17. Majewski DD, Lyons BJ, Atkinson CE, Strynadka NC. Cryo-EM analysis of the SctV cytosolic domain from the Enteropathogenic *E. coli* T3SS injectisome. J Struct Biol. 2020:107660. <https://doi.org/10.1016/j.jsb.2020.107660>
18. Wang H, Avican K, Fahlgren A, Erttmann SF, Nuss AM, Dersch P, Fallman M, Edgren T, Wolf-Watz H. Increased plasmid copy number is essential for *Yersinia* T3SS function and virulence. Science. 2016;353(6298):492–5. <https://doi.org/10.1126/science.aaf7501>
19. Galán JE, Lara-Tejero M, Marlovits TC, Wagner S. Bacterial type III secretion systems: specialized nanomachines for protein delivery into target cells. Annu Rev Microbiol. 2014;68:415–38. <https://doi.org/10.1146/annurev-micro-092412-155725>
20. Chan YC, Wu HP, Chuang DY. Extracellular secretion of Carocin S1 in *Pectobacterium carotovorum subsp. carotovorum* occurs via the type III secretion system integral to the bacterial flagellum. BMC Microbiol. 2009;9(1):18. <https://doi.org/10.1186/1471-2180-9-181>
21. Coburn B, Sekirov I, Finlay BB. Type III secretion systems and disease. Clin Microbiol Rev. 2007;20(4):535–49. <https://doi.org/10.1128/CMR.00013-07>
22. Renault TT, Guse A, Erhardt M. Export mechanisms and energy transduction in type III secretion machines. Curr Top Microbiol Immunol. 2019;20:99–17. https://doi.org/10.1007/82_2019_166
23. Hueck CJ. Type III protein secretion systems in bacterial pathogens of animals and plants. Microbiol Mol Biol Rev. 1998;62:379–433. <https://doi.org/10.1111/j.1365-2958.2006.05301.x>
24. Berg HC, Anderson RA. Bacteria swim by rotating their flagellar filaments. Nature. 1973;245:380–382. <https://doi.org/10.1038/245380a0>
25. Kubori T, Matsushima Y, Nakamura D, Uralil J, Lara-Tejero M, Sukhan A, Galán JE, Aizawa SI. Supramolecular structure of the *Salmonella typhimurium* type III protein secretion system. Science. 1998;280(5363):602–5. <https://doi.org/10.1126/science.280.5363.602>
26. Blocker A, Komoriya K, Aizawa SI. Type III secretion systems and bacterial flagella: insights into their function from structural similarities. Proc Natl Acad Sci USA. 2003;100(6):3027–30. <https://doi.org/10.1073/pnas.0535335100>
27. Cornelis GR. The type III secretion injectisome. Nat. Rev. Microbiol. 2006;4(11):811–25. <https://doi.org/10.1038/nrmicro1526>

28. Galán JE, Wolf-Watz H. Protein delivery into eukaryotic cells by type III secretion machines. *Nature*. 2006;444(7119):567–73. <https://doi.org/10.1038/nature05272>
29. Diepold A. Assembly and post-assembly turnover and dynamics in the type III secretion system. In: Wagner S., Galan J. (eds) *Bacterial type III protein secretion systems*. *Curr Top Microbiol Immunol*. 2019; 8:e1002983–32. https://doi.org/10.1007/82_2019_164
30. Diepold A, Armitage JP. Type III secretion systems: the bacterial flagellum and the injectisome. *Philos Trans R Soc Lond B Biol Sci*. 2015;370(1679):20150020. <https://doi.org/10.1098/rstb.2015.0020>
31. Macnab RM. Type III flagellar protein export and flagellar assembly. *Biochim Biophys Acta Mol Cell Res*. 2004;1694(1–3):207–17. <https://doi.org/10.1016/j.bbamcr.2004.04.005>
32. Erhardt M, Namba K, Hughes KT. Bacterial nanomachines: the flagellum and type III injectisome. *Cold Spring Harb Perspect Biol*. 2010;2(11):a000299. <https://doi.org/10.1101/cshperspect.a000299>
33. Macnab RM. How bacteria assemble flagella. *Annu Rev Microbiol*. 2003;57(1):77–100. <https://doi.org/10.1146/annurev.micro.57.030502.090832>
34. Chevance FF, Hughes KT. Coordinating assembly of a bacterial macromolecular machine. *Nat Rev Microbiol*. 2008;6(6):455–65. <https://doi.org/10.1038/nrmicro1887>
35. Diepold A, Wagner S. Assembly of the bacterial type III secretion machinery. *FEMS Microbiol Rev*. 2014;38(4):802–22. <https://doi.org/10.1111/1574-6976.12061>
36. Sun GW, Chen Y, Liu Y, Tan GY, Ong C, Tan P, Gan YH. Identification of a regulatory cascade controlling Type III Secretion System 3 gene expression in *Burkholderia pseudomallei*. *Mol Microbiol*. 2010;76(3):677–89. <https://doi.org/10.1111/j.1365-2958.2010.07124.x>
37. Denise R, Abby SS, Rocha EP. The evolution of protein secretion systems by co-option and tinkering of cellular machineries. *Trends Microbiol*. 2020;28(5):372–86. <https://doi.org/10.1016/j.tim.2020.01.005>
38. Gantotti BV, Kindle KL, Beer SV. Transfer of the drug-resistance transposon Tn5 to *Erwinia herbicola* and the induction of insertion mutations. *Curr Microbiol*. 1981;6(6):377–81. <https://doi.org/10.1007/BF01567015>
39. Dale C, Young SA, Haydon DT, Welburn SC. The insect endosymbiont *Sodalis glossinidius* utilizes a type III secretion system for cell invasion. *Proc Natl Acad Sci USA*. 2001;98(4):1883–8. <https://doi.org/10.1073/pnas.98.4.1883>
40. Nierman WC, DeShazer D, Kim HS, Tettelin H, Nelson KE, Feldblyum T, Ulrich RL, Ronning CM, Brinkac LM, Daugherty SC, Davidsen TD. Structural flexibility in the *Burkholderia mallei* genome. *Proc Natl Acad Sci USA*. 2004;101(39):14246–51. <https://doi.org/10.1073/pnas.0403306101>
41. Chak KF, Safo MK, Ku WY, Hsieh SY, Yuan HS. The crystal structure of the immunity protein of colicin E7 suggests a possible colicin-interacting surface. *Proc Natl Acad Sci USA*. 1996;93(13):6437–42. <https://doi.org/10.1073/pnas.93.13.6437>
42. Notti RQ, Stebbins CE. The structure and function of type III secretion systems. *Microbiol Spec*. 2016;4(1):1–18. <https://doi.org/10.1128/microbiolspec.VMBF-0004-2015>

43. Matsuda S, Okada R, Tandhavanant S, Hiyoshi H, Gotoh K, Iida T, Kodama T. Export of a *Vibrio parahaemolyticus* toxin by the Sec and type III secretion machineries in tandem. *Nat Microbiol*. 2019;4(5):781–8. <https://doi.org/10.1038/s41564-019-0368-y>
44. Gophna U, Ron EZ, Graur D. Bacterial type III secretion systems are ancient and evolved by multiple horizontal-transfer events. *Gene*. 2003;312:151–63. [https://doi.org/10.1016/S0378-1119\(03\)00612-7](https://doi.org/10.1016/S0378-1119(03)00612-7)
45. Deng W, Marshall NC, Rowland JL, McCoy JM, Worrall LJ, Santos AS, Strynadka NC, Finlay BB. Assembly, structure, function and regulation of type III secretion systems. *Nat Rev Microbiol*. 2017;15(6):323–37. <https://doi.org/10.1038/nrmicro.2017.20>
46. Scheibner F, Hartmann N, Hausner J, Lorenz C, Hoffmeister AK, Büttner D. The type III secretion chaperone HpaB controls the translocation of effector and noneffector proteins from *Xanthomonas campestris* pv. *vesicatoria*. *Mol Plant Microbe Interact*. 2018;31(1):61–74. <https://doi.org/10.1128/MMBR.05017-11>
47. Büttner D. Protein export according to schedule: Architecture, assembly, and regulation of type III secretion systems from plant- and animal-pathogenic bacteria. *Microbiol Mol Biol Rev*. 2012;76(2):262–310. <https://doi.org/10.1094/MPMI-06-17-0138-R>
48. Diepold A, Kudryashev M, Delalez NJ, Berry RM, Armitage JP. Composition, formation, and regulation of the cytosolic c-ring, a dynamic component of the type III secretion injectisome. *PLoS Biol*. 2015;13(1):e1002039. <https://doi.org/10.1371/journal.pbio.1002039>
49. Hu B, Morado D. R., Margolin W., Rohde J. R., Arizmendi O., Picking W. L., Picking W. D., and Liu J. Visualization of the type III secretion sorting platform of *Shigella flexneri*. *Proc Natl Acad Sci USA*. 2015;112:1047–1052. <https://doi.org/10.1073/pnas.1411610112>
50. Hu B, Lara-Tejero M, Kong Q, Galán JE, Liu J. In situ molecular architecture of the *Salmonella* type III secretion machine. *Cell*. 2017;168(6):1065–74. <https://doi.org/10.1016/j.cell.2017.02.022>
51. Wattiau P, Bernier B, Deslee P, Michiels T, Cornelis GR. Individual chaperones required for Yop secretion by *Yersinia*. *Proc Natl Acad Sci USA*. 1994;91(22):10493–7. <https://doi.org/10.1073/pnas.91.22.10493>
52. Iriarte M, Cornelis GR. YopT, a new *Yersinia* Yop effector protein, affects the cytoskeleton of host cells. *Mol Microbiol*. 1998;29(3):915–29. <https://doi.org/10.1046/j.1365-2958.1998.00992.x>
53. Lloyd SA, Forsberg Å, Wolf-Watz H, Francis MS. Targeting exported substrates to the *Yersinia* TTSS: different functions for different signals? *Trends Microbiol*. 2001;9(8):367–71. [https://doi.org/10.1016/S0966-842X\(01\)02100-X](https://doi.org/10.1016/S0966-842X(01)02100-X)
54. Sory MP, Boland A, Lambermont I, Cornelis GR. Identification of the YopE and YopH domains required for secretion and internalization into the cytosol of macrophages, using the *cyaA* gene fusion approach. *Proc Natl Acad Sci USA*. 1995;92(26):11998–2002. <https://doi.org/10.1073/pnas.92.26.11998>
55. Cambronne ED, Cheng LW, Schneewind O. LcrQ/YscM1, regulators of the *Yersinia* yop virulon, are injected into host cells by a chaperone-dependent mechanism. *Mol Microbiol*. 2000;37(2):263–73.

<https://doi.org/10.1046/j.1365-2958.2000.01974.x>

56. Wattiau P, Cornells GR. SycE, a chaperone-like protein of *Yersinia enterocolitica* involved in the secretion of YopE. *Mol Microbiol.* 1993;8(1):123–31. <https://doi.org/10.1111/j.1365-2958.1993.tb01209.x>
57. Cheng LW, Anderson DM, Schneewind O. Two independent type III secretion mechanisms for YopE in *Yersinia enterocolitica*. *Mol Microbiol.* 1997;24(4):757–65. <https://doi.org/10.1046/j.1365-2958.1997.3831750.x>
58. Frithz-Lindsten E, Rosqvist R, Forsberg A. YerA mediates the secretion of the YopE cytotoxin of *Yersinia pseudotuberculosis*. *Contrib Microbiol Immunol.* 1995;13:225–9. PMID: 8833840
59. Khater L, Santos TM, Alegria MC, Docena C, da Silva AC, Ramos CH. In silico identification of potential chaperone genes that belong to type III and type IV secretion systems in *Xanthomonas axonopodis* pv *citri*. *Genet Mol Biol.* 2005;28(2):321–7. <https://doi.org/10.1590/S1415-47572005000200024>
60. Bennett JC, Hughes C. From flagellum assembly to virulence: the extended family of type III export chaperones. *Trends Microbiol.* 2000;8(5):202–4. [https://doi.org/10.1016/S0966-842X\(00\)01751-0](https://doi.org/10.1016/S0966-842X(00)01751-0)
61. Fraser GM, Bennett JC, Hughes C. Substrate-specific binding of hook-associated proteins by FlgN and FliT, putative chaperones for flagellum assembly. *Mol Microbiol.* 1999;32(3):569–80. <https://doi.org/10.1046/j.1365-2958.1999.01372.x>
62. Auvray F, Thomas J, Fraser GM, Hughes C. Flagellin polymerisation control by a cytosolic export chaperone. *J Mol Biol.* 2001;308(2):221–9. <https://doi.org/10.1006/jmbi.2001.4597>
63. Muskotál A, Király R, Sebestyén A, Gugolya Z, Végh BM, Vonderviszt F. Interaction of FliS flagellar chaperone with flagellin. *FEBS letters.* 2006;580(16):3916–20. <https://doi.org/10.1016/j.febslet.2006.06.024>
64. Ozin AJ, Claret L, Auvray F, Hughes C. The FliS chaperone selectively binds the disordered flagellin C-terminal D0 domain central to polymerisation. *FEMS Microbiol Lett.* 2003;219(2):219–24. [https://doi.org/10.1016/S0378-1097\(02\)01208-9](https://doi.org/10.1016/S0378-1097(02)01208-9)
65. Francis, M.S. Type III secretion chaperones: a molecular toolkit for all occasions. In *Handbook of Molecular Chaperones: Roles, Structures and Mechanisms*. P. Durante, and C. L (eds). Hauppauge, NY, USA: Nova Science Publishers. 2010; p. 79–147.
66. Bennett JC, Thomas J, Fraser GM, Hughes C. Substrate complexes and domain organization of the *Salmonella* flagellar export chaperones FlgN and FliT. *Mol Microbiol.* 2001;39(3):781–91. <https://doi.org/10.1046/j.1365-2958.2001.02268.x>
67. Parsot C, Hamiaux C, Page AL. The various and varying roles of specific chaperones in type III secretion systems. *Curr Opin Microbiol.* 2003;6(1):7–14. [https://doi.org/10.1016/S1369-5274\(02\)00002-4](https://doi.org/10.1016/S1369-5274(02)00002-4)
68. Bolivar F, Rodriguez RL, Greene PJ, Betlach MC, Heynker HL, Boyer HW, Crosa JH, Falkow S. Construction and characterization of new cloning vehicles II: a multipurpose cloning system. 1977. *Biotechnology (Reading, Mass.).* 1992;24:153 – 71. PMID: 1422009

69. Sambrook J, Fritsch EF, Maniatis T. Molecular cloning: a laboratory manual. Cold Spring Harbor: Cold spring harbor laboratory press; 1989.
70. Liu YG, Whittier RF. Thermal asymmetric interlaced PCR: automatable amplification and sequencing of insert end fragments from P1 and YAC clones for chromosome walking. *Genomics*. 1995;25(3):674–81. [https://doi.org/10.1016/0888-7543\(95\)80010-J](https://doi.org/10.1016/0888-7543(95)80010-J)
71. Hinton JC, Perombelon MC, Salmond GP. Efficient transformation of *Erwinia carotovora* subsp. *carotovora* and *E. carotovora* subsp. *atroseptica*. *J Bacteriol*. 1985;161(2):786-8.
72. Hanahan D. Studies on transformation of *Escherichia coli* with plasmids. *J Mol Biol*. 1983;166(4):557–80. [https://doi.org/10.1016/S0022-2836\(83\)80284-8](https://doi.org/10.1016/S0022-2836(83)80284-8).
73. Kyte J, Doolittle RF. A simple method for displaying the hydropathic character of a protein. *J Mol Biol*. 1982;157(1):105–32. [https://doi.org/10.1016/0022-2836\(82\)90515-0](https://doi.org/10.1016/0022-2836(82)90515-0)
74. Reusch RN, Hiske TW, Sadoff HL. Poly-beta-hydroxybutyrate membrane structure and its relationship to genetic transformability in *Escherichia coli*. *J Bacteriol*. 1986;168(2):553–62. <https://doi.org/10.1128/jb.168.2.553-562.1986>
75. Chang AC, Cohen SN. Construction and characterization of amplifiable multicopy DNA cloning vehicles derived from the P15A cryptic miniplasmid. *J Bacteriol*. 1978;134(3):1141–56.

Figures

Figure 1

Bacteriocin activity of *P. carotovorum* subsp. *carotovorum*. The bacteriocin production of the test strains was examined by the double-layer method (68). Numbered strains: 1, *Escherichia coli* pJB4JI (containing Tn5)/1830; 2, H-rif-8-6 (wild type, parent); 3, TH22-6 (the transposon Tn5 insertional mutant). The indicator organism used in the agar overlay was *Pcc* strain Ea1068. The result showed that TH22-6 strain loses the low-molecular-weight bacteriocin activity to Ea1068, but the high-molecular-weight bacteriocin kept its activity as manifested in the bacteriocin activity test.

AAGCTTTCTCT	GGATGTTGAT	CTGTGGTTGC	GGGAGATGAA	TGCACAACCTG	CCGGGGATTTC	CTTGGGTGCA	AGTGCCGATT	GACTATTTGG
TTCCGATGGCT	AGATACCATG	CAGTTAGCGT	TTATGTTAGA	AGAAAATACA	GTTTGGTATG	TACGCCATTT	AACCTTGCCG	ATGCTTCCGC
TGCCAGAGAA	ATTACTCTCT	TTGCCTACCC	AACCCTGCCC	GCTATTGTGT	CTTACTTGGC	CAAGTGGGTA	TGCAGAAGGG	GATATTAAGT
CGTTATGGGT	GAAAGCTCTC	CCTTTTACAT	TGCAATATGT	GCTAGGCCAT	AGCCAGTTAT	CGCTCTCTGT	GTAAATGGGG	GCTACCGCTG
GGGATTTTATT	ACTGATTAAG	CATTACTCCC	CTGTGTTAAC	TATCGGGGGG	AAAAGAGCAT	TTCGTTTTAG	TTACCAGAGC	CAAGAGGTTA
TTGTGGAAGA	ACAATTGAAT	GAATTTTCGG	AAGATTATTG	TGATGAAGAA	GAAGTGTGTG	TTGACTGGTC	AAATATTCCA	GTAAATATCG
AATTTGTTCT	TGATAGTCAT	AGCATGGTAT	TGGCCGATTT	AGAAAATATA	AGAGCTGGTA	GCTCGCTGCC	GCTCAGTGGT	GATGCTGAGA
AAAAGGTGAA	AATTTATTTA	AATAGAAAAT	ATTTTGCTCA	TGGAGAATTA	ATTGCGCTGG	AAAATGGTGC	GCTAGCCGTT	GAGATTTATC
AACTTGGTAA	CTTTTCAGGT	GGCGAAAACG	GTTGCTCAGA	TGTTTAAACA	TGATATTTCT	TTAATCGCAT	TATTATCTTT	TTTTACACTG
CTACCCCTTTC	TTATTGCAGG	AGGAACCTGC	TTCATTAAAT	TTTCCATTGT	ATTAGTTATG	GTGCGTAACG	CACTTGGTAT	TCAACAAGTG
CCTTCTACGT	TAACCTTAAA	TGGCGTAGCA	CTTATCCTAA	CGGCATTTGT	TATGATGCCA	ATTTGTCAGC	GTGTTTCTAA	TTATATCGAA
ACTAATCATA	TTGATTTTAA	TGATAGCCGC	TCAGTTACAC	AACTGGTCTG	TCAAGGGTTG	GGGAGCTATC	GGGAGTATTT	GGTGGCTTAT
TCTGATCAGG	ATCTGATTCG	TTTTTTCAAT	AAAGTACAAC	AGAGAAATGA	AGCGCTTCCT	GAACCAGAGA	TTGATGATCA	CGAATTAAGT
GCTTTACCGA	TCATGACGCT	AATGCCTGCC	TACGCATTAA	GTGAAATACA	AAGTGCCTTT	AAAATTGCTT	TTTATCTCTA	TGTTCCCTTT
GTGGTTATTG	ACATGGTGGT	GTCTAGTATA	TTATTGGCTC	TGGGAATGAT	GATGATGAGC	CCAGTGACAA	TATCAATTCC	TATAAAATTG
ATTCTCTTTG	TTGCAATGAA	TGGTTGGAGT	TATTTAAAGCT	AACTGCAGTT	CTTATCGCTT	ACTGACTTGA	TGATATCATG	AATAATATTT
TATTTATCGG	AAACTCGGCT	ATGCTTGTTG	AAACGTTGCC	ATTTGGTTTG	AAAATGTTGG	TTGCTACGGT	CGTTGGTATT	ATCGTCGGGC
TTTTTCAGAC	AGTAATGCAA	ATACAGGAGC	CAACCAATGT	TTTTTATATG	CGGTTTCGAT	CTGTGTTTGC	CAGTATTTTT	ATGTTAATTG
AGTGGTTTTTC	TGCTGAAATG	ATTAATTTCA				GAAACGATAA	AATTATGCTG	TTTGCTGCAT
							M L	F A A F
TCTACCTTAA	TTTTCAGCAC	AATATTCTCA	TTTTTGCTGT	GGCTTATGCG	CGGTTGGCGA	TAGTTTTCTA	CATGTTGCCA	ATATTAGGGG
Y L N	F Q H	N I L I	F A V	A Y A	R L A I	V F Y	M L P	I L G E
AGCGCGTATT	GTCACATCTG	ATAATTAAGA	ACGTGATAAT	TTCTCTTACC	ATTATTGGCT	TATGGCCTTG	CTTTGAGCAG	CTTGTTTCAC
R V L	S H L	I I K N	V I I	S L T	I I G L	W P C	F E Q	L V S P
CAGAACAAGG	CTTGCTAGTT	ATTTTGGTGA	AAGAGTGCCT	CGTTGGCTTA	ATTTTAGCTA	TGTGCTTATG	CTTACCTTTT	TGGGTTGTGA
E Q G	L L V	I L V K	E C V	V G L	I L A M	C L C	L P F	W V V I
TTGGTTTGGG	GGAAATCTT	GATAACCAAC	GTGGTGCAAC	GATTAGCGAC	AGTATCGATC	CAGTTAATGG	TGTGCAAAGC	TCTGTGTTGT
G L G	E I L	D N Q R	G A T	I S D	S I D P	V N G	V Q S	S V L S
CTGGCTTTCT	GAACTTTGCA	TTCGGCGCGA	TCTTCTTTCG	TCAGAATGGC	ATGCGTTTGT	TGATGGAAGC	GATGGCACAA	AGTTATCGTG
G F L	N F A	F G A I	F F A	Q N G	M R L L	M E A	M A Q	S Y R V
TTTTTCCGCG	TGGTAGTGAG	CTTAAAGACT	TAACTGGAT	GGAGGCGGGG	AATTTACTGA	TGGTTTTGGT	ACAGAGCAGT	ATTATGTTGG
F P R	G S E	L K D F	N W M	E A G	N L L M	V L V	Q S S	I M L A
CTGCACCAGT	GTTGATTATA	TTAATGATAG	CGGAAATTTT	ACTTGGTGTT	TTTGCGCGCT	ATTGTCCCCA	GCTCAATTCC	TTTTCACTGT
A P V	L I I	L M I A	E I L	L G V	F A R Y	C P Q	L N S	F S L S
CACTAACTAT	CAAAAGTTTC	ATCGCTTTTA	CTGTTTTTTT	GCTCTACGGC	TTTCAAGCGC	TATCAGAAAA	ACCCTTGAAG	ATGTTTTCTG
L T I	K S F	I A F T	V F L	L Y G	F Q A L	S E K	P L K	M F S V
TGACGATGTT	TCAGCAGTTT	TTATCCTGAT	AGGGAGCCGC	GACGATGGCT	GAAAAACAG	AAAAACCAAC	GAATAAACGT	CGTAAAGACT
T N F	Q H F	L S *						
CGGCAAAGAA	AGGGCAAACA	TTTAAAAGTA	AAGATCTTGT	CACCACAGCT	ATTTTGTTTG	TCGGGATCTA	TTTCTCTCC	CATGCTATCG
ATTTTGGTGC	TTTTTTTTCT	CTTTATTATA	TGGCGTTGTT	ACACAACGCT	GACATGAGTA	TAAATGATTT	TCTAATTGAA	CTCTTGCTTG
TTTTTTTTTACA	TATTACGCTG	CCCTTTATCG	TTGTTTGTAT	ATTAGTTGGG	GCAGGGATGA	CTTTATTGCA	AACCCGATTT	TCTATTGCCA
CAAAAGCAAT	AAAGCTGAAT	TTTAAAGCAT	TGAACCCCGT	AGAGGGTGTG	AAGAAGATTT	TTAATGTTAG	AACAATCAAA	GAGTTTGTGA
AGTCCATCTG	CTATCTTATT	GTGTTTATCT	GTAAGTGTAT	CAGCTTATTA	AATAATGATA	TCAAGCCAAG	CTTTAAACAT	CTATAATGGT
AATATCCCTC	AACTGGTCAG	CGTATGGGTT	TCGCTGGCGT	TTAAAGCAGT	GCTAGTTTTT	A		

Figure 2

Nucleotide sequence of the *sctT* gene with the deduced amino-acid sequence. The *sctT* contains 3031 base pairs.

	10	20	30	40	50	60
SctT-Pcc	-----MLFAAFYLN	FQHNI	IIFAVAYARLAI	VFYMLPILGERVLSHLI	IKNVII	ISLTII
YsaT-Yersinia	MDQQVVGLYVSLYST	FQTGLMTLALAWARI	APVFFFLPFLSSKLLNSGM	IKNCVAVELAL		
YsaT-Sodalis	-----	MIIMLAYVRVGIVFL	MLPLLGGRI	LSHLIVKNM	VIFITLI	
	70	80	90	100	110	120
SctT-Pcc	GLWPCFEQ--LVSPEQGLLV	ILVKECVVGLILAMCL	CLPFWV	VIGLGE	ILDNQR	GATISD
YsaT-Yersinia	GMWPFESTIYELNWEETSL	GEMILYELAIGLVLA	FILSLPFLIANII	GELIDNQR	GATISD	
YsaT-Sodalis	GIWPDIRT--IPNIEQGWLV	LIARECVTGLMLAMIL	SLPFWIASAI	GEVIDNQR	GATISD	
	130	140	150	160	170	180
SctT-Pcc	SIDPVNGVQSSVLSGFLN	FAFGAAIFFAQNGMRL	LMEAMAQSYRV	FPRGSELKDF	NWMEA	
YsaT-Yersinia	TIDPANGVESSEMSVLV	SHIVV-MIFLTQGGMYQ	LAQTFAESYRLL	PFAAGFSHF	DSLPL	
YsaT-Sodalis	SIDPVNGLQSTTMS	SFLSFAFG-AIFMAQGM	ILLMNALVQSYRV	PSGSSLLGAQ	WAVA	
	190	200	210	220	230	240
SctT-Pcc	GNLLMVLVQSSIMLAAPV	LIILMIAEILLGVFARY	CPQLNSFSLSLTI	KSFIAFTV	FLLY	
YsaT-Yersinia	GRWLNQLVVQAI	VLAAPILVTLFITEVAL	GLYSRFCPQLNAF	SLSLTIKSAIA	FMVFLY	
YsaT-Sodalis	GKLVHLLLENSVILA	APTTLITLMLSEMTL	GVFSRYCPQLNPF	SLSLKSFI	AFTIFMFY	
	250	260	270	280	290	300
SctT-Pcc	GFQAALSEKPLKMF	SVTMFQHFLS-----				
YsaT-Yersinia	-FQNEVPTILINMIS	LSPLHDFGVAPQ-				
YsaT-Sodalis	GF-GAMSEPLMLL	FTLTPFYQFVLP	AVGN			

Figure 3

Alignment of the predicted amino acid sequence of the Pcc sctT gene product with the amino acid sequences of YsaT protein in Sodalis and Yersinia. Dashes are inserted to optimize the alignment.

A

B

Figure 4

(A) Bacteriocin Assay for complementation of *sctT* and *flhA*: 1, H-rif-8-6 (wild type); 2, TH22-6 (*sctT*-KO); 3, *flhA*-KO; 4, *sctT*/*sctT*-KO; 5, *flhA*/*flhA*-KO. (B) Transcription analysis of the *sctT*, *flhA*, *caroS1K* and *caroS3K*. (B) Bacteriocin activity assay for the deletion of T3SS-related genes.

A

B

Figure 5

(A) Bacteriocin assay and deletion of various Type-IIIa and Type-IIIb SS related genes. 1, Serratia (Marker); 2, H-rif-8-6 (wild type); 3, *flhB*-KO; 4, *flgH*-KO; 5, *flgF*-KO; 6, *fliR*-KO; 7, *fliE*-KO; 8, *flgG*-KO; 9, *fliG*-KO; 10, *sctD*-KO; 11, *sctQ*-KO; 12, *sctV*-KO; and 13, *sctU*-KO. (B) Transcription analysis of 1, *flhB*; 2, *flgH*; 3, *flgF*; 4, *fliR*; 5, *fliE*; 6, *flgG*; 7, *fliG*; 8, *sctD*; 9, *sctQ*; 10, *sctV*; and 11, *sctU*.

Figure 6

The hypothesized flagellar T3bSS with the injectisome T3SS subunits as chaperones.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryFigure1.docx](#)
- [SupplementaryFigure2.docx](#)
- [SupplementaryFigure3.docx](#)
- [SupplementaryFigure4.docx](#)