Circulating SARS-CoV-2 Variants in Italy, October 2020-March 2021

Alessia Lai (✉ alessia.lai@unimi.it)

University of Milan: Università degli Studi di Milano <https://orcid.org/0000-0002-3174-5721>

Annalisa Bergna

University of Milan: Università degli Studi di Milano

Stefano Menzo

Università Politecnica delle Marche: Università Politecnica delle Marche

Gianguglielmo Zehender

University of Milan: Università degli Studi di Milano

Sara Caucci

Università Politecnica delle Marche: Università Politecnica delle Marche

Valeria Ghisetti

University of Turin: Università degli Studi di Torino

Francesca Rizzo

University of Salerno: Università degli Studi di Salerno

Fabrizio Maggi

University of Insubria: Università degli Studi dell'Insubria

Francesco Cerutti

ASL Città di Torino: ASL Città di Torino

Giorgio Giurato

University of Salerno: Università degli Studi di Salerno

Alessandro Weisz

University of Salerno: Università degli Studi di Salerno

Chiara Turchi

Università Politecnica delle Marche: Università Politecnica delle Marche

Bianca Bruzzone

Hospital San Martino: Ospedale Policlinico San Martino

Francesca Ceccherini Silberstein

University of Rome Tor Vergata: Università degli Studi di Roma Tor Vergata

Nicola Clementi

Università Vita e Salute San Raffaele: Università Vita Salute San Raffaele

Annapaola Callegaro

ASST Papa Giovanni XXIII: Aziende Socio Sanitarie Territoriale Papa Giovanni XXIII

Fabio Sagradi

ASST Cremona: Azienda Socio Sanitaria Territoriale di Cremona

Daniela Francisci

University of Perugia: Università degli Studi di Perugia

Emmanuele Venanzi Rullo

University of Messina: Università degli Studi di Messina

Ilaria Vicenti

University of Siena: Università degli Studi di Siena

Massimo Clementi

Università Vita e Salute San Raffaele: Università Vita Salute San Raffaele

Massimo Galli

University of Milan: Università degli Studi di Milano

Short report

Keywords: SARS-CoV-2 virus, complete genome sequencing, COVID-19 RT-PCR testing, Spike protein, viral variants

Posted Date: June 10th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-591143/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Virology Journal on August 14th, 2021. See the published version at <https://doi.org/10.1186/s12985-021-01638-5>.

Abstract

A growing number of emerging SARS-CoV-2 variants is being identified worldwide, potentially impacting the effectiveness of current vaccines. We report the data obtained in several Italian regions involved in the SARS-CoV-2 variant monitoring from the beginning of the epidemic and spanning the period from October 2020 to March 2021.

Background

Starting from April 2020 when the Cluster 5 [1] variant was first described, multiple SARS-CoV-2 (Severe acute respiratory syndrome coronavirus 2) variants have emerged in different parts of the world. The B.1.1.7 lineage was first identified in the UK from a sample obtained in late September 2020 [2, 3], B.1.351 was identified in October 2020 in South Africa [4], and P.1 was identified in Brazil in December 2020 [5]. These variant strains show multiple changes (deletions and substitutions) in the spike protein (9 in B.1.1.7, 10 in B.1.351, and 12 in P.1) compared with the reference genome Wuhan-Hu-1 sequence (EPI_ISL_406800), some of which affect the receptor-binding domain (RBD) region. The major issue with these variants is their potential to outcompete and rapidly replace formerly prevalent lineages, first in the areas where they likely emerged and subsequently spread in many other countries [6–8]. Increased SARS-CoV-2 diversity raises the concern of escape from pre-existing immunity, elicited either by previous infection or by vaccination.

Main Text

Here, we report data from several Italian centers located in Campania, Lazio, Lombardy, Liguria, Marche, Piedmont, Tuscany, Sicily and Umbria, involved in the SARS-CoV-2 variant monitoring from the beginning of the epidemic and spanning the period from October 2020 to March 2021.

Globally, we analysed data from 3,744 samples obtained by different techniques: RT-PCR variant screening assays (n = 2,095), spike Sanger or next generation sequencing (n = 649) and whole genome sequencing (WGS; n = 1,000). Table 1 shows the data for each month.

Table 1
Data stratified according to months and methodologies.

Month	WGS* (n = 1,000)	Spike (n = 649)	Real Time (n = 2,095)	Total (n = 3,744)
October 2020	137	20	1	158
November 2020	168	31	0	199
December 2020	139	34	0	173
January 2021	291	86	388	765
February 2021	193	214	827	1,234
March 2021	72	264	879	1,215
*WGS. Whole genome sequencing.				

The number of samples considered for variant monitoring increased from 4.2% (158/3744) in October to 32% (1198/3744) in March, due to the increasing interest in viral variants at global level. In the study period, the B.1.1.7 variant significantly increased, growing from 3.5% (6/173) in December, to 88.0% (1054/1198) in March ($p < .001$). Particularly, from the second half of December this lineage was already present in Liguria (n = 1 on the 18th), Campania (n = 3, on the 28th) and Marche (n = 2, on the 31st), with only one known case related to return from UK. The most important increase in its prevalence was observed from the second week of January (20/126, 16.0%) to the end of the same month (35/68, 51.5%) reaching 73.7% (171/232) at the end of February. In parallel, the prevalence of previously circulating variants significantly decreased from 99.4% (157/158) to 9.5% (114/3744). The P.1 lineage was first detected in the second half of January in Campania (n = 1), Lombardy (n = 1) and Umbria (n = 2), subsequently setting at around 2% in February (33/1226) and March (24/1198). Only few B.1.351 and B.1.525 cases were identified, never reaching 1% of the total samples analysed, being firstly detected in Liguria at the end of January and in Campania in mid-February, respectively. The Scottish variant (B.1.258) was firstly observed in Campania, Marche and Piedmont in mid-October, 4 cases were reported in November in Campania (n = 2) and Piedmont (n = 2), thereafter it was only reported in Campania until February reaching a proportion of 18.5% (17/92) of total cases, including its descendant lineages (B.1.258.3, B.1.258.14). In mid-March, two cases of the New York lineage B.1.526 were identified in Marche. Figure 1 shows the main viral variants observed overtime. Concerning other variants, the main circulating lineage was B.1.177 that represented, together with its descendents (B.1.177.4, B.1.177.8, B.1.177.14, B.1.177.15, B.1.177.52, B.1.177.53, B.1.177.75), more than half of cases (50.4%, 54.3%, 66.1%, 71.4% and 59.8% from October to February). Despite limited data regarding lineage assignment in March (n = 28), we observed a 27% prevalence of B.1 and B.1.177 lineages (Fig. 1, Panel B). Clade assignment was available for a larger number of strains compared to lineage because these data was

obtained also by spike sequencing highlighting the clear predominance of 20E(EU1) ranging from 43.2% (68/157) to 67.7% (189/279). Some noteworthy mutations were observed in different lineages: N439K in one lineage A and B.1, E484K in three B.1.177 and A222V + E484K + N501Y in clade 20E.

Conclusion

This study provides insights into the rapid change in the prevalence of SARS-CoV-2 variants over six months in Italy. Interestingly, we observed the first entry of B.1.1.7 in Lombardy in late December but in line with the identification of first retrospective data in UK on 20 September [9–11]. In addition to known variants of concern, we documented other minor variants that could be early warnings of upcoming changes. Continuous monitoring of all variants is mandatory to comprehensively investigate and keep track of virus evolution, particularly along with expanding vaccination still based on original strains that have been largely substituted by novel variants.

Abbreviations

SARS-CoV-2: Severe Acute Respiratory Syndrome Coronavirus 2.

RBD: Receptor-binding domain.

RT-PCR: Real time polymerase chain reaction.

WGS: Whole genome sequencing.

Declarations

Ethics approval

The study was approved by the Ethical Committee of the Sacco Hospital (Prot N. 0012266).

Consent for publication

Not applicable.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This research was supported in part by Regione Campania (POR Campania FESR 2014/2020 grants: “Monitoring the spread and genomic variability of the Covid 19 virus in Campania using NGS technology”, CUP B14I20001980006 and azione 1.5 “GENOMAeSALUTE”, CUP: B41C17000080007) and by Ministero della Salute COVID-2020-12371817. The funding sources had no role in the study design, data collection analyses, interpretation or writing of the report.

Author Contributions

Conceptualization, MC and MG; formal analysis, AL, AB, SM, GZ, SC, VG, FR, FM, FC, GG, AW, CT, BB, FCS, NC, AC, FS, DF, EVR., IV.; investigation, AL, AB; resources, SM, GZ, VG, FM, AW, BB, FSC, MC, MG; writing—original draft preparation, AL, AB; writing—review and editing, AL, AB; visualization and supervision, all authors. All authors have read and agreed to the published version of the manuscript.

Acknowledgment

Members of the SCIRE collaborative group: Claudia Balotta, Maria Gori, Patrizia Bagnarelli, Andreina Baj, Federica Novazzi, Andrea Orsi, Patrizia Caligiuri, Simona Boccotti, Maria Concetta Bellocchi, Loredana Sarmati, Massimo Andreoni, Nicasio Mancini, Elena Criscuolo, Rosa Gallitelli, Sophie Testa, Filippo Dragoni, Maurizio Zazzi.

We thank the members of the NGS and bioinformatics core facilities of the Laboratory of Molecular Medicine and Genomics for WGS data generation and analysis and Gianluigi Franci, Massimiliano Galdiero, Pasquale Pagliano, Giuseppe Portella and Teresa Rocco for their scientific contribution.

References

1. Oude Munnink, B.B.; Sikkema, R.S.; Nieuwenhuijse, D.F.; Molenaar, R.J.; Munger, E.; Molenkamp, R.; et al. Transmission of SARS-CoV-2 on mink farms between humans and mink and back to humans. *Science*. 2021;371, 172-177.
2. Davies, N.G.; Abbott, S.; Barnard, R.C.; Jarvis, C.I.; Kucharski, A.J.; Munday, J.D.; et al. Estimated transmissibility and impact of SARS-CoV-2 lineage B.1.1.7 in England. *Science*. 2021;372, eabg3055.
3. Volz, E.; Mishra, S.; Chand, M.; Barrett, J.C.; Johnson, R.; Geidelberg, L.; et al. Assessing transmissibility of SARS-CoV-2 lineage B.1.1.7 in England. *Nature*. 2021.
4. Tegally, H.; Wilkinson, E.; Giovanetti, M.; Iranzadeh, A.; Fonseca, V.; Giandhari, J.; et al. Detection of a SARS-CoV-2 variant of concern in South Africa. *Nature*. 2021;592, 438-443.
5. Faria, N.R.; Mellan, T.A.; Whittaker, C.; Claro, I.M.; Candido, D.D.S.; Mishra, S.; et al. Genomics and epidemiology of the P.1 SARS-CoV-2 lineage in Manaus; Brazil. *Science*. 2021;eabh2644.
6. Funk, T.; Pharris, A.; Spiteri, G.; Bundle, N.; Melidou, A.; Carr, M.; et al. Characteristics of SARS-CoV-2 variants of concern B.1.1.7; B.1.351 or P.1: data from seven EU/EEA countries; weeks 38/2020 to 10/2021. *Euro Surveill*. 2021;26, 2100348.

7. European Centre for Disease Prevention and Control. SARS-CoV-2 - increased circulation of variants of concern and vaccine rollout in the EU/EEA; 14th update – 15 February 2021. ECDC: Stockholm; 2021.
8. Galloway, S.E.; Paul, P.; MacCannell, D.R.; Johansson, M.A.; Brooks, J.T.; MacNeil, A.; et al. Emergence of SARS-CoV-2 B.1.1.7 Lineage - United States; December 29; 2020-January 12; 2021. MMWR Morb Mortal Wkly Rep. 2021;70, 95-99.
9. Di Giallonardo, F.; Puglia, I.; Curini, V.; Cammà, C.; Mangone, I.; Calistri, P.; et al. Emergence and Spread of SARS-CoV-2 Lineages B.1.1.7 and P.1 in Italy. Viruses. 2021;13, 794.
10. Lo Menzo, S, Marinello, S.; Biasin, M.; Terregino, C.; Franchin, E.; Crisanti, A.; Cattelan, A. The first familial cluster of the B.1.1.7 variant of SARS-CoV-2 in the northeast of Italy. Infection. 2021; 1–5.
11. Rambaut, A.; Loman, N.; Pybus, O.; Barclay, W.; Barrett, J.; Carabelli, A.; et al. Preliminary genomic characterisation of an emergent SARS-CoV-2 lineage in the UK defined by a novel set of spike mutations. ARTIC Network. 2020.

Figures

A

B

Figure 1

SARS-CoV-2 viral variants observed overtime in Italy. Panel A highlights the variants of concern; panel B shows all other circulating lineages.