

Impact of preoperative fecal short chain fatty acids on postoperative infectious complications in esophageal cancer patients

Masaaki Motoori (✉ mmotoori@gh.opho.jp)

Osaka General Medical Center <https://orcid.org/0000-0002-8529-2844>

Koji Tanaka

Osaka University

Keijiro Sugimura

Osaka International Cancer Institute

Hiroshi Miyata

Osaka International Cancer Institute

Takuro Saito

Osaka University

Yasuhiro Miyazaki

Osaka General Medical Center

Kazumasa Fujitani

Osaka General Medical Center

Yukiko Kado

Yakult Central Institute

Takashi Asahara

Yakult Central Institute

Masahiko Yano

Osaka International Cancer Institute

Research article

Keywords: esophageal cancer, esophagectomy, postoperative complications, intestinal environment, organic acids

Posted Date: September 27th, 2019

DOI: <https://doi.org/10.21203/rs.2.15259/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Gastroenterology on March 16th, 2020. See the published version at <https://doi.org/10.1186/s12876-020-01217-y>.

Abstract

Background: The intestinal epithelial barrier allows absorption of dietary nutrients and prevents passage of pathogens and toxins into the body. Severe insults have a negative impact on the intestinal environment, which may decrease intestinal barrier function and cause bacterial translocation. Bacterial translocation, which can cause infectious complications, is the passage of microbes from the gastrointestinal tract across the mucosal barrier to extraintestinal sites. The aim of this study was to investigate the correlation between concentrations of preoperative fecal organic acids and the occurrence of postoperative infectious complications in patients with esophageal cancer.

Methods: Fifty-five patients with esophageal cancer who underwent esophagectomy were enrolled in this study. All patients were administered perioperative synbiotics. Perioperative clinical characteristics and concentrations of preoperative fecal organic acids were compared between patients with or without postoperative infectious complications.

Results: Postoperative infectious complications occurred in 10 patients. In patients with complications, the concentrations of acetic acid and propionic acid were significantly lower than in patients without complications ($p=0.044$ and 0.032 , respectively). The concentration of butyric acid was nonsignificantly lower, while the concentration of lactic acid was nonsignificantly higher in patients with complications. The calculated gap between the concentrations of fecal acetic acid plus propionic acid plus butyric acid minus lactic acid was significantly lower in patients with complications. Multivariate analysis revealed that a low gap between acetic acid plus propionic acid plus butyric acid minus lactic acid was an independent risk factor for postoperative infectious complications ($p=0.027$).

Conclusions : Preoperative fecal concentrations of organic acids had a clinically important impact on the occurrence of postoperative infectious complications in patients with esophageal cancer. To reduce postoperative infectious complications, it may be useful to modulate the intestinal environment and maintain concentrations of fecal organic acids before surgery.

Introduction

Subtotal esophagectomy for esophageal cancer is one of the most invasive gastrointestinal surgeries and is associated with high morbidity and mortality rates, though there have been recent advances in surgical techniques and perioperative management [1, 2]. The intestinal tract is the largest immune organ in the human body. The intestinal epithelial barrier allows absorption of dietary nutrients and prevents passage of pathogens and toxins into the body. Severe insults have a negative impact on the intestinal environment, which may decrease intestinal barrier function and cause bacterial translocation. We previously reported that esophagectomy in patients with esophageal cancer disturbed the intestinal microbiota and decreased the concentrations of organic acids, while the administration of perioperative synbiotics significantly shortened the duration of the postoperative systemic inflammatory response

syndrome and nonsignificantly reduced postoperative infectious complications by maintaining the intestinal microbiota and the concentrations of organic acids [3].

The fermentation of carbohydrates by the intestinal microbiota produces organic acids. Among them, short chain fatty acids (SCFAs) play a variety of important roles in maintaining intestinal mucosal barrier function and preventing bacterial translocation [4]. Bacterial translocation, which can cause infectious complications, is the passage of microbes or their products from the gastrointestinal tract across the mucosal barrier to extraintestinal sites, such as the mesenteric lymph nodes, liver, or bloodstream. Several investigators demonstrated that severe insults, such as major surgery and the severe systemic inflammatory response syndrome, disrupt the intestinal environment, in particular disturbing the intestinal microbiota and decreasing the concentrations of organic acids [3, 5, 6]. However, there is only one study concerning the association between the preoperative intestinal environment and postoperative infectious complications. Yokoyama et al. reported that low preoperative fecal concentrations of SCFAs were significantly associated with the occurrence of postoperative infectious complications in patients undergoing major hepatectomy with extrahepatic bile duct resection [7].

The aim of this study was to investigate the correlation between the preoperative fecal concentrations of organic acids and the occurrence of postoperative infectious complications in patients with esophageal cancer undergoing esophagectomy.

Materials And Methods

Patients

This study included 55 patients with thoracic esophageal cancer who underwent transthoracic esophagectomy at Osaka International Cancer Institute. These patients were included in our previous studies [3, 8]. All patients received gastrointestinal fiberoptic endoscopy and computed tomography (CT) scans of the neck, chest, and upper abdomen for tumor staging according to the TNM classification (7th edition) [9]. Patients underwent transthoracic subtotal esophagectomy with two- or three-field lymphadenectomy and reconstruction using the stomach, jejunum, or colon. An enteral feeding tube was inserted through the stomach or jejunum during the operation. Synbiotics was administered from 5 or more days before surgery to 21 days after surgery to all patients. The administered probiotics were Yakult BL Seichoyaku (Yakult Honsha, Tokyo), containing 1×10^8 live *Bifidobacterium breve* strain Yakult and 1×10^8 live *Lactobacillus casei* strain Shirota/g, and the prebiotics were galacto-oligosaccharides (Oligomate S-HP; Yakult Honsha). Before surgery, Yakult BL Seichoyaku (3 g/day) and galacto-oligosaccharides (15 g/day) were administered orally, and after surgery, Yakult BL Seichoyaku (2 g/day) and galacto-oligosaccharides (10 g/day) were administered through an enteral feeding tube or orally. Enteral feeding using a polymeric formula was initiated on postoperative day 1. Postoperative complications were assessed according to the Clavien-Dindo classification [10]. In this study, postoperative infectious complications were defined as those of grade II or higher.

Fecal bacteriologic and organic acid analysis

Fecal samples were collected 1 day before surgery. For measurement of the number of microbiota, fecal samples were placed into a tube containing RNAlater (Ambion, Austin, TX) and stored at -20°C until use. Total RNA was extracted from feces and the gut microbiota composition was examined using 16S or 23S ribosomal RNA-targeted reverse transcription-quantitative PCR using the Yakult Intestinal Flora-SCAN as described previously [11, 12]. For measurement of the concentrations of organic acids, fecal samples were placed into a vacant tube and stored at -20°C until use. Feces were homogenized in perchloric acid and the concentrations of organic acids were measured using a Waters high-performance liquid chromatography system (Waters 432 Conductive Detector; Waters, Milford, MA) and a Shodex RSpak KC-811 column (Showa Denko, Tokyo, Japan) [13].

Statistical analysis

Statistical analyses were performed using JMP13 (SAS Institute) software. The relationships between postoperative infectious complications and clinical factors, numbers of fecal bacteria, or concentrations of fecal organic acids were evaluated by the chi-square test or the Mann-Whitney U test. Multivariate analysis of infectious complications was performed using multiple logistic regression. Differences were significant at $p < 0.05$.

The study protocol was approved by the Human Ethics Review Committees of Osaka International Cancer Institute and registered in the University Hospital Medical Information Network (<http://www.umin.ac.jp>; registration number ID UMIN000004704 and UMIN000006875). A signed consent form was obtained from each patient.

Results

Patient characteristics and surgical outcome

The clinical characteristics of the patients are presented in Table 1. The clinical disease stages were Stage I in 16 patients, Stage II in 15 patients, Stage III in 20 patients, and Stage IV (due to supraclavicular lymph node metastasis) in four patients. Neoadjuvant chemotherapy or chemoradiotherapy were performed in 36 patients.

Postoperative infectious complications occurred in 10 patients (18%), including pulmonary complications in seven, anastomotic leakage in one, pyothorax in one, and wound infection in two. There was no perioperative mortality.

Table 2 shows the association between preoperative and intraoperative factors and postoperative infectious complications. Postoperative infectious complications were more common in the elderly, but

not significantly so ($p = 0.073$).

Table 3 shows the association between the number of representative fecal obligate anaerobes, administered probiotics, or fecal organic acid concentrations, and postoperative infectious complications. There was no significant difference in the number of representative fecal obligate anaerobes between patients with or without postoperative complications. The number of *Bifidobacterium breve* strain Yakult was nonsignificantly lower in the patients with complication than in those without. By contrast, the concentrations of acetic acid and propionic acid were significantly lower in the patients with complication ($p = 0.044$ and 0.032 , respectively). The concentration of butyric acid was nonsignificantly lower and the concentration of lactic acid was nonsignificantly higher in the patients with complication. The concentration gap between the acetic acid plus propionic acid plus butyric acid minus lactic acid (APB-L gap), was significantly lower in the patients with complication than in those without. ($46.3 \pm 25.1 \mu\text{mol/g}$ of feces vs. $75.2 \pm 34.3 \mu\text{mol/g}$ of feces, respectively, $p = 0.026$), as shown in Figure 1. There was no significant correlation between fecal organic acid concentrations and the number of representative fecal obligate anaerobes (data not shown).

Table 4 shows the multivariate analysis for postoperative infectious complications. The APB-L gap cutoff value was determined using receiver operating characteristics curve analysis. Among factors with $p < 0.2$ in the univariate analysis, the APB-L gap was the only factor significantly associated with postoperative infectious complications ($p = 0.027$).

Discussion

Patients with esophageal cancer who undergo esophagectomy have various risk factors that can negatively impact the intestinal environment, such as malnutrition due to cancer stenosis, preoperative therapy, surgical stress, use of antibiotics, postoperative fasting, and parenteral nutrition. If the preoperative intestinal environment is maintained, postoperative infectious complications may be reduced. This study revealed that concentrations of preoperative fecal SCFAs were significantly associated with the occurrence of postoperative infectious complications. Acetic acid, propionic acid, and butyric acid make up the majority of SCFAs (>95%) [14]. In this study, the patients with complication demonstrated significantly lower concentrations of acetic acid and propionic acid and nonsignificantly lower concentrations of butyric acid than those without complications. The concentration of lactic acid was nonsignificantly higher in the patients with complication, and the APB-L gap was a significant independent risk factor for postoperative infectious complications. These results are consistent with those of the previous study by Yokoyama et al. [7]. In their study of patients undergoing major hepatectomy with extrahepatic bile duct resection, the preoperative fecal concentrations of acetic acid and butyric acid were significantly lower and the concentration of lactic acid was nonsignificantly lower in patients who developed postoperative infectious complications, and the acetic acid plus butyric acid minus lactic acid gap was an independent risk factor for postoperative infectious complications. Lactic acid is digested by obligate anaerobes to produce SCFAs [15]. Yokoyama et al. speculated that a low

acetic acid plus butyric acid minus lactic acid gap may reflect an impaired intestinal environment and thus lead to postoperative infectious complications [7].

SCFAs play various roles in maintaining the intestinal environment, and contribute to the prevention of bacterial translocation. SCFAs stimulate the proliferation and differentiation of intestinal epithelial cells and the secretion of mucus by goblet cells [16, 17]. They also maintain the acidity of the intestine and suppress the growth of harmful bacteria. Acetic acid, which is the most common organic acid, has an antimicrobial effect on harmful bacteria [18] and promotes the defensive functions of host epithelial cells [19]. Furthermore, Hsieh et al. reported that acetic acid strengthened tight junctions in Caco-2 monolayers [20]. Ohata et al. reported that propionic acid and butyric acid induced tight junction permeability in Caco-2 monolayer cells [21]. Butyric acid is the major source of colonic epithelial cells, and induces the differentiation of regulatory T cells and ameliorates chronic intestinal inflammation in mice [22]. Thus, maintaining the concentration of SCFAs prior to iatrogenic gastrointestinal tract damage is useful for preventing bacterial translocation.

Organic acids are mainly produced in the large intestine by obligate anaerobes. In this study, there was no significant difference between the two groups in the number of representative fecal obligate anaerobes, as shown in Table 3. This result is inconsistent with that of the prior study mentioned above [7]. We previously reported that in patients with advanced esophageal cancer, neoadjuvant chemotherapy altered the intestinal microbiota and concentrations of organic acids [8]. In this study, about two-thirds of patients received preoperative chemotherapy or preoperative chemoradiotherapy. When we analyzed the correlation between the preoperative number of representative fecal obligate anaerobes or the fecal concentrations of organic acids and the incidence of postoperative infectious complications only in patients who received preoperative therapy, no significant difference was found in the number of representative fecal obligate anaerobes between the two groups, while the concentrations of acetic acid and propionic acid were significantly lower in the patients with complication than in those without complication ($p = 0.025$ and 0.022 , respectively, data not shown). The concentration of SCFAs is considered to reflect the function of the intestinal microbiota, which may be affected by preoperative therapy. Thus, the concentration of SCFAs but not the number of representative fecal obligate anaerobes was useful for predicting postoperative infectious complications in this study.

Acetic acid, which is the most common SCFA, is mainly generated by *Bifidobacterium* [19, 23]. In this study, the concentration of acetic acid was significantly lower in the patients with complication than in those without complication. This may be because while the number of *Bifidobacterium* did not differ between the two groups, the number of *Bifidobacterium breve* strain Yakult, which was the administered probiotics, was nonsignificantly higher in the complication (-) group. Since *Bifidobacterium breve* strain Yakult can selectively metabolize galacto-oligosaccharide, the production of acetic acid may be increased [24]. In this study, although all patients received synbiotics, including *Bifidobacterium breve* strain Yakult, for at least 5 days before surgery, the number of preoperative fecal *Bifidobacterium breve* strain Yakult was nonsignificantly lower in the patients with complication than in those without complication. It is considered that different patients had different responses to synbiotics. The mechanism of this variation

is unclear. Currently, we administer perioperative synbiotics to all patients with esophageal cancer scheduled to undergo esophagectomy. If a patient has low concentrations of SCFAs after receiving synbiotics, more careful attention should be paid during the perioperative period to prevent postoperative infectious complications.

Our study has several limitations. First, the number of enrolled patients was small. Second, about two-thirds of patients received preoperative chemotherapy or chemoradiotherapy, both of which alter the intestinal environment, while about one-third of patients received no preoperative therapy. In addition, the duration of synbiotic administration varied among individuals. This heterogeneity may have affected the results of our study. Third, there are no data concerning intestinal permeability, which correlates with the degree of bacterial translocation.

Conclusions

Preoperative fecal concentrations of acetic acid and propionic acid were significantly lower in patients who developed postoperative infectious complications than in those who did not. To reduce postoperative infectious complications after esophagectomy in patients with esophageal cancer, it may be useful to modulate the intestinal environment and maintain concentrations of fecal SCFAs.

Abbreviations

SCFAs; short chain fatty acids

APB-L gap; the concentration gap between the acetic acid plus propionic acid plus butyric acid minus lactic acid

Declarations

Ethics approval and consent to participate

The study protocol was approved by the Human Ethics Review Committees of Osaka International Cancer Institute and registered in the University Hospital Medical Information Network (<http://www.umin.ac.jp>; registration number ID UMIN000004704 and UMIN000006875). A signed consent form was obtained from each patient.

Consent for publication

Not applicable.

Availability of data and material

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests

Funding

This study was partly supported by the Osaka Foundation for the Prevention of Cancer and Cardiovascular Diseases. This funding was used for the collection.

Authors' contributions

Study concepts: MM, KT

Study design: MM, KS

Data acquisition: MM, KT, TS, YM, TA, YK

Data analysis and interpretation: MM, HM, TA

Manuscript preparation: MM

Manuscript editing: KF, MY

Manuscript review: MY

All authors have read and approved the manuscript.

Acknowledgments

We wish to express our sincere gratitude for the valuable assistance of Akira Takahashi, Satomi Atobe, Rie Date, and Takako Minami of the Yakult Central Institute in performing the fecal bacteriologic and organic acid analyses.

References

1. Takeuchi H, Miyata H, Gotoh M, et al. A risk model for esophagectomy using data of 5354 patients included in a Japanese nationwide web-based database. *Ann Surg.* 2014;260:259-266.

2. Merkow RP, Bilimoria KY, McCarter MD, et al. Short-term outcomes after esophagectomy at 164 American College of Surgeons National Surgical Quality Improvement Program hospitals: effect of operative approach and hospital-level variation. *Arch Surg*. 2012;147:1009-1016.
3. Tanaka K, Yano M, Motoori M, et al. Impact of perioperative administration of synbiotics in patients with esophageal cancer undergoing esophagectomy: a prospective randomized controlled trial. *Surgery* 2012;152: 832-842.
4. Ohata A, Usami M, Miyoshi M. Short-chain fatty acids alter tight junction permeability in intestinal monolayer cells via lipoxygenase activation. *Nutrition* 2005;21:838-847.
5. Kanazawa H, Nagino M, Kamiya S, et al. Synbiotics reduce postoperative infectious complications: a randomized controlled trial in biliary cancer patients undergoing hepatectomy. *Arch Surg*. 2005;390:104-113.
6. Shimizu K, Ogura H, Goto M, et al. Synbiotics decrease the incidence of septic complications in patients with severe SIRS: a preliminary report. *Dig Dis Sci*. 2009;54:1071-1078.
7. Yokoyama Y, Mizuno T, Sugawara G, et al. Profile of preoperative fecal organic acids closely predicts the incidence of postoperative infectious complications after major hepatectomy with extrahepatic bile duct resection: Importance of fecal acetic acid plus butyric acid minus lactic acid gap. *Surgery*. 2017;162:928-936.
8. Motoori M, Yano M, Miyata H, et al. Randomized study of the effect of synbiotics during neoadjuvant chemotherapy on adverse events in esophageal cancer patients. *Clin Nutr*. 2017;36:93-99.
9. Sobin LH, Wittekind CH. TNM classification of malignant tumors. 7th ed. New York: Wiley-Liss; 2009
10. Dindo D, Demartines N, Clavien PA. Classification of surgical complications: A New Proposal With Evaluation in a Cohort of 6336 Patients and Results of a Survey. *Ann Surg* 2004;240:205-213.
11. Matsuda K, Tsuji H, Asahara T, et al. Establishment of an analytical system for the human fecal microbiota, based on reverse transcription quantitative PCR targeting of multicopy rRNA molecules. *Appl Environ Microbiol*. 2009;75:1961-1969.
12. Fujimoto J, Tanigawa K, Kudo Y, et al. Identification and quantification of viable *Bifidobacterium breve* strain Yakult in human faeces by using strain-specific primers and propidium monoazide. *J Appl Microbiol*. 2011;110:209-217.
13. Kamiya S, Nagino M, Kanazawa H, et al. The value of bile replacement during external biliary drainage: an analysis of intestinal permeability, integrity, and microbiota. *Ann Surg*. 2004; 239:510-517.
14. Cook SI1, Sellin JH. Review article: short chain fatty acids in health and disease. *Aliment Pharmacol Ther*. 1998;12:499-507.
15. den Besten G1, van Eunen K, Groen AK, et al. The role of short-chain fatty acids in the interplay between diet, gut microbiota, and host energy metabolism. *J Lipid Res*. 2013;54:2325-2340.
16. Guarner F, Malagelada JR. Gut flora in health and disease. 2003;361:512-519.

17. Willemsen LE, Koetsier MA, van Deventer SJ, van Tol EA. Short chain fatty acids stimulate epithelial mucin 2 expression through differential effects on prostaglandin E₁ and E₂ production by intestinal myofibroblasts. *Gut*. 2003;52:1442-1447.
18. Hayakawa M, Asahara T, Ishitani T, Okamura A, Nomoto K, Gando S. Synbiotic therapy reduces the pathological gram-negative rods caused by an increased acetic acid concentration in the gut. *Dig Dis Sci*. 2012;57:2642-2649.
19. Fukuda S, Toh H, Hase K, et al. Identification and quantification of viable Bifidobacterium breve strain Yakult in human faeces by using strain-specific primers and propidium monoazide. *Nature*. 2011;469:543-547.
20. Hsieh CY, Osaka T, Moriyama E, Date Y, Kikuchi J, Tsuneda S. Strengthening of the intestinal epithelial tight junction by Bifidobacterium bifidum. *Physiol Rep*. 2015;3: e12327.
21. Ohata A, Usami M, Miyoshi M. Short-chain fatty acids alter tight junction permeability in intestinal monolayer cells via lipoxygenase activation. *Nutrition*. 2005;21:838-847.
22. Furusawa Y, Obata Y, Fukuda S, et al. Commensal microbe-derived butyrate induces the differentiation of colonic regulatory T cells. *Nature*. 2013;504:446-450.
23. Asahara T, Shimizu K, Nomoto K, Hamabata T, Ozawa A, Takeda Y. Probiotic bifidobacteria protect mice from lethal infection with Shiga toxin-producing Escherichia coli O157:H7. *Infect Immun*. 2004;72:2240-2247.
24. Shigehisa A, Sotoya H, Sato T, Hara T, Matsumoto H, Matsuki T. Characterization of a bifidobacterial system that utilizes galacto-oligosaccharides. *Microbiology*. 2015;61:1463-1470.

Tables

Table I. Patient characteristics

Preoperative variables		
Gender	male/female	46/9
Age	median (range)	64 (37-76)
Tumor location	upper/middle/lower	6/30/19
Preoperative therapy	no/yes	19/36
cT factor	T1/T2/T3/T4	12/15/26/2
cN factor	N0/N1/N2/N3	25/19/10/1
cM factor	M0/M1	51/4
cStage	I/II/III/IV	16/15/20/4
Intraoperative variables		
Field of lymph node dissection	two-field/three-field	15/40
Organ used in reconstruction	stomach/jejunum	54/1
Operation time (min)	median (range)	558 (411-755)
Blood loss (ml)	median (range)	945 (370-3525)

Table II. Correlation between preoperative or intraoperative factors and postoperative infectious complications

		Complication (-) (n=45)	Complication (+) (n=10)	p value
Gender	male/female	37/8	9/1	0.55
Age		61.4±8.7	66.8±4.9	0.073
Tumor location	upper, middle / lower	30/15	6/4	0.69
Preoperative therapy	no/yes	17/28	2/8	0.28
cStage	I-II/III-IV	19/26	5/5	0.65
BMI (kg/m ²)		21.8±2.9	21.0±1.6	0.39
Operation time (min)		546±78	589±78	0.15
Blood loss (ml)		1045±517	988±413	0.98

Table III. Correlation between fecal microbiota or organic acids and postoperative infectious complications

	Complication (-) (n=45)	Complication (+) (n=10)	p value
Fecal microbiota (log ₁₀ cells/g of feces)			
<i>Clostr. coccooides</i> group	9.46±0.56	8.65±2.35	0.77
<i>Clostr. leptam</i> subgroup	9.22±1.38	8.51±2.36	0.3
<i>Bacteroides fragilis</i> group	9.29±1.70	8.71±2.00	0.52
<i>Bifidobacterium</i>	9.80±1.12	9.79±0.67	0.35
<i>Atopobium</i> cluster	9.21±0.74	8.80±1.2	0.23
<i>Prevotella</i>	6.58±2.17	5.94±2.38	0.48
Total <i>Lactobacillus</i>	7.56±0.16	7.43±1.18	0.50
<i>Bifidobacterium breve</i> strain Yakult	8.03±1.23	6.76±2.32	0.11
<i>Lactobacillus casei</i> strain Shirota	6.77±2.17	5.93±2.38	0.27
Fecal organic acid (µmol/g of feces)			
Total organic acids	80.9±29.8	56.9±37.0	0.11
Acetic acid	56.0±25.8	36.7±19.6	0.044
Propionic acid	16.0±10.2	9.2±4.8	0.032
Butyric acid	4.5±3.9	2.6±2.2	0.19
Lactic acid	1.3±2.6	2.3±3.4	0.12
Valeric acid	0.88±0.59	0.74±0.28	0.39
Iso-valeric acid	1.34±0.92	0.90±0.32	0.11
Formic acid	0.97±1.5	1.2±1.6	0.57
Succinic acid	2.3±6.0	4.7±8.6	0.67

Table IV. Multivariate analysis of postoperative infectious complications

		Odds ratio (95% CI)	p value
Age	≥70 (vs. ≤69)	1.24 (0.28-5.80)	0.77
Operation time (min)	≥550 (vs. <550)	3.47 (0.79-18.8)	0.11
APB-L gap* (μmol/g of feces)	≤85 (vs. >85)	8.04 (1.24-160.34)	0.027

*: gap between the acetic acid plus propionic acid plus butyric acid minus lactic acid

Figures




Figure 1

APB-L gap in patients with or without postoperative infectious complications