

# Discovery of Genomic Regions Associated With Resistance to Late Wilt Disease Caused by *Harpophora Maydis* (Samra, Sabet and Hing) in Maize (*Zea mays* L.)

**N. C. Sunitha**

University of Agricultural Sciences

**E. Gangappa**

University of Agricultural Sciences

**R. P. Veeresh Gowda**

Bayer CropScience

**Ramesh S** (✉ [ramesh\\_uasb@rediffmail.com](mailto:ramesh_uasb@rediffmail.com))

University of Agricultural Sciences <https://orcid.org/0000-0001-9666-7974>

**Sunil Biradar**

Bayer CropScience AG

**Dharanendra Swamy**

Bayer CropScience AG

**H. B. Hemareddy**

Bayer CropScience AG

---

## Research Article

**Keywords:** Late wilt disease, mapping, QTL × QTL interaction, QTL × Location interaction

**Posted Date:** June 22nd, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-599138/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

---

**Version of Record:** A version of this preprint was published at Journal of Applied Genetics on November 29th, 2021. See the published version at <https://doi.org/10.1007/s13353-021-00672-x>.

## Abstract

Late wilt disease (LWD) caused by *Harpophora maydis* (Samra, Sabet and Hing) is emerging as major production constraint in maize across the world. As a prelude to develop maize hybrids resistance to LWD, genetic basis of resistance was investigated. Two  $F_{2:3}$  mapping populations (derived from CV156670 × 414-33 (P-1) and CV156670 × CV143587 (P-2)) were challenged with LWD at two locations (Kallinayakanahalli and Muppadihatta) during 2017 post-rainy season. Wider range of LWD scores were observed at both locations in both the populations. LWD response was influenced by significant Genotype × location interaction. Six and 56  $F_{2:3}$  progeny families showed resistance level better than resistant parent. 150 and 199 polymorphic SNP markers were used to genotype P-1 and P-2, respectively. Inclusive composite interval mapping was performed to detect significant QTL, QTL × QTL, QTL × Location interaction effects. Three major and four minor QTL controlling LWD resistance were detected on chromosome-1. Position and effect of the QTL varied with the location. Significant di-QTL interactions involving QTL (with significant and/or non-significant effects) located within and between all the ten chromosomes were detected. Five of the seven detected QTL in our study showed significant QTL × location interaction. Though two major QTL (*q-lw-1.5* and *q-lw-1.6*) with lower Q×L interaction effects could be considered as stable, their phenotypic variance is not large enough to deploy them in MAS. Based on these results, strategies to breed maize for resistance to LWD are discussed.

## Introduction

Maize productivity in India is constrained by several prevalent and emerging insect pests and diseases. Of these, post-flowering stalk rot (PFSR) is a complex disease caused by three pathogens, namely, *Fusarium moniliforme* J. (Sheld), *Macrophomina phaseolina* (Tassi) Goid and *Harpophora maydis* (Samra, Sabet and Hing) (Khokhar et al. 2014). Stalk rot caused exclusively by seed-borne and soil-borne pathogen *H. maydis* causes premature wilting symptoms at post-flowering stage, a condition known as Late Wilt Disease (LWD). Typical symptoms of LWD include pre-mature wilting of leaves, discoloration of the stalk followed by stalk tissue disintegration and fibrousness in advanced stages (Samra et al. 1963). LWD has been reported in about 10 countries, with significant economic losses in Egypt (Sabet et al. 1961), India (Payak et al. 1970), Spain and Portugal (Molinero-Ruiz et al. 2010) and Israel (Drori et al. 2013). Based on survey, economic losses of 40% have been reported in Egypt (Labib et al. 1975; Galal et al. 1979), up to 100% in Israel (Degani et al. 2019) and up to 51% in India (Johal et al. 2004) due to LWD. However, 3.5 to 38.4% loss of grain yield attributable to soil inoculation by *H. maydis* has been reported based on an empirical study using a limited number of genotypes in Egypt (El-Naggarr et al. 2015). The Egyptian, Indian, and Hungarian isolates have been reported to differ in their morphology, pathogenicity, and route of infection (Warren 1983). Very recently, we have reported 5.8 to 44.2% grain yield losses attributable to LWD by creating artificial epiphytotic condition through injecting *H. maydis* inoculum into the stem internode (Sunitha et al. 2020) in India.

Genetic intervention is regarded as eco-friendly and cost-effective option to mitigate losses caused by LWD (El-Shafey et al. 1988; Zeller et al. 2000). As a prelude to genetic intervention, stable sources of resistance to *H. maydis* (Shekhar et al. 2010; Rakesh et al. 2016b; Aruna 2017; Mahesh 2019) have been identified. The inheritance of resistance to LWD has been reported as complex with significant genotype × environment interaction (Shekhar et al. 2010; Rakesh 2016; Aruna 2017; Mahesh 2019). Hence, direct selection for LWD resistance is likely to be less effective. Nevertheless, DNA markers could be used as effective surrogates of such traits in maize for which *a priori* identification and validation of closely linked markers is essential. As a step towards this, Quantitative Trait Locus/Loci (QTL) conferring LWD tolerance (Rakesh, 2016) on chromosomes 1,3,5,7 and 10 have been detected. The objectives of our study were to (i) validate the QTL detected by Rakesh (2016) (ii) to detect and estimate size effects of new LWD resistance governing QTL, if any in two different genetic backgrounds and (iii) detect and estimate size effects of their interaction with two spatial environments.

## Material And Methods

### Basic genetic material and development of mapping populations

The basic genetic material consisting of one LWD resistant inbred line, CV156670 and two LWD susceptible inbred lines, 414 – 33 and CV143587, developed and being maintained at Bayer Crop Science (erstwhile Monsanto India Ltd.) were used as parents to develop two connected mapping populations. For intellectual property protection reasons, the pedigree of these inbreds is not included. The two  $F_1$ 's (available with Bayer Crop Science), CV156670 × 414 – 33 and CV156670 × CV143587 were selfed during 2017 summer season to derive  $F_2$  populations. Two  $F_2$  populations derived from connected crosses CV156670 × 414 – 33 and CV156670 × CV143587 were selfed during 2017 rainy season. These  $F_{2:3}$  populations constituted inter-connected mapping populations to detect QTL controlling LWD resistance.

### Phenotyping mapping populations for responses to LWD

The mapping populations were evaluated at two locations, Kallinayakanahalli (KNH) 13.464013°N, 77.519178°E and Muppadihatta (MPG) 13.241161°N, 77.483637°E. 300  $F_{2:3}$  progeny families each from two  $F_2$  populations, parental inbred lines and checks (susceptible and resistant check) were sown in single rows of 4 m length with 0.6 m spacing between rows and 0.2 m spacing between plants in two replications following randomized complete block design (RCBD) during 2017 post-rainy season. Recommended package of practices were followed to establish good crop stand till artificial inoculation. The  $F_{2:3}$  progeny families were screened for responses to LWD using artificial inoculation protocol (Rakesh et al. 2016a).

### Artificial inoculation of *H. maydis* inoculum

Naturally infested LWD stalk specimens collected from commercial maize production fields were used for isolation of pathogens and inoculum preparation. The inoculum concentration was assessed using haemocytometer and was adjusted to  $4 \times 10^6$  spore  $ml^{-1}$ . A hole was made at the second internode on the stem of plants using a proprietary specialized stem jabbers and 2 ml of the *H. maydis* was injected to the holes using a syringe twice at 45 and 55 days after

sowing (DAS). From each progeny family, 20 plants were inoculated. All the recommended production practices were followed except for the application of potassium fertilisers and fungicides to maintain the plants after inoculation.

## Sampling and data recording

The stalks of the artificially inoculated plants were split open at 110 DAS to examine the typical symptoms of LWD. Disease symptoms were manifested by discoloration, disintegration and fibrousness at the inoculated internode. Based on the coverage of disease symptom within an internode and its spread across internodes, the data on LWD severity was recorded on all the inoculated plants using modified 1–9 scale (Table 1) (Rakesh et al. 2016a).

Table 1  
ANOVA of mean LWD scores in two  $F_{2:3}$  populations across two locations

Source of variation	CV156670 × CV143587 (P-1)				CV156670 × CV414-33 (P-2)				
	Degrees of freedom	KNH	MPG	Degrees of freedom	KNH	MPG			
Mean sum of squares	'F' probability	Mean sum of squares	'F' probability	Mean sum of squares	'F' probability	Mean sum of squares	'F' probability		
Between $F_{2:3}$ families	235	0.71**	< 0.001	0.47	0.08	186	1.07	< 0.05	0.71 < 0.001
Within $F_{2:3}$ families	13	0.15		0.17		06	0.32		0.07

## Genotyping $F_{2:3}$ families

Five  $F_{2:3}$  seeds from each plant of two  $F_2$  populations were pooled and genotyped using Monsanto proprietary-SNP markers through TaqMan assay (Semagn et al. 2015). A total of 150 and 199 SNP markers polymorphic between the parents of the two mapping populations were selected from proprietary database (unpublished) for genotyping the two mapping populations derived from CV156670 × 414 – 33 and CV156670 × CV143587, respectively.

## Statistical analysis

### LWD responses

LWD responses from 20 plants of each  $F_{2:3}$  progeny family were recorded. Data were cured based on missing data points, outliers and correlation between replicates (Mailman et al. 2007; Rakesh 2016). Mean LWD scores of  $F_{2:3}$  families were used for statistical analysis after data curation. Analysis of Variance (ANOVA) of  $F_{2:3}$  progeny families was performed to partition the total variation in LWD response scores into those attributable to  $F_{2:3}$  progeny families and within families. Additionally, pooled ANOVA was performed to test for the consistency of the  $F_{2:3}$  progeny families for LWD response across two locations and to quantify the contribution of location and genotype × location interaction. Genotypic data of markers with > 15% missing data and markers showing significant segregation distortion were not considered for statistical analysis.

### Detection of QTL controlling resistance to LWD

Markers showing segregation distortion were removed (Supplementary table 1 and 2). The genotypic and phenotypic data of  $F_{2:3}$  progeny families evaluated in two locations were integrated to detect QTL controlling LWD resistance, initially using single marker analysis. Subsequently, Inclusive Composite Interval Mapping (ICIM) was performed to detect and estimate size effects of QTL and QTL × QTL interactions (Li et al. 2015) controlling LWD resistance at individual locations and across locations (pooling data from both locations). The accuracy of QTL position and significance of size effect of QTL and QTL × Location interaction conferring LWD resistance was determined using data-driven estimates of threshold LOD scores obtained by 1000 permutations (Churchill and Doerge 1994). Similarly, significant QTL × QTL interactions controlling LWD resistance were detected at threshold LOD value of 5.0 and their size effects were estimated. All these statistical analyses were implemented using QTL IciMapping software version 4.0 (Wang et al. 2016). The detected di-QTL interactions associated with resistance to LWD, dominance × dominance di-QTL interactions were interpreted based on theoretical investigations of Kearsey and Pooni (1996).

## Results

### LWD responses of $F_{2:3}$ mapping populations

The LWD response scores of the  $F_{2:3}$  progeny families derived from CV156670 × 414 – 33 (P-1) and CV156670 × CV143587 (P-2) recorded at both KNH and MPG locations were normally distributed (Fig. 1). While LWD response scores of  $F_{2:3}$  progeny families ranged from 3.17–9.00 and from 3.13–8.11 at KNH and MPG, respectively in P-1, the responses ranged from 2.38–8.38 and from 3.12–8.64 at KNH and MPG, respectively in P-2. Wider range of LWD scores at both locations in both the populations indicated sufficient LWD expression. Further, there were considerable numbers of plants towards both resistant and susceptible parents (Fig. 2). Six progeny families including two at KNH and four at MPG in P-1; 56 progeny families including 37 at KNH and 19 at MPG in P-2 showed resistance level better than resistant parent. Similarly, six progeny families including four at KNH and two at MPG in P-1; five progeny families including two at KNH and three at MPG in P-2 showed higher susceptibility than susceptible parent. The  $F_{2:3}$  families derived from both the mapping populations differed significantly across both the locations except P-1 at MPG (Table 1). Further, LWD responses of  $F_{2:3}$  families of both populations interacted significantly with location as evident from significance of genotype × location (G×L) interaction mean squares in pooled ANOVA (Table 2). While

mean squares attributable to location was non-significant, those attributable to genotype × location was significant in P-2. This trend was reversed in P-1 (Table 2).

Table 2  
Pooled ANOVA of mean LWD scores in F<sub>2:3</sub> populations across two locations

Source of variation	CV156670 × CV143587 (P-1)				CV156670 × CV414-33 (P-2)			
	Degrees of freedom	Mean sum of squares	'F' probability	Per cent contribution	Degrees of freedom	Mean sum of squares	'F' probability	Per cent contribution
Genotypes	235	1.52	< 0.001	46.55	186	3.17	< 0.001	77.89
Locations	01	22.41	< 0.001	2.92	01	0.01	0.971	0.00
Genotype × Location (G × L)	235	0.61	0.08	18.68	186	0.44	< 0.001	10.81
Error	470	0.52	-	31.85	372	0.23	-	11.30

## Detection of QTL controlling LWD resistance

### QTL main effects

Seven QTL *i.e.*, two (*q-lw-1.1* and *q-lw-1.2*) in P-1 (Fig. 3) and five (*q-lw-1.3*, *q-lw-1.4*, *q-lw-1.5*, *q-lw-1.6* and *q-lw-1.7*) in P-2 detected on chromosome-1 (Fig. 4) were found associated with LWD resistance with phenotypic variation explained (PVE) ranging from 4.34 to 13.06 % (Table 3). Among these, three major QTL (*q-lw-1.4* and *q-lw-1.6* at MPG; *q-lw-1.5* at KNH) with > 10% PVE were detected in P-2. Among the remaining four minor QTL, two (*q-lw-1.1* and *q-lw-1.2*) were from P-1 and other two (*q-lw-1.3* and *q-lw-1.7*) were from P-2. Further, while five (*q-lw-1.1*, *q-lw-1.3*, *q-lw-1.5*, *q-lw-1.6* and *q-lw-1.7*) of these seven detected QTL showed dominance effects, one (*q-lw-1.4*) showed additive effects in desirable direction (decreasing effects) for LWD resistance (Table 3).

Table 3  
Position and effect of QTL conferring LWD resistance detected individual location-wise and across locations (pooled)

Trait	QTL	Environment	Population	Chromosome	Genomic Position (cM)	Flanking markers		LOD	PVE Q (%)	PVE Q × L (%)	Additive effect	Dominance effect
						Right	Left					
Late Wilt Disease	<i>q-lw-1.1</i>	Kallinayakanahalli	CV156670	01	138.60	Marker-12	Marker-13	4.45	7.92	-	0.44	<b>-0.13</b>
	<i>q-lw-1.2</i>	Muppadighatta	CV143587 (P-1)	112.60	Marker-9	Marker-10	4.56	9.91	-	0.30	0.05	
		Pooled			4.61	8.28	3.80	0.17	0.04			
	<i>q-lw-1.3</i>	Kallinayakanahalli	CV156670		11.40	Marker-151	Marker-152	4.06	4.34	-	0.41	<b>-0.29</b>
	<i>q-lw-1.5</i>	Muppadighatta	CV414-33 (P-2)	9.40	4.22	10.10	-	0.33	<b>-0.31</b>			
		Pooled			8.19	8.85	0.04	0.33	<b>-0.29</b>			
	<i>q-lw-1.4</i>	Kallinayakanahalli		224.40	Marker-179	Marker-180	12.27	11.72	-	<b>-0.02</b>	1.08	
		Pooled			12.34	8.73	6.80	<b>-0.01</b>	0.56			
<i>q-lw-1.6</i>	Muppadighatta		83.40	Marker-160	Marker-161	5.97	13.06	-	0.40	<b>-0.28</b>		
	Pooled			6.23	4.35	1.20	0.24	<b>-0.17</b>				
<i>q-lw-1.7</i>	Pooled		232.40	Marker-180	Marker-181	5.42	5.60	0.31	0.18	<b>-0.38</b>		

### QTL × Location interaction

Pooled analysis across locations indicated significant QTL × location (Q×L) interaction. Of the seven detected QTL, five (*q-lw-1.2*, *q-lw-1.4*, *q-lw-1.5*, *q-lw-1.6* and *q-lw-1.7*) interacted significantly with location with their size effects ranging from 0.04 to 6.80 % PVE (Table 3). Of these five interacting QTL, only *q-lw-1.7* was not detected when individual location-wise analysis was performed. This means that, remaining four (*q-lw-1.2*, *q-lw-1.4*, *q-lw-1.5* and *q-lw-1.6*) of these five QTL detected using pooled analysis were detected only in either of the locations but not in both the locations. These four QTL were also stable for their position but unstable for their effects (decreased size effects) as evident from pooled analysis.

Of the five QTL which showed significant Q × L interaction, one QTL, *q-lw-1.4* (6.80% PVE Q × L) interacted with locations to a greater extent than the other four QTL (Fig. 5). None of the QTL detected through location-wise analysis were common in both locations. Two QTL (*q-lw-1.3* and *q-lw-1.5*) flanked by same pair of markers, namely, Marker-151 and Marker-152 (with inter-marker distance of 23.8 cM) were detected in P-2. While the QTL, *q-lw-1.3* was mapped at 11.40 cM

in KNH (in location-wise analysis), *q-lw-1.5* was mapped at 9.40 cM in MPG (in location-wise as well as pooled analysis). Further, their position and effects differed (Table 3).

## Di-QTL epistasis

A total of 48 significant di-QTL interactions with size effects (in % PVE) ranging from 4.00 to 19.03% were detected across both locations in P-1. Of these, ten were between QTL located within the same chromosome (Fig. 6a). Only nine of these 48 interactions involved QTL with significant effects. Similarly, 52 significant di-QTL epistatic interactions including ten within same chromosome were detected in P-2 with their size effects (in % PVE) ranging from 3.18 to 13.83 % (Fig. 6b). Only four of these 52 significant di-QTL interactions involved QTL with significant effects. Summation of direction and effects of dominance (h) and dominance × dominance (l) effects associated with all di-QTL interaction effects indicated duplicate di-QTL epistasis in both the populations. Duplicate di-QTL epistasis was in desirable direction (dominant decreasing effects) at KNH and MPG in P-1 and P-2, respectively (Table 4).

Table 4  
Direction of dominance effects and all possible dominance × dominance di-QTL interactions controlling resistance to LWD without considering their significance or otherwise

Main QTL effect (h)	Dom × Dom effect (l)	CV156670 × CV143587 (P-1)			CV156670 × CV414-33 (P-2)			Interpretation on type of epistasis
		KNH	MPG	Across locations	KNH	MPG	Across locations	
+	+							Complementary epistasis between dominant increasing effect QTLs
-	-							Complementary epistasis between dominant decreasing effect QTLs
+	-		∏		∏			Duplicate epistasis between dominant increasing effect
-	+	∏		∏		∏	∏	Duplicate epistasis between dominant decreasing effect

## Validation of reported QTL conferring resistance to LWD

Rakesh (2016) reported QTL on chromosome-1 at 103.20 cM (IMD- 10.6 cM), flanked by MONIND14101828 and MONIND14262087. This QTL explained only 7.26 % of LWD resistance in mapping population derived by CV138811 × CV143587. In our study, one of these markers namely, Marker-161 (MONIND14101828) flanking the QTL, *q-lw-1.6* could be validated in P-2 (CV156670 × CV143587). However, this QTL flanked by Marker-160 and Marker-161 was detected at position (83.40 cM) and phenotypic variation (13.06 % PVE at MPG and 4.35 % PVE across locations) different from that reported by Rakesh (2016) (Table 3).

## Discussion

LWD is emerging as a major and serious biotic production constraint with a potential to reduce yield and quality of both grain and fodder (Sunitha et al. 2020; El-Naggarr et al. 2015; Drori et al. 2013; Johal et al. 2004). Developing LWD resistant cultivars is considered as an eco-friendly and sustainable approach to mitigate losses caused by LWD. Precise information on the number and mode of action of genes/QTL controlling resistance to LWD help devise suitable strategies to develop resistant cultivars. The results of our study indicate possible involvement of a large number of genes controlling resistance to LWD as evident from the normal distribution of LWD responses in  $F_{2,3}$  populations. Several previous researchers such as Shehata (1976), Nawar and Salem (1985), Abdel-Snbour and Bekhit (1993) and El-Hosary and El-Fiki (2015) have also reported the involvement of a large number of genes controlling LWD resistance in maize. Further, recovery of individuals that surpass the levels of resistant and susceptible parents indicate (i) presence of alleles controlling resistance even in susceptible parent and those controlling susceptibility even in resistant parent and (ii) complementation of parental alleles controlling resistance. Thus,  $F_{2,3}$  populations serve as potential source for recovering inbred lines with levels of resistance to LWD better than the resistant parent.

Most of the reports (Wu et al. 2020; Zhang et al. 2012; Yang et al. 2010; Suneetha 2016) related to mapping QTL controlling resistance to stalk rot are predominantly on stalk rot caused by *Fusarium* sp. followed by *Macrophomina phaseolina* but not on the stalk rot caused exclusively by *Harpophora maydis* (Late wilt disease). This could be because of co-existence of *Harpophora* with *Fusarium* sp. and/or *Macrophomina* sp. which form a pathogen-complex causing PFSR (Degani et al. 2020; Khokhar et al. 2014). Additionally, soil-borne *Fusarium* sp. and/or *Macrophomina* sp. are known to invade the *H. maydis* infested stalks (Drori et al. 2013). This causes difficulty in phenotyping the symptoms caused exclusively by *H. maydis*. To address the issue of possible confounding effects of pathogens other than *H. maydis*, we used Indian pure isolates of *H. maydis* to inoculate the spores directly into the second internode of the stem (Rakesh et al. 2016) using stem jabbers (specialised injectors) twice at 45 DAS and 55DAS (Sunitha et al. 2020).

*LWD resistance conferring QTL*: Difficulty arising due to complexity of the disease and phenotyping could be the potential causes for sporadic reports on genetic basis of LWD resistance. The only report (Rakesh 2016) on mapping genomic regions controlling LWD resistance in two  $F_{2,3}$  populations (derived using CV138811 and CV143587 as donor and susceptible parents, respectively) resulted in the identification of three major QTL on chromosomes 3, 5 and 10; and six minor QTL on chromosomes 1, 2, 3, 5, 6 and 7. QTL analysis is not only intended to implement MAS for QTL but also for understanding the genetics of the quantitative trait. Hence, all the identified QTL, whether their effects are large or small and with or without environmental sensitivity are informative (Asins 2002). In our study, in addition to three major QTL detected either location-wise or across locations (pooled), four minor QTL were detected. In most of the studies, QTL with small effects go undetected due to bias caused by less-stringent threshold levels. To minimize such bias, we used LOD threshold derived from 1000 permutations (Churchill and Doerge 1994; Beavis 1994) in our study.

**QTL×Location interaction:** Position and effect of the QTL associated with LWD resistance varied significantly with the spatial environments represented by two locations in our study. If a quantitative trait exhibits significant genotype by environment interaction, it then follows that underlying QTL should also display significant interaction with either temporal or spatial environments (Bernardo 2020). In our study, the  $F_3$  progenies of both the crosses displayed significant interaction with the environment represented by two locations for responses to LWD (Table 2). Hence, detected significant interaction of LWD resistance governing QTL with location environments in our study did not surprise us. Several researchers have also reported significant QTL × location interaction associated with resistance response of maize diseases such as Fusarium ear rot (Wu et al. 2020; Robertson-Hoyt et al. 2006), northern leaf blight (Xia et al. 2020; Chen et al. 2016) and rough dwarf disease (Wang et al. 2019). The major QTL (> 10% PVE) detected through location-wise mapping behaved as minor QTL (< 10% PVE) when detected through pooled analysis. We could also observe shift in the position of QTL ( $q-lw-1.3$  and  $q-lw-1.5$ ) within the same flanking marker interval in different locations. However, such QTL despite showing significant G × E interaction, help select genotypes adapted to specific locations (Asins 2002). As expected based on previous reports by Shekhar et al. (2010), we could also observe significant influence of location and genotype × location interaction on the expression of LWD. Significance of G × E interaction for a trait reflects that underlying genomic regions exhibit significant QTL × Location interaction (Bernardo 2020). Among the three major QTL identified by us, although two QTL ( $q-lw-1.5$  and  $q-lw-1.6$ ) showed significant Q×L interaction, the magnitude of interaction effects was relatively low. These QTL with lower Q×L interaction effects could be considered as stable and deployed to implement MAS for LWD resistance after validation. This is because, the efficiency of MAS may get reduced if detected QTL controlling target trait exhibits significant interaction with either the environment or background genotype used (Asins 2002).

**QTL×QTL interaction:** The chances of recovering recombinant inbred lines (RILs) with dominant di-QTL interactions are fewer with any population other than  $F_2$ . However, even a large  $F_2$  mapping population may contain fewer two-QTL RILs limiting the statistical power of detecting di-QTL interactions (Bernardo, 2020). Hence, there are chances that epistatic interactions go undetected. We addressed this limitation by increasing the LOD significance of > 5.0 and considering phenotypic data from replicated trials from  $F_{2,3}$ . Some of the di-QTL interactions which involve QTL with their individual effects being non-significant are also informative, provided their di-QTL interaction effects are of higher magnitude. Under such instance, deployment of QTL alone in MAS without considering its significant di-QTL interaction would be ineffective. Hence, such QTL showing significant epistatic effects of large magnitude need to be introgressed together into recipient genetic background. However, there are chances that the magnitude of the significant di-QTL interaction may get reduced or get enhanced in recipient parent genetic background. The effects of such interacting QTL in recipient genetic background will be known after introgression. The direction of cumulative dominance effects associated with duplicate di-QTL epistasis varied with location. The duplicate di-QTL epistasis with decreasing dominance effects on LWD response is desirable. However, these di-QTL interaction effects need to be confirmed through extensive multi-location evaluation.

#### QTL genetic background interaction

Genetic background has considerable influence on the position and effect of QTL conferring disease resistance (Asins 2002; Awata et al. 2020). The same was evident in our study and none of the QTL was common between both populations, P-1 and P-2 (which differed for susceptible parent genome). However, we could validate only one of the reported markers, MONIND14101828 (Marker-161) flanking the reported QTL (Rakesh 2016) in P-2. While the QTL region having this marker, MONIND14101828 (Marker-161) in common is flanked by inter-marker distance of 11.8 cM in our study, it is flanked by inter-marker distance of 10.6 cM in previous report (Rakesh, 2016). Thus, this region needs fine mapping to identify QTL with narrow inter-marker distance for efficient deployment in MAS.

In summary, three major QTL,  $q-lw-1.4$ ,  $q-lw-1.5$  and  $q-lw-1.6$  were identified from our study. However, PVE% explained by these three QTL is not large enough for use in MAS. Further, only one of these QTL,  $q-lw-1.6$  could be validated. The three major QTL exhibited significant interaction with locations. Considering that the validated QTL reported by us and those reported by previous researcher exhibit significant interactions with locations, further research investigation is essential to identify stable QTL with large PVE for implementing MAS.

## Declarations

**Funding:** This research was funded by Bayer Crop Science (erstwhile Monsanto India Ltd.) under collaborated project with University of Agricultural Sciences, Bengaluru, Karnataka, India including financial support in the form of 'Monsanto student fellowship'.

**Conflict of interest:** The authors declare that they have no conflict of interest.

**Availability of Data and material:** Not applicable

**Code availability:** Not applicable

**Acknowledgement:** Extension of field and lab facilities by Monsanto India Limited under collaborated project with University of Agricultural Sciences, Bengaluru, Karnataka, India are gratefully acknowledged.

**Authors' Contributions:** EG, VRPG and SR conceptualized the work. SNC conducted experiments and analyses. DS and SNC inoculation and screening for late wilt disease. SB, SR and SNC interpreted results. SNC and SR revised the manuscript. EG, SR,VRPG and HBH reviewed and edited the final version.

## References

Abdel-Snbour MS, Bekhit MMM (1993) Genetic variability and diallel analysis of late wilt resistance and some other characters in maize. Zagazig J Agril Res (Egypt)

- Nawar AA, Salem MA (1985) Diallel analysis of the inheritance of late wilt and leaf blight in maize [Egypt]. *Monoufeyah J Agril Res (Egypt)*.
- Aruna K (2017) Identification of high yielding and late wilt resistant maize inbreds. M.Sc. (Agri.) Thesis, Univ Agric Sci, Bangalore.
- Asins MJ (2002) Present and future of quantitative trait locus analysis in plant breeding. *Plant Breed* 121(4):281-291. <https://doi.org/10.1046/j.1439-0523.2002.730285.x>
- Awata LA, Beyene Y, Gowda M, LM S, Jumbo MB, Tongoona P, Danquah E, Ifie BE, Marchelo-Drugga PW, Olsen M, Ogugo V, Mugo S, Prasanna BM (2020) Genetic analysis of QTL for resistance to maize lethal necrosis in multiple mapping populations. *Genes* 11(1): 32. <https://doi.org/10.3390/genes11010032>
- Bernardo R (2020) Breeding for quantitative traits in plants. Third edition Woodbury, MN: Stemma press
- Chen G, Wang X, Long S, Jaqueth J, Li B, Yan J, Ding J (2016) Mapping of QTL conferring resistance to northern corn leaf blight using high-density SNPs in maize. *Mol Breed* 36(1): 4. <https://doi.org/10.1007/s11032-015-0421-3>
- Churchill GA, Doerge RW (1994) Empirical threshold values for quantitative trait mapping. *Genet* 138(3):963-971. <https://doi.org/10.1093/genetics/138.3.963>
- Degani O, Dor S, Abraham D, Cohen R (2020) Interactions between *Magnaportheopsis maydis* and *Macrophomina phaseolina*, the Causes of Wilt Diseases in Maize and Cotton. *Microorganisms* 8(2): 249. <https://doi.org/10.3390/microorganisms8020249>
- Degani O, Dor S, Movshovitz D, Rabinovitz O (2019) Methods for studying *Magnaportheopsis maydis*, the maize late wilt causal agent. *Agronomy* 9(4):181. <https://doi.org/10.3390/agronomy9040181>
- Drori R, Sharon A, Goldberg D, Rabinovitz O, Levy M, Degani O (2013) Molecular diagnosis for *Harpophora maydis*, the cause of maize late wilt in Israel. *Phytopathologia Mediterranea* 52(1): 16–29.
- El-Hosary AAA, El-Fiki IAI (2015) Diallel cross analysis for earliness, yield, its components and resistance to late wilt in maize. *Int J Agric Sci Res* 5:199-210
- El-naggarr AAA, Sabryr AM, Yassin MA (2015) Impact of late wilt disease caused by *Harpophora maydis* on maize yield. *J Biol Chem Env Sci* 10(3): 577- 595
- El-Shafey HA, El-Shorbagy FA, Khalil II, El-Assiuty EM (1988) Additional sources of resistance to the late-wilt disease of maize caused by *Cephalosporium maydis*. *Agric Res Rev* 66:221-230
- Galal AA, El-Rouby MM, Gad AM (1979) Genetic analysis of resistance to late wilt (*Cephalosporium maydis*) in variety crosses of maize. *Zeitschrift für Pflanzenzüchtung* 83: 176–183
- Johal L, Huber DM, Martyn R (2004) Late wilt of corn (maize) pathway analysis: intentional introduction of *Cephalosporium maydis*. In: Pathways Analysis for the Introduction to the U.S. of Plant Pathogens of Economic Importance. U.S. Department of Agriculture, animal and plant health inspection service technical report no. 503025.
- Kearsey MJ, Pooni HS (1996) The genetical analysis of quantitative traits. (Chapman and Hall: New York).
- Khokhar MK, Hooda KS, Sharma SS, Singh V (2014) Post flowering stalk rot complex of maize-present status and future prospects. *Maydica* 59(3): 226-242
- Labib HA, Salem A, Abd El Rahim ME, Abd El Fattah A (1975) DC 19, a new maize hybrid seed resistant to late-wilt disease caused by *Cephalosporium maydis*. *Agricultural Research Review* 53(8):1-4.
- Li S, Wang J, Zhang L (2015) Inclusive composite interval mapping of QTL by environment interactions in biparental populations. *PLoS One* 10(7):e0132414. <https://doi.org/10.1371/journal.pone.0132414>
- Mahesh B (2019) Identification of high yielding and late wilt (*Harpophora maydis*) disease resistant hybrids of maize (*Zea mays* L.). M.Sc. (Agri.) Thesis, Univ Agric Sci, Bangalore.
- Mailman MD, Feolo M, Jin Y, Kimura M, Tryka K, Bagoutdinov R, Hao L, Kiang A, Paschall J, Phan L, Popova N (2007) The NCBI dbGaP database of genotypes and phenotypes. *Nat Genet* 39(10):1181-1186 doi:10.1038/ng1007-1181
- Molinero-Ruiz ML, Melero-Vara JM, Mateos A (2010) *Cephalosporium maydis*, the cause of late wilt in maize, a pathogen new to Portugal and Spain. *Plant Dis* 94(3):379. <https://doi.org/10.1094/PDIS-94-3-0379A>
- Pandey SK, Sarkar DE, Sharma SU (2006) Quantitative Trait Loci (QTLs) for plant disease response. *Proceedings-Indian National Science Academy* 72(3):179
- Payak MM, Lal S, Renfro BL (1970) *Cephalosporium maydis*-a new threat to Maize in India. *Indian Phytopathol* 23(3):562-569
- Rakesh B (2016) Mapping genomic regions controlling late wilt disease resistance caused by *Harpophora maydis* in maize (*Zea mays* L.) Ph.D. Thesis, Univ Agri Sci, Bangalore.
- Rakesh B, Gangappa E, Sonali G, Gowda RV, Swamy SD, Ramesh S, Hemareddy HB, Nagaraju N (2016a) Modified Method of Screening Maize Inbred Lines to Late Wilt Disease Caused by *Harpophora maydis*. *Mysore J Agric Sci* 50:684-690

- Rakesh B, Gangappa E, Sonali G, Ramesh S, Nagaraju N (2016b) Identification of maize (*Zea mays* L.) inbred lines resistant for late wilt disease caused by *Harpophora maydis*. *Mysore J Agric Sci* 50(2):366-368
- Robertson-Hoyt LA, Jines MP, Balint-Kurti PJ, Kleinschmidt CE, White DG, Payne GA, Maragos CM, Molnár TL, Holland JB (2006) QTL mapping for Fusarium ear rot and fumonisin contamination resistance in two maize populations. *Crop Sci* 46(4):1734-1743. <https://doi.org/10.2135/cropsci2005.12-0450>
- Sabet KA, Samra AS, Hingorani MK, Mansour IM (1961) Stalk and root rots of Maize in the United Arab Republic. *Plant Protection Bulletin, FAO* 9(7): 121-125
- Samra AS, Sabet KA, Hingorani MK (1963) Late wilt disease of maize caused by *Cephalosporium maydis*. *Phytopathol* 53: 402-406
- Semagn K, Beyene Y, Babu R, Nair S, Gowda M, Das B, Tarekegne A, Mugo S, Mahuku G, Worku M, Warburton ML (2015) Quantitative Trait Loci Mapping and Molecular Breeding for Developing Stress Resilient Maize for Sub-Saharan Africa. *Crop Sci* 55(4):1449-1459. <https://doi.org/10.2135/cropsci2014.09.0646>
- Shehata AH 1976, Gene action involved in the manifestation of late wilt (*Cephalosporium maydis*) of maize. *Egyptian J Genet Cytol* 5(1): 42-47
- Shekhar M, Kumar S, Sharma RC, Singh R (2010) Sources of resistance against post-flowering stalk rots of maize. *Arch Phytopathol Pflanzenschutz* 43(3): 259-263. doi: 10.1080/03235400701803846
- Suneetha P (2016) Mapping of gene(s) for resistance to post flowering stalk rot in maize (*Zea mays* L.) caused by *Macrophomina phaseolina* (Tassi) Goid (Doctoral dissertation, Professor Jayashankar Telangana State Agricultural University Rajendranagar, Hyderabad)
- Sunitha NC, Gangappa E, Gowda RV, Ramesh S, Biradar S, Swamy D, Hemareddy HB (2020) Assessment of Impact of Late Wilt Caused by *Harpophora maydis* (Samra, Sabet and Hing) on Grain Yield and its Attributing Traits in Maize (*Zea mays* L.). *Mysore J Agric Sci* 54(2):30-36
- Wang J, Li H, Zhang L, Meng L (2016) User's manual of QTL IciMapping ver. 4.1'. Quantitative Genetics group, Institute of Crop Science, Chinese Academy of Agricultural Sciences (CAAS): Beijing/Genetic Resources Program, International Maize and Wheat Improvement Center (CIMMYT): Mexico City.
- Wang X, Yang Q, Dai Z, Wang Y, Zhang Y, Li B, et al. (2019) Identification of QTLs for resistance to maize rough dwarf disease using two connected RIL populations in maize. *PLoS ONE* 14(12): e0226700. <https://doi.org/10.1371/journal.pone.0226700>
- Warren HL (1983) Potential disease problems-late wilt of maize. In *Phytopathology* (Vol. 73, No.5, pp.782-782). 3340 Pilot Knob Road, ST Paul, MN 55121: Amer Phytopathological Soc
- Wu Y, Zhou Z, Dong C, Chen J, Ding J, Zhang X, Mu C, Chen Y, Li X, Li H, Han Y (2020) Linkage mapping and genome-wide association study reveals conservative QTL and candidate genes for Fusarium rot resistance in maize. *BMC Genomics* 21:357 <https://doi.org/10.1186/s12864-020-6733-7>
- Xia H, Gao W, Qu J, Dai L, Gao Y, Lu S, Zhang M, Wang P, Wang T (2020) Genetic mapping of northern corn leaf blight-resistant quantitative trait loci in maize. *Med* 99(31). doi: 10.1097/MD.00000000000021326
- Yang Q, Yin G, Guo Y, Zhang D, Chen S, Xu M (2010) A major QTL for resistance to Gibberella stalk rot in maize. *Theor Appl Genet* 121(4):673-687
- Zeller KA, Jurgenson JE, El-Assiuty EM, Leslie JF (2000) Isozyme and amplified fragment length polymorphisms from *Cephalosporium maydis* in Egypt. *Phytoparasitica* 28(2):121-130
- Zhang D, Liu Y, Guo Y, Yang Q, Ye J, Chen S, Xu M (2012) Fine-mapping of qRfg2, a QTL for resistance to Gibberella stalk rot in maize. *Theor Appl Genet* 124(3):585-596.

## Figures







Figure 1

Frequency distribution for responses to LWD of two F2:3 populations at Kallinayakanahalli and Muppadighatta locations





Figure 2

Patterns of expression of responses to LWD by two F2:3 populations (a. P-1 and b. P-2) across two locations




Figure 3  
 Genome-wide detection of QTL and their effects controlling LWD response mapped in F2:3 of CV156670 (R) × 414-33(S) (P-1) at (a) Kallinayakanahalli (b) Muppadighatta




Figure 4  
 Genome-wide detection of QTL controlling LWD response and corresponding additive and dominance effects mapped in F2:3 of CV156670(R) × CV143587(S) (P-2) at (a) Kallinayakanahalli (b) Muppadighatta





Figure 5  
 Genome-wide detection of QTL controlling LWD response and QTL  $\times$  Location interactions in F2:3 of (a) CV156670(R)  $\times$  414-33(S)/ P-1 (b) CV156670(R)  $\times$  CV143587(S)/ P-2



## Figure 6

Genome-wide detection of significant di-QTL interactions controlling LWD response mapped in F2:3 of (a) CV156670(R) × 414-33(S) /P-1 (b) CV156670(R) × CV143587(S) /P-2.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementarytables.docx](#)