

Alcohol Consumption and Sports-betting Among Young Male Motorcycle Taxi Bodaboda Riders in Urban Southwestern Uganda

Stella Cheryl Nabifo

Mbarara University of Science and Technology Faculty of Medicine

Jonathan Izudi

Mbarara University of Science and Technology Faculty of Medicine

Francis Bajunirwe (✉ fbaj@yahoo.com)

Department of Community Health Mbarara University of Science and Technology P.O.BOX 1410,
Mbarara, Uganda <https://orcid.org/0000-0002-1326-5230>

Research article

Keywords: Sports-betting, alcohol consumption, Uganda, young males

Posted Date: August 25th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-60020/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Version of Record: A version of this preprint was published on February 16th, 2021. See the published version at <https://doi.org/10.1186/s12889-021-10406-7>.

Abstract

Introduction: The popularity of sports-betting is growing globally and may be associated with alcohol consumption among young adults. In this study, we examine the relationship between alcohol consumption plus other substances with sports-betting in a group of young adult males in Uganda.

Methods: We conducted a cross-sectional study and interviewed male motorcycle taxi *bodaboda* riders in the southwestern Uganda town of Mbarara. We asked questions about alcohol consumption, smoking and history and frequency of sports-betting.

Results: We enrolled 401 riders. The median age was 29.0 years, interquartile range 25-32. Seventy four (18.5%) had engaged in sports-betting in the past 30 days. Among those that reported sports-betting, 16(21.6%) engaged daily or almost daily. Alcohol use was significantly associated with gambling with an adjusted risk ratio (aRR) of 2.08(95% CI 1.36, 3.18) among moderate drinkers compared to low or non-drinkers but not among heavy drinkers. Cigarette smoking was significantly associated with sports-betting with an aRR 1.85(95% CI 1.13, 3.01).

Conclusion: Our data show sports-betting is common among these relatively young male motorcycle riders, and is associated with alcohol consumption and cigarette smoking. Interventions to regulate sports-betting should be co-packaged with those to reduce alcohol consumption and cigarette smoking among young adult males.

Introduction

There is a global increase in the popularity of sports-betting and involvement of sports-betting companies in sponsorship and advertisement in popular sports events such as soccer [1]. The number of sports-betting companies involved in shirt sponsorship for football in European leagues has grown exponentially. In the top five European soccer leagues namely England, Germany, Italy, France, and Portugal, shirt sponsorship grew from one deal in the 2002/3 season to 26 in the 2010/11 season [2]. In the hugely popular English Premier League alone, gambling companies that have first team shirt sponsorships grew from four deals in 2008, to six in 2012, and ten in 2017, accounting for half of the 20 English Premier League teams [3].

Soccer is the most popular game in majority of countries in sub Saharan Africa. The European soccer leagues are extremely popular in Africa and therefore soccer sponsors are likely to reach a very wide audience on the continent through their advertisement. The linkage between soccer and sports-betting or gambling advertisement poses potential public health threats to the soccer fans, majority of who are minors or young adults, because of the potential risk of developing problem gambling [4–6] and addiction.

Serious problem gambling also referred to as pathological gambling is recognized as a disorder by the World Health Organization and was included in the Diagnostic and Statistical Manual for Mental

Disorders (DSM-III) in 1980. It was renamed as gambling disorder and classified under the new 'Addictions and Related Disorders' category [7].

The global marketing of sports-betting and alcoholic products together in live sports events such as soccer matches is very common [8] and betting advertisement associates significant moments of sports such as scoring goals with alcohol [9]. Alcohol consumption has been linked to gambling including sports-betting. Several studies have confirmed the association between alcohol consumption and sports-betting [10, 11] and also with smoking or substance abuse [12]. However, most of these studies have been conducted in resource rich settings.

In resource limited settings, because of the popularity of soccer, exposure to sports-betting advertisement is increasing and anecdotal reports suggest that sports-betting may also be on the rise especially among youth in urban and peri-urban areas. Motorcycle taxis commonly referred to as *boda bolas* are operated by a predominantly youthful urban male population, who earn a daily income and may be a soft target for the sports-betting industry. Therefore, the purpose of this study was to establish the frequency and pattern of sports-betting, and the relationship between alcohol consumption and sports-betting among *boda boda* riders in an urban setting in Uganda. We hypothesized that alcohol consumption and cigarette smoking are associated with increased likelihood of sports-betting.

Materials And Methods

Setting

We conducted a cross sectional study in Mbarara municipality located in Mbarara district in western Uganda. The municipality is a mixed urban and peri-urban setting with an estimated population of nearly 100,000 inhabitants. Located 250 km from the capital Kampala, it is the largest urban setting in southwestern Uganda.

Study participants and sampling

The boda boda cyclists are predominantly male, with an insignificant number of female riders. The riders are identifiable by the stage where they operate from. Each stage has a different number of riders ranging from a few to almost 15 in the more crowded areas of the municipality.

We used a multi stage sampling approach to select the stages from which our respondents were picked. In this multi-stage approach, we started at the division (or subcounty), which is the largest geographical unit in the municipality. From each division, we randomly selected one parish, and from each parish we selected three villages. We listed all the stages in the selected villages and randomly selected the stages for participation and selected more stages from the larger divisions. We selected an equal number of cyclists from each stage. Our sample size calculation was adjusted for the design effect arising from potential clustering of riders at a given stage. We collected data using a questionnaire that was translated into the local language for those that were not able to speak English.

Inclusion and exclusion criteria

The *boda boda* cyclists were eligible to participate if they were residing within Mbarara municipality, had been riding for at least 6 months, a period required for them to obtain an official registration from the Municipal commercial officers. Participants were enrolled if they provided individual informed consent to participate in the study. We enrolled participants of 18 to 59 years of age. We did not enroll those who operate on night duty as they were not available for interview during normal working hours.

Measurements

We collected data on social demographics and income. We also collected data on sports-betting, and asked if they had ever been involved in sports-betting, and the frequency of their involvement. The participants in our study do not have access to other forms of gambling other than sports-betting, and therefore sports-betting and gambling are used interchangeably in our description. We collected data on alcohol consumption and used the Alcohol Use Disorders Identification Test (AUDIT) tool, a 10-item screening tool developed by the World Health Organization [13]. We asked participants about their smoking habits, namely whether they had ever smoked and if they had smoked in the past 6 months.

Data analysis

We performed univariate, bivariate, and multivariate analyses. We summarized numeric data using descriptive statistics namely means and standard deviations or medians with interquartile ranges (IQR), and categorical data using frequencies and percentages. We coded sports-betting as a dichotomous outcome. We considered a rider to have engaged in sports-betting if they reported they had been actively involved in sports-betting in the past 30 days. We defined active sports-betting as spending money to complete a transaction in a sports-betting activity. We coded the outcome as "1" when the participant was involved in sports-betting or "0" when not involved.

Alcohol consumption was classified into three categories based on the AUDIT-C score namely low, moderate and high based on the volume of consumption with the smallest consumers in the low category and the largest consumers in the high category. An AUDIT score of 0 to 4 was considered low, 5 to 8 was classified as moderate and any score higher than 8 was considered high or hazardous alcohol use.

We examined categorical variables such as residence by sports-betting status using the Chi-square test. We also examined continuous variables such as age by sports-betting status, and here we used the student's t-test. We regarded test results with probability values (p values) less than 0.05 as statistically significant

The outcome variable was frequent (more than 10%) so the odds ratio (OR) was not an appropriate measure of effect because it would overestimate the strength of the association. [14, 15] Therefore, for the regression analysis, we computed risk ratios (RR) in multivariate analysis, using modified Poisson regression with robust error variance to control for mild violations of the assumptions of Poisson

regression analysis. We reported each RR with 95% confidence intervals (CI). Prior to the multivariate analysis, we checked for multicollinearity and considered variance inflation factor (VIF) greater than 10 as indicator of collinearity.

Human subjects

The study was approved by the Research Ethics Committee at Mbarara University of Science and Technology. Study procedures were adherent to the requirements under the Helsinki declaration. Participants were invited and voluntarily accepted to participate.

Individual informed and written consent was obtained from all participants, and all were adults above 18 years. Consent process was conducted in the local language to enhance understanding.

Results

Socio-demographic characteristics of participants

We enrolled 401 riders and administered interviews to them. The mean age (standard deviation) of the participants was 29.3 ± 5.9 years (median age, 29.0; Interquartile range or IQR 25–32). Table 1 presents a summary of the socio-demographic characteristics of participants stratified by sports-betting status. There was no statistically significant difference in mean age between participants who engaged in sports-betting and those who never: 28.97 ± 5.41 versus 29.42 ± 6.07 , $p = 0.558$. Statistically significant differences by sports-betting were observed with respect to time for end of work day ($p = 0.043$) and alcohol consumption ($p = 0.002$) and smoking status. Riders who end their work day by 7:00 pm are more likely to engage in sports-betting compared to those who end their work day after 7:00 pm. Alcohol consumption and smoking are more frequent among persons who were engaged in sports-betting compared to those who were not. The other variables such as the division of residence, religion, educational level, marital status, number of children, income per day and ownership of a motorcycle taxi did not differ (all $p > 0.05$) between participants who engaged in sports-betting and those that did not. The riders who smoke were more likely to engage in sports-betting, however use of marijuana was not related to sports-betting.

Table 1
Characteristics of motorcycle taxi riders by sports betting status, Mbarara municipality, Uganda

Characteristics	Level	Engaged in sports betting		P value
		No (n = 327)	Yes (n = 74)	
Division of residence	A	110 (33.6)	23 (31.1)	0.185
	B	79 (24.2)	24 (32.4)	
	C	108 (33.0)	17 (23.0)	
	D	30 (9.2)	10 (13.5)	
Mean age (SD)		29.42 (6.07)	28.97 (5.41)	0.558
Religion	Catholic	123 (37.6)	25 (33.8)	0.131
	Protestant	138 (42.2)	38 (51.4)	
	Moslem	45 (13.8)	4 (5.4)	
	Others	21 (6.4)	7 (9.5)	
Educational level	None/primary	204 (62.6)	44 (59.5)	0.618
	Secondary/Tertiary	122 (37.4)	30 (40.5)	
Marital status	Never married	90 (27.5)	18 (24.3)	0.428
	Married	218 (66.7)	54 (73.0)	
	Separated	19 (5.8)	2 (2.7)	
Number of children	≤ 2	211 (64.9)	49 (66.2)	0.977
	3–5	100 (30.8)	22 (29.7)	
	> 5	14 (4.3)	3 (4.1)	
Owns a <i>bodaboda</i>	No	169 (51.8)	40 (55.6)	0.659

Note: 1) * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ at 5% level of significance

		Engaged in sports betting		
	Yes	157 (48.2)	32 (44.4)	
Income per day in dollars	< 5	135 (41.7)	40 (54.1)	0.071
	≥ 5	189 (58.3)	34 (45.9)	
End of work day	Before 7.00 pm	38 (11.8)	16 (21.6)	0.043*
	After 7.00 pm	283 (88.2)	58 (78.4)	
Alcohol consumption	Low	266 (81.3)	49 (66.2)	0.002*
	Moderate	40 (12.2)	21 (28.4)	
	High	21 (6.4)	4 (5.4)	
Ever smoked cigarettes	No	296 (90.5)	52 (70.3)	< 0.001
	Yes	31 (9.5)	22 (29.7)	
Smoked cigarettes in the past 6 months	No	301 (92.0)	58 (78.4)	0.001
	Yes	26 (8.0)	16 (21.6)	
Ever smoked marijuana	No	305 (93.3)	66 (89.2)	0.337
	Yes	22 (6.7)	8 (10.8)	
Ever smoked marijuana in the past six months	No	320 (98.8)	71 (95.9)	0.24
	Yes	4 (1.2)	3 (4.1)	

Note: 1) * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ at 5% level of significance

Frequency And Pattern Of Sports-betting

The results for frequency and pattern of sports-betting are presented in Table 2 below. Of the 401 participants, 74 (18.5%) had engaged in sports-betting in the past 30 days. Among those that reported sports-betting in the past 30 days, 31 or 41.9% engaged less than weekly, 27 (36.5%) engaged weekly while 16 (21.6%) daily or almost daily.

Table 2
Frequency and pattern of sports-betting among motorcycle taxi riders in Mbarara municipality, Uganda

	Category	n (%)
Ever engaged in sports-betting in the past 30 days (n = 401)	No	327 (81.5)
	Yes	74 (18.5)
Pattern of sports-betting in the past 30 days (n = 74)	Daily or Almost daily	16 (21.6)
	Weekly	27 (36.5)
	Less than weekly	31 (41.9)

Factors Associated With Sports-betting

We present the results of the regression analysis in Table 3. In the unadjusted analysis, data show that earning five or more dollars per day was not significantly associated with betting (Unadjusted risk ratio or uRR, 0.67; 95% CI, 0.44–1.01), although marginally significant. Ending work day after 7.00 pm (uRR, 0.55; 95% CI, 0.36–0.93) was associated with lower likelihood of sports-betting. Alcohol consumption was significantly associated with sports-betting. There was a more than 2 fold increase in the likelihood of sports-betting among those in the moderate alcohol consumption category (uRR, 2.21; 95% CI, 1.44–3.41) compared to those in the low consumption category as the referent. The association was not significant among those in the high consumption category (uRR, 1.03; 95% CI, 0.40–2.62). Smoking was significantly associated with sports-betting uRR 2.36, 95% 1.50, 3.71).

After adjusting for potential confounders, riders who end the work day after 7.00 pm were 38% (adjusted risk ratio or aRR, 0.62; 95% CI, 0.39–0.99) less likely to engage in sports-betting compared to those who end the day before 7 pm. We observed an inverted U relationship between alcohol consumption and sports-betting. Riders in the moderate alcohol consumption category were nearly 2 times more likely to engage in sports-betting compared to those in the low consumption category (aRR, 1.86; 95% CI, 1.18–2.94). However, among those in the high consumption category, the relationship was not significant. Smoking remained significantly associated with sports-betting after adjustment for all other variables in the model with aRR 1.85 (95% CI 1.13, 3.01). Income earned per day was not significantly related to sports-betting but those who ended their work day after 7 pm were less likely to engage in sports-betting compared to those who end their work day earlier.

Table 3
Modified poisson regression analysis of the relationship between alcohol and sports-betting among motorcycle taxi riders, Mbarara municipality, Uganda

Modified Poisson regression analysis					
Characteristics	Level	uRR	95% CI	aRR	95% CI
Income per day/dollars	< 5	1		1	
	≥ 5	0.67	(0.44,1.01)	0.74	(0.49,1.11)
End of work day	Before 7.00 pm	1		1	
	After 7.00 pm	0.57*	(0.36,0.92)	0.62*	(0.39,0.99)
Smoked cigarettes in last 6 months	No	1		1	
	Yes	2.36***	(1.50,3.71)	1.85*	(1.13,3.01)
Alcohol consumption	Low	1		1	
	Moderate	2.21***	(1.44,3.41)	1.86**	(1.18,2.94)
	High	1.03	(0.40,2.62)	0.97	(0.37,2.58)

Note: 1) * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ at 5% level of significance; 2) 95% confidence intervals for risk ratio (RR) in brackets; 3) ARR: Adjusted Risk Ratio; 4) URR: Unadjusted Risk Ratio; 5) Risk Ratios are exponentiated coefficients.

Discussion

Our data show that almost 20% of the motorcycle taxi riders have been involved in sports-betting in the most recent month, and that among those that engage in sports-betting, about one fifth of them engage daily or nearly daily. Our data show that riders who stop work earlier are more likely to engage in gambling, but most importantly, alcohol consumption and cigarette smoking were significantly associated with gambling. Income earned per day is unrelated to the likelihood to get engaged in sports-betting. Our study is one of the few from sub Saharan Africa that have examined the association between alcohol consumption and gambling.

The frequency of sports-betting in our study population is comparable to that from a study in the United States [16], although this was a student population. However, the frequency in our study population is only half of that demonstrated in a South Africa population [17]. This study in South Africa was conducted among a general adult population, mostly low income poor population. Our study population consists of low income earners as well but younger. However, the frequency gambling in South Africa is much higher and may be because gambling is well established there, since the 1990s and with a variety of options unlike in Uganda. In our study population, sports-betting was the only form of gambling

available to the riders. Other forms of gambling such as casinos are available in the capital Kampala, but are likely to be accessed by the higher income earners as the costs of gambling in the casinos are likely to be higher.

Our findings of the association between alcohol consumption and gambling are in agreement with those of a study in South Africa among a population of gamblers [18]. The results confirm our primary hypothesis and agree with several other studies outside the African continent. We observed an inverted U association, indication that among the heavy drinkers, there was no association with sports-betting. It is not clear whether alcohol is the chicken or the egg in the pathway and what the causal mechanism may be. Studies suggest alcohol drinking may increase the propensity to gamble [19]. There is a possibility that heavy drinkers lose their income to drinking and have limited resources left to gamble.

Our study also shows a significant association between cigarette smoking and sports-betting. This finding fits our hypothesis and also agrees with findings from other studies [20]. Alcohol and nicotine in the cigarettes are both substances of addiction and their close association with gambling, which is also addictive is not unexpected. Our data adds to the limited existing evidence of the link between the use of alcohol and cigarettes with gambling in sub Saharan Africa [18]. For the continent, these results are significant because of the close association of all these factors with mental health conditions, which are much neglected. Alcohol, substance abuse, anxiety and depression have all been found to be highly prevalent among persons with problem gambling [21]. Marijuana was not commonly used and our data did not provide sufficient opportunity to explore its association with sports-betting.

It is important to note that we conducted our study among motor cycle taxi riders, a sub group of the population that comprises predominantly young male Ugandans. We chose this group because they represent a large proportion of urban male youth who earn a living from a meager daily income. Youth unemployment is very high in Uganda with more than 60% of youth aged between 18 and 35 years facing unemployment. Motorcycle taxis provide an option for them to earn a living. These results are therefore generalizable to a large proportion of out of school youth in Uganda.

Although our study provides unique data from sub Saharan Africa, it has some weaknesses. We did not measure the severity of the gambling. There are well established tools to do this such as the Gambling severity index score. Future studies should consider including the measurement of the grade of severity of gambling. The strength of our study is that it presents data in a field that is under researched in sub Saharan Africa. The finding of an inverted U relationship between alcohol consumption and sports-betting is novel and should be further examined and replicated in other studies. Although several studies have shown the association between substance abuse and gambling [16, 22, 23], very few such studies have been done in resource limited settings.

Conclusions And Recommendations

In conclusion, our study has shown that gambling via sports-betting is very common in this low income category group that represents a large proportion of male youth and young adults in Uganda. There is

concern that sports-betting is on the rise and some of the participants are at risk for developing problem gambling and addiction. Our findings have important implications. There is a need to put in place structures to regulate the users to prevent overindulgence. This will be a challenge because sports-betting services can be accessed via mobile phones by users in the privacy of their homes. In addition, sports-betting is legal in Uganda and many other countries in sub Saharan Africa and the industry is lucrative and contributes revenue to the government coffers through taxation. There is a likelihood that interventions to regulate access to sports-betting may be met with resistance from certain stakeholders that benefit from the business. Lastly, given the close association between sports-betting and alcohol consumption, interventions should be co-packaged to deal with the two public health challenges concurrently.

Declarations

Acknowledgements

This work is part of a Masters of Public Health dissertation for SCN. The authors wish to thank the First Mile Program at the Department of Community Health for providing financial support to collect the data.

Author contributions

FB conceived the idea. SCN and FB supervised the data collection, SCN and JI did the data entry, analysis and drafted the tables. FB did the first draft of the manuscript. All authors approved the final version of the manuscript

Conflicts of interest: None to declare

Human subjects: The study was approved by the Research Ethics Committee at Mbarara University of Science and Technology. Study procedures were adherent to the requirements under the Helsinki declaration. Participants were invited and voluntarily accepted to participate.

Consent: Individual informed and written consent was obtained from all participants, who were all adults above 18 years. Consent process was conducted in the local language to enhance understanding.

Data and/or Code availability: Data are available upon reasonable request and with approval from the Research Ethics committee

References

1. Bunn C, Ireland R, Minton J, Holman D, Philpott M, Chambers S. Shirt sponsorship by gambling companies in the English and Scottish Premier Leagues: global reach and public health concerns. *Soccer society*. 2019;20(6):824–35.
2. Foley-Train J. **Sports betting: Commercial and integrity issues.** *Report prepared for the Association of British Bookmakers, European Gaming and Betting Association.* European Sport Security Association

Remote Gambling Association Retrieved January. 2014;21:2015.

3. Lopez-Gonzalez H, Griffiths MD. Betting, Forex Trading, and Fantasy Gaming Sponsorships-a Responsible Marketing Inquiry into the 'Gambification' of English Football. International journal of mental health addiction. 2018;16(2):404–19.
4. Binde P. **Gambling advertising: A critical research review**. In.: Responsible Gambling Trust; 2014.
5. Hanss D, Mentzoni RA, Griffiths MD, Pallesen S. The impact of gambling advertising: Problem gamblers report stronger impacts on involvement, knowledge, and awareness than recreational gamblers. Psychology of addictive behaviors: journal of the Society of Psychologists in Addictive Behaviors. 2015;29(2):483–91.
6. Binde P, Romild U. Self-Reported Negative Influence of Gambling Advertising in a Swedish Population-Based Sample. Journal of gambling studies. 2019;35(2):709–24.
7. Mayes R, Horwitz AV. DSM-III and the revolution in the classification of mental illness. J Hist Behav Sci. 2005;41(3):249–67.
8. Sharman S, Ferreira CA, Newall PWS. **Exposure to Gambling and Alcohol Marketing in Soccer Matchday Programmes**. *Journal of gambling studies* 2019.
9. Lopez-Gonzalez H, Estévez A, Jiménez-Murcia S, Griffiths MD. Alcohol Drinking and Low Nutritional Value Food Eating Behavior of Sports Bettors in Gambling Advertisements. International journal of mental health addiction. 2018;16(1):81–9.
10. Blankenship J, Starling R, Woodall WG, May PA. Gambling and alcohol use: trends in the state of New Mexico from 1996–1998. Journal of gambling studies. 2007;23(2):157–74.
11. French MT, Maclean JC, Ettner SL. Drinkers and bettors: investigating the complementarity of alcohol consumption and problem gambling. Drug Alcohol Depend. 2008;96(1–2):155–64.
12. Weinberger AH, Franco CA, Hoff RA, Pilver C, Steinberg MA, Rugle L, Wampler J, Cavallo DA, Krishnan-Sarin S, Potenza MN. Cigarette smoking, problem-gambling severity, and health behaviors in high-school students. Addictive behaviors reports. 2015;1:40–8.
13. Babor TF, de la Fuente JR, Saunders J, Grant M. **The Alcohol Use Disorders Identification Test: Guidelines for use in**. Primary care 2001.
14. Schmidt OC, Kohlmann T. When to use the odds ratio or the relative risk? Int J Public Health. 2008;53:165–7.
15. Spiegelman D, Hertzmark E. Easy SAS calculations for risk or prevalence ratios and differences. Am J Epidemiol. 2005;162(3):199–200.
16. Zhai ZW, Duenas GL, Wampler J, Potenza MN. **Gambling, Substance Use and Violence in Male and Female Adolescents**. *Journal of gambling studies* 2020.
17. Dellis A, Spurrett D, Hofmeyr A, Sharp C, Ross D. Gambling participation and problem gambling severity among rural and peri-urban poor South African adults in KwaZulu-Natal. Journal of gambling studies. 2013;29(3):417–33.

18. Skaal L, Sinclair H, Stein DJ, Myers B. Problem Gambling Among Urban and Rural Gamblers in Limpopo Province, South Africa: Associations with Hazardous and Harmful Alcohol Use and Psychological Distress. *Journal of gambling studies*. 2016;32(1):217–30.
19. Barrett SP, Collins P, Stewart SH. The acute effects of tobacco smoking and alcohol consumption on video-lottery terminal gambling. *Pharmacol Biochem Behav*. 2015;130:34–9.
20. Hammond CJ, Pilver CE, Rugle L, Steinberg MA, Mayes LC, Malison RT, Krishnan-Sarin S, Hoff RA, Potenza MN. An exploratory examination of marijuana use, problem-gambling severity, and health correlates among adolescents. *Journal of behavioral addictions*. 2014;3(2):90–101.
21. Sharp C, Dellis A, Hofmeyr A, Kincaid H, Ross D. First Evidence of Comorbidity of Problem Gambling and Other Psychiatric Problems in a Representative Urban Sample of South Africa. *Journal of gambling studies*. 2015;31(3):679–94.
22. Ethier AR, Kim HS, Hodgins DC, McGrath DS. **High Rollers: Correlates of Problematic Cocaine Use Among a Community Sample of Gamblers**. *Journal of gambling studies* 2020.
23. Dufour M, Nguyen N, Bertrand K, Perreault M, Jutras-Aswad D, Morvannou A, Bruneau J, Berbiche D, Roy É. Gambling Problems Among Community Cocaine Users. *Journal of gambling studies*. 2016;32(3):1039–53.