

# Impact of Galcanezumab on Total Pain Burden: Findings From Phase 3 Randomized, Double-Blind, Placebo-Controlled Studies in Patients with Episodic or Chronic Migraine (EVOLVE-1, EVOLVE-2, and REGAIN Trials)

**Jessica Ailani**

Georgetown University

**J Scott Andrews**

Eli Lilly and Company

**Mallikarjuna Rettiganti**

Eli Lilly and Company

**Robert A Nicholson** (✉ [Robert.Nicholson@lilly.com](mailto:Robert.Nicholson@lilly.com))

Eli Lilly and Company

---

## Research article

**Keywords:** Migraine, chronic migraine, burden of illness, galcanezumab, headache, pain, prevention, clinical trial

**Posted Date:** August 20th, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-60145/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published on October 17th, 2020. See the published version at <https://doi.org/10.1186/s10194-020-01190-7>.

# Abstract

## BACKGROUND

Focus on the frequency of migraine pain may undervalue the total burden of migraine as pain duration and severity may present unique, additive burden. A composite measure of total pain burden (TPB; frequency, severity, and duration) of migraine may provide a more comprehensive characterization of pain burden and treatment response in patients with episodic migraine (EM) or chronic migraine (CM). The impact of galcanezumab 120 mg once-monthly injection versus placebo on TPB among patients with EM or CM was analyzed.

## METHODS

Patients from randomized, double-blind, placebo-controlled episodic (6-month) and chronic migraine (3-month) studies received once-monthly subcutaneous injection of galcanezumab 120 mg once-monthly or placebo. Total pain burden for a given month was calculated as severity-weighted duration by multiplying duration (hours) and maximum pain severity (0 = none, 1 = mild, 2 = moderate, 3 = severe) of migraine for each day and summing these over the days in a month. Least square (LS) mean change from baseline in monthly TPB across Months 1–6 (EM) and Months 13 (CM) were compared post hoc using a mixed-model repeated measures model. Correlation of the Migraine Specific Quality of Life Questionnaire (MSQ) and Migraine Disability Assessment Scale (MIDAS) to TPB at baseline was assessed.

## RESULTS

At baseline, the duration of migraine on a given migraine headache day accounted for the greatest unique proportion of variability (EM, 57.4% and CM, 61.1%) to the TPB after adjusting for frequency of migraine headache days and maximum pain severity. The LS mean decreases from baseline in monthly TPB across months were greater with galcanezumab than placebo for patients with EM (68.6 versus 36.2) and CM (102.6 versus 44.4). The average percent reduction of TPB from baseline was significantly greater with galcanezumab compared with placebo in patients with EM (50.8% versus 17.2%) and CM (29.7% versus 11.0%). Total pain burden in patients with EM and CM correlated with MSQ total score ( $r = 0.35$  and  $r = -0.37$ ) and MIDAS ( $r = 0.34$  and  $r = 0.32$ ).

## CONCLUSIONS

Greater reduction in TPB was seen in patients with EM and CM treated with galcanezumab 120 mg once-monthly injection relative to placebo. Discussing TPB supports patient-centric conversations regarding treatment expectations when clinicians are evaluating options for migraine prevention.

# TRIAL REGISTRATION:

ClinicalTrials.gov: #NCT02614183 (I5Q-MC-CGAG; EVOLVE-1), #NCT02614196 (I5QMCCGAH; EVOLVE-2), and #NCT02614261 (I5Q-MC-CGAI; REGAIN) – all 3 trials were registered on 23 November 2015.

## Background

It is well established that migraine can interfere with the occupational, educational, household, family, and social facets of daily life [1–9]. Consideration of the burden of migraine often focuses solely on the frequency of migraine pain; doing so may undervalue the total burden of migraine as pain duration and severity may present unique, additive burden [3, 6, 8, 9]. Total pain burden of migraine has been conceptualized as a composite measure involving frequency, duration, and severity. However, as novel therapies designed for the preventive treatment of migraine have become available, little research has focused on total pain burden. Evaluating this composite measure of pain may provide more comprehensive characterization of the total pain burden of migraine and inform treatment expectations when evaluating options for migraine prevention.

Galcanezumab is a humanized IgG4 monoclonal antibody that binds calcitonin gene-related peptide (CGRP) and prevents its biological activity without blocking the CGRP receptor and is indicated for the preventive treatment of migraine [10]. The efficacy of galcanezumab as a preventive treatment for migraine has been established in patients with episodic or chronic migraine across three randomized, double-blind, placebo-controlled, Phase 3 studies (EVOLVE1, EVOLVE-2, and REGAIN). In those studies, galcanezumab-treated patients with episodic migraine experienced 4.3 to 4.7 fewer migraine headache days/month (versus 2.3 to 2.8 with placebo), and patients with chronic migraine had 4.8 fewer migraine headache days/month (versus 2.7 with placebo) [11–13].

The efficacy of galcanezumab for reduction of migraine frequency is established. However, it is unclear whether galcanezumab impacts total pain burden of migraine over time. Indication of this would provide greater clarity to providers regarding the overall potential clinical value of galcanezumab. In this post hoc study, total pain burden was compared between those taking galcanezumab 120 mg once-monthly injection (with an initial 240 mg loading dose) and placebo among patients with episodic or chronic migraine.

## Methods

### Study design

This post hoc analysis of Phase 3, double-blind, placebo-controlled studies in adult patients utilized data from two, 6-month episodic migraine studies pooled (EVOLVE-1 and EVOLVE-2) and one 3-month chronic migraine study (REGAIN) to analyze the impact of galcanezumab 120 mg once-monthly injection relative to placebo on total pain burden among patients with episodic migraine or chronic migraine [11-13].

Patients enrolled in the studies were 18 to 65 years of age and had a diagnosis of migraine (with or without aura) for at least one year prior to enrollment and onset prior to 50 years of age [11-14].

In all studies, patients entered a prospective baseline period of 30 to 40 days in which they completed an electronic patient-reported outcome diary to record the occurrence of headaches, headache duration, headache features, severity of headache, and use of headache medication. Eligible patients were randomized (1:1:2) to subcutaneous injections of galcanezumab 120 mg/month (following an initial 240 mg loading dose), galcanezumab 240 mg/month, or placebo. During all three studies, patients were permitted to continue acute migraine medications including triptans, ergots, nonsteroidal anti-inflammatory drugs, aspirin, and acetaminophen. Opioid- and barbiturate-containing medications were limited to 3 days/month, and only one corticosteroid injection was allowed during any period. Exclusion criteria of note for all three studies included, but was not limited to, prior exposure to any CGRP antibody or any therapeutic antibody 12 months prior to screening, using opioids or barbiturates more than twice per month, persistent daily headache, cluster headache, head or neck trauma within the past 6 months, possible posttraumatic headache, primary headache, or a medical or psychiatric illness that would preclude study participation [11-13]. The study protocols were reviewed and approved by the appropriate institutional review board for each of the study sites. The studies were conducted according to Good Clinical Practice and the Declaration of Helsinki guidelines. Patients provided written informed consent before undergoing study procedures. The studies are registered with ClinicalTrials.gov (NCT02614183, NCT02614196, and NCT02614261).

## Outcomes and statistical methods

The primary outcome of the EVOLVE-1, EVOLVE-2, and REGAIN studies was the overall mean change from baseline in the number of monthly migraine headache days during the double-blind treatment phase. This post hoc analysis evaluated the galcanezumab 120 mg and placebo treatment arms of the 3 studies. The objective of these analyses was to derive and compare between treatment groups, a “total pain burden” measure that incorporates: *frequency* of migraine headache days in a month, *duration* of migraine headache on a given day, and maximum *severity* of migraine headache on a given day. The total pain burden for a given month (severity-weighted duration) was calculated by multiplying duration (hours) of migraine headache and maximum pain severity (0=none, 1=mild, 2=moderate, 3=severe) for each migraine headache day and summing these over the days in a month. As an example, consider a patient who has 2 days of migraine headache in a month. The patient reports 2 hours of migraine headache on Day 1, which is of mild severity (score = 1) and 3 hours of migraine headache on Day 2, which is of moderate severity (score = 2). The total pain burden score for that month would be calculated as the sum of (2 hours x 1) and (3 hours x 2) which equals 8 severity-weighted hours of total pain burden. The change from baseline in monthly total pain burden measure over the double-blind period was analyzed for both episodic (6 months) and chronic migraine (3 months) studies using a mixed-model repeated measures (MMRM) model. The analysis of the episodic migraine studies included the following fixed effect variables: baseline, treatment, month, study indicator (EVOLVE-1 or EVOLVE-2), pooled region/country (nested within study), and the treatment by month, and baseline by month interaction

effects. For the chronic migraine study, the fixed effect variables included: baseline, treatment, month, pooled country, baseline medication overuse (yes/no), concurrent prophylaxis use (yes/no), the interaction effects of treatment by month, and baseline by month. A marginal unstructured covariance structure was assumed to account for the correlation induced due to repeated measures on patients. The mean percentage change in total pain burden was analyzed using an MMRM model, as described earlier, for the change from baseline analyses. A similar MMRM modeling approach was also used to analyze change from baseline in each of the individual components of frequency of migraine headache days, duration per migraine headache day, and severity.

The Type II squared semi-partial correlation of each individual component within the total pain burden measure at baseline was obtained from the following regression model on the log transformed variables:  $\log(\text{total pain burden}) = \log(\text{monthly migraine headache days}) + \log(\text{mean hours per migraine headache day}) + \log(\text{severity of remaining migraine headache days})$ . These provide the additional or unique proportion of variability explained by each component in its ability to predict total pain burden after adjusting for the other two components.

To explore the construct validity of the total pain burden measure, the Spearman's rank correlation was used to determine the degree of correlation of the Migraine Specific Quality of Life Questionnaire (MSQ) and Migraine Disability Assessment Scale (MIDAS) to total pain burden at baseline. The MSQ is a self-administered instrument that evaluates the physical and emotional limitations of specific concern to patients with migraine [15]. The MIDAS quantifies days with headache-related disability across 5 areas over the last 3 months (90 days) [16].

## Sensitivity analyses

The pain severity scores (0=none, 1=mild, 2=moderate, 3=severe) can be thought of as arbitrary. Therefore, a sensitivity analysis was performed by calculating (and comparing between groups) the total pain burden score using the square root of severity score in the calculation instead of severity (that is, using scores of 1=mild, 1.414=moderate, and 1.732=severe) and using the square of the severity score in the calculation instead of severity (that is, using scores of 1=mild, 4=moderate, and 9=severity). To further understand whether the changes observed in total pain burden can be fully explained by the change observed in frequency of migraine headache days, the original MMRM model analyzing change from baseline in total pain burden was repeated after including an additional time-varying covariate of the change from baseline in the number of monthly migraine headache days.

## General considerations

Baseline demographics and descriptive summaries were provided for continuous variables using means and standard deviation and/or median and quartiles; for categorical variables, frequency and percentages were used. Treatment effects from MMRM models were expressed as least squares (LS)

means and 95% confidence intervals (CI). All statistical tests done were two-sided assuming a significance level of 5%. All statistical analyses were done using SAS<sup>®</sup> Enterprise Guide version 7.1. All analyses done were post hoc and results should be considered exploratory.

## Results

### Patient disposition

A total of 1338 patients with episodic migraine (placebo n=894, galcanezumab 120 mg n=444) and 836 patients with chronic migraine (placebo n=558, galcanezumab 120 mg n=278) were considered for this post hoc analysis. The general baseline demographics and characteristics were balanced between treatment groups in the two episodic and one chronic migraine studies and have been previously well-described [11-13]. Relevant characteristics for the episodic and chronic migraine populations are shown in Table 1. Among the three studies, patients were a mean age of 41 years, and the majority were female, White, and from North America. The mean number of migraine headache days/month among patients with episodic migraine was 9.1 days and among patients with chronic migraine was 19.4 days. At baseline, both populations had a mean MIDAS total score indicative of severe disability and an MSQ total score similar to patients presenting for migraine care. In the prospective baseline period (30 to 40 days), the mean total pain burden across both treatment groups was 124.3 severity-weighted hours for patients with episodic migraine and 320.4 severity-weighted hours for patients with chronic migraine.

### Total pain burden

The LS mean reduction from baseline in monthly total pain burden, was greater in galcanezumab-treated patients compared with placebo-treated patients (Figure 1). Starting at Month 1 in both the episodic and chronic migraine populations, the mean reductions from baseline were greater with galcanezumab compared with placebo and the same pattern was observed in all subsequent months of treatment (all  $p < 0.001$ ) (data not shown). The overall monthly LS mean severity-weighted hours reduction from baseline across Months 1 to 6 for episodic migraine and across Months 1 to 3 for chronic migraine were greater for the galcanezumab groups compared with the placebo groups ( $p < 0.001$ ; Figure 1).

The average percent reduction from baseline in total pain burden in patients with episodic migraine across 6 months was greater with galcanezumab treatment (50.8%) compared with placebo (17.2%); mean difference was 33.6% (95% CI 41.3, 25.9;  $p < 0.001$ ). Likewise, for patients with chronic migraine, the average percent reduction across 3 months was greater with galcanezumab (29.7%) compared with placebo (11.0%); mean difference was 18.7 (95% CI 25.6, 11.7;  $p < 0.001$ ). As illustrated in Supplemental Material animations, for both the episodic and chronic migraine populations treated with galcanezumab, the percentage reduction in total pain burden was significantly greater than placebo starting at Month 1 ( $p < 0.001$ ) and the significantly greater percentage reduction was maintained at each subsequent month ( $p < 0.001$ ).

### Contribution of components to total pain burden

The duration of migraine on a given migraine headache day accounted for the greatest unique proportion of variability (episodic migraine studies, 57.4% and chronic migraine study, 61.1%) to the baseline total pain burden after adjusting for the frequency of migraine headache days and maximum pain severity (Table 2). In other words, the *duration* (hours) of migraine pain on the days when a patient had a migraine headache in a month contributed the most variability to total pain burden rather than the *number* of days with migraine headache in a month.

Results from the sensitivity analyses related to pain severity scaling revealed that total pain burden remained significantly lower for the galcanezumab group compared with placebo even when using the square root of severity scores or the square of severity scores in the calculation of total pain burden. Thus, altering the severity rating scores on the calculation of total pain burden did not considerably alter the findings.

### **Total pain burden correlations to MSQ and MIDAS**

The MSQ and MIDAS scores at baseline were associated to total pain burden in the episodic ( $r = -0.35$  and  $r = 0.34$ , respectively;  $p < 0.0001$ ) and chronic migraine ( $r = -0.37$  and  $r = 0.32$ , respectively;  $p < 0.0001$ ) populations.

## **Change from baseline of individual components**

The LS mean change for the individual components of total pain burden overall monthly across Months 1 to 6 for episodic migraine and Months 1 to 3 for chronic migraine are shown in Table 3. For both populations, the reductions from baseline in the frequency of, duration per, and the severity of the migraine headache days were significantly greater with galcanezumab treatment compared to placebo.

The sensitivity analysis conducted to explore the residual impact on total pain burden, even after controlling for the reduction of migraine headache day frequency, demonstrated that for both episodic and chronic migraine populations, the reduction in total pain burden was still greater in the galcanezumab groups compared with the placebo groups. In other words, galcanezumab treatment had impact on the other components of total pain burden (duration and severity) and supports findings from the analysis of the individual components of total pain burden (Table 3). Shown in Figure 2 is the total pain burden (monthly severity-weighted duration in hours) adjusted for change in migraine headache days for patients with episodic and chronic migraine across months in the double-blind phase. The reduction in total pain burden was significantly greater in the galcanezumab groups compared with the placebo groups.

## **Discussion**

Treatment with galcanezumab 120 mg once-monthly injection (following a 240 mg initial loading dose) resulted in greater reduction of total pain burden relative to placebo. This study, combined with previous

studies demonstrating the benefit of galcanezumab for reducing monthly migraine headache days, provides a more holistic view of the effect galcanezumab can have in patients with episodic or chronic migraine.

A composite measure of total pain burden produces a measure that may be more aligned to the personal pain experience in migraine relative to solely assessing frequency, duration, and severity as independent components. The focus on pain frequency as the sole metric for evaluating treatment “benefit” may be driven, in part, by regulatory bodies’ recommendations/requirements regarding what outcomes “matter” when considering the potential benefits of preventive treatment for migraine. Clinically, total pain burden may better reflect what clinicians and patients discuss regarding the individual’s pain experience and could prove useful to further patient-centric discussions regarding treatment expectations when clinicians are evaluating options for migraine prevention. In this current analysis, all three components of total pain burden independently contributed to total pain burden; duration of migraine headache per day and number of monthly migraine headache days were the two most influential components. The current findings suggest that assessing total pain burden in a research or clinical setting allows for a more robust evaluation of the potential benefit of a preventive treatment for migraine (in this case, galcanezumab) in reducing the overall pain experience.

The evidence presented here suggests that galcanezumab could benefit a patient beyond reduction in migraine headache days per month. This observation may be especially relevant for those with chronic migraine as these patients will likely to continue to experience migraine headache more frequently than those with episodic migraine. If those remaining migraine headache days feature pain that lasts fewer hours and/or is less severe, then patients may be experiencing benefit that goes beyond those identified through unidimensional frequency counts.

The results support the hypothesis that total pain burden is a composite of various components that contribute unique information. Further, total pain burden is not simply a proxy to the number of migraine headache days since the reduction of total pain burden was greater with galcanezumab relative to placebo even after accounting for the reduction in monthly migraine headache days. This implies that the patient is experiencing an added benefit beyond experiencing fewer migraine headache days per month. Moreover, the positive correlation of total pain burden with MSQ and MIDAS further supports the construct validity of this composite measure and that disability is significantly correlated with total pain burden.

## **Limitations**

This study provides data regarding total pain burden in which published research pertaining to migraine is sparse. The outcomes from this study are limited by the post hoc nature of the evaluation and the clinical trial setting that restricts generalizability to the real-world population of patients with migraine. Another limitation of note may be the 4point scale used to assess severity and its use in calculating the

total pain burden given that a 4-point scale may have less sensitivity in detecting changes in severity than an 11point scale.

The study outcomes are strengthened by the large sample size for both the populations of patients with episodic and chronic migraine and by the controlled nature of the study that allows for comparisons between groups providing greater confidence in considering the veracity of observed outcomes. Additionally, in the current analysis of total pain burden, we calculated severity-weighted hours such that 1 hour of moderate pain was considered equally burdensome as 2 hours of mild pain (similarly, this calculation considered 1 hour of severe pain as equally burdensome as 3 hours of mild pain). Others may consider different weighting algorithms to be more representative of total pain burden. However, our sensitivity analysis showed that changes in severity weights did not significantly alter the results when assessing the impact of galcanezumab on total pain burden.

## Conclusions

Greater reduction in total pain burden was seen in patients with episodic or chronic migraine treated with galcanezumab 120 mg once-monthly injection relative to placebo. This allows for a more robust evaluation of the potential benefit of preventive treatments (e.g., galcanezumab) in reducing the overall pain experience. Moreover, the discussion of total pain burden is well situated to support patient-centric discussions regarding treatment expectations when clinicians are evaluating options for migraine prevention.

## Abbreviations

<b>CGRP</b>	<b>Calcitonin gene-related peptide</b>
CI	Confidence intervals
EVOLVE-1, EVOLVE-2	6-month, randomized, placebo-controlled studies (identical design) of galcanezumab in patients with episodic migraine
LS	Least squares
MIDAS	Migraine Disability Assessment Scale
MMRM	Mixed-model repeated measures
MSQ	Migraine Specific Quality of Life Questionnaire
REGAIN	3-month randomized, placebo-controlled study of galcanezumab in patients with chronic migraine

## Declarations

Data were presented in part as a poster presentation at the 72nd American Academy of Neurology (AAN) Virtual 2020 Meeting; May 18, 2020.

Ethics approval and consent to participate: Stated in Study Design Section of Methods, “*Study protocols were reviewed and approved by the appropriate institutional review board for each of the study sites. The studies were conducted according to Good Clinical Practice and the Declaration of Helsinki guidelines. Patients provided written informed consent before undergoing study procedures.*”

Consent for publication: Not applicable

Availability of data and materials: The datasets generated and/or analyzed during the current study are not publicly available due intellectual asset protection but are available from the corresponding author upon reasonable request.

Competing interests:

Jessica Alani, *CME programing*: Miller Communications (CME content and speaker fee), Avent (CME content and speaker fee), Peer View (CME speaker fee), Forefront (CME content and speaker fee); *Consultant*: (Honoraria for independent consulting) Alder, Amgen, Allergan, Electrocore, Eli Lilly and Company, Promius, Teva, Impel, Satsuma, Zosano, Revance, Alpha Sites Consulting, Neurodiem; *Research Support*: American Migraine Foundation, Allergan, Biohaven, Eli Lilly and Company; *Speaker’s Bureau*: (Honoraria for promotional speaking) Allergan, Amgen, Electrocore, Eli Lilly and Company, Promius, Teva; *Other*: Current pain and headache reports, section editor.

J. Scott Andrews, Employee of Eli Lilly and Company, and/or one of its subsidiaries, Indianapolis, IN, USA

Mallikarjuna Rettiganti, Employee of Eli Lilly and Company, and/or one of its subsidiaries, Indianapolis, IN, USA

Robert A. Nicholson, Employee of Eli Lilly and Company, and/or one of its subsidiaries, Indianapolis, IN, USA

Study funding: Eli Lilly and Company, Indianapolis, IN, USA funded the design of the study and the collection, analysis, and interpretation of data and the medical writing and editorial support the manuscript.

Author contributions: All authors made substantial contributions to the study conception and design. SW, MR, and RN made major contributions to data collection and MW performed the statistical analysis. All authors were major contributors in the writing and critical review and interpretation of the manuscript. All authors read and approved the final manuscript.

Acknowledgments: Medical writing support was provided by Millie Hollandbeck (Synchrogenix, a Certara Company) and was funded by Eli Lilly and Company, Indianapolis, IN, USA.

## References

1. Hawkins K, Wang S, Rupnow MF. Indirect cost burden of migraine in the United States. *J Occup Environ Med.* 2007;49:368-74.
2. Blumenfeld A, Varon SF, Wilcox TK, Buse DC, Kawata AK, Manack A, et al. Disability, HRQoL and resource use among chronic and episodic migraineurs: results from The International Burden of Migraine Study. *Cephalalgia.* 2011;31:301-5.
3. Payne KA, Varon SF, Kawata AK, Yeomans K, Wilcox TK, Manack A, et al. The International Burden of Migraine Study (IBMS): study design, methodology, and baseline cohort characteristics. *Cephalalgia.* 2011;31:1116-30.
4. Bloudek LM, Stokes M, Buse DC, Wilcox TK, Lipton RB, Goadsby PJ, et al. Cost of healthcare for patients with migraine in five European countries: results from The International Burden of Migraine Study (IBMS). *J Headache Pain.* 2012;13:361-78.
5. Buse DC, Manack AN, Fanning KM, Serrano D, Reed ML, Turkel CC, Lipton RB. Chronic migraine prevalence, disability, and sociodemographic factors: results from the American Migraine Prevalence and Prevention Study. *Headache* 2012.52:1456-70.
6. Adams AM, Serrano D, Buse DC, Reed ML, Marske V, Fanning KM, Lipton RB. The impact of chronic migraine: the Chronic Migraine Epidemiology and Outcomes (CaMEO) Study methods and baseline results. *Cephalalgia.* 2015;35:563-78.
7. GBD 2015 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet.* 2016;388:1545-602.
8. Dodick DW. Migraine. *Lancet.* 2018;391:1315-30.
9. Martelletti P, Schwedt TJ, Lanteri-Minet M, Quintana R, Carboni V, Diener HC, et al. My Migraine Voice survey: a global study of disease burden among individuals with migraine for whom preventive treatments have failed. *J Headache Pain.* 2018;19:115.
10. Emgality [package insert]. Indianapolis, IN: Eli Lilly and Company; 2018.
11. Detke HC, Goadsby PJ, Wang S, Friedman DI, Selzler KJ, Aurora SK. Galcanezumab in chronic migraine: the randomized, double-blind, placebo-controlled REGAIN study. *Neurology.* 2018;91:e2211-21.
12. Skljarevski V, Matharu M, Millen BA, Ossipov MH, Kim BK, Yang JY. Efficacy and safety of galcanezumab for the prevention of episodic migraine: results of the EVOLVE-2 phase 3 randomized controlled clinical trial. *Cephalalgia.* 2018;38:1442-54.
13. Stauffer VL, Dodick DW, Zhang Q, Carter JN, Ailani J, Conley RR. Evaluation of galcanezumab for the prevention of episodic migraine: the EVOLVE-1 randomized clinical trial. *JAMA Neurol.* 2018;75:1080-88.
14. Headache Classification Committee of the International Headache Society (IHS). The International Classification of Headache Disorders, 3rd edition (beta version). *Cephalalgia* 2013;33:629-808.
15. Jhingran P, Osterhaus JT, Miller DW, Lee JT, Kirchoetfer L. Development and validation of the Migraine Specific Quality-of-Life questionnaire. *Headache.* 1998;38:295-302.

16. Stewart WF, Lipton RB, Kolodner K, Liberman J, Sawyer J. Reliability of the migraine disability assessment score in a population-based sample of headache sufferers. *Cephalalgia* 1999;19:107-14.

## Tables

Table 1. Summary of Baseline Patient Demographics and Characteristics

<b>Baseline Demographics and Disease Characteristics<sup>a</sup></b>	<b>Episodic Migraine N=1773</b>	<b>Chronic Migraine N=1113</b>
Age (years), mean (SD)	41.3 (11.4)	41.0 (12.1)
Gender (female), n (%)	1499 (84.6)	946 (85.0)
Race (white), n (%)	1333 (75.2)	879 (79.1)
Ethnicity (not Hispanic or Latino), n (%)	1341 (78.9)	788 (75.3)
Region, North America, n (%)	1304 (73.6)	641 (57.6)
Duration of migraine disease, mean (SD)	20.3 (12.4)	21.1 (12.8)
Migraine headache days/month, mean (SD)	9.1 (3.0)	19.4 (4.5)
MIDAS total score, mean (SD)	33.1 (28.7)	67.2 (57.3)
MSQ total score, mean (SD)	58.0 (16.6)	44.9 (18.0)
	<b>N=1338</b>	<b>N=836</b>
Total monthly pain severity-weighted duration in hours <sup>b</sup>	124.3 (94.0)	320.4 (223.8)

Abbreviations: MIDAS = Migraine Disability Assessment; MSQ = Migraine-Specific Quality of Life Questionnaire version 2.1; SD=standard deviation.

<sup>a</sup>Includes pooled data from two 6-month episodic migraine studies and one 3-month chronic migraine study; includes both galcanezumab 120 mg and 240 mg dose groups.

<sup>b</sup>Includes placebo and galcanezumab 120 mg and 240 mg dose groups.

Note: Daily severity-weighted duration was calculated as duration (in hours) x pain severity (none=0, mild=1, moderate=2, severe=3).

Table 2. Additional Proportion of Variability of the Three Components That Contributed to Total Pain Burden at Baseline

<b>Additional Proportion of Variability That Contributed to Total Pain Burden at Baseline<sup>a</sup></b>	<b>Episodic Migraine<sup>b</sup></b>	<b>Chronic Migraine<sup>c</sup></b>
Frequency of monthly migraine headache days	22.6%	13.2%
Duration (hours per migraine headache days)	57.4%	61.1%
Severity of remaining migraine headache days	4.9%	4.8%

<sup>a</sup> Semi-partial provide the unique (or additional) proportion of variability explained by the particular variable in its ability to predict the outcome after accounting for the other 2 components.

<sup>b</sup>Includes pooled data from two parallel 6-month studies.

<sup>c</sup>Includes data from a 3-month study.

Table 3. Individual Components of Total Pain Burden - Changes From Baseline

	<b>Episodic Migraine</b>			<b>Chronic Migraine</b>		
	Overall Monthly Across Months 1 to 6			Overall Monthly Across Months 1 to 3		
Pain burden individual components, LS mean (SE)	Placebo (n=872)	Galcanezumab (n=435)	LS Mean Difference (95% CI)	Placebo (n=535)	Galcanezumab (n=273)	LS Mean Difference (95% CI)
Number of migraine headache days	-2.6 (0.2)	-4.5 (0.2)**	-2.0 (-2.4, -1.6)**	-2.7 (0.4)	-4.8 (0.4)**	-2.1 (-2.9, -1.3)**
Hours per migraine headache day	0.1 (0.1)	-0.6 (0.1)**	-0.7 (-0.9, -0.4)**	0.1 (0.1)	-0.5 (0.2)**	-0.7 (-1.0, -0.3)**
Severity of remaining migraine headache days <sup>a</sup>	-0.2 (0.02)	-0.2 (0.02)*	-0.04 (-0.1, -0.0)*	-0.1 (0.02)	-0.2 (0.02)**	-0.1 (-0.1, -0.03)**

Abbreviations: CI=confidence interval; LS=least squares; SE=standard error.

\*p<0.05 versus placebo.

\*\*p≤0.001 versus placebo.

<sup>a</sup> Severity measured as: none=0, mild=1, moderate=2, severe=3.

## Figures





Figure 1

Total pain burden (monthly severity-weighted duration in hours) for patients with episodic migraine and chronic migraine. Greater decreases in total pain burden (severity-weighted hours) were observed with galcanezumab treatment Months 1 to 6 overall for patients with episodic migraine and Months 1 to 3 overall for patients with chronic migraine.



**Figure 2**

Total pain burden sensitivity analysis: total monthly severity-weighted duration in hours adjusted for change in migraine headache days for patients with episodic migraine and chronic migraine. Greater decreases in total pain burden (severity-weighted hours) were observed with galcanezumab treatment Months 1 to 6 overall for patients with episodic migraine and Months 1 to 3 overall for patients with chronic migraine.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [AilaniGalcanezumabPainBurdenAnimationChronicMigraine.mp4](#)
- [AilaniGalcanezumabPainBurdenAnimationEpisodicMigraine.mp4](#)