

Daily Mean Temperature and HFMD: Risk Assessment and Attributable Fraction Identification in Ningbo China

Rui Zhang

Chinese Center for Disease Control and Prevention

Zhehan Lin

China Population Communication Center

Zhen Guo

Institute of Medical Information / Medical Library, CAMS & PUMC

Zhaorui Chang

Chinese Center for Disease Control and Prevention

Ran Niu

National Institute for Nutrition and health, Chinese Center for Disease Control and Prevention

Yu Wang

National Institute of Environmental Health, Chinese Center for Disease Control and Prevention

Songwang Wang

Chinese Center for Disease Control and Prevention

Yonghong Li (✉ liyonghong@nieh.chinacdc.cn)

National Institute of Environmental Health, Chinese Center for Disease Control and Prevention

<https://orcid.org/0000-0001-5534-1644>

Research

Keywords: Hand, Foot and Mouth Disease (HFMD), Meteorological factors, Temperature, Distributed lag non-linear model (DLNM), Ningbo

Posted Date: August 24th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-60179/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Journal of Exposure Science & Environmental Epidemiology on February 5th, 2021. See the published version at

<https://doi.org/10.1038/s41370-021-00291-y>.

Daily Mean Temperature and HFMD: Risk Assessment and Attributable Fraction Identification in Ningbo China

Rui Zhang¹

Email: zhangrui@chinacdc.cn

Zhehan Lin²

Email: cpccyoung2017@163.com

Zhen Guo³

Email: guo.zhen@imicams.ac.cn

Zhaorui Chang¹

Email: changzr@chinacdc.cn

Ran Niu⁴

Email: niuran@nih.chinacdc.cn

Yu Wang⁵

Email: wangyu@nieh.chinacdc.cn

Songwang Wang^{1,*}

Email: wangws@chinacdc.cn

Yonghong Li^{5,*}

Email: liyonghong@nieh.chinacdc.cn

1 Information Center, Chinese Center for Disease Control and Prevention, Beijing

102206, China.

2 China Population Communication Center, Beijing 100013, China.

3 Institute of Medical Information / Medical Library, CAMS & PUMC, Beijing
100020, China

4 National Institute for Nutrition and health, Chinese Center for Disease Control and
Prevention, Beijing 100050, China.

5 National Institute of Environmental Health, Chinese Center for Disease Control and
Prevention, Beijing 100021, China.

Corresponding authors at: National Institute of Environmental Health, Chinese
Center for Disease Control and Prevention, No 7. Panjiayuan Nanli, Chaoyang
District, Beijing 100021, China. **Tel.:** +86-0150930270, **Fax:** +86-0150930146.
(Yonghong Li).

Email:

liyonghong@nieh.chinacdc.cn (Yonghong Li),

wangws@chinacdc.cn (Songwang Wang).

Abstract

Background:

Hand, foot and mouth disease (HFMD) remains a significant public health issue, especially in developing countries. Many studies have reported the association between environmental temperature and HFMD. However, the results are highly heterogeneous in different regions. In addition, there are few studies on the attributable risk of HFMD.

Objectives:

The study aimed to assess the association between temperature and HFMD incidence and to evaluate the attributable burden of HFMD in Ningbo China.

Methods:

The research used daily incidence of HFMD from 2014 to 2017 and distributed lag non-linear model (DLNM) to investigate the effects of daily mean temperature (Tmean) on HFMD incidence from lag 0 to 30 days, after controlling potential confounders. The lag effects and cumulative relative risk (CRR) were analyzed. Attributable fraction (AF) of HFMD incidence due to temperature was calculated. Stratified analysis by age and gender were also conducted.

Results:

The Significant associations between Tmean and HFMD incidence were observed in Ningbo for lag 0-30. Two peaks were observed at both low (5°C-11°C) and high

(16°C-29°C) temperature scales. For low temperature scale, the highest CRR was 2.22% (95% CI: 1.61-3.07) at 7°C on lag 0-30. For high temperature scale, the highest CRR was 3.54% (95% CI: 2.58-4.88) at 24°C on lag 0-30. The AF of low and high temperature was 5.23% (95% CI: 3.10-7.14) and 39.55% (95% CI: 30.91-45.51), respectively. There was no significant difference between gender- and age-specific AFs, even though the school-age and female children had slightly higher AF values.

Conclusions:

The result indicates that both high and low temperatures were associated with daily incidence of HFMD, and more burdens were caused by heat in Ningbo.

Key words

Hand, Foot and Mouth Disease (HFMD); Meteorological factors; Temperature; Distributed lag non-linear model (DLNM); Ningbo

1. Background

Hand, foot and mouth disease (HFMD) is a common infectious disease in China. It was first reported in New Zealand in 1957 along with the isolation of Coxsackie virus as the pathogen [1]. This emerging infectious disease was then named as Hand, foot and mouth disease according to its typical symptoms in 1959 [2]. Infants and children under 6 years old are more likely to get this disease, though elder children and adults could also be infected. HFMD can be caused by various enteroviruses, including Coxsackie virus A16 (CV-A16) and Enterovirus 71 (EV71). Main clinical symptoms of HFMD include fever, mouth ulcers, and vesicles on the hands and feet. In most cases, the disease is mild and self-limiting. But a small proportion can rapidly develop neurological and systemic complications, including brainstem encephalitis and poliomyelitis-like acute flaccid paralysis [3].

Previous studies have shed light on the effect of meteorological factors such as temperature, relative humidity and altitude on HFMD [4-9]. However, many studies have shown that the association between the incidence of HFMD and seasons or temperatures varied with different regions. Some areas have the peak once a year, such as in summer in Japan [10, 11], and in autumn in Finland [12]. Two peaks may occur in other regions, such as in summer and autumn in the United Kingdom [13, 14] and Belgium [15]. The incidence trends of HFMD are different in North and South China. In northern China, HFMD outbreaks in June among cities, such as Beijing [16, 17] and Hefei [8]. In several cities of southern China, like Kunming [18] and Guangdong [19, 20], the HFMD outbreaks in May and October-November. These

discrepancies could arise from various local climatic conditions, differences in socioeconomic status, and the demographic characteristics of different regions. Our understanding of the impact of seasonal and meteorological variables on disease transmission remains limited. Further research is required into the effects of meteorological variables on the incidence of HFMD.

At present, attributable risk has been used to study the quantitative dependence and regularity between risk factors and morbidity and mortality of infectious diseases. It has transformed factors such as lifestyle, physical, meteorological and pollution indicators into measurable indicators for predicting the risk of disease or death in the future. An understanding of the temperature contributing to disease burden is critical for determining research priorities and informing national health policy. However, the contribution of temperature to the incidence of HFMD burden has not been quantified in Ningbo.

Therefore, this research aims to analyze the association between the daily mean temperature and the incidence of HFMD in Ningbo, as well as evaluate the attributable fraction of HFMD due to temperature. Thus, reliable basis for the prevention and control of HFMD in Ningbo could be provided.

2. Methods

2.1. Geography

Ningbo locates in the Southern China and ranges in latitude from $28^{\circ}51'$ to $30^{\circ}33'N$ and in longitude from $120^{\circ}55'$ to $122^{\circ}16'E$. It bounds on the east by the

East China Sea and Zhoushan Archipelago, on the north by Hangzhou Bay, across which it faces Jiaxing and Shanghai, on the west by Shaoxing, and on the south by Taizhou. Its land area is 9,186km², while oceanic territory amounts to 9,758km².

2.2. Climate

Ningbo has a humid subtropical climate with four distinctive seasons, characterized by hot, humid summers and chilly, cloudy and dry winters (with occasional snow). The mean annual temperature is 17.15°C, with monthly daily averages ranging from 5.3°C in January to 28.8°C in July. The city receives an average annual rainfall of 1,430mm and is affected by the plum rains of the Asian monsoon in June, when average relative humidity also peaks [21].

2.3. Incidence Data

The daily incidence data of HFMD in Ningbo from January 2014 to December 2017 were collected from the official website of The Data-center of China Public Health Science [22]. The diagnostic criteria of HFMD were based on the clinical criteria set by the Hand, Foot and Mouth Disease Control and Prevention Guide published by the National Health Commission of the People's Republic of China [23]. To identify vulnerable populations, the data was reclassified by gender (male, female) and age (0-3, 4-5, ≥6 years old). The meteorological data were collected from the China Meteorological Administration, including daily maximum, minimum and mean temperatures (°C), and daily mean relative humidity (RH, %) and daily precipitation (mm).

2.4. Data analysis

A distributed lag nonlinear model (DLNM) with quasi-Poisson regression was applied to evaluate the association between the daily mean temperature and the incidence of HFMD in Ningbo. The DLNM model was defined by the following formula [24]:

$$\log[E(Y_t)] = \alpha + cb(Tmean_t, lag = 30) + ns(time_i, df = 7/year) + ns(RH, 3) + \beta DOW_t$$

where Y_t represents the expected number of HFMD incidence on day t . α is the intercept. $Tmean_t$ is the daily mean temperature on day t . cb refers to the crossbasis function, which specifies the exposure-lag-response relationship simultaneously in the exposure-response and lag-response dimensions; a quadratic B spline (bs) with 4 degrees of freedom (df) was used for the exposure-response relationship and natural cubic splines (ns) with 4 degrees of freedom for the lag-response relationship. The lag day up to 30 days reflects the maximum lag day of the temperature effect. A smooth function of time with 7 degrees of freedom per year was used in the model to control the seasonality and long-term trends. Mean relative humidity (RH), as potential confounding variables, was modeled as a natural cubic spline (ns) with 3 degrees of freedom. DOW stands for day of the week, which was entered as a categorical variable, and β is the coefficient of DOW [24-27].

The reference temperature for the DLNM analysis was identified according to the relationship between daily mean temperature and the incidence of HFMD, that is, the temperature, at which the incidence of HFMD had the relative lower risk was

identified as the reference temperature for this study. Cumulative relative risks (CRR) for specific temperature scales under certain lag days were calculated to assess the effects of temperature on the incidence of HFMD.

Attributable fraction (AF) was calculated to evaluate the incidence burden of HFMD caused by temperature. There are two ways to define AF within the dlnm framework. One is backward perspective, the other is forward perspective. Backward perspective is a commonly used explanation of lag effect in research, but its calculation process is more complicated than forward perspective. The principle and calculation of forward perspective are relatively simple, but the actual risk may be underestimated [27]. In this study, we used backward perspective to calculate AF. The backward attributable fraction $b - AF_{x,t}$ at time t were obtained by the following formulas [27]:

$$b - AF_{x,t} = 1 - \exp\left(-\sum_{l=l_0}^L \beta_{x_{t-l},l}\right)$$

l_0 and L correspond to minimum and maximum lags, respectively.

Finally, we calculated the empirical confidence interval values through Monte Carlo simulations [28] and the related 2.5th and 97.5th percentiles of multivariate normal distribution were interpreted as 95% empirical confidence intervals [27].

Stratified analyses by gender and age were also conducted. The statistical significance tests of gender- and age-specific differences were conducted by using the formula below [29]:

$$(\widehat{Q}_1 - \widehat{Q}_2) \pm 1.96\sqrt{S\widehat{E}_1^2 + S\widehat{E}_2^2}$$

Where \widehat{Q}_1 and \widehat{Q}_2 are the estimates for the two categories, and \widehat{SE}_1 and \widehat{SE}_2 are their respective standard errors.

2.5. Sensitivity analysis

We conducted the collinear diagnosis to see if there is collinearity among RH and precipitation. Sensitivity analysis was performed by changing the df of RH, adjusting and without adjusting RH and precipitation, respectively.

Data analysis was conducted using R software 3.6.2 with the package 'dlnm' [24]. The 'dlnm' package was used for fitting DLNM. For all statistical tests, statistical significance as a two-tailed $P < 0.05$.

3. Results

3.1. Descriptive analysis

Descriptive statistics for the number of HFMD cases based on gender and age, as well as meteorological variables including daily mean temperature, daily mean relative humidity and daily precipitation are summarized in [Table 1](#). A total of 129,897 HFMD cases from January 2014 to December 2017 were included in our analyses, of which 76,846 (59.16%) were male and 53,051 (40.84%) were female. The group aged 0-3 years, 4-5 years and aged ≥ 6 years accounted for 76.43%, 17.19% and 6.38%, respectively. The mean value of daily mean temperature was 17.46°C, the standard deviation was 8.42°C, the maximum value was 32.9°C, and the minimum value was -4.5°C. The mean value of daily mean relative humidity was 79.75%, the standard deviation was 11.18%, the maximum value was 100%, and the minimum value was

34%. The mean value of daily precipitation was 5.0mm, the standard deviation was 14.38mm, the maximum value was 276.2mm, and the minimum value was 0mm.

Table 1. Descriptive statistics of daily incidence of HFMD cases and meteorological factors in Ningbo, 2014–2017

	Total	Mean±SD	Min	Max	Proportion
Daily mean Incidence	129,897	88.91±76.84	1	479	
Sex					
Male	76,846	52.6±45.76	0	302	59.16%
Female	53,051	36.31±31.71	0	183	40.84%
Age					
0-3	99,284	67.96±56.29	1	309	76.43%
4-5	22,331	15.28±16.83	0	124	17.19%
≥6	8,282	5.67±6.13	0	46	6.38%
Daily mean temperature	-	17.46±8.42	-4.5	32.9	-
Daily mean relative humidity	-	79.75±11.18	34	100	-
Daily precipitation		5.0±14.38	0	276.2	-

Figure 1. Decomposition of additive time series of the daily incidence of HFMD in Ningbo from January 2014 to December 2017

[Fig. 1](#) shows the decomposition analysis of additive time series of the daily incidence cases of HFMD in Ningbo from January 2014 to December 2017, which included the trend of the observed cases, the long-term and seasonal trends and the random variation. The long-term trend showed that the overall incidence of HFMD in Ningbo presented a downward trend from 2014 to 2015, followed by slowly rising in 2016 and then decreasing in 2017. The seasonal lines showed a strong seasonality that a bimodal seasonal pattern was observed in Ningbo, which was characterized by peaks in HFMD incidence in the summer (June) and early winter (November). And the random line showed the randomness of the data. It was indicated that the dataset of the incidence of HFMD during 2014-2017 in Ningbo was a typical time series and the meteorological variables might be associated with HFMD.

3.2. Associations between temperature and HFMD

[Fig. 2](#) shows the CRR of daily HFMD incidence associated with daily mean temperature on lag 0-30. Analysis of the crude relationship revealed that the potential overall CRR of HFMD had an approximately M-shape with two peaks during 5°C-11°C and 16°C-29°C which was consistent with the bimodal seasonal trend. It was demonstrated in [Fig. 2](#) that the lowest point between the two peaks was 14°C, so we chose 14°C as the reference temperature. The highest risk was at 7°C during 5°C-11°C and at 24°C during 16°C-29°C, so 7°C and 24°C was chosen as the

representative low and high temperatures to calculate relative risk (RR) and CRR, respectively.

Figure 2. Cumulative relative risk (CRR) of daily HFMD incidence associated with daily mean temperature on lag 0-30 (reference temperature: 14°C)

Fig. 3 shows the RR of daily incidence of HFMD associated with high and low temperatures. It indicated that the temperature-HFMD association was nonlinear, immediate and lasted throughout 30 days lag periods. The RR of daily incidence of HFMD at 7°C were more prominent over shorter 10 days lag periods with the highest RR of 1.08 (95% CI: 1.06-1.10) on lag 3 (Table 2) and diminished over time, whereas the effects at 24°C were persistent over longer 30 days lag periods with the highest RR of 1.06 (95% CI: 1.04-1.07) on lag 14 (Table 2).

Figure 3. Relative risk of daily incidence of HFMD associated with temperatures (a: 24°C, b: 7°C) on lag 0-30 (reference temperature: 14°C)

Table 2. Relative risks and cumulative relative risks of HFMD incidence

	RR		CRR	
	7°C, lag 3	24°C, lag 14	7°C, lag 0-30	24°C, lag 0-30
Overall	1.08(1.06, 1.10)	1.06(1.04, 1.07)	2.22(1.61, 3.07)	3.54(2.58, 4.88)
Sex				

Male	1.08(1.06, 1.11)	1.05(1.03, 1.07)	2.50(1.77, 3.55)	3.12(2.22, 4.40)
Female	1.07(1.04, 1.09)	1.07(1.05, 1.09)	1.89(1.31, 2.71)	4.23(2.94, 6.07)
Age				
0-3	1.07(1.05, 1.09)	1.05(1.04, 1.07)	2.04(1.47, 2.84)	3.31(2.40,4.59)
4-5	1.11(1.07, 1.14)	1.08(1.06, 1.11)	3.64(2.19, 6.03)	5.08(3.08, 8.36)
≥6	1.07(1.03, 1.12)	1.07(1.04, 1.10)	1.84(0.93, 3.63)	3.88(2.00, 7.55)

It was shown in [Table 2](#) that the CRR at 7°C from lag 0-30 was 2.22 (95% CI: 1.61-3.07) and the CRR at 24°C from lag 0-30 was 3.54 (95% CI: 2.58-4.88). The results displayed that both gender- and age-specific difference were statistically insignificant. At the low temperature of 7°C, the CRR of male at was 2.50 (95% CI: 1.77-3.55) and that of female was 1.89 (95% CI: 1.31-2.71). The CRR of group aged 0-3 years, 4-5 years and ≥6 years was 2.04 (95% CI: 1.47-2.84), 3.64 (95% CI: 2.19-6.03) and 1.84 (95% CI: 0.93-3.63), respectively. At the high temperature of 24°C, the CRR of male was 3.12% (95% CI: 2.22-4.40), and that of female was 4.23% (95% CI: 2.94-6.07). The CRR of group aged 0-3 years, 4-5 years and ≥6 years was 3.31% (95% CI: 2.40-4.59), 5.08% (95% CI: 3.08-8.36) and 3.88% (95% CI: 2.00-7.55), respectively.

3.3. Attributable fraction of HFMD due to temperature

[Table 3](#) presents the AF of HFMD due to temperatures for the overall group as well as gender- and age-specific subgroups. It was shown that the AFs were higher for hot temperatures (16°C-29°C) than those for cold temperatures (5°C-11°C). The estimated

overall AF was 45.50 (95% CI: 36.21-52.48). The AF due to cold temperature (5°C-11°C) was 5.23 (95% CI: 3.10-7.14) and the AF due to hot temperature (16°C-29°C) was 39.55 (95% CI: 30.91-45.51). The AF of male and female was 42.98 (95% CI: 31.51-51.23) and 48.26 (95% CI: 39.09-54.77), respectively. The group aged 4-5 years (AF: 53.34, 95% CI: 40.91-60.96) and ≥ 6 years (50.59, 95% CI: 26.93-62.89) seemed to have higher AF values than the group aged 0-3 years (43.44, 95% CI: 34.14-51.41), but there was no significant difference. The similar trends of the age-specific AFs due to cold and hot temperatures were also observed.

Table 3. Attributable fraction of HFMD incidence

Group	Overall	5°C-11°C	16°C-29°C
Overall	45.50(36.21, 52.48)	5.23(3.10, 7.14)	39.55(30.91, 45.51)
Sex			
Male	42.98(31.51, 51.23)	5.92(3.60, 7.79)	36.61(27.02, 44.29)
Female	48.26(39.09, 54.77)	4.20(1.55, 6.47)	43.15(34.21, 48.94)
Age			
0-3	43.44(33.14, 51.41)	4.67(2.41, 6.59)	38.31(29.88, 44.79)
4-5	53.34(40.91, 60.96)	8.36(5.25, 10.81)	44.16(34.93, 51.28)
≥ 6	50.59(26.93, 62.89)	3.99(-1.79, 7.50)	43.74(26.05, 56.20)

3.4. Sensitivity Analysis

The result of collinear diagnosis showed that there was no serious collinearity among daily mean RH and daily precipitation (correlation coefficient is 0.35). The results (see [Table S1](#)) remained almost unchanged and statistically significant, indicating the stability of the models.

4. Discussion

We assessed the association between daily mean temperature and the daily incidence of HFMD in Ningbo and two peaks were discovered during specific cold (5°C-11°C) and hot (16°C-29°C) temperatures. What's more, to our best knowledge, for the first time we evaluated the AF of HFMD incidence due to temperature in Ningbo and found that much more disease burden of HFMD could be attributed to high temperatures than that attributed to cold temperatures. Stratified analysis implied that the occurrence of HFMD of school-age children seemed to be more sensitive to temperatures than the group aged 0-3 years.

In this study, we found that the incidence of HFMD in Ningbo has semiannual peaks, including a higher peak in summer and the lower one in early winter. The result is consistent with the previous studies which demonstrated that HFMD outbreaks once a year in high latitudes regions but twice a year in tropical and subtropical regions [[2](#), [10](#), [19](#)].

Several previous studies have shed light on the effect of meteorological factors such as temperature on HFMD [[4-9](#), [19](#), [30](#)]. However, the results have shown that the association between the incidence of HFMD and temperature varied with different

regions [25, 30-35]. Chen et al. [36] found that the occurrence of HFMD epidemics was bimodal in Wuhan, one peak occurred when the monthly average temperature was below 15°C during autumn-winter and the other occurred when the monthly average temperature exceeds 25°C in summer. The study in Guangzhou [19] also found that the association between the daily incidence of HFMD and the daily mean temperature increased rapidly when the daily mean temperature was below 25°C but flattened above 25°C. While our result showed that the risk of HFMD increased during both low temperature from 5°C to 11°C and high temperature from 16°C to 29°C of daily mean temperatures, but declined at both extremely cold (<5°C) and hot (>29°C) temperature. The inconsistency of these findings can be partly attributable to the diversity of methodologies and data sources, but it also implies that the temperature-HFMD relationship might be modified by some location-specific variables. The heterogeneity across studies is still poorly understood which hinder a more comprehensive characterization of the temperature-HFMD relationship.

As some previous studies suggested, our results imply that the effect of temperature on HFMD may involve with complex mechanisms. Some evidential studies found that temperature could affect the breeding, growth, and transmission of pathogens, as well as human behavior [37]. First, HFMD can be transmitted either by exposure to infectious individuals or contaminated environment. During the suitable temperature (such as 5°C-11°C and 16°C-29°C), individuals are more likely to have outdoor activities, thereby suitable temperature may increase the chances that susceptible individuals contact with infectious individuals or contaminated environment.

Conversely, people will reduce outdoor activities and take some protective measures at extreme temperatures. Second, both extremely cold and hot temperatures can reduce the reproduction of pathogens. Temperature is considered as the main factor determining enterovirus inactivation in the environment [38]. According to the previous studies in the laboratory, EV-A71 and CVA16 are lack of a thermostatic mechanism, and their reproduction and survival rates are strongly affected by fluctuations in temperature. The EV-A71 survives at 20°C slightly better than at 25°C [19], and at 36°C is more active than at 39°C [39]. Therefore, extremely temperature may shorten the survival time of enterovirus in the environment. All of above might explain the two peaks during specific suitable temperature scales and the dropping down of risk at extreme temperatures.

Measures of attributable risk are an integral part of epidemiological analyses, particularly when aimed at the planning and evaluation of public health interventions [26, 27, 32]. It was presented in our results that the overall attributable fraction of HFMD incidence due to hot temperatures (39.55%) was significantly higher than that due to cold temperatures (5.23%). This results indicated that when the temperature is between 16°C and 29°C in summer in Ningbo, public-health authorities should prepare fully to respond to a possible epidemic of HFMD, including increasing access to health-care resources, the distribution of scientific knowledge to the public, medical staff and public health personnel, the availability of essential medical equipment, active disease surveillance, and the design of other more specific control measures to mitigate the risk of disease transmission.

It was displayed in our study that the number of the HFMD cases of group aged 0-3 years was the highest. However, the attributable fractions of group aged 4-5 years and ≥ 6 years were both higher than the AF of group aged 0-3 years, which meant that the occurrence of HFMD of the school-age children be more susceptible to the ambient temperature. Our results indicated that more attention should also be paid to the school-age children especially by strengthening the hand hygiene and the monitor of symptoms and diseases in kindergarten and primary school especially during periods with specific ambient temperatures.

Although difference of temperature effects between males and females were not statistically significant, the daily incidence of HFMD tended to be higher in males than in females. This result is consistent with other studies [2, 19]. One possible reason is that boys may be more active and thus have more chances to infect the disease.

Some limitations of this paper should be acknowledged. First, we used ambient temperature data monitored at a fixed location as most previous epidemiological studies, rather than individual real exposure data, which could lead to measurement errors for exposure. Second, as the availability of the data, we did not control the effect of air pollution on the incidence of HFMD. Third, our study focused on Ningbo, a southern city in China. The findings need to be verified in more other cities or regions. However, our results warrant further research on climate change and the incidence of HFMD and provide scientific evidence for the planning of control and prevention of HFMD in Ningbo.

5. Conclusions

Global climate change has a profound impact on infectious diseases including HFMD. This study can help local health departments to provide early warnings to control the risk before reaching its peak and implement effective interventions to reduce the burden of HFMD.

Declarations

- Ethics approval and consent to participate

Since no primary data collection was undertaken, no patient or public was involved, no formal ethical assessment or informed consent was required. All data were collected from the official website and all data were fully anonymized.

- Consent for publication

Not applicable

- Availability of data and materials

Everyone can apply the daily incidence data of HFMD in Ningbo from January 2014 to December 2017 through the official website of The Data-center of China Public Health Science (<http://www.phsciencedata.cn/Share/en/index.jsp>).

- Competing interests

The authors declare that they have no competing interests.

- Funding

This work was supported by the Special Foundation of Basic Science and Technology Resources Survey from Ministry of Science and Technology of China

[2017FY101201, 2017FY101207] and National Science and Technology Major Project [2017ZX10305501-009-001].

- Authors' contributions

Rui Zhang, Songwang Wang and Yonghong Li were major contributors in writing the manuscript. Zhehan Lin and Zhen Guo analyzed daily HFMD incidence associated with daily mean temperature. Zhaorui Chang, Ran Niu and Yu Wang calculated the CRR and AF of HFMD incidence due to temperature. All authors read and approved the final manuscript

- Acknowledgements

Not applicable

Rui Zhang¹, Zhehan Lin², Zhen Guo³, Zhaorui Chang¹, Ran Niu⁴, Yu Wang⁵, Songwang Wang^{1,*}, Yonghong Li⁵

References

1. Robinson CR, Doane FW, Rhodes AJ: **Report of an outbreak of febrile illness with pharyngeal lesions and exanthem: Toronto, summer 1957; isolation of group A Coxsackie virus.** *Can Med Assoc J* 1958, **79**(8):615-621.
2. Zhu Q, Hao Y, Ma J, Yu S, Wang Y: **Surveillance of hand, foot, and mouth disease in mainland China (2008-2009).** *Biomed Environ Sci* 2011, **24**(4):349-356.
3. Ooi MH, Wong SC, Lewthwaite P, Cardoso MJ, Solomon T: **Clinical features, diagnosis, and management of enterovirus 71.** *Lancet Neurol* 2010, **9**(11):1097-1105.
4. Zhang X, Xu C, Xiao G: **Spatial heterogeneity of the association between temperature and hand, foot, and mouth disease risk in metropolitan and other areas.** *Sci Total Environ* 2020, **713**:136623.
5. Ji XY, Huang LY, Song J, Fei CN, Liu J, Liu H: **Short-term effects of meteorological factors, air pollution, and sunspot on childhood hand, foot, and mouth disease in Tianjin, China: a new time series regression, 2014-2018.** *Environ Sci Pollut Res Int* 2020.
6. Guo T, Liu J, Chen J, Bai Y, Long Y, Chen B, Song S, Shao Z, Liu K: **Seasonal Distribution and Meteorological Factors Associated with Hand, Foot, and Mouth Disease among**

- Children in Xi'an, Northwestern China.** *Am J Trop Med Hyg* 2020, **102**(6):1253-1262.
7. Gu S, Li D, Lu B, Huang R, Xu G: **Associations between ambient air pollution and daily incidence of pediatric hand, foot and mouth disease in Ningbo, 2014-2016: a distributed lag nonlinear model.** *Epidemiol Infect* 2020, **148**:e46.
 8. Deng J, Gao X, Xiao C, Xu S, Ma Y, Yang J, Wu M, Pan F: **Association between diurnal temperature range and outpatient visits for hand, foot, and mouth disease in Hefei, China: a distributed lag nonlinear analysis.** *Environ Sci Pollut Res Int* 2020.
 9. Bo Z, Ma Y, Chang Z, Zhang T, Liu F, Zhao X, Long L, Yi X, Xiao X, Li Z: **The spatial heterogeneity of the associations between relative humidity and pediatric hand, foot and mouth disease: Evidence from a nation-wide multicity study from mainland China.** *Sci Total Environ* 2020, **707**:136103.
 10. Onozuka D, Hashizume MJSotTE: **The influence of temperature and humidity on the incidence of hand, foot, and mouth disease in Japan.** 2011, **s 410–411**(none):119-125.
 11. Sumi A, Toyoda S, Kanou K, Fujimoto T, Mise K, Kohei Y, Koyama A, Kobayashi N: **Association between meteorological factors and reported cases of hand, foot, and mouth disease from 2000 to 2015 in Japan.** *Epidemiol Infect* 2017, **145**(14):2896-2911.
 12. Blomqvist S, Klemola P, Kaijalainen S, Paananen A, Simonen ML, Vuorinen T, Roivainen M: **Co-circulation of coxsackieviruses A6 and A10 in hand, foot and mouth disease outbreak in Finland.** *J Clin Virol* 2010, **48**(1):49-54.
 13. Bendig JW, Fleming DM: **Epidemiological, virological, and clinical features of an epidemic of hand, foot, and mouth disease in England and Wales.** *Commun Dis Rep CDR Rev* 1996, **6**(6):R81-86.
 14. Sinclair C, Gaunt E, Simmonds P, Broomfield D, Nwafor N, Wellington L, Templeton K, Willocks L, Schofield O, Harvala H: **Atypical hand, foot, and mouth disease associated with coxsackievirus A6 infection, Edinburgh, United Kingdom, January to February 2014.** *Euro Surveill* 2014, **19**(12):20745.
 15. Druyts-Voets E: **Epidemiological features of entero non-poliovirus isolations in Belgium 1980-94.** *Epidemiol Infect* 1997, **119**(1):71-77.
 16. Liu XF, Sun XM, Sun XW, Yang YQ, Huang CH, Wen H: **Epidemiological study on hand, foot and mouth disease in Tongzhou District, Beijing, 2013-2017.** *J Int Med Res* 2019, **47**(6):2615-2625.
 17. Dong W, Li X, Yang P, Liao H, Wang X, Wang Q: **The Effects of Weather Factors on Hand, Foot and Mouth Disease in Beijing.** *Sci Rep* 2016, **6**:19247.
 18. Ma S, Zhang Y, Du C, Yang T, Liu Q, Pan Y, Chen J, Shi H, Sun Q, Liu L *et al*: **Dynamic Constitution of the Pathogens Inducing Encephalitis in Hand, Foot and Mouth Disease in Kunming, 2009-2011.** *Jpn J Infect Dis* 2015, **68**(6):504-510.
 19. Chen C, Jiang Q, Song Z, Li Y, Wang H, Lu Y, Wang D, Li M, Li T: **Influence of temperature and humidity on hand, foot, and mouth disease in Guangzhou, 2013-2017.** *J Int Med Res* 2020, **48**(6):300060520929895.
 20. Zhang H, Yang L, Li L, Xu G, Zhang X: **The epidemic characteristics and spatial autocorrelation analysis of hand, foot and mouth disease from 2010 to 2015 in Shantou, Guangdong, China.** *BMC Public Health* 2019, **19**(1):998.
 21. Lin D, Guo H, Wang J, Shi PJJöBNU: **Precipitation and temperature changes during high-temperature periods in Ningbo,Zhejiang Province in the last 50 years:case of**

- Yinzhou Station. 2015.
22. **The Data-center of China Public Health Science** [<http://www.phsciencedata.cn/Share/en/index.jsp>]
 23. **Hand, Foot and Mouth Disease Control and Prevention Guide.** In.; 2008.
 24. Gasparrini A: **Distributed Lag Linear and Non-Linear Models in R: The Package dlnm.** *J Stat Softw* 2011, **43**(8):1-20.
 25. Su XC, Yibin; Wang, Yu; Liu,Yue; Li, Na; Li, Yonghong; Yao, Xiaoyuan: **Regional Temperature-Sensitive Diseases and Attributable Fractions in China.** *International Journal of Environmental Research and Public Health* 2020, **17**(1).
 26. Liu Z, Tong MX, Xiang J, Dear K, Perspectives PBJEH: **Daily Temperature and Bacillary Dysentery: Estimated Effects, Attributable Risks, and Future Disease Burden in 316 Chinese Cities.** 2020, **128**(5):057008.
 27. Gasparrini A, Leone M: **Attributable risk from distributed lag models.** *BMC Medical Research Methodology* 2014, **14**(1):55.
 28. Sander GJJJoE: **Interval estimation by simulation as an alternative to and extension of confidence intervals.** 2004(6):1389-1397.
 29. Zanobetti AZA, Epidemiology JSJAJo: **Individual-Level Modifiers of the Effects of Particulate Matter on Daily Mortality.** 2006, **163**(9):849-859.
 30. Xu C, Zhang X, Wang L, Zhou Y, Xiao G, Liao J: **Effects of temperature fluctuations on spatial-temporal transmission of hand, foot, and mouth disease.** *Sci Rep* 2020, **10**(1):2541.
 31. Xing W: **Epidemiological characteristics of hand-foot-and-mouth disease in China, 2008-2012 %J Lancet Infectious Diseases.** 2014, **14**(4):308.
 32. Qi H, Li Y, Zhang J, Chen Y, Guo Y, Xiao S, Hu J, Wang W, Zhang W, Hu Y *et al*: **Quantifying the risk of hand, foot, and mouth disease (HFMD) attributable to meteorological factors in East China: A time series modelling study.** *Sci Total Environ* 2020, **728**:138548.
 33. Gu J, Liang L, Song H, Kong Y, Ma R, Hou Y, Zhao J, Liu J, He N, Zhang Y: **A method for hand-foot-mouth disease prediction using GeoDetector and LSTM model in Guangxi, China.** *Sci Rep* 2019, **9**(1):17928.
 34. Du Z, Lin S, Marks T, Zhang W, Deng T, Yu S, Hao Y: **Weather effects on hand, foot, and mouth disease at individual level: a case-crossover study.** *BMC Infect Dis* 2019, **19**(1):1029.
 35. Coates SJ, Davis MDP, Andersen LK: **Temperature and humidity affect the incidence of hand, foot, and mouth disease: a systematic review of the literature - a report from the International Society of Dermatology Climate Change Committee.** *Int J Dermatol* 2019, **58**(4):388-399.
 36. Chen B, Sumi A, Toyoda S, Hu Q, Zhou D, Mise K, Zhao J, Kobayashi N: **Time series analysis of reported cases of hand, foot, and mouth disease from 2010 to 2013 in Wuhan, China.** *BMC Infect Dis* 2015, **15**:495.
 37. Wang P, Goggins WB, Chan EY: **Hand, Foot and Mouth Disease in Hong Kong: A Time-Series Analysis on Its Relationship with Weather.** *PLoS One* 2016, **11**(8):e0161006.
 38. Bertrand I, Schijven JF, Sanchez G, Wyn-Jones P, Ottoson J, Morin T, Muscillo M, Verani

- M, Nasser A, de Roda Husman AM *et al*: **The impact of temperature on the inactivation of enteric viruses in food and water: a review.** *J Appl Microbiol* 2012, **112**(6):1059-1074.
39. Arita, Virology MJoG: **Temperature-sensitive mutants of enterovirus 71 show attenuation in cynomolgus monkeys.** 2005, **86**(5):1391-1401.

Figures




Figure 1

Decomposition of additive time series of the daily incidence of HFMD in Ningbo from January 2014 to December 2017




Figure 2

Cumulative relative risk (CRR) of daily HFMD incidence associated with daily mean temperature on lag 0-30 (reference temperature: 14°C)




Figure 3

Relative risk of daily incidence of HFMD associated with temperatures (a: 24°C, b: 7°C) on lag 0-30 (reference temperature: 14°C)

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [TableS1.docx](#)
- [TableS1.docx](#)