A New Paradigm for South Pacific Climate Variability and Predictability

Jiale Lou (✉ jiale.lou@noaa.gov)

NOAA <https://orcid.org/0000-0002-4014-5242>

Terence O'Kane

CSIRO

Neil Holbrook

University of Tasmania <https://orcid.org/0000-0002-3523-6254>

Article

Keywords: climate variability, atmospheric science, ocean science, South Pacific

Posted Date: July 19th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-602270/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Communications Earth & Environment on October 21st, 2021. See the published version at <https://doi.org/10.1038/s43247-021-00295-4>.

A New Paradigm for South Pacific Climate Variability and Predictability

Jiale Lou^{1,2,*†}, Terence J. O’Kane^{3,4}, Neil J. Holbrook^{1,4}

¹Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, TAS, Australia

²ARC Centre of Excellence for Climate System Science, University of Tasmania, Hobart, TAS, Australia

³CSIRO Oceans and Atmosphere, Hobart, TAS, Australia

⁴ARC Centre of Excellence for Climate Extremes, University of Tasmania, Hobart, TAS, Australia

*Current affiliations – Cooperative Institute for Research in the Environmental Sciences, University of Colorado Boulder, Boulder, CO, USA

† Physical Sciences Laboratory, NOAA, Boulder, CO, USA

Corresponding author address:

Jiale Lou

NOAA Physical Sciences Laboratory, 325 Broadway, Boulder, CO, 80305

e-mail: jiale.lou@noaa.gov

Summary

Pacific climate variability is largely understood based on El Niño–Southern Oscillation (ENSO), the North Pacific focused Pacific decadal oscillation (PDO) and/or the whole of Pacific region interdecadal Pacific oscillation – which respectively represent the dominant modes of interannual and decadal climate variability. However, the role of the South Pacific, including atmospheric drivers and cross-scale interactions between interannual and decadal climate variability, has received considerably less attention. Here we propose a new paradigm for South Pacific climate variability whereby the Pacific-South American (PSA) mode, characterised by two mid-tropospheric modes (PSA1 and PSA2), provides coherent noise forcing that acts to excite multiple spatiotemporal scales of oceanic responses in the upper South Pacific Ocean ranging from seasonal to decadal. While PSA1 has long been recognised as highly correlated with ENSO, we find that PSA2 is critically important in generating a sea surface temperature (SST) quadrupole pattern in the extratropical South Pacific. This sets up a precursor that optimally determines the predictability and evolution of SST 9 months in advance of the peak phases of both the leading South Pacific SST mode and ENSO. Our results show that the atmospheric PSA mode is the key driver of oceanic variability in the South Pacific subtropics. Importantly, we anticipate that South Pacific Ocean climate variability is potentially predictable on seasonal to interannual time scales. We expect this new knowledge to be of potential benefit to marine conservation, fisheries and aquaculture management practices that utilise climate forecasts for their operations and planning.

Pacific decadal variability has been historically understood in terms of the North Pacific-focused Pacific decadal oscillation (PDO ^[1]) and by the basin-scale interdecadal Pacific oscillation (IPO ^[2]). However, neither the PDO nor the IPO explicitly isolate the South Pacific contribution. Analogous to the PDO, the South Pacific decadal oscillation (SPDO ^[3, 4]) has been identified and described as the South Pacific centre of action contributing to the entire basin-wide Pacific decadal variability. Refs ^[4, 5] show that the SPDO can be viewed as a combination of different dynamical processes operating on different timescales. Those processes include the atmospheric forcing, tropical El Niño–Southern Oscillation (ENSO) teleconnections, and the internal oceanic dynamics. Importantly, the atmospheric forcing associated with the Pacific-South American pattern 1 (PSA1) has been identified as the main stochastic driver of the SPDO ^[4] (cf Fig.1 (left)). ENSO has been shown to impact extratropical variability via the atmospheric bridge ^[6] and oceanic pathways ^[7]. Recent studies point out that extratropical ocean dynamics can feed back to the tropical Pacific via a sea surface temperature (SST) quadrupole pattern ^[8] or a South Pacific meridional mode ^[9-11]. These oceanic processes act as extratropical precursors that might potentially guide the predictability and evolution of ENSO.

In this study, we propose a new paradigm for South Pacific climate variability whereby the pair of atmospheric eastward-propagating PSA modes can excite extratropical South Pacific Ocean responses on multiple time scales ranging from seasonal to decadal. Although the South Pacific Ocean responds to fast-varying atmospheric forcing on distinct timescales, the resultant spatial SST features remain difficult to distinguish from the background state. This suggests that the characterisation and identification of predictable signals, and their sources, require careful separation of the climate signal and relevant noise processes. This remains a necessary but challenging problem.

The atmospheric Pacific-South American mode

The PSA mode is represented by an eastward-propagating wave train extending from eastern Australia to Argentina, characterised in mid-tropospheric geopotential height in terms of two invariant empirical orthogonal function (EOF) patterns with associated principal component (PC) time series (PSA1 and PSA2; see methods section) whose phases are nearly in quadrature with each other and whose explained variances are of nearly equal amplitude ^[12, 13]. In combination, they produce the single propagating PSA mode. The PSA mode is known to strongly influence the Antarctic cryosphere ^[14], wind and significant wave heights across the Southern Hemisphere oceans ^[13], and the South American monsoon system including rainfall ^[15] and weather and climate extremes over Brazil ^[16].

The respective PSA1 pattern has been widely recognised as being highly correlated with ENSO, where previous studies argue that it results in part as an atmospheric response to ENSO ^[15, 17, 18]. While regression analysis shows that there is indeed a very close relationship between PSA1 and ENSO (Fig. 1a), the origin of this relationship remains an active area of research ^[19, 20]. Meanwhile, the connection between the atmospheric PSA2 and tropical SST variability is less clear. Ref ^[15] argues that PSA2 is responsible for the quasi-biennial component of ENSO (Fig. 1b). Although ENSO-PSA correlations are widely documented, the processes by which ENSO and the PSA mode are dynamically connected, and, in particular, how the PSA mode feeds back to the tropics, remain unclear.

Diverse views exist on the mechanisms that link the atmospheric PSA mode with ENSO. Some studies ^[15, 17, 21, 22] argue that the ENSO-PSA connection is directly linked via atmospheric Rossby wave propagation. However, other studies ^[20, 23, 24] point out that synoptic scale Rossby waves are primarily generated and trapped locally within the Southern Hemisphere subtropical and polar jet

streams. Unlike the teleconnection between the Madden-Julian oscillation and North Atlantic oscillation ^[25], there is little direct dynamical evidence in terms of Rossby wave source or wave activity flux (see, for example, refs ^[19, 20, 26]) to suggest that poleward propagating large-scale tropical Pacific Rossby waves, potentially initiated by ENSO variability, teleconnect the PSA to ENSO. Even if sufficient Rossby wave sources were to be generated in the equatorial Pacific, ray tracing theory and wave activity flux calculations make it readily apparent that they are blocked by a reflecting barrier associated with the presence of the subtropical jet, and low latitude wave breaking, from propagation to the midlatitudes. Instead, Ref ^[27] argues that the PSA mode is linked to ENSO via a direct modulation of the midlatitude jets by the thermal winds generated by tropical convection that is highly correlated with ENSO. This indirectly influences the interannual variability of coherent synoptic features forming within these midlatitude waveguides where local Rossby waves sources are prevalent.

Oceanic signal driven by the PSA mode

Analysis of South Pacific SST shows that the first two modes of SST variability, referred to as the SPDO ^[3, 4] and South Pacific quadrupole SST pattern ^[8], are primarily driven by the variability associated with the atmospheric PSA1 and PSA2 patterns (Figs. 1 e and f). Using a univariate first-order autoregressive (AR1) model ^[28, 29], we found that the integrated atmospheric PSA1 and PSA2 variability could explain a significant fraction of the variance of the first two SST modes in the South Pacific Ocean (Figs. 1 e and f; $R=0.76$ and 0.71 , respectively; significant at $>99\%$ level; see the methods section for details).

A second integration of the AR1 model (following a similar approach to refs ^[4, 30]) – with the first integration being from the atmosphere to the surface ocean, and the second integration from the surface ocean to the subsurface ocean – reveals that the leading SST mode is further reddened by the extratropical upper ocean (Fig. 1 e). The integrated SPDO signal (blue curve in Fig. 1 e) resembles the time evolution of the leading vertically averaged temperature (VAT) mode through the upper 300 m of the ocean (see the methods section), with a temporal correlation of $R=0.88$ (statistically significant at the 99% level). To first order, we conclude that the South Pacific Ocean integrates the fast-varying atmospheric “noise” forcing to produce a “reddened” oceanic signal, with a pronounced increase in amplitude of the spectrum at low frequencies and decreased amplitude at high frequencies, where the main potentially predictable signal is generated on the time scales of the ocean dynamics.

Oceanic noise driven by the PSA mode

The extratropical South Pacific not only responds via reddening processes to the fast-varying atmosphere but also via coherence resonances ^[31]. Specifically, the forcing due to coherent synoptic scale disturbances in the atmosphere associated with the PSA imprints onto the surface ocean further enhancing internal SST variability at its preferred frequency. To examine this mechanism, we generalised the univariate AR1 model to a higher-dimensional multivariate field through inclusion of SST anomalies in the tropical and South Pacific oceans using a linear inverse model ^[32] (LIM; see methods section). Previous studies have shown that the tropical and South Pacific oceans can be approximated as a stochastically forced linear system, where different dynamical processes are represented by distinct damping time scales via the decomposition of the LIM ^[5, 32-35].

Fig. 2 shows the fastest damped SST mode pair with a damping time scale of two months. This pair of complex patterns (Figs. 2 a and b) constitutes a single propagating wave train with a complete cycle of 26 months. Along with the propagating features, the spatial patterns of the fastest damped modes also bear a strong resemblance to the atmospheric PSA1 and PSA2 patterns. Projecting the corresponding time series of the real and imaginary components of the fastest damped mode onto the monthly 500-hPa geopotential height (Z500) anomalies produces regression maps (Figs. 2 c and d) that closely resemble the atmospheric PSA1 and PSA2 with spatial correlations of 0.74 and 0.69 (statistically significant at the 95% level), respectively, in the South Pacific region. This suggests that the high-frequency atmospheric PSA fluctuations can excite SST modes whose frequencies are subject to the atmospheric drivers. In contrast to the reddening processes that act to excite the potentially predictable low frequency ocean signal, the pair of most damped SST modes are correlated with the atmospheric PSA forcing that acts as a stochastic forcing on the SST system.

The optimal growth and extratropical precursor

Previous studies have extensively discussed the tropical and extratropical precursors of ENSO [8, 10, 32, 36, 37]. Despite there being little direct evidence, in terms of atmospheric dynamics, to show that the atmospheric PSA mode can directly modulate ENSO evolution in the tropics [27], the indirect influence has nevertheless been widely identified in both observations [11] and model simulations [36, 38] via the “atmosphere → extratropical ocean → tropical ocean” pathway. One way in which this indirect influence may occur is through a “seasonal footprinting” mechanism [39], where the atmospheric forcing drives an extratropical anomalous SST “footprint” in the boreal

spring, which persists through the boreal summer, and sustains wind stress anomalies in the tropics that are crucial to initiate ENSO events.

The LIM approach enables an objective determination of the optimal initial perturbations that maximise, for example, ENSO and SPDO growth. This provides an ideal framework through which to investigate the dynamical precursors and predictability of the peak phases of ENSO and the SPDO. Unlike lead-lag correlations that have been widely applied to identify ENSO precursors [8, 9, 11], the LIM represents a multivariate linear stochastically forced model and provides a dynamical approximation (see methods section) that satisfies conditionally causal^[40] relationships between the optimal precursors and their peak phases.

The interactions between the non-normal damping SST modes can give rise to a transient amplification of the variance of the deterministic SST system at a preferred temporal growth scale. Such transient amplification is useful in sampling and interpreting errors in initial conditions^[41] and explains the actual variance growth in the system^[32]. Transient amplification of monthly tropical and South Pacific SST anomalies allows a specific set of initial perturbations to develop into the peak phases that are found to exist at 6- to 10-month lead times [5, 9, 32, 33, 42]. Our LIM experimental results show that the optimal growth time of SST anomalies in the tropical and South Pacific oceans (see methods section) is 9 months. These results imply the existence of linear growth events that act to maximise ENSO and SPDO development leading to enhanced predictability if the initial perturbations can be sufficiently well specified.

The spatial patterns of the initial SST perturbations in the tropical Pacific and South Pacific are shown in Figs. 3 a and c, respectively. These initial patterns co-evolve over nine months into the optimal final peak phases in Figs. 3 b and d, which are closely associated with ENSO and the SPDO with pattern correlations of 0.99 and 0.97 (statistically significant at the 99% level),

respectively. The tropical ENSO precursor (Fig. 3 a) has been discussed in previous studies [32, 33, 37] and may be associated with the recharge-discharge mechanism as first described by Jin [43]. The extratropical precursor (Fig. 3 c) resembles a quadrupole structure similar to the second South Pacific SST mode with the pattern correlation of 0.85 (significant at the 95% level). The initial and final spatial patterns (Figs. 3 c and d) were projected onto the monthly SST anomalies to reconstruct the time series. The reconstructed time series of the optimal initial perturbations and final peak phases in the South Pacific (Figs. 3 e and f) are highly correlated with the second and first South Pacific SST modes with temporal correlations of 0.86 and 0.98 (statistically significant at the 99% level), respectively. This suggests that the South Pacific quadrupole SST pattern, which is primarily driven by the atmospheric PSA2, is the optimal local (linear) precursor that maximises the SPDO growth. Given the almost synchronous manifestation of feature relationships between ENSO and the SPDO [3, 4], these results imply that the extratropical SST precursor of ENSO is also related to the quadrupole SST pattern in the South Pacific Ocean. Our results support the “seasonal footprinting” mechanism whereby the atmospheric PSA2 excites the South Pacific quadrupole SST anomalies, which then persist through the boreal summer to guide ENSO growth.

A new paradigm for South Pacific climate variability and predictability

Fig. 4 summarises the atmospheric PSA mode and its South Pacific Ocean responses based on the combined results from the AR1 and LIM investigations. Our results suggest that the eastward-propagating PSA mode provides an important source of atmospheric forcing to excite the extratropical South Pacific Ocean responses operating on multiple time scales via reddening processes and coherence resonances. To first order, the close relationships between the integrated PSA1 and PSA2 patterns and the leading South Pacific SST modes support the general concept

that fast atmospheric variations are a critically important excitation source of low frequency oceanic variability. The leading integrated subsurface temperature mode, that resembles the spatial pattern of its SST counterpart, can be regarded as a cumulative response to atmospheric PSA1 forcing. That is, the inclusion of extratropical subsurface processes further reddens the SST variations thereby enhancing the most persistent and potentially predictable signal. Furthermore, the atmospheric PSA1 and PSA2 patterns can excite SST variations via coherence resonances such that the resultant fastest damping SST modes are slaves to the atmospheric forcing but synchronised to the spatiotemporal features of the real and imaginary components of the propagating PSA.

The extratropical SST precursor of ENSO and SPDO growth is strongly associated with the South Pacific quadrupole SST pattern. The set of initial conditions related to this quadrupole pattern can optimally determine how the SST anomalies will evolve along its deterministic (linear) trajectory and lead to the SPDO and ENSO peaks over the following nine months, and from which the linear predictability intrinsic to the SST system can be inferred. While the influence of the atmospheric PSA mode on tropical SST variability is less apparent than the influence of the tropical SST variability on the PSA mode ^[8, 20, 27], our results indicate that the South Pacific quadrupole SST pattern acts as an oceanic bridge that links the atmospheric PSA forcing to ENSO dynamics centred in the tropics, and represents the most probable pathway for midlatitude atmospheric variability to influence tropical SST via an indirect “seasonal footprinting” mechanism. Although the actual observed nonlinear dynamics are more complicated than the tangent linear dynamics that form the basis of the LIM propagator, the essential dynamics of the system are captured. The proposed paradigm provides a new mechanistic framework to understand the dynamics and predictability of

221 South Pacific climate variability and to interpret extratropical atmospheric drivers and the oceanic
 222 responses on multiple time scales.

223
 224 **Fig. 1 | Atmospheric forcing of the leading two SST modes in the South Pacific Ocean. a, b,**
 225 The PSA1 (a) and PSA2 (b) patterns are obtained by regressing the monthly near-global Z500
 226 anomalies (NCEP-NCAR; contour) and monthly SST anomalies (ACCESS-O; shade) onto the
 227 PSA1 and PSA2 time series, respectively. (c) and (d), The corresponding normalised PSA1 and
 228 PSA2 time series. (e) and (f), The time series of the leading two SST modes of variability in the
 229 South Pacific Ocean (ACCESS-O; black) are reconstructed (red) using an AR1 model forced by

the PSA1 and PSA2 time series. The blue curve in (e) indicates the second integration forced by the SST SPDO time series. The units are in standard deviations (s.d.). The significance of the temporal correlations is estimated taking account of the effective number of degrees of freedom due to serial correlation^[44].

Fig. 2 | The fastest damped SST modes. a, b, The imaginary (a) and real (b) parts of the fastest damped SST modes are obtained using a reduced-order linear inverse model (see the text for details). The pair of SST modes shown has the fastest damping time scale of 2.3 months. c, d, the maps are obtained by regressing the monthly Z500 anomalies (NCEP-NCAR) onto the time series of the imaginary (c) and real (d) parts of the fastest damped modes.

Fig. 3 | The optimal evolution of the tropical and South Pacific SST. The optimal initial structures in the (a) tropical Pacific and (c) South Pacific are obtained using a linear inverse model (see the text for details), which linearly co-evolve over 9 months into their peak phases in the (b) tropical Pacific and (d) South Pacific. The spatial patterns shown in a-d are normalised according to the variance in each domain. e and f, the reconstructed time series of the (e) optimal initial structure and (f) peak phase in the South Pacific (black) are compared with the time series of the leading two South Pacific SST modes (red).

Fig. 4 | Schematic of new paradigm for South Pacific Ocean variability and predictability.

The top panels are the same as shown in Figs. 2 a and b indicating the fastest damped mode pair.

The middle panels are the second and third modes of the monthly Z500 anomalies (NCEP-NCAR) in the southern hemisphere. The left panels of the bottom layer are the leading EOF modes of the monthly SST anomalies and VAT anomalies in the South Pacific Ocean, respectively. The right panel of the bottom layer is the same as shown in Fig. 3 c indicating the optimal precursor of the SPDO (illustrated by the green arrow). Red arrows indicate reddening processes. Blue arrows indicate coherence resonances. Grey arrows indicate the propagating features of the corresponding modes.

260 **References**

- 261 1. Mantua, N. J. & Hare, S. R. The Pacific decadal oscillation. *J. Oceanogr.* **58**, 35-44 (2002).
- 262 2. Power, S. et al. Inter-decadal modulation of the impact of ENSO on Australia. *Clim. Dynam.* **15**,
- 263 319-324 (1999).
- 264 3. Chen, X. & Wallace J. M. ENSO-like variability: 1900–2013*. *J. Climate* **28**, 9623-9641(2015).
- 265 4. Lou, J., Holbrook, N. J. & O’Kane T. J. South Pacific decadal climate variability and potential
- 266 predictability. *J. Climate* **32**, 6051-6069 (2019).
- 267 5. Lou, J., O’Kane, T. J. & Holbrook, N. J. A linear inverse model of tropical and South Pacific
- 268 seasonal predictability. *J. Climate* **33**, 4537-4554 (2020).
- 269 6. Taschetto, A.S. et al. ENSO atmospheric teleconnections, *El Niño Southern Oscillation in a*
- 270 *Changing Climate* 309-335 (2020).
- 271 7. Sprintall, J. et al. ENSO oceanic teleconnections, *El Niño Southern Oscillation in a Changing*
- 272 *Climate* 337-359 (2020).
- 273 8. Ding, R., Li, J. & Tseng, Y.-h. The impact of South Pacific extratropical forcing on ENSO and
- 274 comparisons with the North Pacific. *Clim. Dynam.* **44**, 2017-2034 (2015).
- 275 9. Zhao, Y. & Di Lorenzo, E. The impacts of extra-tropical ENSO precursors on tropical Pacific
- 276 decadal-scale variability. *Sci. Rep.* **10**, 3031 (2020).
- 277 10. Zhang, H., Clement, A. & Di Nezio, P. The South Pacific meridional mode: a mechanism for
- 278 ENSO-like variability. *J. Climate* **27**, 769-783 (2014).
- 279 11. You, Y. & Furtado, J. C. The role of South Pacific atmospheric variability in the development of
- 280 different types of ENSO. *Geophys. Res. Lett.* **44**, 7438-7446 (2017).
- 281 12. Lau, K.M., Sheu, P. J. & Kang, I.S. Multiscale low-frequency circulation modes in the global
- 282 atmosphere. *J. Atmos. Sci.* **51**, 1169-1193 (1994).
- 283 13. Echevarria, E.R. et al. Influence of the Pacific-South American modes on the global spectral
- 284 wind-wave climate. *J. Geophys. Res. Oceans* **125**, e2020JC016354 (2020).
- 285 14. Irving, D. & Simmonds, I. A new method for identifying the Pacific–South American pattern and
- 286 its influence on regional climate variability. *J. Climate* **29**, 6109-6125 (2016).
- 287 15. Mo, K.C. & Paegle, J. N. The Pacific-South American modes and their downstream effects. *Int. J.*
- 288 *Climatol.* **21**, 1211-1229 (2001).
- 289 16. Rehbein, A. et al. Severe weather events over southeastern Brazil during the 2016 dry season.
- 290 *Adv. Meteorol.* **2018**, 4878503 (2018).
- 291 17. Karoly, D. J. Southern Hemisphere circulation features associated with El Niño–Southern
- 292 Oscillation events. *J. Climate* **2**, 1239-1252 (1989).
- 293 18. Cai, W. et al. Climate impacts of the El Niño–Southern Oscillation on South America. *Nat. Rev.*
- 294 *Earth Environ.* **1**, 215-231(2020).
- 295 19. Karoly, D. J., Plumb, R.A. & Ting, M. Examples of the horizontal propagation of quasi-stationary
- 296 waves. *J. Atmos. Sci.* **46**, 2802-2811(1988).
- 297 20. Li, X. et al. A Rossby wave bridge from the tropical Atlantic to west Antarctica. *J. Climate* **28**,
- 298 2256-2273 (2015).
- 299 21. Renwick, J. A. & Revell, M. J. Blocking over the South Pacific and Rossby wave propagation.
- 300 *Mon. Weather Rev.* **127**, 2233-2247 (1999).
- 301 22. Cai, W. et al. Teleconnection pathways of ENSO and the IOD and the mechanisms for impacts on
- 302 Australian rainfall. *J. Climate* **24**, 3910-3923 (2011).
- 303 23. Ambrizzi, T. & Hoskins, B.J. Stationary rossby-wave propagation in a baroclinic atmosphere. *Q.*
- 304 *J. R. Meteorol. Soc.* **123**, 919-928 (1997).
- 305 24. Ambrizzi, T., Hoskins, B.J. & Hsu, H.-H. Rossby wave propagation and teleconnection patterns
- 306 in the austral winter. *J. Atmos. Sci.* **52**, 3661-3672 (1995).
- 307 25. Lin, H. & Brunet, G. Extratropical response to the MJO: nonlinearity and sensitivity to the initial
- 308 state. *J. Atmos. Sci.* **75**, 219-234 (2018).

26. O’Kane, T. J. et al. On the dynamics of persistent states and their secular trends in the waveguides of the Southern Hemisphere troposphere. *Clim. Dynam.* **46**, 3567-3597 (2015).
27. O’Kane, T. J., Monselesan, D. P. & Risbey, J. S. A multiscale reexamination of the Pacific–South American pattern. *Mon. Weather Rev.* **145**, 379-402 (2017).
28. Frankignoul, C. & Hasselmann, K. Stochastic climate models, Part II Application to sea-surface temperature anomalies and thermocline variability. *Tellus* **29**, 289-305 (1977).
29. Hasselmann, K. Stochastic climate models Part I. Theory. *Tellus* **28**, 473-485 (1976).
30. Di Lorenzo, E. & Ohman, M. D. A double-integration hypothesis to explain ocean ecosystem response to climate forcing. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 2496-9 (2013).
31. Pierini, S. Low-frequency variability, coherence resonance, and phase selection in a low-order model of the wind-driven ocean circulation. *J. Phys. Oceanogr.* **41**, 1585-1604 (2011).
32. Penland, C. & Sardeshmukh, P. D. The optimal growth of tropical sea surface temperature anomalies. *J. Climate* **8**, 1999-2024 (1995).
33. Capotondi, A. & Sardeshmukh, P. D. Optimal precursors of different types of ENSO events. *Geophys. Res. Lett.* **42**, 9952-9960 (2015).
34. Newman, M. Interannual to decadal predictability of tropical and North Pacific sea surface temperatures. *J. Climate*, **20**, 2333-2356 (2007).
35. Penland, C. & Magorian, T. Prediction of Niño 3 sea surface temperatures using linear inverse modeling. *J. Climate* **6**, 1067-1076 (1993).
36. Liguori, G. & Di Lorenzo, E. Separating the North and South Pacific meridional modes contributions to ENSO and tropical decadal variability. *Geophys. Res. Lett.* **46**, 906-915 (2019).
37. Penland, C. & Matrosova, L. Studies of El Niño and interdecadal variability in tropical sea surface temperatures using a nonnormal filter. *J. Climate* **19**, 5796-5815 (2006).
38. Zhang, H. et al. Equatorial signatures of the Pacific meridional modes: Dependence on mean climate state. *Geophys. Res. Lett.* **41**, 568-574 (2014).
39. Vimont, D. J., Wallace, J.M. & Battisti, D. S. The seasonal footprinting mechanism in the Pacific: Implications for ENSO. *J. Climate* **16**, 2668-2675 (2003).
40. Granger, C. W. J. Some recent development in a concept of causality. *J. Econom.* **39**, 199-211 (1988).
41. Moore, A. M. & Kleeman, R. The dynamics of error growth and predictability in a coupled model of ENSO. *Q. J. R. Meteorol. Soc.* **122**, 1405-1446 (1996).
42. Newman, M., Alexander, M. A. & Scott, J. D. An empirical model of tropical ocean dynamics. *Clim. Dynam.* **37**, 1823-1841(2011).
43. Jin, F.-F. An equatorial ocean recharge paradigm for ENSO. Part I: Conceptual model. *J. Atmos. Sci.* **54**, 811-829 (1997).
44. Davis, R.E. Predictability of sea surface temperature and sea level pressure anomalies over the North Pacific Ocean. *J. Phys. Oceanogr.* **6**, 249-266 (1976).
45. Kalnay, E. et al. The NCEP/NCAR 40-year reanalysis project. *Bull. Am. Meteorol. Soc.* **77**, 437-472 (1996).
46. O’Kane, T. J. et al. Storm tracks in the Southern Hemisphere subtropical oceans. *J. Geophys. Res. Oceans* **119**, 6078-6100 (2014).
47. von Storch, H. et al. Principal oscillation patterns: A review. *J. Climate* **8**, 377-400 (1995).
48. Lou, J., O’Kane, T.J. & Holbrook, N. J. A linear inverse model of tropical and South Pacific climate variability: Optimal structure and stochastic forcing. *J. Climate* **34**, 143-155 (2021).

Methods

The reanalysis data used here were monthly detrended anomalies of the 500-hPa geopotential heights (Z500) from the National Centers for Environmental Prediction/National Center for Atmospheric Research (NCEP-NCAR) ^[45]. The monthly detrended sea surface temperature (SST) anomalies were generated using an atmosphere-forced ocean model, the Australian Community Climate and Earth-System Simulator-Ocean (ACCESS-O). The model configuration is described in ref ^[46]. The original 360×300 tripolar ACCESS-O model grid has been remapped to a regular $2.5^\circ \times 2.5^\circ$ grid. The reanalysis data and model simulations were linearly detrended at each grid point, with monthly anomalies calculated by removing the climatological monthly mean.

The atmospheric PSA1 and PSA2 patterns were defined by computing the second and third empirical orthogonal functions/principal components (EOFs/PCs) of the monthly Z500 anomalies in the Southern Hemisphere, respectively. The leading two South Pacific SST modes, which are referred to as the South Pacific decadal oscillation (SPDO) and the South Pacific quadrupole SST pattern, were obtained by computing the first and second EOFs/PCs of the monthly SST anomalies in the South Pacific Ocean domain from (20°S-70°S; 120°E-60°W). The subsurface temperature variability was characterised by spatiotemporal variations of vertically averaged temperatures (VAT) through the upper 300-m of the ocean, that includes the mixed layer and thermocline across most regions. The subsurface SPDO is derived from the first EOF/PC of the VAT in the South Pacific Ocean (20°S-70°S; 120°E-60°W). Statistical significance tests of the correlation coefficients calculated in this study took account of the effective number of degrees of freedom due to serial correlation in the time series, following the method of ref ^[44].

378 The oceanic reddening response to atmospheric PSA forcing was quantified using a doubly
379 integrated first-order autoregressive (AR1) model. A univariate AR1 model can be written as

$$380 \quad \frac{dx}{dt} = lx + \xi, \quad (1)$$

381 where x is a time series and $-1/l$ is associated with the damping time scale. This univariate AR1
382 process only has one degree of freedom and is unable to oscillate when the damping coefficient l
383 is positive ^[47]. Predictability of the time series x is largely limited by the damping scale $-1/l$. ξ
384 in Eq. (1) represents a high-frequency noise forcing term.

385 The optimal damping time scales are 6 months and 5 months, respectively, when the time series
386 of the PSA1 and PSA2 patterns were specified as the atmospheric forcing of the first two South
387 Pacific SST modes in the first AR1 integration from the atmosphere to the surface ocean. In the
388 second integration from the surface ocean to the subsurface ocean, the optimal damping time scale
389 is 15 months when the time series of the SST SPDO was specified as the forcing of the leading
390 South Pacific VAT mode.

391 The pair of fastest damped noise mode and optimal initial perturbations were estimated using a
392 generalised AR1 model, here referred to as the linear inverse model (LIM) ^[32], or alternatively
393 termed principal oscillation pattern (POP) analysis ^[47]. The LIM can be written in the form of a
394 linear stochastic differential equation:

$$395 \quad \frac{d\mathbf{X}}{dt} = \mathbf{L}\mathbf{X} + \boldsymbol{\xi} \quad (2)$$

396 where the evolution of the state vector $\mathbf{X}$ can be expressed as the sum of the deterministic dynamics,
397 $\mathbf{L}\mathbf{X}$, that constitute “slow” processes and the stochastic forcing term, $\boldsymbol{\xi}$, that constitutes “fast”
398 processes. In this study, we defined the model state vector $\mathbf{X}$ as

$$\mathbf{X} = \begin{bmatrix} \mathbf{SST}_{\text{TP}} \\ \mathbf{SST}_{\text{SP}} \end{bmatrix}, \quad (3)$$

where we considered monthly SST anomalies from the tropical Pacific (i.e., $\mathbf{SST}_{\text{TP}}$ in the region 20°S-20°N, 120°E-60°W) and South Pacific Ocean (i.e., $\mathbf{SST}_{\text{SP}}$ in the region 22.5°S-70°S, 120°E-60°W). We retained the leading 4 tropical Pacific and 6 South Pacific SST EOFs/PCs in the LIM to reduce the spatial degrees of freedom and ensure linearity and stability of the SST system. The robustness of the LIM has been demonstrated in the refs ^[5, 48], showing that the choices of using the leading 10 to 16 EOFs/PCs to construct the state vector $\mathbf{X}$ and the different combinations of the leading EOFs/PCs used from the different regions give similar results.

The dynamical operator $\mathbf{L}$ in Eq. (2) encapsulates the evolution of the state vector $\mathbf{X}$ and can be estimated from

$$\mathbf{L} = \tau_0^{-1} \ln[\mathbf{C}(\tau_0)\mathbf{C}(0)^{-1}], \quad (4)$$

where $\mathbf{C}(\tau_0)$ and $\mathbf{C}(0)$ are the time-lagged and zero-lagged (cross-)covariance matrices of the state vector $\mathbf{X}$. That is, $\mathbf{C}(\tau_0) = \langle \mathbf{X}(t + \tau_0)\mathbf{X}^T(t) \rangle$ and $\mathbf{C}(0) = \langle \mathbf{X}(t)\mathbf{X}^T(t) \rangle$, where the angle brackets here denote an ensemble average or a time average over all t for variables with stationary statistics. Here, $\tau_0 = 1$ month was used to ensure the stability of the LIM. Previous studies ^[5, 48] examined the sensitivity of using different choices of τ_0 to estimate the dynamical operator $\mathbf{L}$, finding that the LIM is insensitive to τ_0 ranging from 1 month to 6 months.

For a system described by Eq. (2), the most probable evolution at time $t + \tau$ can be determined by

$$\mathbf{X}(t + \tau) = \mathbf{G}(\tau)\mathbf{X}(t) = e^{\mathbf{L}\tau}\mathbf{X}(t), \quad (5)$$

given a state vector $\mathbf{X}(t)$.

The damping modes can then be diagnosed by applying eigen-decomposition to the dynamical operator $\mathbf{L}$ (i.e., $\mathbf{L}\mathbf{p} = \lambda\mathbf{p}$). Since the $\mathbf{L}$ matrix is not symmetric, some or all of its eigenvalues λ and eigenvectors $\mathbf{p}$ are complex. The damping time scales $-1/\sigma$ and/or the oscillatory periods $2\pi/\omega$ of the damping modes can be directly estimated from the corresponding eigenvalues $\lambda = \sigma + i\omega$, where σ and ω are the real and imaginary parts of λ , respectively. In this study, the pair of complex damping modes that has the fastest damping time scale has been discussed.

The transient growth over any time interval τ can be written as the norm of the final state $\mathbf{X}(t)$ divided by the norm of the initial state $\mathbf{X}(0)$:

$$\gamma^2 = \frac{\|\mathbf{X}(t)\|^2}{\|\mathbf{X}(0)\|^2} = \frac{\mathbf{X}^T(t)\mathbf{X}(t)}{\mathbf{X}^T(0)\mathbf{X}(0)} = \frac{\mathbf{X}^T(0)\mathbf{G}^T(\tau)\mathbf{G}(\tau)\mathbf{X}(0)}{\mathbf{X}^T(0)\mathbf{X}(0)}. \quad (6)$$

The optimal growth that maximises the amplification of the variance can be estimated as the leading eigenvalue γ_1^2 of $\mathbf{G}^T(\tau)\mathbf{G}(\tau)$. The leading eigenvalues at different time intervals [i.e., $\gamma_1^2(\tau)$] then form a trajectory that indicates the modal interferences and variance changes of the system in the absence of stochastic forcing. Since the dynamical operator of the LIM is nonnormal (i.e., $\mathbf{L}^T\mathbf{L} \neq \mathbf{L}\mathbf{L}^T$), it allows the resultant damping modes to interact with each other and contribute to the transient amplification of the variance without the presence of stochastic forcing. This provides a perspective to, for example, interpret errors in the initial conditions and understand the error growth of the system. In the present LIM, the largest transient amplification of the SST variance was found at 9 months.

The optimal precursor indicates the initial condition that maximises the growth of the state vector over a specified time interval τ ($\tau = 9$ months in this case). The optimal precursor is derived as the corresponding leading eigenvector of $\mathbf{G}^T(\tau)\mathbf{G}(\tau)$ when $\tau = 9$ months is specified.

Data availability

The atmospheric data used in this work are freely available from <https://psl.noaa.gov/data/gridded/data.ncep.reanalysis.pressure.html>. The model configurations and model outputs used in this work are available on request from TJO. The AR1 and LIM codes are available on request from JL.

Acknowledgements

JL is grateful for a PhD scholarship provided by the ARC Centre of Excellence for Climate System Science (CE110001028), a University of Tasmania tuition fee scholarship, and a CSIRO-UTAS Quantitative Marine Science top-up scholarship [including support from the Australian Commonwealth Scientific Research Organisation (CSIRO) Postgraduate scheme]. TJO was supported by the CSIRO Decadal Climate Forecasting Project (<https://research.csiro.au/dfp>). NJH acknowledges funding support from the ARC Centre of Excellence for Climate Extremes (CE170100023) and the Australian Government National Environmental Science Programme Climate Systems Hub.

Author contributions

JL conceived the work, carried out the data analyses and wrote most of the paper. TJO supplied the ACCESS-O model outputs. TJO and NJH contributed to the interpretation of the model results and analysis, and also to the text.

462 **Competing interests**

463 The authors declare no competing interests.

464