Resilience, Occupational Burnout, and Parenting Stress in Nurses Caring for COVID-2019 Patients

Saeid Bashirian

Hamadan University of Medical Sciences School of Public Health

Mostafa Bijani

Fasa University of Medical Science

Seyed Reza Borzou

Hamadan University of Medical Sciences School of Nursing and Midwifery

Salman Khazaei

Hamadan University of Medical Sciences Medical School

Fateme Mohammadi (✉ mohammadifateme47@yahoo.com)

Hamadan University of Medical Sciences School of Nursing and Midwifery

Research article

Keywords: occupational burnout, resilience, parenting stress, nurses, coronavirus, COVID 2019

Posted Date: October 29th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-60538/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Caring of patients with Corona Virus Infection Disease (COVID 2019) may have had an impact on the occupational burnout, resilience, and parenting of the nurses.

Purpose: Evaluation occupational burnout, resilience, and parenting stress in nurses caring for COVID 2019 patients.

Methods: The present study is a cross-sectional study Base on Strengthening the Reporting of Observational Studies in Epidemiology Statement (STROBE). 420 nurses cared for COVID 2019 patients in 5 hospitals selected via convenience sampling. Each participant completed the scales online. Data were analyzed in SPSS v. 22.

Results: Occupational burnout has a negative correlation with resilience and a positive correlation with parenting stress and can predict 61.32% of changes in the occupational burnout variance of nurses.

Conclusion: the resilience, parenting stress, marital status, number of children, employment status, and gender predicated High percentage of the nurses' occupational burnout variance. The nurse managers should use these findings to provide appropriate environments for nurses, that to develop more comprehensive plans for support of nurses for the current and future crises.

Background

One of the most recent emerging diseases is Corona Virus Infection Disease (COVID 2019) which started in China in 2019, very quickly became a pandemic, and has caused a large number of deaths in many countries so far [1]. The initial symptoms of COVID 2019 are similar to those of the influenza, but the infection gradually develops and causes severe cardiopulmonary and renal disorders and can eventually cause death [2]. Because COVID 2019 is a little-known disease, the genetic structure of the virus constantly changes, new symptoms are continuously recorded in the infected, and, most importantly, there is not a definite cure or specialized vaccine for it, over 8.3 million people in the world have contracted the disease in the past three months, over 224,000 of whom have lost their lives as a result [3, 4]. This is while, the Iran outbreak of COVID 2019 is complex, At the beginning of the pandemics Iran was the third country with the highest number of reported COVID 2019 cases after China, and Italy, but, Iran has been heavily hit by the virus is now facing the third wave of the COVID 2019 outbreak compared with other countries that are currently facing the second wave. Although the majority of Iranians wear masks, but inappropriate economic situation has resulted to very few restrictions on work, activity and travel in all cities of Iran, therefore large numbers of people use public transportation every day, and the streets is crowded, In addition all employees are at work every day and telecommuting from home is very low [5]. These factors have caused about 530,500 people in Iran have infected with COVID 2019 and over 30,500 of whom have lost their lives [5, 6].

The high fatality rate of COVID-19 worldwide has caused many people to experience considerable tension with adverse effects on their social activities and psychological security [7]. Due to the special nature of their occupation, nurses are at higher risk of suffering serious physical and psychological harms, including stress and depression. It is obvious that these psychological crises can influence nurses' personal and family responsibilities as well, subject them to occupational burnout, and adversely affect their performance of their parental duties [8, 9]. In January 2020, WHO declared the COVID-19 pandemic to be a public health emergency of international concern (PHEIC) [10]. According to the same organization, nurses play a significant role in healthcare and treatment systems in caring for and improving the health of patients infected with the coronavirus [1, 8]. However, caring for COVID-19 patients has subjected nurses to great psychological tension and work stress, thus the need for taking measures to protect nurses' psychological and physical well-being [1]. Accordingly, Stuijzand et al. (2020) state that healthcare professionals during epidemic and pandemic ([Severe Acute Respiratory Syndrome\(SARS\)](#), [Middle East respiratory syndrome coronavirus\(MERS\)](#), [Ebola Virus Disease](#)), experience a lot of mental health problems including psychological distress, insomnia, alcohol/drug misuse, and symptoms of posttraumatic stress disorder (PTSD), depression, anxiety, anger and burnout [11]. Also, Lehmann et al. (2020) represent that health care professionals reported Depression, stress, social isolation, and job fatigue during Ebola outbreak. Social isolation has been largely due to parenting stress and concerns about the transmission of the disease to their children and other family members, On the other hand, although job stress and fatigue can affect their tolerance, but they have overcome these tensions and work pressures with team working and observing the principles of infection control [12]. A study conducted by Liu et al. (2020) confirms that, compared to other members of medical staff, nurses experience higher levels of anxiety and depression in caring for COVID-19 patients and face more difficulties in the performance of not only their professional duties, but family responsibilities and daily activities [7]. Similarly, he study of Roy (2020) shows that nurses are experiencing high degrees of stress, anxiety, depression, and post-traumatic stress disorder (PTSD) in the COVID-19 crisis. Thus, it is essential that the executive administrators of healthcare systems and the crisis committees of hospitals consider the status of nurses' psychological well-being, anxiety, depression, and occupational burnout [13]. The present study is an attempt at measuring occupational burnout, resilience, and parenting stress in nurses who care for COVID-19 patients in Iran.

1. Methods

1.1 Study design and setting

This study was a cross-sectional research investigation conducted Base on Strengthening the Reporting of Observational Studies in Epidemiology Statement (STROBE) from February to June 2020. The two following aims were examined in study "evaluation of occupational burnout, resilience, and parenting stress in nurses who care for COVID-19 patients" and "investigating the relationship between occupational burnout with resilience, parenting stress and demographic characteristics in nurses who care for COVID-19 patients".

1.2 Participants and sampling

In this study sample size has been estimated 630 samples according to the study of Roy et al. With $\beta = 80\%$ and $\alpha = 0.05$ and taking into account the 10% drop in each group. The participants were selected via convenience sampling. So nurses, who provided care to COVID-2019 patients in 5 hospitals affiliated with University of Medical Sciences in the west of Iran, were invited and selected via convenience sampling to participate in the study. The inclusion criteria were: being willing to participate, being in practice in one of the hospitals assigned for Covid-19 patients, and having at least one month of work experience in caregiving wards for patients with COVID2019. The subjects who failed to answer more than half of the items on their questionnaires or did not turn in their questionnaires were excluded. The subjects were asked to complete and submit the questionnaires—a personal (demographic) characteristics questionnaire, an occupational burnout scale, resilience scale, and, if they were parents, the parenting stress scale—online. The researchers sent emails and reminder messages to the participants, so that, the majority of the questionnaires (90%) were gathered in June. 420 of the subjects completed and returned the questionnaires via e-mail or a social network. Thus, the response rate was 66.7%. The nurses' reasons for not participating in this study were High prevalence of COVID2019 in Iran, heavy workload and intensive shifts, also infected with COVID2019.

1.3 Measurements

1.3.1 The occupational burnout scale

Developed by Maslach in 1996, the Maslach Burnout Inventory is the most commonly used scale for measuring occupational burnout [14]. The scale consists of 22 items which address three dimensions of occupational burnout: 9 items for emotional exhaustion, 5 items for depersonalization, and 8 items for personal accomplishment. The items measure the frequency and degree of burnout on an 8-point Likert scale: from 0 (very little) to 7 (very high). In the domain of emotional exhaustion, scores of 27 and above indicate severe emotional exhaustion, scores of 16 and below indicate slight emotional exhaustion, and scores of between 17 and 26 indicate average emotional exhaustion. In the domain of depersonalization, scores of above 13 indicate severe depersonalization and in the domain of personal accomplishment, scores of below 31 indicate low levels of personal accomplishment. The internal reliability of this scale to equal a Cronbach's alpha of 0.80 [14]. In their study, Baher et al. (2014) find the Cronbach's alpha of the scale to be 0.79 [15].

1.3.2 The self-report Caregiver Burden Inventory

The Caregiver Burden Inventory was developed by Novak and Guest in Canada in 1988 to measure caregivers' burden. The scale consists of 24 items which assess caregivers' burden in five domains: time-dependence burden (5 items (1, 2, 3, 4, 5)), developmental burden (5 items 6, 7, 8, 9, 10), physical burden (4 items 11, 12, 13, 14), social burden (5 items 15, 16, 17, 18, 19), and emotional burden (5 items (20, 21, 22, 23, 24)). Completion of the scale requires approximately 15 minutes. The items are scored on a 5-point Likert scale ranging from 0 (Not at all) to 4 (Absolutely). The score range is between 0 and 96: 0 to 32

indicate low burden, 33 to 64 indicate moderate burden, and 65 to 96 indicate great burden [16]. In their study conducted in Iran, Hormozi et al. report the reliability of the inventory to be 0.82, which is a satisfactory value [17].

1.3.3 The Parenting Stress Index-Short Form (PSI-SF)

Created by Abidin in 1995, the Parenting Stress Index comprises 3 subscales: parental distress (items 1 through 12), parent-child dysfunctional interactions (items 13 through 24), and child characteristics (items 25 through 36). The scale measures parental stress on a 5-point Likert scale ranging from 1 (=completely disagree) to 5 (=completely agree). [18]. In their studies, Yeh (2001) and Reitman (2002) report the reliability of the scale to be 0.89 and 0.90 respectively [19, 20]. Assessing the reliability and validity of the scale in Iran, Shirzadi finds the reliability of the various dimensions of the index to be between 0.59 and 0.86, which are acceptable values [21].

1.4 Statistical methods

After data were collected, they were analyzed using descriptive statistics (frequency, percentage, mean, and standard deviation) in SPSS v. 22. To investigate the relationship between occupational burnout on the one hand and resilience and parenting stress on the other, the researchers applied the chi-square test, independent t-test, and ANOVA. Significance level was set at 0.05. Subsequently, the variables of demographics, resilience, and parenting stress which were found to correlate with occupational burnout ($p < 0.25$) were entered into multiple linear regression with the backward technique. Before executing the analysis of multiple linear regressions, the researchers examined the assumptions of normality of data, homogeneity of variance, and independence of residuals.

2. Results

2.1 Demographic Information

Of the 420 nurses who participated in the study, 73.8% were female and 26.2% were male. The range of the participants' ages was between 23 and 55 years with the mean of 35.24 ± 3.56 years. The majority of the participants 73.8% were married, 54.8% of whom had two children. Also, most of the participants had a bachelor of science in nursing 63.1%, were contractual employees 50.5%, had 11 years of work experience, and worked 32 rotating shifts per month. The findings of the study showed that there was a statistically significant relationship between occupational burnout on the one hand and marital status, number of children, employment status, and gender on the other.

Table1. The participants' demographic characteristics and occupational burnout scores				
Demographic variables		Number (%)	Occupational burnout Means $\pm$SD	p-value
Age (years)	23-33	197 (46.9)	38 $\pm$ 1.21	0.324
	34-44	154 (36.7)	37 $\pm$ 1.96	
	45-55	69 (16.4)	37 $\pm$ 1.43	
Gender	Female	300 (71.4)	41 $\pm$ 2.12	0.042
	Male	120 (28.6)	38 $\pm$ 1.87	
Marital status	Single	110 (26.2)	32 $\pm$ 2.76	0.033
	Married	310 (73.8)	38 $\pm$ 1.54	
Number of children	None	12 (3.9)	35 $\pm$ 2.32	0.031
	1	80 (25.8)	38 $\pm$ 1.61	
	2	170 (54.8)	39 $\pm$ 2.14	
	More than 2	48 (15.5)	42 $\pm$ 1.87	
Education	Associate	73 (17.4)	31 $\pm$ 1.32	0.451
	Bachelor	265 (63.1)	32 $\pm$ 1.05	
	Master	82 (19.5)	32 $\pm$ 1.08	
Employment status	Permanent	208 (49.5)	33 $\pm$ 2.69	0.039
	Contractual	212 (50.5)	38 $\pm$ 1.57	
Work experience	Less than 10 years	150 (35.7)	36 $\pm$ 2.96	0.310
	10-20 years	197 (46.9)	37 $\pm$ 1.52	
	20-30 years	73 (17.4)	37 $\pm$ 2.24	

2.2 Resilience, occupational burnout and parenting stress in the participants

The resilience means score of the nurses who participated in the present study was 32.33 $\pm$ 2.57, the occupational burnout means score was 32.33 $\pm$ 2.57. Also, the parenting stress mean score of the 310 nurses who were married and had children was found to be 17.53 $\pm$ 1.58 during the COVID-19 crisis (Table 2).

Table2. The means and standard deviations of the participants' resilience, occupational burnout and parenting stress scores			
Variable	Dimension	Means $\pm$SD per dimension	Total Means $\pm$SD
Resilience	Time-dependence burden	14.32 $\pm$ 1.42	32.33 $\pm$ 2.57
	Developmental burden	15.76 $\pm$ 2.18	
	Physical burden	10.42 $\pm$ 1.53	
	Social burden	16.78 $\pm$ 1.34	
	Emotional burden	10.11 $\pm$ 1.28	
Occupational burnout	Emotional exhaustion	48 $\pm$ 3.22	32.33 $\pm$ 2.57
	Depersonalization	23 $\pm$ 2.13	
	Personal accomplishment	39 $\pm$ 2.38	
Parenting stress	Parental stress	18 $\pm$ 1.09	17.53 $\pm$ 1.58
	dysfunctional interactions	17 $\pm$ 2.53	
	Child characteristics	15 $\pm$ 1.12	

2.3 The relationship between the participants' resilience, occupational burnout, and parenting stress

The findings of the study show that there is a strong and inverse correlation between occupational burnout and resilience in nurses who care for COVID-19 patients ($p<0.001$, $r=-0.70$). Also, a more strong and inverse correlation p was found to exist between occupational burnout and resilience in married nurses than single nurses. The results showed a positive correlation between the occupational burnout and parenting stress scores of the nurses with children ($p<0.001$, $r=0.74$) (Table 3).

Table 3. The relationship between resilience, occupational burnout, and parenting stress in married and single nurses and the entire participants			
Occupational burnout	Resilience (married nurses)	$r= -0.71$	$p<0.001$
	Resilience (single nurses)	$r= -0.69$	$p<0.001$
	Resilience (total)	$r= -0.70$	$p<0.001$
Occupational burnout	Parenting stress (married nurses)	$r= 0.74$	$p<0.001$

2.4 The predictor variables of occupational burnout in nurses who care for COVID-19 patients

The variable of resilience, parenting stress, number of children, marital status, employment status, and gender which had a p -value of smaller than 0.25 were entered into multiple linear regressions with the backward technique. These variables remained in the model and accounted for about 61.32% of the occupational burnout variance in the nurses who provided care to COVID-19 patients (Table 4).

Table 4. The predictor variables of occupational burnout in nurses who care for COVID-19 patients					
Variable	Unstandardized coefficients		standardized coefficients	T	P-value
	B	Standard deviation	β		
Resilience	-0.634	2.53	-0.643	-3.32	0.001
Parenting stress	0.568	2.49	0.572	2.78	0.001
Number of children	0.113	2.28	0.441	2.54	0.012
Marital status	0.298	2.54	0.301	3.87	0.024
Employment status	0.382	2.34	0.214	1.98	0.022
Gender	0.331	1.89	0.186	1.64	0.039
Adjusted R2: 61.32%					

3. Discussion

This study showed nurses have reported high levels of occupational burnout low levels resilience and high levels tension in their relations with their children. Though a few studies have addressed the work stress, knowledge, and awareness of nurses who care for COVID-19 patients, there are not any studies of resilience, occupational burnout, or parenting stress in this group of caregivers. Therefore, the researchers had to use articles which measure resilience, occupational burnout, or parenting stress in nurses who care for patients with other specific diseases. The occupational burnout means score of the nurses was found to be 32.33 ± 2.57 , which indicates high burnout. Occupational burnout in nurses is a hot topic in nursing

associations: studies show that the physical and psychological pressures of caring for patients lead to occupational burnout in the long run and, thus, nurses' length of service, work shifts, and salary must be redefined according to the type of patients they care for and the units where they work [22]. However, most of these studies report the occupational burnout of nurses with 5 years' or more work experience to be average, which is not consistent with the findings of the present study [22-24]. This discrepancy can be attributed to the sudden emergence of COVID-19 which is highly infectious and has significantly increased nurses' workload and caused them to have to work longer shifts and stay away from their families in the past few months. The results of the present study show a statistically significant difference between the married and single participants in terms of their occupational burnout scores ($p < 0.033$). This difference may be due to the fact that, in addition to stress in the workplace, married nurses, especially those with children, are subject to greater psychological tension ($p < 0.031$) as a result of disruption in their performance of their duties as spouses and parents [22, 25-27]. The results of the study also show that contractual nurses ($p < 0.039$) and female nurses ($p < 0.041$) suffer from greater occupational burnout. Job insecurity, lower salaries, and the lower physical-psychological resilience of female nurses can account for the statistically significant difference between the occupational burnout scores of contractual and permanent nurses and male and female nurses [22, 27]. Similarly, the study of Salahian et al shows that nurses' employment status and gender have an impact on their occupational burnout [22]. In the present study, the resilience means score of the nurses was found to be 32.33 ± 2.57 , which is considered low [28, 29]. In contrast, other studies report average to high levels of resilience for nurses, even for those who provide care to critically or terminally ill patients. The discrepancy can be attributed to lack of a definite treatment for COVID-19 and the high speed and rate of the infection which have subjected nurses to severe physical and psychological tension with extremely adverse effects on their resilience [28-30]. The parenting stress means score of 298 of the participants was found to be 17.53 ± 1.58 . There are studies on parenting stress in working mothers or parents with a chronically ill child [31, 32]; however, there has not been much research into parenting stress in nurses. This lack of research may be due to the fact that nurses who work in rotating shifts can spend at least a few hours a day with their children and attend to their upbringing. But in the COVID-19 crisis, many nurses have to work longer hours caring for the infected and, due to the high infectiousness of the disease, would rather reside in hospitals temporarily or have their relatives take away their children to their own homes for the safety of their children. Stress in the workplace, fear of transmitting the infection to their children, having to stay away from their children, and inability to monitor their children's upbringing have subjected nurses with children to extra psychological tension and, consequently, increased occupational burnout in them.

The results of the present study show a strong and inverse correlation between the participants' occupational burnout with resilience, and a strong and direct correlation between the participants' occupational burnout with parental stress. In addition the variable of resilience, parenting stress, number of children, marital status, employment status, and gender described about 61.32% of the occupational burnout variance in the nurses who provided care to COVID-2019 patients. Study was not found that had evaluated and predicted the relationship between occupational burnout with resilience, parenting stress and demographic characteristic in nurses at the same time. Therefore, other studies were used that

examined the relationship between occupational burnout with resilience or occupational burnout with parenting stress or occupational burnout with demographic factors. In this regard, Amini et al. stated that there is an inverse relationship between occupational burnout with resilience of nurses [28]. Also, Zou et al. expressed occupational burnout was negatively related to resilience [33], these findings are consistent with the results of present study. In addition [Salahian](#) et al. stated nurses who worked in the infectious diseases ward reported more occupational burnout than nurses in other wards and job stress, role clarity, and workload predicted about 36.5% of the nurses' occupational burnout variance[22]. However nurses in present study provided care to COVID-2019 patients considering rapid transmission of the disease and their high workload, they have reported more occupational burnout and less resilience, which is consistent with the results of present study.

Rafiee et al, and Ahmadi et al. expressed individual characteristics of nurses (gender, work experience) and workplace are effective on nurses' occupational burnout, which is consistent with the present study [34, 35].

Finally, the results of the study show that nurses who provide care to COVID-19 patients and have had to work more shifts in recent months are affected by higher levels of occupational burnout, have less resilience, and are experiencing more tension in their relationship with their children than before the emergence of the pandemic. In view of the persistence of the COVID-19 crisis in Iran and the world in coming months, it is mandatory that health administrators take effective measures to reduce nurses' occupational burnout and enhance their physical and psychological well-being.

One of the major limitations of the present study was the relatively low return rate of the questionnaires via e-mail which could have been due to the hectic work schedules of nurses in the crisis. Moreover, the variables addressed in the present study were measured over a 6-month period—it is suggested that future studies assess occupational burnout, resilience, and parenting stress in nurses who care for COVID-19 patients in coming months and years in other societies and larger samples in order to acquire a more accurate understanding of nurses' occupational burnout in in this crisis. Health administrators and policy-makers can use these findings to develop more comprehensive plans for the current and future crises.

4. Conclusion

In the present study, the nurses who care for COVID-19 patients were found to be suffering from high levels of occupational burnout. Also, the results showed that nurses' occupational burnout correlates with their resilience, parenting stress, marital status, number of children, employment status, and gender: these variables predicated 61.32% of the subjects' occupational burnout variance. The nurse managers and policy makers of health organizations should examine and use the findings of this study in order to provide appropriate environments for nurses, also to develop more comprehensive plans for support of nurses for the current and future crises.

Declarations

Ethics approval and consent to participate

The institutional review board of the medical universities located in the west of Iran provided ethics approval (approval number: 9904242498). Also at the beginning of each interview, the researcher introduced herself and explained the goals of the study and assured that all information would remain confidential and that they could withdraw from the study at any time. The researchers provided the opportunity for participants to inform the researcher about their withdrawal from the study at any stage of the research and assured. Finally, the written consent was obtained from study participants.

Consent for Publication

Not applicable

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request

Competing interests

The authors declare that they have no competing interests.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not for profit sectors.

Authors' contributions

FM, SB, MB, and SRB were involved in the conception of the study and designed the study. They are responsible for data collection. Then FM, MB, SKH and SB analyzed data. FM, MF, SRB, SKH, SB and BT drafted the primary manuscript, revised and approved the final manuscript.

Acknowledgment

The present article is extracted from a research project registered under the ethical code (1399.194) at Hamadan University of Medical Sciences, Hamadan, Iran. The authors would like to appreciate Hamadan University of Medical Sciences for financially supporting this research. The researchers would like to express their gratitude to the authorities at Hamadan School of Nursing and Midwifery, the pre-hospital emergency care personnel who participated in the study, and all the other individuals who cooperated with them.

References

1. Organization WH: **Coronavirus disease 2019 (COVID-19): situation report, 72.** 2020.
2. Wu Z, McGoogan JM: **Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72 314 cases from the Chinese Center for Disease Control and Prevention.** *Jama* 2020, **323**(13):1239-1242.
3. Coronavirus N: **Situation Report–22 (World Health Organization, 2020).** In.; 2019.
4. Lai C-C, Shih T-P, Ko W-C, Tang H-J, Hsueh P-R: **Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and corona virus disease-2019 (COVID-19): the epidemic and the challenges.** *International journal of antimicrobial agents* 2020:105924.
5. Nemati M, Ebrahimi B, Nemati F: **Assessment of Iranian nurses' knowledge and anxiety toward COVID-19 during the current outbreak in Iran.** *Archives of Clinical Infectious Diseases* 2020, **15**(COVID-19).
6. **252 patients died and 3890 new patients were identified overnight in the country**
7. Liu S, Yang L, Zhang C, Xiang Y-T, Liu Z, Hu S, Zhang B: **Online mental health services in China during the COVID-19 outbreak.** *The Lancet Psychiatry* 2020, **7**(4):e17-e18.
8. Greenberg N, Docherty M, Gnanapragasam S, Wessely S: **Managing mental health challenges faced by healthcare workers during covid-19 pandemic.** *bmj* 2020, **368**.
9. Lai J, Ma S, Wang Y, Cai Z, Hu J, Wei N, Wu J, Du H, Chen T, Li R: **Factors associated with mental health outcomes among health care workers exposed to coronavirus disease 2019.** *JAMA network open* 2020, **3**(3):e203976-e203976.
10. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, Wang B, Xiang H, Cheng Z, Xiong Y: **Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus–infected pneumonia in Wuhan, China.** *Jama* 2020, **323**(11):1061-1069.
11. Stuijtzand S, Deforges C, Sandoz V, Sajin C-TC-T, Jaques C, Elmers J, Horsch A: **Psychological Impact of an Epidemic/Pandemic on the Mental Health of Healthcare Professionals: A Rapid Review.** *BMC Public Health* 2020, **20**(1230):1-18.
12. Lehmann M, Bruenahl CA, Löwe B, Addo MM, Schmiedel S, Lohse AW, Schramm C: **Ebola and psychological stress of health care professionals.** *Emerging Infectious Diseases* 2015, **21**(5):913.
13. Roy D, Tripathy S, Kar SK, Sharma N, Verma SK, Kaushal V: **Study of knowledge, attitude, anxiety & perceived mental healthcare need in Indian population during COVID-19 pandemic.** *Asian Journal of Psychiatry* 2020:102083.
14. Maslach C, Jackson SE, Leiter MP, Schaufeli WB, Schwab RL: **Maslach burnout inventory**, vol. 21: Consulting psychologists press Palo Alto, CA; 1986.
15. Baher A: **The relationship between emotional intelligence and sense of self-efficacy and burnout among staff of Azad University of Tabriz.** *Training and Evaluation (Education)* 2009, **2**(7):99-119.
16. Novak M, Guest C: **Application of a multidimensional caregiver burden inventory.** *The gerontologist* 1989, **29**(6):798-803.

17. Sakar H, Mahtab A-K, Farshad S, Fahimeh T, Mirzadeh FS, Hossien F: **Validation study: The Iranian version of caregiver abuse screen (CASE) among family caregivers of elderly with dementia.** *Journal of gerontological social work* 2019, **62**(6):649-662.
18. Abidin R: **Parenting stress index** . Lutz, FL: **Psychological Assessment Resources**. In.: Inc; 1995.
19. Reitman D, Currier RO, Stickle TR: **A critical evaluation of the Parenting Stress Index-Short Form (PSI-SF) in a head start population.** *Journal of Clinical Child and Adolescent Psychology* 2002, **31**(3):384-392.
20. Yeh CH, Chen ML, Li W, Chuang HL: **The Chinese version of the parenting stress index: a psychometric study.** *Acta Paediatrica* 2001, **90**(12):1470-1477.
21. Shirzadi P, Framarzi S, Ghasemi M, Shafiee M: **Investigating Validity and reliability of Parenting Stress Index–short form among Fathers of normal child under 7 years old.** *Rooyesh-e-Ravanshenasi Journal (RRJ)* 2015, **3**(4):91-110.
22. Salahian A, Oreizi H, Babamiri M, Asgari A: **The predictor factors of burnout syndrome in Isfahan nurses.** 2012.
23. Sandeep K, Sandhyarani B, Rao ES, Sharma H: **A Study to Assess Knowledge regarding Perceived Causes of Burnout Syndrome among Staff Nurses at selected Hospitals of Bareilly.** *International Journal of Advances in Nursing Management* 2019, **7**(4):351-354.
24. Velando-Soriano A, Ortega-Campos E, Gómez-Urquiza JL, Ramírez-Baena L, De La Fuente EI, Cañadas-De La Fuente GA: **Impact of social support in preventing burnout syndrome in nurses: A systematic review.** *Japan Journal of Nursing Science* 2020, **17**(1):e12269.
25. da Silva AA, Sanchez G, Barbosa NMS, de Oliveira MZ: **Predictor variables for burnout among nursing professionals.** *REVISTA DE PSICOLOGÍA (LIMA)* 2019.
26. da Silva Azevedo D, Martins Ferraz MM, Alencar Ferreira RdS, Caetano Lira JA, da Silva Azevedo D, Ribeiro Amorim SM, Portela Veloso LU: **RISK OF BURNOUT SYNDROME IN MENTAL HEALTH NURSES.** *Journal of Nursing UFPE/Revista de Enfermagem UFPE* 2019, **13**.
27. Sofology M, Efstratopoulou M, Dunn T: **Predicting burnout syndrome in Greek mental health professionals.** *Journal of Social Service Research* 2019, **45**(1):142-149.
28. Amini F: **The relationship between resiliency and burnout in nurses.** *Journal of research development in nursing and midwifery* 2013, **10**(2):94-102.
29. Edward K-I, Hercelinskyj G: **Burnout in the caring nurse: learning resilient behaviours.** *British journal of nursing* 2007, **16**(4):240-242.
30. Torgheh M, Alipour A: **Effect of humour on burnout and resiliency of nurses.** *Journal of Holistic Nursing And Midwifery* 2015, **25**(2):57-64.
31. Anthony LG, Anthony BJ, Glanville DN, Naiman DQ, Waanders C, Shaffer S: **The relationships between parenting stress, parenting behaviour and preschoolers' social competence and behaviour problems in the classroom.** *Infant and Child Development: An International Journal of Research and Practice* 2005, **14**(2):133-154.

32. van der Geest IM, van den Heuvel-Eibrink MM, Passchier J, van den Hoed-Heerschop C, Pieters R, Darlington ASE: **Parenting stress as a mediator of parents' negative mood state and behavior problems in children with newly diagnosed cancer.** *Psycho-Oncology* 2014, **23**(7):758-765.
33. Zou G, Shen X, Tian X, Liu C, Li G, Kong L, Li P: **Correlates of psychological distress, burnout, and resilience among Chinese female nurses.** *Industrial health* 2016:2015-0103.
34. Ahmadi O, Azizkhani R, Basravi M: **Correlation between workplace and occupational burnout syndrome in nurses.** *Advanced biomedical research* 2014, **3**(44):1-4.
35. Rafiee F, Oskouie F, Nikraves M: **Key factors in nurses' reaction to Burnout: A qualitative study.** *Razi Journal of Medical Sciences* 2007, **13**(53):83-94.