

Population-based Study of Long-term Anticoagulation for Treatment and Secondary Prophylaxis of Venous Thromboembolism in Men With Prostate Cancer in Sweden

Yanina Balabanova (✉ yanina.lenz@bayer.com)

Bayer AG

Bahman Farahmand

Bayer AB

Pär Stattin

Uppsala University

Hans Garmo

Uppsala University

Gunnar Brobert

Bayer AB

Research Article

Keywords: Anticoagulants, Bleeding, Prostate Cancer, Venous Thromboembolism

Posted Date: June 30th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-608457/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at BMC Urology on February 2nd, 2022. See the published version at <https://doi.org/10.1186/s12894-022-00967-z>.

Abstract

Background: Epidemiological data on anticoagulation for venous thromboembolism (VTE) in prostate cancer are sparse. We aimed to investigate associations between anticoagulation duration and risks of VTE recurrence after treatment cessation and major on-treatment bleeding in men with prostate cancer in Sweden.

Methods: Using nationwide prostate cancer registry and prescribing data, we followed 1413 men with VTE and an outpatient anticoagulant prescription following prostate cancer diagnosis. Men were followed to identify cases of recurrent VTE, and hospitalized major bleeding. We calculated adjusted hazard ratios (HRs) with 95% confidence intervals (CIs) to quantify the association between anticoagulation duration (reference ≤ 3 months) and recurrent VTE using Cox regression. We estimated 1-year cumulative incidences of major bleedings from anticoagulation initiation.

Results: The outpatient anticoagulation prescribed was parenteral (64%), direct oral anticoagulant (31%), and vitamin K antagonist (20%). Median duration of anticoagulation was 7 months. Adjusted HRs (95% CI) for off-treatment recurrent pulmonary embolism (PE) were 0.32 (0.09–1.15) for >3–6 months' duration, 0.21 (0.06–0.69) for >6–9 months and 0.16 (0.05–0.55) for >9 months; corresponding HRs for deep vein thrombosis (DVT) were 0.67 (0.27–1.66), 0.80 (0.31–2.07), and 1.19 (0.47–3.02). One-year cumulative incidences of intracranial, gastrointestinal and urogenital bleeding were 0.9%, 1.7%, 3.0% during treatment, and 1.2%, 0.9%, 1.6% after treatment cessation.

Conclusion: The greatest benefit in reducing recurrent VTE risk was with >9 months anticoagulation for PE and >3–6 months for DVT. Risks of major bleeding were low not being substantially different to the risks after treatment cessation.

Introduction

Venous thromboembolism (VTE) is the second most common cause of death in patients with cancer after disease progression,[1] and around 15–20% of all patients with VTE have cancer.[1–3] Prostate cancer is the most commonly diagnosed cancer in men aged ≥ 55 years worldwide.[4] Although prostate cancer is associated with a low risk of VTE relative to other cancer types, the risk of VTE has been estimated to be three-fold higher in men with prostate cancer than among men of the same age without cancer in the general population.[5] Owing to the high prevalence of prostate cancer, VTE is commonly seen among affected men in clinical practice,[1] hence the importance of prescribing effective agents for treatment and secondary prophylaxis.

For the majority of patients with cancer-associated thrombosis, clinical guidelines recommend long-term anticoagulant therapy – usually with low-molecular weight heparin (LMWH) or a direct oral anticoagulant (DOAC) – to help prevent VTE recurrence. Some guidelines advocate at least 3 months of anticoagulant therapy,[6] and others recommend 6 months[7] or a minimum of 6 months,[8, 9] and there is currently no consensus regarding therapy beyond 6 months. The pivotal clinical trials that shaped guideline development were limited to 3–6 months duration,[10] and observational data on the topic are limited. Determining the optimal duration of anticoagulant therapy requires balancing the reduced risk of VTE recurrence against the increased risk of bleeding, which is also raised in patients with cancer. Whether prolonged anticoagulation beyond 6 months affords a favourable benefit–risk ratio is a question that remains unanswered.

The aim of this study was to evaluate the type and duration of long-term anticoagulant therapy in men with prostate cancer and VTE, the association between duration of treatment and risk of VTE recurrence after treatment cessation, and risk of major bleedings on- and off-treatment. Of note is that we refrain from using the term 'cancer-associated VTE' because of the difficulty in determining the true likely cause of the thrombosis using our study design that used secondary collected data.

Methods

Data source

We used data from the Prostate Cancer data Base Sweden (PCBaSe) 4.0, a research database comprising information from the National Prostate Cancer Register in Sweden and other linked national health registers. Details of PCBaSe have been published previously.[11, 12] Briefly, the database contains information on 180,000 men diagnosed with prostate cancer between 1998 and 2016 and captures 98% of all newly diagnosed cases of prostate cancer in Sweden since 1998. For this study, we used data from the National Patient Registry,[13] which captures hospital inpatient/outpatient diagnoses using International Classification of Diseases (ICD–10) codes (with a look-back period to 1998), and the Prescribed Drug Registry (with a look-back period to 2005),[14] which records medications dispensed from all Swedish pharmacies using Anatomical Therapeutic Classification (ATC) codes. We also obtained data from the Swedish longitudinal integration database for health insurance and labour market Register for information on education level and marital status, and data from the Registries of Immigration and Emigration. The study protocol was approved by the Research Ethics Authority, Uppsala (Dnr 2019 – 01319).

Identification of men with prostate cancer and VTE

A flowchart depicting the identification of men with prostate cancer and VTE is shown in Fig. 1. The source population included all men in PCBaSe 4.0 with a first diagnosis of prostate cancer between 1 January 2007 (about 18 months after the Prescribed Drug Registry became available) and 31 December 2016, with at least 1 year of observational follow-up. From this source population, we included all men with a first inpatient/outpatient diagnosis of VTE (index VTE) recorded as a primary diagnosis in the National Patient Registry, and with a prescription for an anticoagulant in the Prescribed Drug Registry any time after the index VTE (N = 2404). We excluded men with either a past record of VTE or atrial fibrillation and those with a prescription for oral anticoagulant treatment any time before the index VTE, or for parental anticoagulant treatment in the 2 weeks before the index VTE (see **Supplementary Table 1** for codes). For our analyses, we retained only men whose index VTE was recorded between 2013 and 2017, to cover the period from which DOACs were approved in Sweden.

Anticoagulant treatment for VTE

We classified the type of anticoagulant(s) dispensed during the four weeks after the index VTE date as either parenteral anticoagulant, vitamin K antagonist (VKA) or DOAC; men could have been prescribed more than one type of anticoagulant during this period, either as a bridge between parenteral and oral agents or because they switched between drug class. The duration of an individual prescription was calculated based on all dispensed prescriptions any time after index VTE using information on pack size and the numbers of packs dispensed. Total anticoagulant exposure was determined by summing the lengths of individual consecutive prescriptions. A gap of more than 30% of supplied days was considered to represent the cessation of anticoagulant exposure. To check the validity of the computer algorithm used to determine treatment exposure, prescription start/end dates records for each patient were manually reviewed by one author (BF). We categorized exposure duration as ≤ 3 months, > 3 to 6 months, > 6 to 9 months and > 9 months irrespective of anticoagulant type.

Co-variates

We obtained information on sociodemographics (including educational and marital status), comorbidities and comedications before the index VTE. The Charlson comorbidity index was determined at the time of prostate cancer diagnosis.

Recurrent VTE

To identify cases of off-treatment recurrent VTE, we followed-up men from the date that their anticoagulation treatment ended until the earliest of the following: the first subsequent entry of a VTE diagnosis as the main diagnosis, death, the date of emigration, or the end of available follow-up (data lock point) in PCBaSe (31 December 2017). For this analysis, the study population was restricted to men whose treatment cessation occurred at least 1-year before the data-lock date to ensure adequate potential follow-up time to identify VTE events.

Major bleeding

We performed separate follow-ups to identify cases of major bleeding as the main cause of hospitalisation, one for each bleeding type (intracranial, gastrointestinal, urogenital or 'other bleeding' (see **Supplementary Table 2** for codes). We followed-up men from the start of anticoagulation therapy until the earliest of the following: the first subsequent entry of the major bleeding, death, the date of emigration, or the end of follow-up. We recorded whether the bleeding event occurred on-treatment or off-treatment (after ending of the anticoagulant treatment).

Statistical analysis

We summarised sociodemographics, clinical characteristics and duration of anticoagulant exposure using frequency counts and percentages for categorical variables, and with medians with interquartile range (IQR) for continuous variables. One-year cumulative incidences of recurrent VTE and major bleeding were calculated by dividing the number of incident cases during the first year of follow-up by the number of men at the start of follow-up (patients were not censored at death or low-to follow-up), with 95% confidence intervals computed using the binominal distribution. For major bleeding, cumulative incidence estimates were also stratified by type of bleeding. Incidence rates of recurrent VTE and major bleeding over the whole period of observation were calculated as the number of VTE/major bleeding events divided by the total person-years of follow-up, with 95% confidence intervals (CIs) determined using the Poisson distribution. We used Cox proportional hazards regression models to compute hazard ratios (HRs) with 95% CIs for associations between duration of anticoagulant exposure and off-treatment recurrent VTE adjusted for confounders and using ≤ 3 months duration as the reference group. We present age-adjusted HRs based on the finding that no additional variable made any notable change to the risk estimate when included into the model. SAS version 9.4 was used for all analyses.

Results

Baseline characteristics of patients with incident VTE

A total of 1413 men with prostate cancer experienced a VTE and were dispensed anticoagulant therapy from a pharmacy: 50% (700/1413) had a pulmonary embolism (PE), of which 68% were hospitalised, and 45% (640/1413) had a deep vein thrombosis (DVT), of which 12% were hospitalised (5%, 73/1413 had a non-specified VTE). Clinical and sociodemographic characteristics are shown in **Supplementary Tables 3–5**. Median age was 73 years (IQR 68–79) at VTE diagnosis. Median time from prostate cancer diagnosis to VTE diagnosis was 3 years (IQR 1–6). Approximately 20% of patients experienced their VTE within 1 year after prostate cancer diagnosis. Common comorbidities included hypertension (67%), cerebrovascular disease (10%), diabetes (14%) and hyperlipidaemia (12%), and just over 10% of patients had a history of urogenital bleeding. Commonly used medications at baseline (any time before index VTE) included antihypertensives (67%), statins (35%) and antiplatelets (30%). No major differences were seen in the clinical profiles of men with DVT vs. those with PE.

Anticoagulation

The vast majority (96%) of men started their outpatient anticoagulant therapy within the four weeks after their VTE diagnosis. Men received their first prescription at a median of 2 days after their VTE (range 1–6 days). Almost two-thirds

(64%) were prescribed parenteral anticoagulation, 31% a DOAC and 20% a VKA (Fig. 2). The substantial decline between 2013 and 2017 in the use of parental anticoagulation (83–53%) and VKAs (45–4%) corresponded with a substantial increase in DOAC use (5–51%). The prevalence of cardiovascular disease was slightly higher among men who received a VKA/DOAC than among those prescribed parenteral anticoagulation. A history of bleeding, in particularly urogenital bleeding, was more common among men who received parenteral anticoagulation. The median duration of use was 7 months (IQR 3–13). Fifty seven per cent of men were prescribed > 6 months' anticoagulation, and a quarter were prescribed > 1 year of anticoagulation (**Supplementary Table 6**). In general, anticoagulation was longer for PE (median 8 months, IQR 5–15) than for DVT (median 6 months, IQR 3–9). Over two-thirds (67%) of men treated for PE received outpatient anticoagulation for more than > 6 months, with over a third (35%) treated for > 1 year (**Supplementary Table 6**).

Recurrent VTE and major bleeding

Five hundred and ten men were included in the analysis of recurrent VTE. Incidence rates of recurrent VTE over the whole of follow-up are shown in **Supplementary Table 7a**. The incidence rate of recurrent VTE per 100 person-years was 19.2 (95% CI: 11.2–30.8) in men with \leq 3 months' therapy, 13.1 per 100 person-years (95% CI: 7.3–21.5) for > 3–6 months, 14.7 per 100 person-years (95% CI: 8.9–22.6) for > 6–9 months, and 15.0 per 100 person-years (95% CI: 8.9–23.7) for > 9 months. As shown in Table 1, compared with men with \leq 3 months' anticoagulation for the index PE, adjusted HRs for recurrent VTE were 0.32 (95% CI: 0.09–1.12) for > 3–6 months therapy, 0.26 (95% CI: 0.08–0.82) for > 6–9 months, and 0.18 (95% CI: 0.15–0.60) for > 9 months. Corresponding HRs relating to treatment of the index DVT were 0.67 (95% CI: 0.27–1.66) for > 3–6 months, 0.80 (95% CI: 0.31–2.07) for > 6–9 months, and 1.19 (95% CI: 0.47–3.02) for > 9 months.

Table 1

HRs (95% CIs) for recurrent VTE after the cessation of anticoagulation for the index VTE by duration of use.

Duration of anticoagulant therapy	DVT			PE			VTE*		
	Incident cases /person- yrs	Crude RR (95% CI)	Adjusted RR [†] (95% CI)	Incident cases /person- yrs	Crude RR (95% CI)	Adjusted RR [†] (95% CI)	Incident cases /person- yrs	Crude RR (95% CI)	Adjusted RR [†] (95% CI)
≤ 3 months N = 102	11 / 68	1.00	1.00	5 / 13	1.00	1.00	17 [†] / 88	1.00	1.00
> 3 to 6 months N = 127	9 / 75	0.74 (0.31–1.79)	0.71 (0.29–1.74)	6 / 35	0.47 (0.14–1.55)	0.32 (0.09–1.12)	15 / 115	0.68 (0.34–1.37)	0.66 (0.33–1.33)
> 6 to 9 months N = 150	9 / 60	0.93 (0.39–2.24)	0.99 (0.41–2.43)	11 / 68	0.45 (0.16–1.29)	0.26 (0.08–0.82)	20 / 137	0.77 (0.40–1.46)	0.76 (0.40–1.45)
> 9 months N = 131	9 / 39	1.41 (0.58–3.40)	1.39 (0.56–3.44)	8 / 75	0.30 (0.10–0.90)	0.18 (0.05–0.60)	18 [†] / 120	0.78 (0.40–1.52)	0.74 (0.38–1.44)
*Includes 73 men with VTE not specified as either DVT/PE, either ICD-10 I809 or I82)									
[†] Adjusted for age, time from prostate cancer to VTE diagnosis, and the following comorbidities before the index VTE: cardiovascular disease, hypertension, diabetes, major bleeding, cerebrovascular disease, and use of statins or NSAIDs before the index VTE.									
[†] One case was classified as neither DVT or PE, but as 'other VTE'.									
Notes: Analysis included 510 men with VTE and at least 1 year of follow-up after the end of therapy). Anticoagulants included LMWH, VKAs and DOACs.									
CI, confidence interval; DVT, deep vein thrombosis; DOAC, direct oral anticoagulant; HR, hazard ratio; LMWH, low-molecular weight heparin; NSAID, non-steroidal anti-inflammatory drug; PE, pulmonary embolism; VKA, vitamin K antagonist; VTE, venous thromboembolism									

One-year cumulative incidences of recurrent VTE were 17% for ≤ 3 months, 12% >3–6 months, 13% for > 6–9 months, and 14% for > 9 months (**Supplementary Table 8**). Cumulative incidence of recurrent VTE was highest in men with ≤ 3 months anticoagulation at all timepoints during the year after treatment cessation. Cumulative incidences at 60 days after anticoagulation cessation were 7% for men who received ≤ 3 months' duration, 4% for > 3–6 months or > 6–9 months' therapy, and 2% for > 9 months' therapy (Fig. 3); corresponding estimates at 180 days were 14%, 8%, 9%, and 7%, respectively, and at 360 days, they were 17%, 12%, 13%, and 14%, respectively.

Incidence rates of major bleeding rates were low; rates over the whole of follow-up are shown in **Supplementary Table 7b**. For one-year cumulative incidences, on-treatment estimates were slightly higher or similar to the respective off-treatment estimates. For urogenital bleeding these were 3.0% (on-treatment) compared with 1.6% (off-treatment), 1.7% (on-treatment) and 0.9% (off-treatment) for gastrointestinal bleeding, and 0.9% (on-treatment) and 1.2% (off-treatment) for intracranial bleeding (Fig. 4; **Supplementary Table 9**). The cumulative incidence of most types of major bleeding increased most rapidly over the first 90 days and increased more steadily thereafter (Fig. 4).

Discussion

In this large population-based study of men with prostate cancer in Sweden, the average duration of outpatient anticoagulant therapy for the treatment and secondary prophylaxis of VTE was 7 months; however, duration of treatment varied between patients. A fifth of men received anticoagulation for ≤ 3 months, while a quarter received ≥ 1 year duration of therapy. For cases of PE, anticoagulation was, on average, prescribed for a longer period. A reduced risk of recurrent VTE was seen among patients prescribed $> 3-6$ months therapy (vs ≤ 3 months therapy), with > 9 months treatment appearing to offer the greatest benefit for PE, and $> 3-6$ months offering affording the greatest benefit for DVT. Risks of major bleeding were low and did not appear to be substantially higher during the on-treatment period than after treatment cessation. This was especially apparent for intracranial bleeding, while for urogenital bleeding, the risk was somewhat increased during the on-treatment period.

Previous studies on this topic included patients across several cancer sites with different anticoagulation requirements and have evaluated VTE risks using different definitions. The average duration of anticoagulation among our study cohort (7 months) is similar to that seen among cancer patients in the international RIETE registry (6 months).[15] In US claims database studies, the average length of anticoagulation for VTE has been higher. Between 2007 and 2011, Kaatz *et al*^[16] found that cancer patients were anticoagulated for VTE (mostly warfarin) for an average of 10 months, while for the period 2007–2014, Khorana *et al*^[17] reported a median duration of 8 months for warfarin/rivaroxaban therapy among patients with cancer. We observed a significant increase in DOAC use for VTE among men in our study (reaching 51% in 2017), with a corresponding decline in use of VKAs (4% in 2017) and of parenteral anticoagulants (53% in 2017), albeit the latter was still used by over half of men long-term. This rise in DOAC use is consistent with temporal trends in other cancer populations, and reflect the current guideline recommendations.^[18] It is important to note that our data did not enable us to determine whether a recurrent VTE event was ‘cancer-associated’, nor was this the aim of the study – our focus was on all men with prostate cancer who were diagnosed with a VTE at any time point following cancer diagnosis, irrespective of treatment. In men with prostate cancer, it is challenging to determine whether a VTE event is cancer-associated because this type of cancer can have slow progression, and patients may require different treatments depending on disease stage; thrombotic activity could therefore be triggered at different times in different patients. Furthermore, it is noteworthy that only about 20% of men in our study were diagnosed with a VTE within a year of cancer diagnosis.

Owing to advances in early screening and treatment, life expectancy of men with prostate cancer can be relatively long when compared with other cancer types, and timely diagnosis and treatment of VTE in these men is important because it can impact prognosis and survival. Yet, anticoagulant treatment decisions are challenging because these patients are characteristically of older age and with comorbidities that may increase thromboembolism and bleeding risks. Our findings reflect the adoption of an individualised approach to treating VTE in this heterogenous population. Notwithstanding the necessity for personalized treatment approaches, a higher risk of recurrent VTE was seen in men treated with ≤ 3 months’ anticoagulation vs longer durations, consistent with findings among cancer patients in the claims database study by Khorana *et al* 2019.^[19] The small possible increase in recurrent DVT risk observed in men treated for > 9 months is likely to be an artefact from an inherent increased in this subset of patients.

A major strength of our study is the focus on a single cancer type. Selection bias was minimized because the cohort included unselected men with prostate cancer of different stages/grades, across all ages and including those with comorbidities. Sweden has a tax-funded public healthcare system. Data completeness in the Swedish Prescribed Drug Register is virtually complete because Swedish pharmacies are required to participate by law. We therefore captured treatment patterns in virtually all men with prostate cancer and VTE in the country, none of whom were lost to follow-up. Although drugs issued in hospital are not captured in PCBaSe, we were interested in long-term anticoagulation, where

the minimum recommended duration is 3 months. Furthermore, the proportion of men hospitalised for their VTE was low (68% for PE and 12% for DVT). We included both inpatient and outpatient VTE events, based on previous knowledge that a high proportion of VTE events in cancer patients occur in the ambulatory setting.[3] The main study limitation is the likelihood of residual confounding such as VTE severity. Other factors that influence VTE risk, such as cancer stage/grade, cancer treatment (including surgery), age and comorbidity profile, all of which can affect VTE risk, [20, 21] may also have influenced treatment decisions, with the duration of treatment dependent the individual's risk profile. We were unable to validate the VTE diagnoses as the results of imaging procedures are not routinely recorded in the Patient Register. Additionally, anticoagulant duration may have been misclassified because information on adherence was unavailable, and this may have impacted on our VTE recurrence/bleeding analyses. Lastly, some of our analyses may have been underpowered to detect significant differences between anticoagulant duration categories.

In conclusion, our study provides useful information on the benefits and risks of anticoagulation therapy beyond 6 months' duration for VTE in men with prostate cancer, on which there is currently no consensus. Our results suggest the greatest potential for the reducing VTE recurrence occurs with > 9 months treatment for PE and > 3–6 months for DVT. Risks of major bleeding were low overall and did not differ substantially for events that occurred during anticoagulation and those that occurred after anticoagulation cessation (off-treatment events), particularly for intracranial bleeding. Further epidemiological studies are needed to confirm/refute our findings and help guide clinical decision making in routine practice. Other beneficial areas for further research in this field include the effects of specific anticoagulants on relevant clinical outcomes and gaining a better understanding of the prescribing choices.

Abbreviations

ATC, Anatomical Therapeutic Classification

CI, confidence interval

DOAC , direct oral anticoagulant

HR, hazard ratio

IQR, inter-quartile range

LMWH, low-molecular weight heparin

PCBaSe, Prostate Cancer data Base Sweden

VKA, vitamin K antagonist

VTE, venous thromboembolism

Declarations

Ethical approval and consent to participate

The study was carried out in accordance with the Helsinki Declaration. There was no requirement for informed consent by the Research Ethics Authority, Uppsala – the ethics committee that approved the study protocol (Dnr 2019-01319). The Research Ethics Authority Uppsala approved the use of the opt-out informed consent in NPCR for which there is no collection of signed consent.

Consent for publication

Not applicable.

Availability of data and materials

The datasets analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

YL is an employee of Bayer AG. BF and GB are employees of Bayer AB. PS declares that Region Uppsala has, on behalf of NPCR, made agreements on quarterly reports from Patient-overview Prostate Cancer (part of NPCR) with Astellas, Janssen, and Bayer, as well as research projects with Astellas, Bayer, and Janssen. HG declares no conflicts of interest.

Funding

This study was funded by Bayer AG. The funder had no role in any aspect of the study apart from those performed by YL, GB and BF who were all employees of Bayer at the time the study was carried out.

Authors' contributions

YB designed the study, interpreted the patient data, and reviewed manuscript drafts. BF designed the study, analyzed and interpreted the patient data, and reviewed manuscript drafts. GL designed the study, interpreted the patient data, and reviewed manuscript drafts. PS obtained the data, interpreted the patient data and reviewed manuscript drafts. HG obtained the data, interpreted the patient data and reviewed manuscript drafts. All authors read and approved the final manuscript.

Acknowledgements

We thank Susan Bromley from EpiMed Communications Ltd (Abingdon, UK) for medical writing assistance funded by Bayer AG, the study funder, and in accordance with Good Publication Practice. This project was made possible by the continuous work of the National Prostate Cancer Register of Sweden (NPCR) steering group: Pär Stattin (chair), Ingela Franck Lissbrant (deputy chair), Johan Styrke, Camilla Thellenberg Karlsson, Lennart Åström, Hampus Nugin, Stefan Carlsson, Marie Hjälms-Eriksson, David Robinson, Mats Andén, Ola Bratt, Magnus Törnblom, Johan Stranne, Jonas Hugosson, Maria Nyberg, Olof Akre, Per Fransson, Eva Johansson, Gert Malmberg, Hans Joelsson, Fredrik Sandin, and Karin Hellström.

References

1. Iorga RA, Bratu OG, Marcu RD, Constantin T, Mischianu DLD, Socea B, Gaman MA, Diaconu CC: Venous thromboembolism in cancer patients: Still looking for answers (Review). *Exp Ther Med* 2019, 18(6):5026–5032.
2. Lee AY: Management of thrombosis in cancer: primary prevention and secondary prophylaxis. *Br J Haematol* 2005, 128(3):291–302.
3. Streiff MB: Association between cancer types, cancer treatments, and venous thromboembolism in medical oncology patients. *Clin Adv Hematol Oncol* 2013, 11(6):349–357.
4. Ferlay J, Ervik M, Lam F, et al. Global Cancer Observatory: Cancer Today . Available at: <https://gco.iarc.fr/today>. Accessed 27 April 2021
5. Cronin-Fenton DP, Søndergaard F, Pedersen LA, Fryzek JP, Cetin K, Acquavella J, Baron JA, Sørensen HT: Hospitalisation for venous thromboembolism in cancer patients and the general population: a population-based cohort study in Denmark, 1997-2006. *Br J Cancer* 2010, 103(7):947–953.

6. National Comprehensive Cancer Network: Cancer-Associated Venous Thromboembolic Disease. 2020.
7. Mandalà M, Falanga A, Roila F: Management of venous thromboembolism (VTE) in cancer patients: ESMO Clinical Practice Guidelines. *Ann Oncol* 2011, 22 Suppl 6:vi85–92.
8. Farge D, Frere C, Connors JM, Ay C, Khorana AA, Munoz A, Brenner B, Kakkar A, Rafii H, Solymoss S *et al*: 2019 international clinical practice guidelines for the treatment and prophylaxis of venous thromboembolism in patients with cancer. *The Lancet Oncology* 2019, 20(10):e566-e581.
9. Key NS, Khorana AA, Kuderer NM, Bohlke K, Lee AYY, Arcelus JI, Wong SL, Balaban EP, Flowers CR, Francis CW *et al*: Venous Thromboembolism Prophylaxis and Treatment in Patients With Cancer: ASCO Clinical Practice Guideline Update. *J Clin Oncol* 2020, 38(5):496–520.
10. Wang TF, Li A, Garcia D: Managing thrombosis in cancer patients. *Res Pract Thromb Haemost* 2018, 2(3):429–438.
11. Hagel E, Garmo H, Bill-Axelsson A, Bratt O, Johansson JE, Adolfsson J, Lambe M, Stattin P: PCBaSe Sweden: a register-based resource for prostate cancer research. *Scand J Urol Nephrol* 2009, 43(5):342–349.
12. Van Hemelrijck M, Garmo H, Wigertz A, Nilsson P, Stattin P: Cohort Profile Update: The National Prostate Cancer Register of Sweden and Prostate Cancer data Base—a refined prostate cancer trajectory. *Int J Epidemiol* 2016, 45(1):73–82.
13. Socialstyrelsen. The National Patient Register. <https://www.socialstyrelsen.se/en/statistics-and-data/registers/register-information/the-national-patient-register/> Accessed 27 April 2021
14. Socialstyrelsen. The Swedish Prescribed Drug Register. <https://www.socialstyrelsen.se/en/statistics-and-data/registers/register-information/the-swedish-prescribed-drug-register/> Accessed 27 April 2021
15. Farge D, Trujillo-Santos J, Debourdeau P, Bura-Riviere A, Rodriguez-Beltrán EM, Nieto JA, Peris ML, Zeltser D, Mazzolai L, Hij A *et al*: Fatal Events in Cancer Patients Receiving Anticoagulant Therapy for Venous Thromboembolism. *Medicine* 2015, 94(32):e1235–e1235.
16. Kaatz S, Fu A-C, AbuDagga A, LaMori J, Bookhart BK, Damaraju CV, Tan H, Schein J, Nutescu E: Association between anticoagulant treatment duration and risk of venous thromboembolism recurrence and bleeding in clinical practice. *Thrombosis Research* 2014, 134(4):807–813.
17. Khorana AA, McCrae KR, Milentijevic D, Fortier J, Nelson WW, Laliberte F, Crivera C, Lefebvre P, Yannicelli D, Schein J: Current practice patterns and patient persistence with anticoagulant treatments for cancer-associated thrombosis. *Res Pract Thromb Haemost* 2017, 1(1):14–22.
18. Angelini DE, Elson P, McCrae K, Khorana AA: Anticoagulant Prescription Patterns for Cancer Associated Thrombosis: New Insights into the Use of Direct Oral Anticoagulants. *Blood* 2017, 130(Supplement 1):4725-4725.
19. Khorana AA, McCrae KR, Milentijevic D, Fortier J, Nelson WW, Laliberte F, Crivera C, Lefebvre P, Schein J: Duration of anticoagulant therapy and VTE recurrence in patients with cancer. *Support Care Cancer* 2019, 27(10):3833–3840.
20. Khorana AA, Cohen AT, Carrier M, Meyer G, Pabinger I, Kavan P, Wells PS: Prevention of venous thromboembolism in ambulatory patients with cancer. *ESMO Open* 2020, 5(6):e000948.
21. Van Hemelrijck M, Garmo H, Holmberg L, Bill-Axelsson A, Carlsson S, Akre O, Stattin P, Adolfsson J: Thromboembolic events following surgery for prostate cancer. *Eur Urol* 2013, 63(2):354–363.

Supplementary Tables

Supplementary Tables are not available with this version.

Figures




Figure 1

Identification of men with prostate cancer and VTE treated with anticoagulant therapy. *ICD-10 codes I82, I801, I802, I803, I809 as main diagnosis. AF, arial fibrillation; DOAC, direct oral anticoagulant; VKA, vitamin K antagonist; VTE, venous thromboembolism




Figure 2

Type of anticoagulant(s) prescribed during the four weeks after the index VTE among the 1413 men with prostate cancer and VTE (2013–2017) stratified by type of VTE (DVT/PE). Note: Men were eligible to have received more than one type of anticoagulant treatment during this four-week time period. In total, an outpatient prescription for an anticoagulant during the four weeks after the index VTE was issued to 95.5% of men, with 4.5% of men issued an anticoagulant prescription after four weeks. DOAC, direct oral anticoagulant; DVT, deep vein thrombosis; PE, pulmonary embolism; VKA, vitamin K antagonist; VTE, venous thromboembolism




Figure 3

Cumulative incidence (%) of recurrent VTE after the cessation of anticoagulant treatment by duration of therapy and time since cessation of therapy. Note: Analysis included 510 men with VTE and at least 1 year of follow-up after the cessation of therapy. VTE, venous thromboembolism




Figure 4

Cumulative incidence (%) of on-treatment and off-treatment major bleeding of anticoagulant therapy. Note: On-treatment analysis included all 1413 men with VTE, and off-treatment analysis include 936 patients. VTE, venous thromboembolism