Evaluation of Historical CMIP6 Model Simulations and Future Projections of Temperature Over the Pan-Third Pole Region

Xuewei Fan

Beijing Normal University

Qingyun Duan (✉ qyduan@hhu.edu.cn)

Hohai University

Chenwei Shen

Beijing Normal University

Yi Wu

Beijing Normal University

Chang Xing

Beijing Normal University

Research Article

Keywords: CMIP6, Temperature, the Pan-Third Pole, Climate change

Posted Date: July 13th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-619314/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Environmental Science and Pollution Research on December 1st, 2021. See the published version at <https://doi.org/10.1007/s11356-021-17474-7>.

Evaluation of historical CMIP6 model simulations and future projections of temperature over the Pan-Third Pole region

Xuewei Fan¹, Qingyun Duan^{*1,2}, Chenwei Shen¹, Yi Wu¹, Chang Xing¹

¹*State Key Laboratory of Earth Surface Processes and Resource Ecology, Faculty of Geographical Science, Beijing Normal University, Beijing 100875, China*

²*College of Hydrology and Water Resources, Hohai University, Nanjing 210098, China*

Abstract: The Pan-Third Pole (PTP) region, which encompasses the Eurasian highlands and their surroundings, has experienced unprecedented, accelerated warming during the past decades. This study evaluates the performance of historical simulation runs of the Coupled Model Intercomparison Project (CMIP6) in capturing spatial patterns and temporal variations observed over the PTP region for mean and extreme temperatures. In addition, projected changes in temperatures under four Shared Socioeconomic Pathway (SSP) scenarios (SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5) are also reported. Four indices were used to characterize changes in temperature extremes: the annual maximum value of daily maximum temperature (TX_x), the annual minimum value of daily minimum temperature (TN_n), and indices for the percentage of warm days (TX90p) and warm nights (TN90p). Results indicate that most CMIP6 models generally capture the characteristics of the observed mean and extreme

*Corresponding author: Qingyun Duan
Email: qyduan@hhu.edu.cn

temperatures over the PTP region, but there still are slight cold biases in the Tibetan Plateau. Future changes of mean and extreme temperatures demonstrate that a strong increase will occur for the entire PTP region during the twenty-first century under all four SSP scenarios. Between 2015 and 2099, ensemble area-averaged annual mean temperatures are projected to increase by 1.24 °C/100 yr, 3.28 °C/100 yr, 5.57 °C/100 yr, and 7.40 °C/100 yr for the SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5 scenarios, respectively. For TXx and TNn, the most intense warming is projected in Central Asia. The greatest number of projected TX90p and TN90p will occur in the Tibetan Plateau and Southeast Asia subregions, respectively.

Keywords

CMIP6; Temperature; the Pan-Third Pole; Climate change

Article Highlights:

- CMIP6 models perform well in reproducing the spatial and temporal variability of the mean and extreme temperatures over the PTP region.
- The performance of the multi-model mean is superior to that of most individual models.
- CMIP6 models reported a strong increase in mean and extreme temperatures over the PTP region during the 21st century.

Introduction

Warming of the climate system is unequivocal, a fact that has drawn overwhelming attention from the public, governments, and academic communities in recent decades. The warming has exerted profound, worldwide impacts on human life (Cheng et al., 2018; Sun et al., 2019), agricultural production (Liu et al., 2019; Liu et al., 2021; Tigchelaar et al., 2018), land use (Seneviratne et al., 2018), and natural ecosystems (Gou et al., 2020; Gou et al., 2021; Kong et al., 2020; Miao et al., 2019; Zheng et al., 2021). Among the areas expected to be sensitive to this warming is the region known as the Third Pole, which includes the Tibetan Plateau and the mountains surrounding it. Extending westward and northward from the Third Pole, the Pan-Third Pole (PTP) region is the core region of the “The Belt and Road” initiative promoted by the Chinese government, which has been building a new platform for international cooperation among more than 70 countries in Asia, Africa, and Europe. It covers more than 20 million km² and supports over 3 billion people with its resources. The PTP region is among the regions in the world most vulnerable to the impact of climate change, since it has the world's highest elevations and hosts the largest mass of glaciers and snow cover outside the polar regions (Wang et al., 2020). Under global warming, the PTP region, and especially the Tibetan Plateau, has experienced rates of warming twice the global average over the last 50–60 years (Deng et al., 2017; Pepin et al., 2019; You et al., 2019). Furthermore, the projected warming of some areas of the PTP region will exceed 4 °C above pre-industrial levels by 2100, which far exceeds the 2 °C goal set by the Paris Agreement of the United Nations Framework Convention on Climate

Change (Yao et al., 2017). The warming in the PTP region is causing Earth system changes characterized by intensive interactions among the processes of the atmosphere, hydrosphere, cryosphere, and biosphere, and is resulting in environmental threats such as glacier retreat, ice collapse, glacial lake expansion, and frequent glacier lake outburst flood (Yang et al., 2014; Yao et al., 2019). These changes may have impacts on the regional and global hydrologic cycle, thereby hindering socioeconomic development in countries along the routes of the Belt and Road Initiative. A deeper understanding of climate changes in the PTP region can inform science-based adaptation strategies to reduce climate risks.

Global climate models (GCMs) have become a major and vital tool for projecting future changes in climate; their reliability depends on their ability to reproduce historical and current climate features. To this end, the World Climate Research Programme has promoted a set of experiments known as the Coupled Model Intercomparison Project (CMIP) since the 1990s, which have delivered systematic and high-quality simulations for better understanding past climate changes and making projections and uncertainty estimates of the future (Annan and Hargreaves, 2011; Meehl et al., 2000). Studies using simulations from the fifth phase of CMIP (CMIP5; Taylor et al., 2012) have advanced our understanding of regionally heterogeneous climate warming (Bannister et al., 2017; Ongoma et al., 2018; Sun et al., 2017; Sun et al., 2020), highlighting strong warming trends in the high latitudes of the Northern Hemisphere and moderate warming trends in the middle latitudes (Feng et al., 2014). Several studies have investigated the performance of CMIP models for the PTP region.

For example, Dong et al. (2018) assessed the performance of CMIP5 historical simulations and projected future changes under three Representative Concentration Pathways (RCP2.6, RCP4.5, and RCP8.5) over critical Belt and Road regions based on data from 22 models. They found the most significant areas of warming are expected in Kazakhstan and the northern part of the south Belt and Road region. Kamworapan and Surussavadee (2019) evaluated the performances of forty CMIP5 models for simulating climatological temperature and precipitation for Southeast Asia and suggest the use of an ensemble they called 6-GCM-Ensemble for climate studies and projections. Jia et al. (2019) comprehensively assessed the performance of 33 CMIP5 models by an improved-score-based and multiple-criteria method and demonstrated that most models could capture the seasonal temperature patterns.

The sixth phase of CMIP (CMIP6; Eyring et al., 2016) is now in progress, and it features updates to parameterizations, the addition of new physical processes, and somewhat higher resolution compared to CMIP5 (Eyring et al., 2019). CMIP6 contains future scenario experiments named the Scenario Model Intercomparison Project (ScenarioMIP; O'Neill et al., 2016; O'Neill et al., 2020), which produce projections for a new set of emissions and land use scenarios based on the Shared Socioeconomic Pathways (SSPs; Riahi et al., 2017). The SSPs describe plausible future changes in societal aspects such as demographic, technological, economic, social, governance, and environmental factors (O'Neill et al., 2017). The CMIP6 models provide a new opportunity to examine the climate system and make regional projections of the climate under new scenarios, including for the PTP region.

Before assessing projections into the future, it is essential to evaluate the credibility of CMIP6 simulations with respect to observations on regional scales. Recent studies have already investigated the performance of CMIP6 in simulating regional historical climate and achieved positive results (Fan et al., 2020b; Grose et al., 2020; Li et al., 2021; Lovino et al., 2021; Srivastava et al., 2020). However, a comprehensive assessment relative to historical observations over the PTP region has not yet been performed, especially for looking at the climate extremes. Similarly, projected climate changes under the new SSP scenarios have not yet been reported. Motivated by the above issues, this study aims to (1) evaluate the performance of the CMIP6 models in simulating near-surface mean and extreme temperatures over the PTP region during the historical period and (2) assess future climate change in the PTP region based on CMIP6 model simulations for the 21st century.

The paper is organized as follows: Section 2 briefly describes the study area and introduces the data and methodology used in this study. The performance of CMIP6 models in reproducing the observed temperature over the PTP region is assessed in Section 3. Section 4 investigate the projected changes of temperature over the different subregions of PTP in the 21st century. Finally, Section 5 presents the main conclusions.

Data and methods

Study area

The PTP region is located in one of the most fragile and rapidly developing regions of the Earth, around the Third Pole, which mainly includes the Tibetan Plateau and the

northern intracontinental arid region of Asia, extending to the Caucasus Mountains in the west and the western Loess Plateau in the east (Yao et al., 2017). Figure 1 shows the areal extent considered in this study (35°E–125°E and 0°N–60°N) and its three subregions (Tibetan plateau (TP), Southeast Asia (SEA), and Central Asia (CA)). Home to three billion people, the PTP region hosts a substantial part of the “Silk Road Economic Belt” initiative proposed and advanced by China. The climate in the PTP region ranges from arid continental to humid tropical, primarily influenced by alternating predominance of the westerlies in winter and the Asian monsoon in summer. In turn, the PTP region exerts profound thermal and dynamic effects on atmospheric circulation, thus affecting the climate of Asia, the Northern Hemisphere, and beyond (Kang et al., 2019).

CMIP6 datasets and observations

We obtained monthly simulations of near-surface temperatures and daily minimum and maximum near-surface temperatures from the CMIP6 database (<https://esgf-node.llnl.gov/search/cmip6/>) using the historical simulations (1850–2014) and future scenario simulations (2015–2099) from ScenarioMIP. ScenarioMIP recommends four Tier 1 simulation protocols, reflecting different SSPs that result in different radiative forcing magnitudes by 2100: SSP1 - 2.6 (a low-forcing “sustainability” pathway; +2.6 W/m²), SSP2 - 4.5 (a medium-forcing “middle - of - the - road” pathway; +4.5 W/m²), SSP3 - 7.0 (a medium - to high - forcing pathway; +7.0 W/m²), and SSP5 - 8.5 (a high - forcing pathway; +8.5 W/m²). Table 1 provides the basic

information of 16 GCMs that were involved to analyze the changes in annual mean temperature. Only the last 12 GCMs in Table 1 were used to analyze the changes in temperature extremes when considering the availability of daily data in historical experiment and four SSP scenarios.

The reference mean temperature dataset used in this study is the gauge-based gridded Climatic Research Unit (CRU) TS v. 4.03 (Harris et al., 2020), which covers the period 1901–2018 with a $0.5^{\circ} \times 0.5^{\circ}$ spatial resolution (available at https://crudata.uea.ac.uk/cru/data/hrg/cru_ts_4.03/). The dataset is derived by interpolating monthly climate anomalies from extensive networks of weather station observations, and it has been subject to extensive quality control measures. Numerous studies have utilized CRU datasets for temperature-related analysis and reported its capability in simulating temperature in various regions of globe (Ahmed et al., 2020; Fan et al., 2020a; Li et al., 2018; Osborn et al., 2021). All monthly simulations of CMIP6 models were bilinearly interpolated to a common $0.5^{\circ} \times 0.5^{\circ}$ grid to keep the resolution consistent with CRU data.

The HadEX3 dataset (Dunn et al., 2020) is used to investigate observed changes in temperature extremes. This dataset is the recent global land-based climate extremes data developed by the World Meteorological Organization's Expert Team on Climate Change Detection and Indices (ETCCDI), available on a $1.875^{\circ} \times 1.25^{\circ}$ longitude-latitude grid for the period from 1901 to 2018 (<https://www.metoffice.gov.uk/hadobs/hadex3>).

Methods

Calculation of extreme temperature indices

Four indices of temperature extremes (Table 2) from the ETCCDI were selected to assess historical and future changes related to daily maximum temperatures (TX) and daily minimum temperatures (TN) from 1951 to 2099. There are three hot indices (annual maximum value of TX, TXx; the percentage of warm days, TX90p; and the percentage of warm nights, TN90p) and one cold index (annual minimum value of TN, TNn), which together can characterize the intensity and frequency of temperatures extremes. These indices have been widely used to investigate the observed and projected changes in extreme temperatures (Li et al., 2019; Wehner, 2020; Zhou et al., 2014). We calculated all indices at the models' native resolution using the R package *climindex.pcic.ncdf* and regridded the outputs to a common $1^{\circ} \times 1^{\circ}$ grid by bilinear interpolation to balance the resolution of HadEX3. The HadEX3 indices are also regridded to a $1^{\circ} \times 1^{\circ}$ grid. When calculating the TX90p and TN90p indices, the same base period (1961–1990) from HadEX3 was applied to the CMIP6 models.

Model performance metric

Taylor diagrams were employed to quantify the degree of correspondence between model simulations and observations, taking into consideration mean temperature climatology patterns, which are shown by spatial correlation coefficients, root mean square error (RMSE), and the ratio of standard deviations (Taylor, 2001). When the

correlation coefficient and the standard deviation are close to 1 and the RMSE is close to 0, this is indicative of the best match between observation and model simulation. Additionally, we compared the average bias of the spatial pattern of the mean state of temperature between the observational data and the CMIP6 ensembles.

To evaluate the performance of the CMIP6 models in simulating temperature extremes for the PTP region, we construct a “portrait” diagram based on the relative model RMSE (RMSE') of the climatology pattern for the period 1951–2014 (Sillmann et al., 2013). This diagram provides a concise graphical overview of the model performance for various temperature extreme indices. First, we calculated RMSEs for all individual models relative to the HadEX3 data. Then, the RMSEs for all models were collected to derive the relative model error for each model, RMSE', defined as:

$$RMSE' = \frac{RMSE - RMSE_{median}}{RMSE_{median}} \quad (1)$$

where $RMSE_{median}$ is the median of RMSEs for individual models. Negative values of RMSE' indicate that the corresponding model performs better than the majority (50%) of models.

Evaluations of historical temperature simulations

Evaluations of historical mean temperatures

The spatial distributions of annual mean temperatures from the 16 CMIP6 models and CRU observations over the PTP region are shown in Figure 2. Generally, the observed annual mean temperature exhibits a decreasing gradient from the north to south with the lowest temperatures occurring in the high-elevation TP. All the CMIP6

models reproduce the spatial distribution of annual temperature reasonably well and are able to capture the lowest temperatures in the TP. However, there are consistent cold biases in the TP and SEA for most models. Additionally, some models tend to overestimate the annual temperatures in South Asia, especially MIROC6 and MIROC-ES2L. The multi-model mean results are notably similar to observations in most regions, and the biases are relatively smaller than those of most individual models.

Next, we use bias calculations and Taylor diagrams to quantitatively assess the performance of models in simulating the spatial pattern of annual mean temperature. The length of each bar in Figure 3a shows the spread in regional average temperature biases simulated by the CMIP6 models relative to the CRU observations. We found that the overall biases of the PTP region vary between -2.35°C and 2.45°C . Among the three subregions, the TP region exhibits the largest negative biases, while the CA region shows large positive biases. The bias of the CMIP6 multi-model mean is below 2°C , which indicates smaller differences compared with the corresponding observed data than most models produce in simulating spatial patterns. Figure 3b depicts the Taylor diagram of the model simulations against observations, which summarizes the degree of correspondence between the observed and simulated fields. All 16 CMIP6 models are in line with observations, with correlation coefficients (dotted radial lines) all above 0.9. Also, most of the models exhibit a ratio of the standard deviations that is close to 1 and the centered pattern RMSE difference range was 0.17–0.30. This indicates that the CMIP6 models perform reasonably well in simulating the spatial distribution of annual mean temperatures over the PTP region. With respect to the statistical parameters in the

Taylor diagram, CESM2-WACCM, CESM2, EC-Earth3, EC-Earth3-Veg, and GFDL-ESM4 exhibit superior performance compared with other models. However, the CMIP6 ensemble average outperforms many of the individual models, as indicated by its closer distance to the reference point on the Taylor diagram.

Figure 4 shows 10-year moving average values for annual mean temperature for the ensemble of the 16 models and for the observations. The analysis shows that the observed annual mean temperature lies within the 5th–95th percentile range of CMIP6 multi-model ensembles, implying that there is consistency between the observed record and the CMIP6 models. Additionally, the CMIP6 historical simulations can reproduce the observed annual temperature warming trends in the PTP region and its three subregions, although with different magnitudes for bias.

Evaluations of historical temperature extremes

The spatial distributions of the four annual mean temperature extreme indices (TXx, TNn, TX90p, and TN90p) from HadEX3 and the CMIP6 multi-model mean during 1970–1999 over the PTP region are shown in Figure 5. The warm extreme indices calculated from both the observations and the model simulations show the decreasing trend from north to south, while the cold extreme index shows the opposite change, indicating that the CMIP6 multi-model ensemble simulates the spatial distribution of temperature extremes effectively. We also investigate the spatial variability of climatological annual mean temperature extreme indices for 12 individual models, and the results are shown in Figures S1 to S4. For TXx, the multi-model mean and each

model display consistent positive biases in most regions but show a negative bias in the TP. MIROC6 shows a larger warm bias than other models, but a smaller bias in the TP region. Among the four extreme indices, the CMIP6 models perform best in simulating TNn, with considerable consistency across models. However, most models underestimate the observed TNn of the TP. The TX90p for the CMIP6 models and the multi-model mean display substantially positive biases of about 2% over most parts of the PTP region, especially in the northern parts of the PTP region and South Asia. Several models (especially EC-Earth3-Veg, EC-Earth3, CanESM5, GFDL-ESM4, and IPSL-CM6A-LR, and MRI-ESM2-0) simulate high TN90p over parts of the PTP region. These biases are also found in the CMIP6 multi-model mean, but with smaller magnitude. In general, the four extreme indices simulated by the CMIP6 models are basically biased in the TP region, which may be due to the lack of observation stations and complex terrain in this region, with the coarse resolutions of the climate models leading to difficulties in the model simulation (You et al., 2018; Zhu and Yang, 2020).

A portrait diagram is a powerful tool for evaluating individual models against a reference dataset. Figure 6 presents a portrait diagram for the chosen temperature extreme indices, displaying the relative magnitude of spatially averaged RMSE and the average RMSE for all indices (RMSE_all) in the top row. The cold colors indicate when a model's performance is better than others', on average, and the warm colors indicate when its performance is worse. In general, most models shown exhibit reasonable skill in representing the temperature-based indices at the annual scale, while the performance of several models deviates greatly from that of the median model with respect to

HadEX3 (e.g., MIROC6 for TXx, CanESM5 and IPSL-CM6A-LR for TNn, Earth3-Veg for TN90p and TN90p). Models that perform relatively well for these four indices include MIROC-ES2L, MRI-ESM2-0, UKESM1-0-LL, CNRM-CM6-1, and CNRM-ESM2-1. Based on the RMSE_all (top row), MIROC-ES2L shows the best performance, with negative relative RMSEs for all four indices, followed by MRI-ESM2-0, UKESM1-0-LL, and CNRM-ESM2-1, which exhibit relatively small RMSE' of around 0. The performance of the multi-model mean is superior to that of most individual models due to substantial reduction of the systematic errors in individual models (Kim et al., 2020; Sillmann et al., 2013).

The temporal evolution of the regional averaged indices over the PTP region in the models and HadEX3 is shown in Figure 7 for the time period 1951–2014, and the time series of the three subregions are shown in Figures S5–S7. Note that we use HadEX3 to mask the CMIP6 models to avoid effects related to changes in the spatial coverage of the HadEX3 dataset over time. We found that the HadEX3 temperature indices generally lie within the CMIP6 model spread, with similar variability in many cases. Some exceptions are TNn for whole PTP region and its subregions (Figures 7, S5–S7), and TXx before 1960s for the TP region (Figure S6). For all four indices, there are similar warming trends in the models and HadEX3 for the entire PTP region and the three subregions (except TXx in TP, which may be due to the lack of observed stations in TP). For the extreme indices based on the TX (TXx and TX90p), the trend simulation of the CMIP6 multi-model average generally overestimates the warming trend of the observations, while for the extremes indices based on the TN (TNn and TN90p), the

CMIP6 multi-model average shows a lower increase compared to the HadEX3 dataset.

Projected temperatures in the 21st century

Projected changes in mean temperatures

In this section, we present the projected changes of temperature over the PTP region in the twenty-first century under the scenarios of SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5. We found that the performance of the model is inconsistent for different temperature variables. However, the multi-model means generally exhibit better performance than most individual models, so we use the multi-model mean for projections of the future. Figure 8 shows the spatial distributions of climatological changes in mean temperature in terms of multi-model ensemble means for the near-term (2025–2049), mid-term (2050–2074) and long-term (2075–2099) periods of the 21st century, compared to the baseline period (1970–1999). Projected changes in annual mean temperature suggest increasingly widespread temperature increases under the four future scenarios across the whole PTP region. The largest increases of temperature are predicted for the northern parts of the PTP region—Central Asia and the Tibetan Plateau. In the near term, the different emission scenarios do not lead to dramatically different temperature responses and the multi-model mean projects a less than 3 °C increase relative to the reference period under the four scenarios. The mid-term period can be regarded as a transition period during which the different temperature responses for the SSP2-4.5, SSP3-7.0, and SSP5-8.5 scenarios become increasingly noticeable,

with maximum values of 3.87 °C, 4.55 °C, and 5.28 °C, respectively. However, the temperature changes under the low-forcing sustainability pathway (SSP1-2.6 scenario) are relatively small throughout the 21st century, with increases generally remaining within 3 °C. By the end of the 21st century (2075–2099), changes under the SSP5-8.5 are much larger than other scenarios. The increase in the annual mean temperature under the high-forcing pathway (SSP5-8.5) will exceed 6 °C over most of the PTP region and will exceed 7 °C over the Tibetan Plateau and Central Asia.

The time series of annual mean temperature in the historical (1901–2014) and projected periods (2015–2099) in the PTP region, SEA, TP, and CA are illustrated in Figure 9. The simulations of the multi-model mean show that the annual mean temperature of the PTP region and all three subregions will increase remarkably over the 21st century under all scenarios. On average, the temperature over whole PTP region will rise by 1.24 °C/100 yr, 3.28 °C/100 yr, 5.57 °C/100 yr, and 7.40 °C/100 yr for the SSP1-2.6, SSP 2-4.5, SSP3-7.0, and SSP5-8.5 scenarios, respectively. The projected warming trends under SSP5-8.5 are considerably higher than those under SSP1-2.6 in all the considered regions. The highest warming trends occur in the CA region under all four scenarios, following by the TP region, while the lowest warming trends appear in SEA. The strong warming trends in the entire PTP region, the CA region, and the TP region are all higher than the global land average (about 7.20 °C/100 yr under SSP5-8.5) (Fan et al., 2020a), which will cause greater threats in the vulnerable ecological systems of these regions. Additionally, we found that the warming trend slows down or even declines after 2050 under SSP1-2.6 for the PTP region and its three subregions.

For the model uncertainty of the mean temperature projections, the results suggested that the uncertainty increases with time in the 21st century, and the uncertainty under high-forcing pathways is larger than it is under medium- and low-forcing pathways.

Projected changes in temperature extremes

Figure 10 presents the spatial distributions of projected changes in temperature extreme indices over the PTP region during the long-term future period at the end of the 21st century (2075–2099). The spatial distributions of changes in the near-term (2025–2049) and mid-term (2050–2074) periods are shown in Figures S8 to S11. Projected changes in the TX90p and TN90p are shown in absolute values, and not as differences relative to the reference period as for the TXx and TNn. This is because the percentile indices represent exceedance rates (in %) relative to the 1961–1990 base period, which has been used as the baseline for future changes. All four extreme temperature indices are projected to show prominent increases in the PTP region, with stronger warming under the SSP5-8.5 scenario (Figures S8–S11). The most intense warming for TXx and TNn is mainly projected in the CA region, with magnitudes of around 8 °C and 14 °C, respectively. A pronounced increase in warm days (TX90p) is projected over most of the TP region under all SSP scenarios (around 80% for SSP5-8.5), and the index value in the SEA region will also increase greatly by the end of 21st century. For warm nights (TN90p), the projected strongest increases occur in the SEA region, where the exceedance rate in some areas reaches 99.6%. The robustness of the projected increases of these four indices over the entire PTP region suggests a potential

risk of intensified temperature extremes to natural and social systems under the accelerated emission scenarios. Nevertheless, consistent with the changes in mean temperatures described above, the extreme temperature indices also show little difference with time under the SSP1-2.6 scenario, which indicates the effectiveness of anticipated climate mitigation and adaptation strategies associated with this scenario.

To identify the inter-annual trend under different scenarios, the time series of regional average annual temperature extremes indices over the three subregions and the entire PTP region during 1951–2099 are shown in Figure 11. In general, the CMIP6 models exhibit increasing trends in annual TXx, TNn, TX90p, and TN90p over the PTP region and its three subregions in the 21st century. The increase in TXx is lower than that in TNn for all four CMIP6 SSPs. For the PTP region, the multi-model mean increases in TXx and TNn, respectively, that are projected by the end of the 21st century are 1.22 °C/100 yr and 1.73 °C/100 yr in SSP1-2.6, 3.44 °C/100 yr and 4.35 °C/100 yr in SSP2-4.5, 5.72 °C/100 yr and 7.46 °C/100 yr in SSP3-7.0, and 7.58 °C/100 yr and 10.03 °C/100 yr in SSP5-8.5. By the end of the 21st century, the greatest warming trends under the four scenarios for TXx and TNn are projected in the CA region (8.40 °C/100 yr and 12.86 °C/100 yr in SSP5-8.5, respectively). The smallest annual increases in TXx and TNn are projected in the SEA region under all four SSPs (the value for both indices is 6.49 °C/100 yr in SSP5-8.5). The increasing trends of TN90p are greater than that of warm days (TX90p). Toward the end of the 21st century, the warming trends for TX90p and TN90p over the entire PTP region are 11.94%/100 yr and 14.18%/100 yr for SSP1-2.6, 31.08%/100 yr and 38.54%/100 yr for SSP2-4.5,

49.67%/100 yr and 60.27%/100 yr for SSP3-7.0, and 63.49%/100 yr and 72.80%/100 yr for SSP5-8.5. The strongest warming for TX90p occurs in the SEA region (70.67%/100 yr) followed by warming in the TP region of about 70.46%/100 yr for SSP5-8.5. The highest increase for TN90p (about 78.64%/100 yr for SSP5-8.5 by year 2099) occurs in the TP region, followed by the SEA region (75.21%/100 yr). In conclusion, the warming trends of the nighttime extremes (TNn and TN90p) are larger than those of the daytime extremes (TXx and TX90p), which is consistent with the results for CMIP5 in previous studies (Yin et al., 2019; You et al., 2018), probably because the water vapor and radiative feedbacks are enhanced at lower air temperatures (Ohmura, 2012; You et al., 2018).

Summary and conclusions

In this work, a comprehensive evaluation of CMIP6 ensembles over the Pan-Third Pole region was performed to assess their performance in simulating spatial patterns and temporal variability of mean temperatures and four indices of temperature extremes, which constitute the most important parameters for modeling climate-related changes to the terrestrial water cycle. From the results of the spatial pattern analysis of the mean and extreme temperatures, we found that most CMIP6 simulations are in fairly good agreement with CRU or HadEX3 observations in many areas, but model replication of observed temperature patterns over the Tibetan Plateau is problematic. The multi-model ensemble mean is found to be superior to most CMIP6 model simulations, overall.

Then, temperature projections for the twenty-first century were estimated for four

integrated scenarios of socioeconomic development and greenhouse gas emissions, SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5. The multi-model ensemble of CMIP6 models reveals a continuous increase in the annual mean temperature and four extremes indices over the PTP region during the twenty-first century under all four SSP scenarios. The northern parts of PTP (Central Asia and the Tibetan Plateau) are projected to experience the largest increases in future mean temperature (exceeding 7 °C) by the end of the twenty-first century under SSP5-8.5 relative to the reference periods (1970–1999). For TXx and TNn, the most intense warming will occur in the CA region, with magnitudes of around 8 °C and 14 °C, respectively. The pronounced increase of warm days (TX90p) is projected to be around 80% over the TP region and under the SSP5-8.5 scenario, and the greatest number of projected warm nights (TN90p) occur in the SEA region, where the exceedance rate in some areas reaches 99.6%.

By the end of the twenty-first century, the annual mean temperature averaged over the PTP region is projected to increase for the SSP1-2.6, SSP 2-4.5, SSP3-7.0, and SSP5-8.5 scenarios by 1.24 °C/100 yr, 3.28 °C/100 yr, 5.57 °C/100 yr, and 7.40 °C/100 yr, respectively. For temperature extremes, we found that the increasing trends in indices based on TN are greater than the increases in indices based on TX for all four CMIP6 SSPs. The CA region is projected to exhibit the greatest warming trends for TXx and TNn across the four scenarios (8.40 °C/100 yr and 12.86 °C/100 yr for SSP5-8.5, respectively). The strongest warming during the twenty-first century for TX90p and TN90p occur during SSP5-8.5 in the SEA region (70.67%/100 yr) and the TP region (78.64%/100 yr), respectively. Finally, we note that the projected changes in

mean and extreme temperatures are stronger under the highest emissions scenario;
future heat risk in the Pan-Third Pole region would be mitigated by reducing
greenhouse gas emissions.

Acknowledgments

We acknowledge the World Climate Research Program's Working Group on Coupled
Modeling, which is responsible for CMIP, and thank climate modeling groups (listed in
Table 1 of this paper) for producing and making available their model output. This
research received support from the Strategic Priority Research Program of the Chinese
Academy of Sciences (No. XDA20060401) and the National Natural Science
Foundation of China (No. 51979004).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the
corresponding author on reasonable request.

Funding

This work was financially supported by the Strategic Priority Research Program of the
Chinese Academy of Sciences (No. XDA20060401) and the National Natural Science
Foundation of China (No. 51979004).

Author information

Affiliations

State Key Laboratory of Earth Surface Processes and Resource Ecology, Faculty of
Geographical Science, Beijing Normal University, Beijing 100875, China

Xuewei Fan, Qingyun Duan, Chenwei Shen, Yi Wu & Chang Xing

College of Hydrology and Water Resources, Hohai University, Nanjing 210098, China

Qingyun Duan

Contributions

Qingyun Duan and Xuewei Fan contributed to the study conception and design. Xuewei Fan and Chenwei Shen performed the experiments and analysis. Material preparation, data collection and analysis were performed by Yi Wu & Chang Xing. The manuscript was written by Xuewei Fan and edited by Qingyun Duan. All authors read and approved the manuscript.

Corresponding author

Correspondence to Qingyun Duan.

Ethics declarations

Consent to participate

All authors have participated and approved the final version of the manuscript.

Consent to publish

All authors have approved the final version of the manuscript and have given their consent for publication.

Conflict of interest

The authors declare no competing interests.

References

- Ahmed K, Sachindra DA, Shahid S, Iqbal Z, Nawaz N, Khan N (2020) Multi-model ensemble predictions of precipitation and temperature using machine learning algorithms. *Atmos Res* 236: 104806. <https://doi.org/10.1016/j.atmosres.2019.104806>
- Annan JD, Hargreaves JC (2011) Understanding the CMIP3 multimodel ensemble. *J Clim* 24(16): 4529-4538. <https://doi.org/10.1175/2011JCLI3873.1>
- Bannister D, Herzog M, Graf H-F, Hosking JS, Short CA (2017) An assessment of recent and future temperature change over the Sichuan basin, China, using CMIP5 climate models. *J Clim* 30(17): 6701-6722. <https://doi.org/10.1175/JCLI-D-16-0536.1>
- Cheng J, Xu Z, Bambrick H, Su H, Tong S, Hu W (2018) Heatwave and elderly mortality: An evaluation of death burden and health costs considering short-term mortality displacement. *Environ Int* 115: 334-342. <https://doi.org/10.1016/j.envint.2018.03.041>
- Deng H, Pepin NC, Chen Y (2017) Changes of snowfall under warming in the Tibetan

504 Plateau. J Geophys Res-Atmos 122(14): 7323-7341.
505 <https://doi.org/10.1002/2017JD026524>

506 Dong T-Y, Dong W-J, Guo Y, Chou J-M, Yang S-L, Tian D, Yan D-D (2018) Future
507 temperature changes over the critical Belt and Road region based on CMIP5
508 models. Adv Clim Chang Res 9(1): 57-65.
509 <https://doi.org/10.1016/j.accre.2018.01.003>

510 Dunn RJH, Alexander LV, Donat MG et al (2020) Development of an updated global
511 land in situ-based data set of temperature and precipitation extremes: HadEX3. J
512 Geophys Res-Atmos 125(16): e2019JD032263.
513 <https://doi.org/10.1029/2019JD032263>

514 Eyring V, Bony S, Meehl GA, Senior CA, Stevens B, Stouffer RJ, Taylor KE (2016)
515 Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6)
516 experimental design and organization. Geosci Model Dev 9(5): 1937-1958.
517 <https://doi.org/10.5194/gmd-9-1937-2016>

518 Eyring V, Cox PM, Flato GM et al (2019) Taking climate model evaluation to the next
519 level. Nat Clim Change 9(2): 102-110. [https://doi.org/10.1038/s41558-018-0355-](https://doi.org/10.1038/s41558-018-0355-y)
520 [y](https://doi.org/10.1038/s41558-018-0355-y)

521 Fan X, Duan Q, Shen C, Wu Y, Xing C (2020a) Global surface air temperatures in
522 CMIP6: historical performance and future changes. Environ Res Lett 15(10):
523 104056. <https://doi.org/10.1088/1748-9326/abb051>

524 Fan XW, Miao CY, Duan QY, Shen CW, Wu Y (2020b) The performance of CMIP6
525 versus CMIP5 in simulating temperature extremes over the global land surface. J

526 Geophys Res-Atmos 125(18): e2020JD033031.
 527 <https://doi.org/10.1029/2020JD033031>
 528 Feng S, Hu Q, Huang W, Ho C-H, Li R, Tang Z (2014) Projected climate regime shift
 529 under future global warming from multi-model, multi-scenario CMIP5
 530 simulations. Global Planet Change 112: 41-52.
 531 <https://doi.org/10.1016/j.gloplacha.2013.11.002>
 532 Gou J, Miao C, Duan Q, Tang Q, Di Z, Liao W, Wu J, Zhou R (2020) Sensitivity
 533 Analysis-Based Automatic Parameter Calibration of the VIC Model for
 534 Streamflow Simulations Over China. Water Resour Res 56(1): e2019WR025968.
 535 <https://doi.org/10.1029/2019WR025968>
 536 Gou JJ, Miao CY, Samaniego L, Xiao M, Wu JW, Guo XY (2021) CNRD v1.0: a high-
 537 quality natural runoff dataset for hydrological and climate studies in China. Bull
 538 Amer Meteorol Soc 102(5), E929-E947. [https://doi.org/10.1175/BAMS-D-20-](https://doi.org/10.1175/BAMS-D-20-0094.1)
 539 0094.1
 540 Grose MR, Narsey S, Delage FP et al (2020) Insights From CMIP6 for Australia's
 541 Future Climate. *Earth's Future*, 8(5): e2019EF001469.
 542 <https://doi.org/10.1029/2019EF001469>
 543 Harris I, Osborn TJ, Jones P, Lister D (2020) Version 4 of the CRU TS monthly high-
 544 resolution gridded multivariate climate dataset. Sci Data 7(1): 109.
 545 <https://doi.org/10.1038/s41597-020-0453-3>
 546 Jia K, Ruan Y, Yang Y, You Z (2019) Assessment of CMIP5 GCM Simulation
 547 Performance for Temperature Projection in the Tibetan Plateau. Earth Space Sci

548 6(12): 2362-2378. <https://doi.org/10.1029/2019EA000962>

549 Kamworapan S, Surussavadee C (2019) Evaluation of CMIP5 Global Climate Models
550 for Simulating Climatological Temperature and Precipitation for Southeast Asia.
551 Adv Meteorol 2019: 1067365. <https://doi.org/10.1155/2019/1067365>

552 Kang S, Zhang Q, Qian Y (2019) Linking atmospheric pollution to cryospheric change
553 in the Third Pole region: current progress and future prospects. Natl Sci Rev 6(4):
554 796-809. <https://doi.org/10.1093/nsr/nwz031>

555 Kim Y-H, Min S-K, Zhang X, Sillmann J, Sandstad M (2020) Evaluation of the CMIP6
556 multi-model ensemble for climate extreme indices. Weather Clim Extreme 29:
557 100269. <https://doi.org/10.1016/j.wace.2020.100269>

558 Kong DX, Miao CY, Wu JW, Zheng HY, Wu SH (2020) Time lag of vegetation growth
559 on the Loess Plateau in response to climate factors: Estimation, distribution, and
560 influence. Sci Total Environ 744, 140726.
561 <https://doi.org/10.1016/j.scitotenv.2020.140726>

562 Li D, Zhou T, Zou L, Zhang W, Zhang L (2018) Extreme high-temperature events over
563 east Asia in 1.5°C and 2°C warmer futures: analysis of NCAR CESM low-
564 warming experiments. Geophys Res Lett 45(3): 1541-1550.
565 <https://doi.org/10.1002/2017GL076753>

566 Li L, Yao N, Li Y, Liu D L, Wang B, Ayantobo OO (2019) Future projections of extreme
567 temperature events in different sub-regions of China. Atmos Res 217: 150-164.
568 <https://doi.org/10.1016/j.atmosres.2018.10.019>

569 Li Y, Yan D, Peng H, Xiao S (2021) Evaluation of precipitation in CMIP6 over the

570 Yangtze River Basin. Atmos Res 253: 105406.
 571 <https://doi.org/10.1016/j.atmosres.2020.105406>

572 Liu B, Martre P, Ewert F et al (2019) Global wheat production with 1.5 and 2.0°C above
 573 pre-industrial warming. Glob Change Biol 25(4): 1428-1444.
 574 <https://doi.org/10.1111/gcb.14542>

575 Liu X, Liu Y, Liu Z, Chen Z (2021) Impacts of climatic warming on cropping system
 576 borders of China and potential adaptation strategies for regional agriculture
 577 development. Sci Total Environ 755: 142415.
 578 <https://doi.org/10.1016/j.scitotenv.2020.142415>

579 Lovino MA, Pierrestegui MJ, Müller OV, Berbery EH, Müller GV, Pasten M (2021)
 580 Evaluation of historical CMIP6 model simulations and future projections of
 581 temperature and precipitation in Paraguay. Clim Change 164(3): 46.
 582 <https://doi.org/10.1007/s10584-021-03012-4>

583 Meehl GA, Boer GJ, Covey C, Latif M, Stouffer RJ (2000) The Coupled Model
 584 Intercomparison Project (CMIP). Bull Amer Meteorol Soc 81(2): 313-318.
 585 <https://www.jstor.org/stable/26215108>

586 Miao C, Duan Q, Sun Q, Lei X, Li H (2019) Non-uniform changes in different
 587 categories of precipitation intensity across China and the associated large-scale
 588 circulations. Environ Res Lett 14(2): 025004. [https://doi.org/10.1088/1748-](https://doi.org/10.1088/1748-9326/aaf306)
 589 [9326/aaf306](https://doi.org/10.1088/1748-9326/aaf306)

590 Ohmura A (2012) Enhanced temperature variability in high-altitude climate change.
 591 Theor Appl Climatol 110, 499–508. <https://doi.org/10.1007/s00704-012-0687-x>

O'Neill BC, Tebaldi C, van Vuuren DP et al (2016) The Scenario Model Intercomparison Project (ScenarioMIP) for CMIP6. *Geosci Model Dev*, 9(9): 3461-3482. <https://doi.org/10.5194/gmd-9-3461-2016>

O'Neill BC, Carter TR, Ebi K (2020) Achievements and needs for the climate change scenario framework. *Nat Clim Change* 10(12): 1074-1084. <https://doi.org/10.1038/s41558-020-00952-0>

O'Neill BC, Kriegler E, Ebi KL et al (2017) The roads ahead: Narratives for shared socioeconomic pathways describing world futures in the 21st century. *Global Environ Chang* 42: 169-180. <https://doi.org/10.1016/j.gloenvcha.2015.01.004>

Ongoma V, Chen H, Gao C (2018) Projected changes in mean rainfall and temperature over East Africa based on CMIP5 models. *Int J Climatol* 38(3): 1375-1392. <https://doi.org/10.1002/joc.5252>

Osborn TJ, Jones PD, Lister DH, Morice CP, Simpson IR, Winn JP, Hogan E, Harris IC (2021) Land Surface Air Temperature Variations Across the Globe Updated to 2019: The CRUTEM5 Data Set. *J Geophys Res-Atmos* 126(2): e2019JD032352. <https://doi.org/10.1029/2019JD032352>

Pepin N, Deng H, Zhang H, Zhang F, Kang S, Yao T (2019) An examination of temperature trends at high elevations across the Tibetan Plateau: The use of MODIS LST to understand patterns of elevation-dependent warming. *J Geophys Res-Atmos* 124(11): 5738-5756. <https://doi.org/10.1029/2018JD029798>

Riahi K, van Vuuren DP, Kriegler E et al (2017) The Shared Socioeconomic Pathways and their energy, land use, and greenhouse gas emissions implications: An

614 overview. Global Environ Chang 42: 153-168.
615 <https://doi.org/10.1016/j.gloenvcha.2016.05.009>

616 Seneviratne SI, Wartenburger R, Guillod BP (2018) Climate extremes, land–climate
617 feedbacks and land-use forcing at 1.5°C. Philos T R Soc A 376(2119): 20160450.
618 <https://doi.org/10.1098/rsta.2016.0450>

619 Sillmann J, Kharin VV, Zhang X, Zwiers FW, Bronaugh D (2013) Climate extremes
620 indices in the CMIP5 multimodel ensemble: Part 1. Model evaluation in the
621 present climate. J Geophys Res-Atmos 118(4): 1716-1733.
622 <https://doi.org/10.1002/jgrd.50203>

623 Srivastava A, Grotjahn R, Ullrich PA (2020) Evaluation of historical CMIP6 model
624 simulations of extreme precipitation over contiguous US regions. Weather Clim
625 Extreme 29: 100268. <https://doi.org/10.1016/j.wace.2020.100268>

626 Sun QH, Miao CY, AghaKouchak A, Mallakpour I, Ji DY, Duan QY (2020) Possible
627 Increased frequency of ENSO-related dry and wet conditions over some major
628 watersheds in a warming climate. Bull Amer Meteorol Soc 101(4): E409 – E426.
629 <https://doi.org/10.1175/BAMS-D-18-0258.1>

630 Sun Q, Miao C, Hanel M, Borthwick AGL, Duan Q, Ji D, Li H (2019) Global heat stress
631 on health, wildfires, and agricultural crops under different levels of climate
632 warming. Environ Int 128: 125-136. <https://doi.org/10.1016/j.envint.2019.04.025>

633 Sun Q, Miao C, Qiao Y, Duan Q (2017) The nonstationary impact of local temperature
634 changes and ENSO on extreme precipitation at the global scale. Clim Dyn 49(11):
635 4281-4292. <https://doi.org/10.1007/s00382-017-3586-0>

636 Taylor KE (2001) Summarizing multiple aspects of model performance in a single
 637 diagram. *J Geophys Res-Atmos* 106(D7): 7183-7192.

638 Taylor KE, Stouffer RJ, Meehl GA (2012) An Overview of CMIP5 and the Experiment
 639 Design. *Bull Amer Meteorol Soc* 93(4): 485-498.
 640 <https://doi.org/10.1029/2000JD900719>

641 Tigchelaar M, Battisti DS, Naylor RL, Ray DK (2018) Future warming increases
 642 probability of globally synchronized maize production shocks. *P Natl Acad Sci*
 643 *Usa* 115(26): 6644. <https://doi.org/10.1073/pnas.1718031115>

644 Wang T, Chen Y, Gan Z, Han Y, Li J, Huang J (2020) Assessment of dominating aerosol
 645 properties and their long-term trend in the Pan-Third Pole region: A study with 10-
 646 year multi-sensor measurements. *Atmos Environ* 239: 117738.
 647 <https://doi.org/10.1016/j.atmosenv.2020.117738>

648 Wehner MF (2020) Characterization of long period return values of extreme daily
 649 temperature and precipitation in the CMIP6 models: Part 2, projections of future
 650 change. *Weather Clim Extreme* 30: 100284.
 651 <https://doi.org/10.1016/j.wace.2020.100283>

652 Yang K, Wu H, Qin J, Lin C, Tang W, Chen Y (2014) Recent climate changes over the
 653 Tibetan Plateau and their impacts on energy and water cycle: A review. *Glob Planet*
 654 *Change* 112: 79-91. <https://doi.org/10.1016/j.gloplacha.2013.12.001>

655 Yao T, Chen F, Cui P, Ma Y, Xu B, Zhu L, Zhang F, Wang W, Ai L, Yang X (2017) From
 656 Tibetan plateau to third Pole and pan-third Pole. *Bull Chin Acad Sci* 32, 924–
 657 931(in Chinese). <https://doi.org/10.16418/j.issn.1000-3045.2017.09.001>

658 Yao T, Xue Y, Chen D (2019) Recent Third Pole's Rapid Warming Accompanies
659 Cryospheric Melt and Water Cycle Intensification and Interactions between
660 Monsoon and Environment: Multidisciplinary Approach with Observations,
661 Modeling, and Analysis. Bull Amer Meteorol Soc 100(3): 423-444.
662 <https://doi.org/10.1175/BAMS-D-17-0057.1>

663 Yin H, Sun Y, Donat MG (2019) Changes in temperature extremes on the Tibetan
664 Plateau and their attribution. Environ Res Lett 14(12): 124015.
665 <https://doi.org/10.1088/1748-9326/ab503c>

666 You Q, Jiang Z, Wang D, Pepin N, Kang S (2018) Simulation of temperature extremes
667 in the Tibetan Plateau from CMIP5 models and comparison with gridded
668 observations. Clim Dyn 51(1): 355-369. [https://doi.org/10.1007/s00382-017-](https://doi.org/10.1007/s00382-017-3928-y)
669 [3928-y](https://doi.org/10.1007/s00382-017-3928-y)

670 You Q, Zhang Y, Xie X, Wu F (2019) Robust elevation dependency warming over the
671 Tibetan Plateau under global warming of 1.5 °C and 2 °C. Clim Dyn 53(3): 2047-
672 2060. <https://doi.org/10.1007/s00382-019-04775-4>

673 Zheng HY, Miao CY, Jiao JY, Borthwick AGL (2021) Complex relationships between
674 water discharge and sediment concentration across the Loess Plateau, China. J
675 Hydrol 596, 126078. <https://doi.org/10.1016/j.jhydrol.2021.126078>

676 Zhou B, Wen QH, Xu Y, Song L, Zhang X (2014) Projected changes in temperature and
677 precipitation extremes in China by the CMIP5 multimodel ensembles. J. Clim
678 27(17): 6591-6611. <https://doi.org/10.1175/JCLI-D-13-00761.1>

679 Zhu Y-Y, Yang S (2020) Evaluation of CMIP6 for historical temperature and

680 precipitation over the Tibetan Plateau and its comparison with CMIP5. Adv Clim

681 Chang Res 11(3): 239-251. <https://doi.org/10.1016/j.accre.2020.08.001>

682

683

Table 1 List of 16 CMIP6 models in this study and their spatial resolutions

Model name	Modeling center	Spatial resolution
BCC-CSM2-MR	Beijing Climate Center, China	320×160
CAMS-CSM1-0	Chinese Academy of Meteorological Sciences, China	320×160
CESM2-WACCM	National Center for Atmospheric Research, Climate and Global Dynamics Laboratory, United States	288×192
CESM2	National Center for Atmospheric Research, Climate and Global Dynamics Laboratory, United States	288×192
CNRM-CM6-1	National Centre for Meteorological Research, France	256×128
CNRM-ESM2-1	National Centre for Meteorological Research, France	256×128
CanESM5	Canadian Centre for Climate Modelling and Analysis, Environment and Climate Change Canada, Canada	128×64
EC-Earth3-Veg	EC-Earth Consortium, Europe	512×256
EC-Earth3	EC-Earth Consortium, Europe	512×256
FGOALS-g3	LASG, Institute of Atmospheric Physics, Chinese Academy of Sciences, China	180×80
GFDL-ESM4	National Oceanic and Atmospheric Administration, Geophysical Fluid Dynamics Laboratory, United States	288×180
IPSL-CM6A-LR	Institut Pierre Simon Laplace, France	144×143
MIROC-ES2L	JAMSTEC (Japan Agency for Marine-Earth Science	128×64

and Technology), AORI (Atmosphere and Ocean
Research Institute, The University of Tokyo), NIES
(National Institute for Environmental Studies), and R-
CCS (RIKEN Center for Computational Science),
Japan

MIROC6	JAMSTEC, AORI, NIES and R-CCS, Japan	256×128
MRI-ESM2-0	Meteorological Research Institute, Japan	320×160
UKESM1-0-LL	Met Office Hadley Centre, United Kingdom	192×144

685

686

Table 2 Definitions of the extreme temperature indices used in this study

Label	Index Name	Index Definition	Units
TXx	Max TX	Annual maximum value of daily maximum temperature	°C
TNn	Min TN	Annual minimum value of daily minimum temperature	°C
TX90p	Warm days	Percentage of days when the daily maximum temperature is above the 90th percentile for the base period 1961–1990	%
TN90p	Warm nights	Percentage of days when the daily minimum temperature is above the 90th percentile for the base period 1961–1990	%

Figure Captions

Figure 1 Location map of the study area. The blue rectangles represent the latitude and longitude ranges of the three subregions: Central Asia (CA), Tibetan Plateau (TP), and Southeast Asia (SEA).

Figure 2 Spatial distributions of annual mean temperatures from 16 CMIP6 models, ensemble averages, and CRU observations over the PTP region for 1970–1999 average.

Figure 3 Area-averaged annual biases of temperature relative to the CRU observational dataset for each CMIP6 ensemble mean during 1970–1999 over the PTP region and its three subregions, (a); the light purple bars represent the maximum and minimum biases of the 16 GCMs, and the horizontal line in dark purple represents the bias of the multi-model mean. Taylor diagram displaying statistics of climatological annual mean temperature generated from 16 GCMs and CRU in the PTP region during 1970–1999, (b); the vertical axis indicates standard deviation ratios, the numbers along the arc are the spatial correlations, and the green circles centered at CRU measure RMSE difference relative to the standard deviation of the observations.

Figure 4 Time series of 10-year moving average annual surface mean temperature from the CMIP6 models and CRU observational dataset over the PTP region and its three subregions during 1901–2014 relative to the period 1970–1999 (blue line and shading: CMIP6; black line: CRU). The trends are calculated for the observations and the CMIP6 ensemble mean during 1901–2014. The shading indicates the ensemble spread (range between the 5th and 95th quantiles).

Figure 5 Spatial distributions of annual mean max TX (TXx), min TN (TNn), warm days (TX90p), and warm nights (TN90p) for HadEX3 and the CMIP6 ensemble mean over the PTP region for the 1970–1999 average.

Figure 6 The portrait diagram of relative RMSEs for the 1970–1999 climatologies of temperature extreme indices simulated by the CMIP6 models with respect to the HadEX3 dataset. (Note that the models were masked by HadEX3 due to the insufficient spatial coverage of HadEX3.) The RMSEs are spatially averaged over the PTP region and the top row indicates the mean relative RMSE across all indices for a particular model.

Figure 7 Time series of the 10-year moving average annual mean max TX (TXx), min TN (TNn), warm days (TX90p), and warm nights (TN90p) from the CMIP6 models and the HadEX3 observational dataset over the PTP region during 1951–2014. For TXx and TNn, the time series shows anomalies with respect to the reference period 1970–1999; For TX90p and TN90p, the time series are displayed as absolute exceedance rate). The trends are calculated for the observations and the CMIP6 ensemble mean during 1951–2014. The shading indicates the ensemble spread (range between the 5th and 95th quantiles).

Figure 8 Spatial distribution of changes in annual mean temperatures over the PTP region in the near-term (2025–2049), mid-term (2050–2074), and long-term (2075–2099) periods of the 21st century, relative to 1970–1999, under the SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5 scenarios.

Figure 9 Time series of changes in annual mean temperature over the PTP region and its three subregions during 1901–2099 relative to the period 1970–1999. The black, green, blue, red, and purple curves represent the results of the CMIP6 ensemble mean for the historical period and for the SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5 scenarios, respectively. The shaded areas are the spreads from the 5th to the 95th percentiles of the annual mean temperatures.

Figure 10 Spatial distributions of projected changes in annual mean max TX (TXx), min TN (TNn), warm days (TX90p), and warm nights (TN90p) over the period 2075–2099 relative to the reference period 1970–1999 under the SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5 scenarios. Note that TX90p and TN90p are displayed as absolute exceedance rate.

Figure 11 Time series of changes in annual mean max TX (TXx), min TN (TNn), warm days (TX90p), and warm nights (TN90p) over the PTP region and its three subregions during 1901–2099 relative to the period 1970–1999. (Note that the time series for TX90p and TN90p are displayed as an absolute exceedance rate). The black, green, blue, red, and purple curves represent the results for the CMIP6 ensemble mean for the historical period and for the SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5 scenarios, respectively. The shaded areas are the spreads from the 5th to the 95th percentiles of the annual mean temperature extreme indices.

Figure 1 Location map of the study area. The blue rectangles represent the latitude and longitude ranges of the three subregions: Central Asia (CA), Tibetan Plateau (TP), and Southeast Asia (SEA).

Figure 2 Spatial distributions of annual mean temperatures from 16 CMIP6 models, ensemble averages, and CRU observations over the PTP region for 1970–1999 average.

Figure 3 Area-averaged annual biases of temperature relative to the CRU observational dataset for each CMIP6 ensemble mean during 1970–1999 over the PTP region and its three subregions, (a); the light purple bars represent the maximum and minimum biases of the 16 GCMs, and the horizontal line in dark purple represents the bias of the multi-model mean. Taylor diagram displaying statistics of climatological annual mean temperature generated from 16 GCMs and CRU in the PTP region during 1970–1999, (b); the vertical axis indicates standard deviation ratios, the numbers along the arc are the spatial correlations, and the green circles centered at CRU measure RMSE difference relative to the standard deviation of the observations.

Figure 4 Time series of 10-year moving average annual surface mean temperature from the CMIP6 models and CRU observational dataset over the PTP region and its three subregions during 1901–2014 relative to the period 1970–1999 (blue line and shading: CMIP6; black line: CRU). The trends are calculated for the observations and the CMIP6 ensemble mean during 1901–2014. The shading indicates the ensemble spread (range between the 5th and 95th quantiles).

Figure 5 Spatial distributions of annual mean max TX (TXx), min TN (TNn), warm days (TX90p), and warm nights (TN90p) for HadEX3 and the CMIP6 ensemble mean over the PTP region for the 1970–1999 average.

Figure 6 The portrait diagram of relative RMSEs for the 1970–1999 climatologies of temperature extreme indices simulated by the CMIP6 models with respect to the HadEX3 dataset. (Note that the models were masked by HadEX3 due to the insufficient spatial coverage of HadEX3.) The RMSEs are spatially averaged over the PTP region and the top row indicates the mean relative RMSE across all indices for a particular model.

Figure 7 Time series of the 10-year moving average annual mean max TX (TXx), min TN (TNn), warm days (TX90p), and warm nights (TN90p) from the CMIP6 models and the HadEX3 observational dataset over the PTP region during 1951–2014. For TXx and TNn, the time series shows anomalies with respect to the reference period 1970–1999; For TX90p and TN90p, the time series are displayed as absolute exceedance rate). The trends are calculated for the observations and the CMIP6 ensemble mean during 1951–2014. The shading indicates the ensemble spread (range between the 5th and 95th quantiles).

Figure 8 Spatial distribution of changes in annual mean temperatures over the PTP region in the near-term (2025–2049), mid-term (2050–2074), and long-term (2075–2099) periods of the 21st century, relative to 1970–1999, under the SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5 scenarios.

Figure 9 Time series of changes in annual mean temperature over the PTP region and its three subregions during 1901–2099 relative to the period 1970–1999. The black, green, blue, red, and purple curves represent the results of the CMIP6 ensemble mean for the historical period and for the SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5 scenarios, respectively. The shaded areas are the spreads from the 5th to the 95th percentiles of the annual mean temperatures.

Figure 10 Spatial distributions of projected changes in annual mean max TX (TXx), min TN (TNn), warm days (TX90p), and warm nights (TN90p) over the period 2075–2099 relative to the reference period 1970–1999 under the SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5 scenarios. Note that TX90p and TN90p are displayed as absolute exceedance rate.

Figure 11 Time series of changes in annual mean max TX (TXx), min TN (TNn), warm days (TX90p), and warm nights (TN90p) over the PTP region and its three subregions during 1901–2099 relative to the period 1970–1999. (Note that the time series for TX90p and TN90p are displayed as an absolute exceedance rate). The black, green, blue, red, and purple curves represent the results for the CMIP6 ensemble mean for the historical period and for the SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5 scenarios, respectively. The shaded areas are the spreads from the 5th to the 95th percentiles of the annual mean temperature extreme indices.

830

Supporting Information for

831

832

Evaluation of historical CMIP6 model simulations and future

833

projections of temperature over the Pan-Third Pole region

834

835

Contents of this file

836

Figures S1 to S11

837

Figure S1 Spatial distributions of annual mean max TX (TXx) for the CMIP6 models over the PTP for the 1970–1999 average.

Figure S2 Spatial distributions of annual mean min TN (TNn) for the CMIP6 models over the PTP for the 1970–1999 average.

Figure S3 Spatial distributions of annual mean warm days (TX90p) for the CMIP6 models over the PTP for the 1970–1999 average.

847

848 Figure S4 Spatial distributions of annual mean warm nights (TN90p) for the CMIP6

849 models over the PTP for the 1970–1999 average.

Figure S5 Time series of 10-year moving average annual mean max TX (TXx), min TN (TNn), warm days (TX90p), and warm nights (TN90p) from the CMIP6 models and HadEX3 observational dataset over the SEA region during 1951–2014 (for TXx and TNn, the time series shows anomalies with respect to the reference period 1970 to 1999; for TX90p and TN90p, the time series are displayed as absolute exceedance rates). The trends are calculated for the observations and the CMIP6 ensemble mean during 1951–2014. The shading indicates the ensemble spread (range between the 5th and 95th quantiles).

859

860 Figure S6 Same as Figure S5, but for the TP region.

861

862 Figure S7 Same as Figure S5, but for the CA region.

863

Figure S8 Spatial distributions of changes in annual mean max TX (TXx) over the PTP regions in near-term (2025–2049), mid-term (2050–2074), and long-term (2075–2099) periods of the 21st century, relative to 1970–1999, under the SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5 scenarios.

869

870 Figure S9 Same as Figure S8, but for min TN (TNn).

871

872 Figure S10 Same as Figure S8, but for warm days (TX90p).

Figure S11 Same as Figure S8, but for warm nights (TN90p).