Preliminary Geological and Geochemical Study of Gold and Base Metals Mineralization in The Surupa Area, Borena Zone, Southern Ethiopia

Waktola Merera (✉ benolmaro4@gmail.com)

Department of Geology, Bule Hora University

Original article

Keywords: Gold mineralization, Mineral paragenesis, rare earth element, Base metals, Precambrian basement rocks, Petrography and Geochemistry

Posted Date: July 20th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-620878/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: The southern Ethiopian Precambrian shield terrain largely of sutured medium to high-grades assemblages of Precambrian crystalline basement consists of metamorphic complex and associated plutonic rocks which belongs to northern and southern continuation of the Mozambique belt and the Arabian-Nubian Shield. The paper reported the information on geology, alteration pattern, ore minerals and geochemical investigation for gold and base metals and to delineate potential area of gold and base metals mineralization. The purpose of the study was to carry out a preliminary investigation on geological and geochemical study for gold and base metals mineral potentials around Surupa area.

Results: The study area consists of heterogeneous sequences of medium to high-grade poly-deformed gneisses with igneous and sedimentary precursors and structurally affected by three phases of deformation like foliations, mesoscale folds, veins and fractures/joints respectively. The rocks of the area characterized by its medium to high-potassium calc-alkaline rocks of high silicon dioxide and displaying predominance of potassium oxide over sodium oxide. The positive and negative correlation with major oxides exhibited indicate the fractionation nature of the rocks during magmatic differentiation.

Conclusions: Based on the result, nickel and cobalt are highly anomalous elements indicating the source for the mineralization controlled by different structures such as fractures and joints. Ore mineral assemblages of pyrite, goethite, magnetite and hematite characterizes the mineralization of the study area. Generally, all those integrated studies confirmed that mineral potential of the study area is mainly controlled by the nature of the rock and associated structures.

1. Introduction

The Precambrian basement rocks of the southern Ethiopia are sources of most of the economic metallic mineral deposits that include primary and secondary enriched deposits of gold, platinum group elements (PGE), nickel, tantalum, base metals, industrial minerals such as phosphate, iron ore, gemstones and also decorative and dimension stones (Ministry of Mines and Energy, 2009). Based on geological set up of the country and possible occurrence of gold and base metal deposits, several studies have been carried out by the governmental and private companies reported their occurrence from different parts of the country in the low-grade basement rocks of greenschist facies (Tadesse, 2004). These includes; brittle–ductile shear-zone related mesothermal gold deposits in Lega Dembi, Megado and Sakaro area; Au, Ag Alluvial–eluvial placers in Adola Belt to Bore; Brittle–ductile shear-zone related mesothermal gold deposits in Adi Dairo-Indallilo area(northern Ethiopia); Au–Cu–Pb–Zn in Ashashire; Au–Cu in Bomo; v) Au–Cu–Pb in Digati; ductile shear zone-hosted mesothermal gold mineralization in south Okote area; vii) mesothermal type gold in south-west Akebo area, and epithermal gold in the rift valley and Afar regions, among others (Tadesse, 2004).

The biggest developed mine in the country is that of southern Ethiopia's Lega Dembi gold mine which remains the only mesothermal gold deposit under mining operations. It is related to the shear-zone

hosted hydrothermal quartz veins within the Neoproterozoic metamorphosed volcano-sedimentary succession of greenschist to amphibolites facies metamorphism (Billay et al., 1997; Tadesse, 2004). It is estimated to have a reserve of about 260 tons of gold with an average grade of 6ppm (Billay et al., 1997) and over 43 tones and 65.94 tones with average grade of 3.7ppm at nearby areas Sakaro and Tulu Kabi around Lega-Dembi respectively (Ministry of Mines and Energy, 2009).

Accordingly, the study area is located very close to the well-known Adola gold field within the southern Ethiopia greenstone belt and there is an extensive artisanal mining activity by the local people for gold. However, no systematic study were conducted in terms of their geology, alteration pattern, ore mineral assemblages, chemistry and petrography of the host rock responsible for gold and base metals mineralization.

The purpose of this paper is to provide information on geology, alteration pattern, ore minerals and geochemical investigation for gold and base metals and to delineate potential area of gold and base metals mineralization.

2. Materials And Methods

A. Study area

The study area covers an area of about 50 sq.km in Surupa Woreda of Borena zone, Oromia Region, Southern Ethiopia, which is bounded by, coordinates of 412000mE to 438000mE longitude and 560000mN to 586000mN latitude (Fig. 1). Topographically, it is characterized by very rugged to gentle sloping from northwest-southeast direction and ranges from 2000 to 1600 meters above mean sea level. The region experiences of a bi-modal monsoon rainfall type, where 60% of the 300-900mm annual rainfall occurs during March to May and 40% between September and November. Heavy cloud cover, vapor and occasionally short showers characterize the period from June to September, while the main dry season occurs from November to March with high evaporation. The highland areas get the mean annual precipitation in the range of 600 to 900 mm whereas the lowland areas receive the lowest annual precipitation in the range of 300 to 500 mm.

Furthermore, the minimum and maximum temperatures of the area fall within the range of 13.3 to 17°C and 18 to 25.1°C, respectively. Generally, the lowland areas tend to be relatively hotter than the highland areas.

B. Samples and Methods

Representative whole rock samples from different lithological units were collected and submitted to Geological Survey of Ethiopia, Addis Ababa for sample preparation by crushing to 70% less than 2mm, riffled to split off 1kg and the splits were pulverized to 75 microns. The pulverized samples were shipped to ALS Services, Ireland laboratory for whole rock analysis to represent combination of ICP-MS and ICP-AES. Trace elements including Ba, Ce, Cr, Cs, Dy, Er, Eu, Ga, Gd, Hf, Ho, La, Lu, Nb, Nd,

Pr, Rb, Sm, Sn, Sr, Ta, Tb, Th, Tm, U, V, W, Y, Yb and Zr were analyzed by Inductively Coupled Plasma-Mass Spectrometry (ICP-MS) to understand the magma source and to discriminate tectonic setting of the rocks of the area. However, Inductively Coupled Plasma-Atomic Emission Spectrometry (ICP-AES) used for analyzing major elements in their oxide forms including SiO₂, Al₂O₃, Fe₂O₃, CaO, MgO, Na₂O, K₂O, Cr₂O₃, TiO₂, MnO, and P₂O₅.

Additionally, about sixty stream sediment samples were collected mostly from first order streams and analyzed for Au, Pb, Co, Cu, Zn and Ni by Atomic Absorption Spectrophotometer (AAS) at geochemical laboratory of Geological Survey of Ethiopia, Addis Ababa.

About nine thin sections and five polished sections have been prepared at Geological Survey of Ethiopia (GSE), Addis Ababa and analyzed at both Mekelle and Bahir Dar Universities for respectively. The fresh parts of the rock samples were sliced by the rock cutter and polished by DAP-U polishing machine to obtain the required size not greater than 4 cm to fit the mould size, which has a maximum diameter of 4cm. The rock samples were ground mechanically using carbide abrasive until the slice reaches the conventional thickness of 30 microns. This is carried out using petro discs with 400-grit aluminum oxide powder.

3. Result And Discussion

Regionally, the basement rocks of Southern Ethiopia including the study area is found in the transitional zone with imprints of Pan-African tectonothermal events (Vail, 1976; Kazmin et al., 1978; Davidson, 1983). The Precambrian rocks of Southern Ethiopia are heterogeneous sequences of high-grade poly-deformed gneisses with igneous and sedimentary origin and mainly represented by Moyale Belt, Adola Belt, and Bulbul Belt comprising the known placer and primary gold mineralization. These rocks are subjected to middle to upper amphibolite facies metamorphism with localized granitization and migmatization and pockets of granulites. The study area is the northern extension of the Moyale Belt and very close to the known Adola belt.

Based on the dominant mineral assemblages and their textures, the rocks are named as feldspar-plagioclase-biotite gneiss, feldspar-plagioclase-biotite-muscovite gneiss, feldspar-quartz-plagioclase-granitic gneiss, feldspar-plagioclase gneiss, plagioclase-pyroxene schist and actinolitic hornblende-quartz-plagioclase-schist. The nature of alignment, texture, size and type of minerals identified in the rocks implies that the rocks have experienced middle to high-grade amphibolite facies metamorphism (Fig. 2). The mineral assemblage from this rock indicates that the protolith is possibly intermediate rock and corresponding geochemistry (Appen1, Table 1, and MtR1) plotting in the intermediate field (Fig. 3).

3.1 Stream Sediment Geochemistry

3.1.1 Univariate data analysis

The distributions of the element concentrations in the stream sediments are observed to be strongly related to specific spatial and environmental factor like hydrothermal alteration zones, geological structures and lithologies drained by streams in the area (Figs. 4 and 5). Concentrations of nickel, cobalt and zinc are direct related and highly anomaly in the northwestern part of the study area mainly clustered on the metagranite and biotite gneiss rocks, which indicates the source for the mineralization controlled by different structures such as fractures and joints (Fig. 5B, E & F). However, the gold result from the stream sediment riches 0.116ppm indicates low concentration relative other base metals. The reason for the low content of Au in the stream sediment is, most of the stream sediments are not properly channelized, and hence accumulation of heavy elements like gold is randomly not stream channel controlled.

3.1.2 Multivariate data analysis

The Zn-Ni-Cu association represented by the first principal component (PC1) is interpreted to be a multi-element geochemical signature reflecting a possible combination of lithologic and chemical controls. Co has weak correlation in this principal component, which suggests these elements are depleted in the mineralized parts of the above stated lithological controls (Table 2).

The second principal component (PC2) (Table 2) represents antipathetic behavior between the negatively loaded Cu-Co and positively loaded Au-Pb. PC2 is possibly due to the leaching of elements from parent rocks enriched in Pb & Au and depleted in Cu & Co. Therefore, PC2 are more likely in the mineralized parts of the rocks of the study area. The variations explained by the PCs can be spatially outlined by calculating the principal component scores. The principal components are calculated according to the formula (George and Carter, 1989) as follow:

$$Sci = \sum_{j=1}^n Icj Zij \quad \text{For } i = 1, 2, 3, n \text{ Samples}$$

Where Sci = scores for sample I on component c; Icj = loadings on element j on component c; Zij = concentration of element j for sample i.

The geochemical investigation in the study area has proved successful in locating the base metal anomalies, associations and potential areas for mineral deposits. Most of the high anomalies recognized from the stream sediments correspond with metagranite and biotite gneiss rocks of north and southwestern part of the study area. The association in Zn-Ni-Cu more reflects to the underlying lithology of metagranite and biotite gneiss unit and the association of Au-Pb is related to base metal mineralization.

3.1.3 Whole rock Geochemistry

Rock geochemical data of the study area show different variation in their geochemistry. Based on the analysis, most of major elements plotted against SiO₂ show systematic negative correlation with TiO₂, Fe₂O₃, MgO, CaO, and MnO, which can be explained by magma differentiation. The concentration of K₂O

and Na_2O increase with increasing of SiO_2 concentration even if there are scattered and irregular distributions patterns (Fig. 6). This indicates normal fractionation igneous trends, which display a continuous gradation in chemical composition from mafic to felsic indicating the parental magma differentiation. The Na_2O and K_2O shows a wider data spread and small data scatter which are positively correlated with SiO_2 .

Table.1: Major (wt. %) and trace (ppm) element concentrations

Rock Name	Biotite gneiss			Metagranite				Amphibolite Schist	
S/Code	MtR1	MtR2	MtR3	Mt4	MtR5	MtR6	MtR7	MtR8	MtR9
Major Oxides (ICP-AES) (wt. %)									
SiO2	57.2	68.4	71.73	65.07	73.9	73.7	74.4	55.03	68.4
TiO2	0.37	0.44	0.53	0.83	0.17	0.04	0.17	0.18	0.57
Al2O3	13.35	15.16	13.79	15.5	13.91	14.85	14.05	15.46	16.05
Fe2O3	9.42	4.47	3.84	4.29	2.19	1	2	9.1	4.89
MgO	8.44	1.81	0.65	1.32	0.46	0.04	0.34	5.87	1.98
CaO	8.57	3.3	1.86	2.37	1.61	0.3	1.06	7.45	1.65
Na2O	2.35	4.23	2.98	4.47	3.14	2.34	3.59	3.68	2.48
K2O	0.18	1.79	4.18	4.07	4.63	8.82	4.82	1.78	3.66
MnO	0.16	0.06	0.06	0.05	0.03	0.01	0.04	0.15	0.08
P2O5	0.03	0.18	0.21	0.42	0.11	0.03	0.03	0.12	0.13
LOI	1.29	1.15	0.45	0.58	0.43	0.45	1.32	1.15	1.58
Total	101.36	100.99	100.28	98.97	100.58	101.58	101.82	99.97	101.47
Trace Elements (ICP-MS)(ppm)									
V	267	62	32	61	28	4.9	13	183	57
Cr	280	160	148	19	128	10	20	44	70
Co	34	43	51	44	132	0.9	4	50	11
Ni	73	9	52	56	41	7	10	81	33
Cu	44	183	67	54	40	8	8	112	32
Zn	62	84	53	90	96	5	27	39	69
Rb	1.4	16	15	24	92	137.5	203	110	101.5
Sr	67.1	404	160	854	271	105.5	165.5	251	272
Y	9.5	9	33	8	9	5.5	7.1	29	23.3
Zr	19	154	295	337	244	8	105	157	169
Nb	0.3	5	6	11	105	2.6	8.7	6	7.5
Cs	0.28	0.5	1	4.3	6.2	2.99	5.85	3.3	2.96

Rock Name	Biotite gneiss			Metagranite				Amphibolite Schist	
Ba	25.1	262	956	1849	922	221	460	174	997
Rare Earth Elements (ICP-MS)(ppm)									
La	1.3	7	1.2	13.8	4.9	1.5	13.7	6.4	0.34
Ce	2.7	9.3	2.8	15.8	6.9	2.2	24.1	5.3	61.8
Pr	0.45	1.2	0.2	2.1	0.5	0.27	2.6	0.6	6.96
Nd	2.2	6.2	1.1	10.3	2.5	1.1	9.3	3.3	25.6
Sm	0.73	2.1	0.5	3.3	0.9	0.38	1.42	1.1	4.75
Eu	0.31	0.8	0.2	1.3	0.4	0.16	0.39	0.4	1.1
Gd	1.24	2.9	0.5	4.4	1.3	0.6	1.42	1.4	4.77
Tb	0.22	0.6	0.1	0.8	0.2	0.14	0.19	1.7	0.69
Dy	1.62	3.9	0.6	5.4	1.8	1.08	1.25	1.7	4.14
Ho	0.36	0.8	0.1	1.1	0.4	0.19	0.25	0.3	0.83
Er	0.89	2.5	0.4	3.3	1.3	0.46	0.53	1	2.55
Tm	0.14	0.4	0.1	0.5	0.2	0.07	0.11	0.1	0.38
Yb	1.03	2.4	0.5	3.2	1.2	0.28	0.72	0.9	2.5

Based on the idea of (Cox et al. 1979), biotite gneiss and metagranite rock units of the area are abundantly falls in the felsic fields and some analysis of amphibolite schist is flanked by intermediate field, which may be due to the compositional variation of the clasts (Fig. 3). The discrimination between tholeiitic and calc-alkaline series interpreted by AFM plot (Irvine and Baragar 1971), (Fig. 7) emphasis that the rocks the study area typically occupy the calc-alkaline affinity indicating their (Na₂O + K₂O) rich in nature. In addition, on SiO₂-FeO_t/MgO plot (Miyashiro 1974) and SiO₂-K₂O plot (Peccerillo and Taylor, 1976), (Fig. 7) most of the samples occupy the calc-alkaline series field and high-K calc-alkaline series respectively.

Table 2
Principal Component analysis result

Element	Principal components	
	PC1	PC2
Au	0.026	0.694
Cu	0.434	-0.227
Zn	0.550	0.106
Pb	0.301	0.605
Co	0.405	-0.280
Ni	0.499	-0.105

A-(alkalis :($\text{Na}_2\text{O} + \text{K}_2\text{O}$), F-($\text{FeO} + \text{Fe}$), and M-(MgO), (B and C) SiO_2 vs. $\text{Fe}_2\text{O}_3/\text{MgO}$ and K_2O of the high K Calc-alkaline rock series further into low-K tholeiitic series of the study area.

The Chondrite normalized REE abundances plot (Boynton, 1984) (Fig. 8), shows all the rock samples of the area has moderate to high negative Eu anomalies which indicate removal of feldspar from the melt by crystal fractionation or the partial melting of a rock in which feldspar is retained in the source (Rollinson, 1993).

According to the ideas of Rollinson (1993), the REE patterns of rocks show high enrichment in LREE and high depletion in HREE when the melting percentage decreases. However, the REE pattern of Surupa rocks shows only slight variation between the abundances of LREE and HREE, which indicates moderate to high percentage melting.

Most samples of the area shows enrichment in the incompatible large ion lithophile elements (LILE: Ce, Ba, Th and Rb) relative to high field strength elements (HFSE: Zr, Y, Eu, and Yb). Large negative Ti anomaly indicates the fractionation of Ti-magnetite. The enrichment of Ba, K and depletion of Nb signifying contamination during fractional crystallization (Fig. 9).

Tectonic discrimination diagram suggest that the rocks have a tendency towards Calc-alkaline basalt (CAB) tectonic setting, which can be characteristic of subduction related magmatism.

5. Conclusion

The study area consists of heterogeneous sequences of medium to high-grade poly-deformed gneisses with igneous and sedimentary precursors and structurally affected by three phases of deformation (D1 to D3) like foliations, mesoscale folds, veins and fractures/joints respectively. Mineral assemblage, gneissose and some schistose texture properties suggest that the rocks of the study area have experienced medium to high grade amphibolite facies metamorphism and shows development of different type of alteration related to metamorphism and hydrothermal activity based on their abundance

specifically silicification, oxidation, chloritization and epidotization. Based on the result, nickel and cobalt are highly anomalous elements indicating the source for the mineralization controlled by different structures such as fractures and joints. Ore mineral assemblages of pyrite, goethite, magnetite and hematite characterizes the mineralization of the study area. Generally, all those integrated studies confirmed that mineral potential of the study area is mainly controlled by the nature of the rock and associated structures.

Declarations

Dear Sir, I would like to submit the manuscript entitled “Preliminary Geological and Geochemical Study of Gold and Base Metals Mineralization in the Surupa Area, Borena Zone, Southern Ethiopia to be considered for a publication of original article in the Energy, Sustainability and Society Journal website.

The study area, which is located very close to the well-known Adola gold field within the southern Ethiopia greenstone belt have recently found gold by artisanal mining activity and local people were experienced by crushing the quartz and country rock for primary gold mining.

Daniest mining PLC. (2015) have also reported that they have identified occurrence of gold and base metals during first year of systematic exploration, which involved geological mapping and geochemical sampling. However, there was no any systematic exploration conducted in the area except the Daniest Mining Company, which was collected soil sampling besides to the geological mapping and does not have analyze for their trace element. Therefore, this study has been designed to provide a better understanding of the geology, and geochemical investigation, and as well to search other possible targets of exploration activity.

I, **Waktola Merera**, a resident of Ambo town of Oromia state of Ethiopia, lecturer at Bule Hora University of Geology department, do hereby declare that this manuscript is original, has not been published before and is not currently being considered for publication elsewhere.

The data I used to accomplish this work were mostly primary data analyzed in different laboratory of Geological survey of Ethiopia as well as Ireland laboratory for rock geochemistry and all data are currently available with me.

I know of no conflicts of interest associated with this publication, and there has been no significant financial support for this work that could have influenced its outcome. As corresponding Author, I confirm that the manuscript has been read and approved for submission by other people invited for their comments.

Acknowledgement

The author is extremely grateful to the people from the study area for their time, hospitality and cooperation's. The author is very much indebted to Dr. Kassa Amare for his detailed critical reading and

constructive comments as well as for the better thoughts he offered. The author also expresses his appreciation and gratitude to all those who gave their support and assistance in researching and writing this paper and who provided him with information, documents and materials.

References

1. Abbate E, Bruni P, Sagri M (2015) *Geology of Ethiopia: A Review and Geomorphological Perspectives*. Springer Netherlands, 33–64 pp
2. Abdelsalam M, Stern R (1996) Sutures and shear zones in the Arabian-Nubian Shield. *J Afr Earth Sc* 23:289–300
3. Avigad D, Gvirtzman Z (2009) Late Neoproterozoic rise and fall of the northern Arabian–Nubian shield: The role of lithospheric mantle delamination and subsequent thermal subsidence. *Tectonophysics* 477(3–4):217–228
4. Avigad D, Stern RJ, Beyth M, Miller N (2007) Detrital zircon U–Pb geochronology of Cryogenian diamictites and Lower Paleozoic sandstone in Ethiopia (Tigray): age constraints on Neoproterozoic glaciation and crustal evolution of the southern Arabian Nubian Shield. *Precambrian Res* 154:88–106
5. Berhe S (1990) Ophiolites in Northeast and East Africa: implications for Proterozoic crustal growth. *Journal of Geological Society of London* 147:647–657
6. Beyth M, Avigad D, Wetzel H, Matthews A, Berhe S (2003) Crustal Exhumation and indications for snowball Earth in the East African Orogen: north Ethiopia and east Eritrea. *Precambr Res* 123:187–201
7. Billay et al (1997) The Geology of the Lega Dembi gold deposit, southern Ethiopia: implications for Pan-African Gold exploration. *Miner Deposita* 32:491–504
8. Birru T, Zhdanov EP (1991) Geological prospecting and exploration for primary gold and other minerals in Agere Mariam, Arero and Bulbul areas (southern Ethiopia), Ethiopian Institute of Geological Survey, Addis Ababa, Ethiopia, v. 2, PP. 1-368
9. Blasband B, White S, Brooijmans P, Boorder HD, Visser W (2000) Late Proterozoic extensional collapse in the Arabian–Nubian Shield, *Journal of the Geological Society, London*, 157: 615–628
10. Boynton, 1984, Nakamura N, 1974. Determination of REE, Ba, Fe, Mg, Na and K in carbonaceous and ordinary chondrites. *Geochemical et Cosmochimica Acta*, 38, 757–775
11. Best MG (2003) *Igneous and Metamorphic petrology*, 2nd edition. Blackwell Science Ltd, a Blackwell Publishing company
12. Chater AM (1971) The geology of the Megado region of Southern Ethiopia. Ph.D. thesis, Department of Earth Science, University of Leeds, United Kingdom, PP. 1–33
13. Daniest Mining PLC (2015) Reconnaissance geological and geochemical report on surupa gold exploration. Southern Ethiopia

14. Davidson A (1983) Reconnaissance geology and geochemistry of parts of Illubabor, Kefa, Gemu Gofa, and Sidamo, Ethiopia. The Omo river project, No. 2. Ethiopian Institute of Geological Surveys, Addis Ababa, Bull, pp 1–129
15. Ethiopian Meteorological, Agency (2012) Recorded Rainfall and Temperature Data of Yabelo Districts in Borana Rangeland; Ethiopian Meteorological Agency. Addis Ababa, Ethiopia, Unpublished data
16. Fritz H, Abdelsalam M, Ali KA, Bingen B, Collins AS, Fowler AR, Hauzenberger CA, Johnson PR, Kusky TM, Macey P, Muhongo S, Stern RJ, Viola G (2013) Orogen styles in the East African Orogen: A review of the Neoproterozoic to Cambrian tectonic evolution. *Joun Afr Earth Sc* 86:65–106
17. Garrett R, Nichol I (1969) Factor analysis in the interpretation of regional geochemical stream sediment data. *QColoSchMine* 64:245–264
18. Getaneh Assefa, Giovanni M, Paola DJ, Roberto Valer (1981) Plate Tectonics and Metallogenic processes in Ethiopia (preliminary report). Unpublished report. Ethiopian Institute of Geological Surveys, Ethiopia, pp 861–867
19. Genzebu W, Hasse N, Yemane T (1997) Geology of Agere Mariam map sheet, Memoir 8, Geological Survey of Ethiopia, pp. 1-118
20. Genzebu W, Hassen N, Yemane T (1994) Geology of the Ageremariam area (NB37-10). Ethiopian Institute of Geological Surveys, Addis Ababa, (Memoir 8), pp. 1-112
21. George H, Bonham-Carter GF (1989) Spatial modeling of geological data for gold exploration, Star Lake area, Saskatchewan. In: F. P. Afterberg and G.F. Bonham-Carter (Eds.) *Statistical Applications in the Earth Sciences*. Geological Survey of Canada Open File Report 93–392, 23 pp
22. Gilboy AF (1970) The geology of the Gariboro region of southern Ethiopia. Ph.D. thesis, Department of Earth Science, University of Leeds, United Kingdom, pp. 1-176
23. Hussein B (1999) The geology, structure and geochemistry of crystalline rocks of the Moyale area, Southern Ethiopia: Implications for the tectogenesis of Precambrian basement, Band 50, Tubinger, Germany
24. Irvine TN, Baragar WRA (1971) A guide to the chemical classification of the common volcanic rocks. *Canadian Journal of Earth Science* 8:523–548
25. Jelenc AD (1966) Mineral occurrences of Ethiopia, Ministry of Mines, Addis Ababa, pp. 1-720
26. Johnson P, Woldehaimanot B (2003) Development of the Arabian-Nubian Shield: perspectives on accretion and deformation in the northern East African Orogen and the assembly of Gondwana. *Supercontinent Assembly and breakup*. Geological Society of London, Special Publication, 206, 289–325
27. Johnson P, Andresen A, Collins A, Fowler H, Fritz W, Ghebreab T, Kusky R, Stern (2011) Late Cryogenian–Ediacaran history of the Arabian Nubian Shield: A review of depositional, plutonic, structural, and tectonic events in the closing stages of the northern East African Orogeny. *J Afr Earth Sc* 61:167–232
28. Kazmin V (1971) Precambrian of Ethiopia *Nature* 230:176–177

29. Kazmin V (1972) Geology of Ethiopia, unpublished report. Ethiopian Institute of Geological Surveys, Addis Ababa
30. Kazmin V, Shiferaw A, Balcha T (1978) The Ethiopian Basement: Stratigraphy and possible manner of evolution. *Geol Rundsch* 67:531–546
31. Kennedy WQ (1964) The structural differentiation of Africa in the Pan-African ($\pm$ 500 m.y.) Tectonic Episode. *Leeds Res Inst African Geol Ann Report* 8:48–49
32. Kröner A, Stern RJ (2004) Pan African Orogeny. *Encyclopedia of Geology* 1:1–12
33. Kozyrev V, Kebede G, Safonov Y, W/Micheal, Gurbanovich B, Tekle G, Kaitukov T, Arijapov A (1985) Regional Geological and exploration work for gold and other minerals in the Adola gold fields. *Ethiopian Mineral Exploration Development Corporation Bulletin*, Vol. 2, Addis Ababa, pp. 1-260
34. Le Bas M, Le Maitre R, Streckeisen A, Zanettin B (1986) Chemical classification of volcanic rocks based on the total alkali-silica diagram. *J Petrol* 27:745–750
35. Meert J (2003) A Synopsis of Events related the Assembly of Eastern Gondwana. *Tectonophysics* 362:1–40
36. Mengesha Tefera., Chernet T and Workineh Haro.(1996). Explanation of the Geological map of Ethiopia. Ethiopian Institute of Geological Survey. Note No. 621-101-122
37. Ministry of Mines and Energy (2009). National report on mining to the United Nation Commission on sustainable development (UNCSD), New York. Unpublished technical report, MOME, Addis Ababa, Ethiopia
38. Miyashiro A (1974) Volcanic rock series in island arcs and active continental margins. *Am J Sci* 274:321–355
39. Pearce J, Cann J (1973) Tectonic setting of basic volcanic rocks determined using trace element analyses. *Earth Planetary Science Letters* 19:290–300
40. Rose A, Hawkes H, Webb J (1979) *Geochemical in Mineral Exploration*. (2ndEd.). Academic Press, London, 657pp
41. Rollinson HR (1993) *Using Geochemical Data: Evaluation, Presentation, and Interpretation*. Pearson Education limited, England
42. Shackleton RM (1993) Tectonics of the Mozambique Belt in East Africa. In: Prichard HM, Alabaster T, Harris NBW, Neary CR (eds) *Magmatic Processes and Plate Tectonics*, 76. Geological Society Special Publication, pp 345–362
43. Stein M, Goldstein SL (1996) From plume head to continental lithosphere in Arabian Nubian Shield. *Nature* 382:773 – 778
44. Stern RJ (1994) Neoproterozoic (900–550 Ma) arc assembly and continental collision in the East African Orogen: implications for consolidation of Gondwanaland. *Ann Rev Earth Planet Sci* 22:319–351
45. Stern RJ, Johnson PR, Kröner A, Yibas B (2004) Neoproterozoic Ophiolites of the Arabian-Nubian Shield. *Precambrian Geology* 13:95–128

46. Sinclair AJ (1974) Selection of threshold values in geochemical data using probability graphs. *JGeochemical Exploration* 3:238
47. Sun S, McDonough W (1989) Chemical and isotopic systematic of oceanic basalts: implications for mantle composition and processes: In *Sunders, A.D., and Norry, M.J. Magmatism in ocean basins*, Geological Society Special Publication, 42, 313–345
48. Tadesse S (2004) Genesis of the shear zone related gold vein mineralization of the Lega Dembi gold deposit, Adola gold field, Southern Ethiopia. *Gondwana Res* 7(2):481–488
49. Tenczer V, Hauzenberger C, Fritz H, Wallbrecher E, Muhongo S, Mogessie A, Hoinkes G, Loizenbauer J. & Bauernhofer A (2005) The metamorphic evolution of the Mozambique Belt in Central Tanzania: new petrological and geochronological data. *Geophysical Research Abstracts*, 7
50. Tsige L, Abdelsalam MG (2005) Neoproterozoic–Early Paleozoic gravitational tectonic collapse in the southern part of the Arabian-Nubian Shield: the Bulbul Belt of southern Ethiopia. *Precambrian Res* 138:297–318
51. Vail JR (1976) Outline of the geochronology and tectonic units of the basement complex of northeast Africa. *Proceeding of Royal Society of London, A* 350, 127–141
52. Warden AJ, Horkel AD (1984) The Geological Evolution of the NE-Branch of the Mozambique Belt (Kenya, Somalia, Ethiopia). *Mitt, österr. geol Ges* 77:161–184
53. Worku H (1996) Geodynamic development of the Adola Belt (southern Ethiopia) in the Neoproterozoic and its control on gold mineralization. Ph.D.thesis, Technical University of Berlin, Wissenschaftliche Schriftenreihe Geologie und Bergbau Band 5, pp. 1-156

Figures

Figure 1

Location map of the study area

Figure 2

Micro-picture of metagranite all under Crossed polarized Light (XPL) for different sample showing different minerals. Bt-biotite, Musc-muscovite, Qtz-quartz, Plg-plagioclase, K-spar-K-feldspar, Grt-garnet and Opq-Opaque.

Figure 3

SiO_2 (wt. %) Vs. $\text{Na}_2\text{O} + \text{K}_2\text{O}$ (wt. %) on the (Cox et al, 1979) diagram were plotted to discriminate the nomenclature of normal igneous rocks.

Figure 4

Histograms of stream sediments in Surupa area (a-f). All elements show positive skewness. All elements are reported in ppm.

Figure 5

Geochemical zonation maps of elements in stream sediments of the study area

Figure 6

Harker variation diagram for selected major elements as a function of SiO₂ wt. %.

Figure 7

Discrimination diagram (A) The AFM triangular variation diagram for Surupa rocks; A-(alkalis :($\text{Na}_2\text{O} + \text{K}_2\text{O}$), F-($\text{FeO} + \text{Fe}$), and M-(MgO), (B and C) SiO_2 vs. $\text{Fe}_2\text{O}_3/\text{MgO}$ and K_2O of the high K Calc-alkaline rock series further into low-K tholeiitic series of the study area.

Spider plot – REE chondrite (Boynton 1984)

Figure 8

Chondrite normalized rare earth element patterns (Boynton 1984)

Figure 9

Primitive mantle normalized spider diagram (Sun and McDonough (1989))

Figure 10

A map showing mineral potential of gold and base metals stream sediment sample result.