Usefulness of Lung Ultrasound in Older Patients with Asymptomatic COVID-19 Pneumonia: Bicentric 3-month Follow-up Study

Chukwuma Okoye

University of Pisa

Valeria Calsolaro

Azienda Ospedaliera Universitaria Pisana

Alessandra Fabbri

Azienda Ospedaliera Universitaria Pisana

Riccardo Franchi

Azienda Ospedaliera Universitaria Pisana

Rachele Antognoli

Azienda Ospedaliera Universitaria Pisana

Ludovica Zisca

Presidio Ospedaliero "Anna Torrigiani"

Camilla Bianchi

Presidio Ospedaliero "Anna Torrigiani"

Alessia Maria Calabrese

University of Pisa

Sara Rogani

University of Pisa

Fabio Monzani (✉ fabio.monzani@med.unipi.it)

University of Pisa

Research Article

Keywords: COVID-19, lung ultrasound, symptoms, follow-up, oldest

Posted Date: July 13th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-625269/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

BACKGROUND: Clinical and prognostic differences between symptomatic and asymptomatic older patients with COVID-19 are of great interest since frail patients often show atypical presentation of illness. Lung Ultrasound (LUS) has been proven to be a reliable tool for detecting early-phase COVID-19 pulmonary alterations. The current study aimed to compare LUS score and 3-month overall mortality between asymptomatic and symptomatic older patients with COVID-19, according to frailty status.

METHODS: Bicentric prospective study on older patients with asymptomatic and symptomatic COVID-19 disease. Patients were stratified according to LUS score tertiles and Clinical Frailty Scale categories. Survival rate was assessed by telephone interviews 3 months after discharge.

RESULTS: 64 symptomatic (24 women, aged 80.0 ± 10.8 years) and 46 asymptomatic (31 women; aged 84.3 ± 8.8 years) were consecutively enrolled. LUS score resulted an independent predictor of 3-month mortality [OR 2.27 (CI 95%: 1.09–4.8), $p=0.03$], and the highest mortality rate was observed in symptomatic and asymptomatic pre-frail and frail patients (70.6% and 66.7%, respectively) with greater LUS abnormalities (3rd tertile).

CONCLUSIONS: LUS identified an acute interstitial lung involvement in most of the older asymptomatic patients. Mortality rate progressively increased according to clinical frailty and LUS score degree, resulting a reliable prognostic tool in both symptomatic and asymptomatic patients.

Background

Severe Acute Respiratory Syndrome caused by Coronavirus, etiological agent of COVID-19 associated pneumonia, has been identified all over the world since, in December 2019, SARS-CoV-2 has been isolated in China for the first time(1). The clinical spectrum of SARS-CoV-2 ranges from asymptomatic infection to a multiorgan inflammatory disease that can degenerate in ARDS (acute respiratory distress syndrome). A major challenge to containing the spread of SARS-CoV-2 is that asymptomatic and pre-symptomatic individuals are infectious, and up to 50% of cases may be attributable to transmission from asymptomatic or pre-symptomatic people(2). To date, few studies have sought at comparing clinical characteristics and pulmonary findings between asymptomatic and symptomatic patients(3,4); a recent meta-analysis proved that more than half of the individuals with SARS-CoV-2 infection without any symptoms presented with CT abnormalities; however, the mean age was low (31.0 ± 23.8 years), and few cases of asymptomatic older patients were described(5). In particular, clinical status of older patients with asymptomatic COVID-19 is of great interest, since frail patients often show non-typical COVID-19 clinical presentation; in a retrospective study by Bianchetti et al(6) regarding older patients with dementia, classical COVID-19 symptoms as dyspnoea or fever were recorded at ED admission in less than one half inpatients. Noteworthy, frail patients are the most at risk for poor outcome, facing a three times higher risk of in-hospital mortality compared to non-frail counterparts(7). The Office for National Statistics (ONS) in England and Wales reported a 79% increase in total deaths in care homes from 2 March to 12 June compared to 2015–2019, higher in nursing homes than residential homes(8). Moreover, a recent study by Rutten et al.(9) reported that, despite symptomatology overlapped between nursing-home residents with and without COVID-19, those with COVID-19 were 3 times more likely to die within 30 days. For these reasons, early detection of the severity of pulmonary involvement is crucial in older patients with COVID-19 diagnosis, regardless their clinical presentation. The American College of Radiology recommends against the use of CT as a screening or initial imaging study to diagnose Covid-19(10); whereas lung ultrasonography has been proven to represent a valid alternative in the early staging of the disease(11). LUS has proven to exhibit a good concordance with HRCT(12) by evaluating the degree of pulmonary involvement via the amount of focal or confluent B-lines and the presence of thickened pleural line with spared areas. The aim of the current study is to evaluate lung ultrasound artefacts in older patients hospitalized for symptomatic SARS-CoV-2 pneumonia, as compared to asymptomatic subjects with positive COVID-19 nasopharyngeal swab (NS). Secondary endpoint is to assess the possible relationship between LUS score and short-term mortality rate in symptomatic and asymptomatic patients.

Methods

We enrolled all the patients older than 65 years consecutively admitted to the Geriatrics Unit of our tertiary care hospital from April 10th to December 12th 2020, with severe SARS-CoV-2 pneumonia, as defined by NIH criteria(13). Thus, all the patients had SpO₂ below 94% on room air at sea level, respiratory rate above 30 breaths/min, PaO₂/FiO₂ ratio (i.e., partial pressure arterial oxygen/fraction of inspired oxygen ratio) below 300 mmHg, or over 50% lung infiltrates. Exclusion criteria were: *i*) critical illness requiring invasive ventilation at admission, *ii*) presence of concomitant pulmonary oedema at admission, *iii*) pre-existing interstitial diseases. All the symptomatic patients underwent HRCT immediately before ward admission, as part of the Emergency Department (ED) evaluation route of SARS-CoV-2 pneumonia. Bedside LUS was independently performed by two skilled clinicians (C.O., D.G.) at admission (within 12 hrs), blinded at HRCT results.

Furthermore, we enrolled asymptomatic, older patients admitted to a COVID-19 low care unit from May 18th to December 12th, 2020. All the subjects were previously nursing home residents, temporarily transferred for observation after having two consecutive PCR nasopharyngeal swabs positive for SARS-CoV-2 infection, with a PCR cycle threshold (Ct) value ≤ 30 . In this setting, all the patients met the NIH criteria for asymptomatic/pre-symptomatic SARS-CoV-2 infection¹³; hence, patients showing any respiratory symptoms, fever, or peripheral O₂ saturation below 94% in air room during the hospital stay, were excluded from the final analysis. LUS was performed by two skilled clinicians (C.O., L.Z.) within the 10th to 14th day after positive NS. At ward admission, the presence of positive history of chronic obstructive pulmonary disease, chronic heart failure, diabetes and arterial hypertension was recorded. Comprehensive geriatric assessment (CGA) was performed using the following scales: Activities of Daily Living (ADL)(14), Instrumental ADL(15), Short Portable Mental Status Questionnaire (SPMSQ)(16), Cumulative Illness Rating Scale (CIRS)(17). The presence of frailty was defined according to the Rockwood Clinical Frailty Scale(18) as follows: patients with score 1–3 were categorized as “Robust”; those with score 4–5 as “Pre-Frail”, and those with score 6–9 as “Frail”.

In both the study settings, a convex and a linear covered probe, 3.5 to 7.5 MHz (Esaote Medical System), were used for chest ultrasound examination. For patients with severe mobility limitations, two operators were concomitantly involved, according to current guidelines(19). As for the scanning scheme, in the

absence of a standardized score for COVID-19 patients, we used a previously validated score(20), namely LUS score for monitoring aeration, ranging from 0 to 3 points in each of the 16 scanned zones. Score 0: predominant A-lines or less than 3 separated B-lines. Score 1: at least 3 B-lines or coalescent B-lines occupying $\leq 50\%$ of the screen without a clearly irregular pleural line. Score 1p: at least 3 B-lines or coalescent B-lines occupying $\leq 50\%$ of the screen with a clearly irregular pleural line. Score 2: coalescent B-lines occupying $> 50\%$ of the screen without a clearly irregular pleural line. Score 2p: coalescent B-lines occupying $> 50\%$ of the screen with a clearly irregular pleural line. Score 3: large consolidations (at least > 1 cm). The final score is obtained by summing up the scores of each area and dividing the sum for the total of zone recorded (Supplemental Fig. 3). All the patients (symptomatic and asymptomatic) were categorized according to LUS score tertiles as follows: T1 (Tertile 1) = LUS score ≤ 0.5 , T2 (Tertile 2) = LUS score $0.6-1.4$; T3 (Tertile 3) = LUS Score ≥ 1.5 . Sensitivity analysis comparing ultrasonographic findings using HRCT as gold standard was performed to confirm LUS accuracy in symptomatic patients.

Each patient of the two cohorts gave written informed consent to participate in the study; in case of patient's inability, the legally authorized delegate provided informed consent. The study protocol complied with the Declaration of Helsinki and was approved by the local Ethic Committee (n° protocol: CEAVNO-2020-17241). Three-month mortality rate of both symptomatic and asymptomatic patients was assessed by telephone interview.

Statistical Analysis

Statistical analysis was performed by using SPSS 27.0 statistical software package (SPSS Inc., Chicago, IL) and GraphPad Prism 9. Continuous variables were expressed as mean $\pm$ standard deviation or, for non-normally skewed distributions, as median and interquartile (IQR), and categorical data were expressed as percentage. Mann-Whitney U-test and χ^2 exact tests were used for multiple comparisons. LUS score tertiles were considered as 3-ordinal-variables, and a Cochran-Armitage test was used for multiple comparison for the mortality outcome, utilizing a p for trend. LUS diagnostic performance was assessed by calculating specificity, sensitivity, negative predictive value (NPV) and positive predictive value (PPV), using HRCT as gold standard. Proportion of agreement (P_0) was utilized to represent the concordance between HRCT and LUS. In order to evaluate the association between LUS score and 3-months all-cause mortality, a logistic univariate regression analysis was carried out, using LUS score as continuous numerical dependent variable. Multivariate logistic regression analysis was also calculated after adjusting for age, sex, history of heart failure (HF), Chronic obstructive pulmonary disease (COPD) and obesity. Statistical significance was considered for $p < 0.05$.

Results

A total of 122 older patients with SARS-CoV-2 infection were initially enrolled; 12 of them were subsequently excluded as described in Fig. 1. Thus, 110 patients (mean age 82.3 ± 10.1 years) were finally included into the study: 64 (59.2%) with symptomatic SARS-CoV-2 pneumonia from the acute Geriatrics ward setting and 46 (41.8%) asymptomatic subjects from the COVID-19 low-care setting. Baseline characteristics of the study population are detailed in Table 1.

Table 1
Clinical characteristics of symptomatic and asymptomatic older patients with SARS-CoV-2 Infection.

	Whole cohort N = 110	Symptomatic N = 64	Asymptomatic N = 46	<i>p</i>
Gender (F%)	55 (50.0)	24 (37.5)	31 (67.4)	<0.01
Age (years)	82.3 ± 10.2	80.8 ± 10.8	84.3 ± 8.8	0.04
ADL (median, IQR)	4 (0–6)	6 (2–6)	2 (0–6)	<0.01
SPMSQ (median, IQR)	2 (0–8)	1 (0–5)	4 (0–10)	0.06
CIRS- c (mean ± SD)	1.80 ± 1.5	2.4 ± 1.4	1.4 ± 0.9	<0.001
Clinical Frailty Scale				<0.001
- Frail (%)	43 (39.1)	22 (34.4)	21 (45.6)	
- Pre-Frail (%)	24 (21.8)	7 (10.9)	17 (37.0)	
- Robust (%)	43 (39.1)	35 (54.7)	8 (17.4)	
Chronic respiratory diseases (%)	16 (31.4)	7 (26.9)	9 (36)	0.48
Chronic Heart Failure (%)	19 (37.2)	10 (40)	9 (34.6)	0.06
Arterial Hypertension (%)	32 (62.5)	18 (69.2)	14 (56)	0.18
Diabetes Mellitus (%)	10 (19.7)	7 (28)	3 (11.5)	0.17
Obesity (%)	10 (19.7)	6 (24)	4 (15.3)	0.39
WBC/mm ³ (median, IQR)	7043(4000)	6710(3810)	5180(4170)	0.655
Lymphocyte count/mm ³ (median, IQR)	1390(780)	995(692)	1610(2375)	0.101
Hs-CRP mg/dl (median, IQR)	6.44(8.20)	6.09 (10.3)	2.10 (3.70)	<0.001
Data are expressed as mean ± standard deviation, median (interquartile range) and number (%) as appropriate. Significant <i>p</i> values are marked in bold. ADL: Activities of Daily Living, CIRS-C: Cumulative Illness Rating Scale-Comorbidity, SPMSQ: Short Portable Mental Status Questionnaire. WBC: White blood count. Hs-CRP: High-sensitivity C-reactive protein.				

At LUS examination, at least three B-lines artefacts (78.3%) and irregular thickened pleural line with spared areas (73.6%) resulted the predominant pattern of abnormalities observed (Fig. 2). B-lines were commonly observed (88.9%), usually bilateral (78.8%), in symptomatic patients and, although less frequently (67.7%, *p* < 0.01 vs symptomatic), with a lower bilateral distribution (34.8%, *p* < 0.001 vs symptomatic) in asymptomatic patients. Pleural effusion was found in 20 (31.1%) symptomatic and 9 (18.7%) asymptomatic patients (*p* = 0.005).

Overall, mean LUS score resulted significantly higher in symptomatic than asymptomatic patients (1.34 ± 0.69 vs 0.60 ± 0.50, *p* < 0.001) (Fig. 3). By sensitivity analysis, B-lines artefacts were found in 54 out of 57 (95.1%) patients with ground glass opacities at HRCT (*P*_o = 95%) with 100% sensitivity (95%CI: 92.4–100%) and 40% specificity (95%CI: 0.1–82.9); PPV 95.1% (95%CI: 84.6–98.8) and NPV 100%, yielding a diagnostic accuracy of 95.8% (95%CI: 78.9–99.9). All the cases of pulmonary consolidations and pleural effusion detected by LUS were confirmed at HRCT examination (*P*_o=100%).

Survival analysis

Nine symptomatic patients died during hospital stay; further 14 symptomatic and 11 asymptomatic patients died at 3-month follow-up. Thus, the overall mortality rate was 35.9% in symptomatic and 23.9% in asymptomatic patients (*p* = 0.17). Moving from the lowest to the highest LUS score tertile, a significant, progressive increase of mortality rate was observed in the whole cohort of patients (15.4%, 28.6% and 50.0%, respectively; *p* for trend = 0.001). The mortality trend remained significant among asymptomatic (11.5%, 35.3% and 66.7%, respectively; *p* for trend = 0.013) but not symptomatic patients (23.1%, 22.2% and 48.5%, respectively; *p* for trend = 0.057) (Fig. 4).

According to CFS, pre-frail and frail patients showed an increased mortality typically in the highest tertile of LUS score compared to T1 and T2, both in asymptomatic (66.7% vs 36.3% and 30%, respectively) and symptomatic patients (70.6% vs 25.0% and 28.6%), as shown in Supplemental Table 1. Furthermore, LUS score emerged as an independent predictor of 3-months all-cause mortality at multivariate analysis after adjusting for age, sex, BMI, diabetes, heart failure and COPD (unadjusted OR = 2.29, CI95%: 1.11–4.9, *p* = 0.028, adjusted OR = 2.27, CI95%: 1.09–4.8, *p* = 0.030).

Discussion

In our study, more than two-thirds of asymptomatic older patients with SARS-CoV-2 infection showed signs of pulmonary involvement evaluated by lung ultrasound. Asymptomatic patients were more likely female, with a lower burden of comorbidity, usually showing unilateral interstitial involvement. As expected, overall mortality rate of asymptomatic patients was lower than symptomatic patients but, moving from the lowest to the highest LUS score tertile, their mortality rate showed a significant, progressive increase, typically in frail and pre-frail patients. Although older individuals suffering from COVID-19

represent the prototype of patients most at risk for adverse outcomes, to our knowledge, no previous studies have attempted to assess LUS artefacts and their relationship with a mid-term mortality risk in asymptomatic oldest patients.

It is well-established that a proportion of patients with COVID-19 without symptoms show HRCT signs of acute pneumonia; however, less is known about the prognostic impact of these findings in older frail patients, since they usually face disproportionate non-typical symptoms and an increase excess mortality after hospital discharge which is not completely understood(8)(9). In the current study we hypothesized a crucial role of asymptomatic pneumonia evaluated with LUS, in the mid-term course of the disease in this particular class of frailest individuals with SARS-CoV-2 infection.

Lung ultrasound has been proved to be an important tool for interstitial lung evaluation of COVID-19 pneumonia and in the follow-up of the disease(21). In COVID-19, we typically see various grades of multiple B-lines with patchy distribution. B-lines can be separated or coalescent, including pictures of sonographic 'white lung'(22). In agreement with Volpicelli et al(11), in our study at least three B lines and a fragmented pleural line were the most frequent ultrasonographic findings, both in asymptomatic and symptomatic patients. In our study, asymptomatic patients were more likely to present unilateral involvement, as first described by Shi et al.(3), who found unilateral, multifocal ground glass opacities at Chest CT. This finding is particularly of interest in the low care hospitals and nursing homes, where an accurate risk stratification is crucial to decide which older patients need to be transferred to a hospital or could be treated in the same setting. As expected, we found a significantly higher LUS score of aeration in symptomatic patients compared to asymptomatic pairs. In other words, patients with respiratory symptoms, showed a higher number of diffuse, bilateral B-lines compared to the asymptomatic ones, thus confirming an increased interstitial pulmonary damage.

In the current study we found a high prevalence of comorbidities, especially cardiovascular and chronic respiratory diseases. Consistent with previous studies(23), patients with severe COVID-19 manifestations showed a higher prevalence of cardiovascular disorders, compared to asymptomatic, although not reaching the statistical significance. Somewhat surprisingly, asymptomatic patients were older and frailer than those symptomatic; however, this result needs to be contextualized in the clinical setting where the asymptomatic patients came from. Indeed, asymptomatic patients were residents of a nursing home temporarily transferred to a COVID-19 low care unit for clinical observation, while symptomatic patients were hospitalized in an acute care geriatrics unit. Although the prognostic role of multimorbidity and frailty is still debated(24), as a fact, older people living in nursing homes are commonly oldest with various degree of disability, needing assistance in basic activities of daily living.

Given that older individuals may present non-specific LUS pattern, such as localized B lines or pleural thickening due to pre-existing fibrosis(25)(26), we performed a concordance analysis by estimating the correlation of LUS findings and HRCT features in the acute geriatric ward. As expected, we found a high proportion of agreement in terms of B-lines, pulmonary consolidations and pleural effusions, consistent with a recent paper by Nouvenne et al(12), in which the significant correlation between HRCT and LUS findings in COVID-19 was firstly reported.

During the three-month post-discharge follow-up, fourteen symptomatic and eleven asymptomatic patients deceased, yielding a worrying residual mortality of 24%, significantly higher than previous reports(27)(28), where a 2–5% mortality was recorded. However, in our study we aimed at focusing on the outcomes in the oldest individuals and patients severely disabled living in nursing homes, thus explaining this substantial difference. Noteworthy, after categorizing LUS score by tertiles, patients with the highest LUS score had two and three-times higher mortality compared to those of the lowest tertile, in symptomatic and asymptomatic patients, respectively. This result is not surprising since various studies have reported the post COVID-19 progression to lung fibrosis in older patients(29)(30), and a recent study by Somani et al(28), showed that the most common cause for re-admission at hospital after COVID-19 was respiratory distress, typically in comorbid patients.

Asymptomatic patients reported a lower overall mortality, facing most often a reduced interstitial damage detected by LUS score and decreased levels of inflammatory markers compared to symptomatic counterparts. Indeed, the overall lower mortality of asymptomatic patients largely depends on the fact that the majority were in the lowest tertile with half mortality than those with symptoms. On the other hand, mortality risk of asymptomatic patients overcame that of symptomatic in both the second and third LUS score tertiles (35.3% vs 22.2% and 66.7% vs 48.5%, respectively). One of the most interesting results emerging from our study is the demonstration of the complex interaction between frailty and morpho-functional pulmonary impairment in COVID-19 patients, regardless their clinical presentation. Hence, we recorded a higher mortality in pre-frail and frail patients in both symptomatic and asymptomatic patients as compared to robust individuals; this result is in line with a recent multi-centre cohort study reporting an increased likelihood of adverse outcomes in patients with CFS score from 4 to 9(31). Interestingly, we found that a bilateral diffuse interstitial damage may act as a mortality-multiplying risk factor in frail patients, doubling the 3-months mortality moving from the lowest to the higher LUS score tertile.

In conclusion, the current study offers some important insight into the clinical status of asymptomatic COVID-19 older patients, highlighting the prognostic role of LUS evaluation in these individuals, regardless their clinical presentation. Given that frail patients often show non-typical COVID-symptoms and are the most at risk for poor outcomes, the utilization of LUS could guide the clinician in the decision-making process and direct an early-stage COVID-19 therapeutic management. However, the generalization of these results is subject to certain limitations. Firstly, as previously mentioned, due to a small sample size, the findings might not be conclusive; therefore, further prospective analyses in larger cohorts of older patients are needed to confirm these results in asymptomatic patients. Secondly, the study population accounts for two different clinical settings and, some significant differences between the two populations (i.e., age and ADL score) might be biased. Thirdly, in agreement with the current guidelines, we did not perform HRCT to asymptomatic patients; therefore, despite the exclusion of patients with a known history of chronic interstitial disease or pulmonary fibrosis, we cannot exclude an overdiagnosis of COVID-19 pneumonia in asymptomatic subjects, being the B-lines an indirect, non-specific sign of interstitial involvement. Notwithstanding, in the concordance analysis carried out in symptomatic patients, we confirmed a high proportion of agreement between the two techniques.

Conclusions

The current study targets on the usefulness of a LUS-based follow-up in the frailest older patients with a recent SARS-CoV2 infection regardless symptoms or clinical presentation, and it brings attention on possible drawbacks of the traditional classification of “asymptomatic or pre-symptomatic SARS-CoV2 infection” in the advanced age. Indeed, signs of acute interstitial injury were detected by lung ultrasound in more than two-thirds of older asymptomatic patients. Moreover, mortality rate progressively raised with the increasing degree of interstitial pulmonary involvement, especially in pre-frail and frail patients. These findings raise a warning regarding the actual clinical status of asymptomatic older subjects and, suggest bed side ultrasound lung examination as an effective tool to guide clinical workup and patients' care.

Declarations

Ethical approval and consent to participate: Each patient gave written informed consent to participate in the study; in case of patient's inability, the legally authorized delegate provided informed consent. The study protocol complied with the Declaration of Helsinki and was approved by the Tuscany Regional Ethical Committee for Clinical Spérimentation (n° protocol: CEAVNO-2020-17241).

Availability of data and materials: The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request

Competing interests: The authors declare that they have no competing interests

Funding sources: This research did not receive any funding from agencies in the public, commercial, or not-for-profit sectors.

Authors' contributions: CO and FM performed the study concept and desing. RA, SR, AC, RF, CB were contributor for acquisition of data. AF, CO, LZ performed the lung ultrasound examination. VC, AC performed the analysis and interpretation of data. CO, FM, VC were responsible for the drafting of the manuscript. FM, VC provided the critical revision of the manuscript for important intellectual content. All authors read and approved the final manuscript.

Acknowledgments: The authors wish to thank Dr. Pasquale Morano for assistance in the definition of the study protocol, all the nurses and staff of the Presidio Ospedaliero “Anna Torrigiani” for the help given through the project. The authors wish also to thank all patients who participated to the study, Dr. Tommaso Riccioni, Dr. Lucia Visaggi, Dr. Daniela Guarino and Dr. Antonio Polini for the important contribution in the study. All the colleagues gave the permission to be named.

References

1. Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med* [Internet]. 2020 Feb 20 [cited 2020 Dec 2];382(8):727–33. Available from: <https://www.nejm.org/doi/full/10.1056/NEJMoa2001017>
2. Kimball A, Hatfield KM, Arons M, James A, Taylor J, Spicer K, et al. Asymptomatic and Presymptomatic SARS-CoV-2 Infections in Residents of a Long-Term Care Skilled Nursing Facility — King County, Washington, March 2020. *MMWR Morb Mortal Wkly Rep* [Internet]. 2020 Apr 3 [cited 2020 Dec 2];69(13):377–81. Available from: http://www.cdc.gov/mmwr/volumes/69/wr/mm6913e1.htm?s_cid=mm6913e1_w
3. Shi H, Han X, Jiang N, Cao Y, Alwalid O, Gu J, et al. Radiological findings from 81 patients with COVID-19 pneumonia in Wuhan, China: a descriptive study. *Lancet Infect Dis* [Internet]. 2020 Apr 1 [cited 2020 Dec 2];20(4):425–34. Available from: <https://pubmed.ncbi.nlm.nih.gov/32105637/>
4. Bandirali M, Sconfienza LM, Serra R, Brembilla R, Albano D, Pregliasco FE, et al. Chest radiograph findings in asymptomatic and minimally symptomatic quarantined patients in codogno, Italy during COVID-19 pandemic [Internet]. Vol. 295, *Radiology*. Radiological Society of North America Inc.; 2020 [cited 2020 Dec 2]. Available from: <https://pubs.rsna.org/doi/abs/10.1148/radiol.2020201102>
5. Kronbichler A, Kresse D, Yoon S, Lee KH, Effenberger M, Shin J II. Asymptomatic patients as a source of COVID-19 infections: A systematic review and meta-analysis. *Int J Infect Dis* [Internet]. 2020 Sep 1 [cited 2020 Dec 2];98:180–6. Available from: <https://doi.org/10.1016/j.ijid.2020.06.052>
6. Bianchetti A, Rozzini R, Guerini F, Boffelli S, Ranieri P, Minelli G, et al. Clinical Presentation of COVID19 in Dementia Patients. *J Nutr Heal Aging* [Internet]. 2020 Jun 1 [cited 2021 May 10];24(6):560–2. Available from: <https://pubmed.ncbi.nlm.nih.gov/33256958/>
7. Blomaard LC, van der Linden CMJ, van der Bol JM, Jansen SWM, Polinder-Bos HA, Willems HC, et al. Frailty is associated with in-hospital mortality in older hospitalised COVID-19 patients in the Netherlands: the COVID-OLD study. *Age Ageing* [Internet]. 2021 Jan 30 [cited 2021 May 14];50(3). Available from: <https://pubmed.ncbi.nlm.nih.gov/33951156/>
8. Deaths registered weekly in England and Wales, provisional - Office for National Statistics [Internet]. [cited 2021 May 10]. Available from: <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/bulletins/deathsregisteredweeklyinenglandandwalesprovisio>
9. Rutten JJS, van Loon AM, van Kooten J, van Buul LW, Joling KJ, Smalbrugge M, et al. Clinical Suspicion of COVID-19 in Nursing Home Residents: Symptoms and Mortality Risk Factors. *J Am Med Dir Assoc* [Internet]. 2020 Dec 1 [cited 2021 May 10];21(12):1791-1797.e1. Available from: <https://pubmed.ncbi.nlm.nih.gov/33256958/>
10. American College of Radiology. ACR recommendations for the use of chest radiography and computed tomography (CT) for suspected COVID-19 infection. [Internet]. 2020. Available from: <https://www.acr.org/Advocacy-and-Economics/ACR-Position-Statements/Recommendations-for-Chest-Radiography-and-CT-for-Suspected-COVID19-Infection>.
11. Volpicelli G, Lamorte A, Villén T. What's new in lung ultrasound during the COVID-19 pandemic. *Intensive Care Med* [Internet]. 2020 Jul 1 [cited 2020 Dec 2];46(7):1445–8. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7196717/>
12. Nouvenne A, Zani MD, Milanese G, Parise A, Baciarello M, Bignami EG, et al. Lung Ultrasound in COVID-19 Pneumonia: Correlations with Chest CT on Hospital admission. *Respiration* [Internet]. 2020 Aug 1 [cited 2020 Dec 2];99(7):617–24. Available from: <https://www.karger.com/Article/FullText/509223>

13. COVID-19 Treatment Guidelines Panel. Coronavirus Disease 2019 (COVID-19) Treatment Guidelines. National Institutes of Health. Available at <https://www.covid19treatmentguidelines.nih.gov/>. Accessed 3/15/2021. 2021.
14. Katz S, Ford AB, Moskowitz RW, Jackson BA, Jaffe MW. Studies of Illness in the Aged: The Index of ADL: A Standardized Measure of Biological and Psychosocial Function. *JAMA* [Internet]. 1963 Sep 21;185(12):914–9. Available from: <https://doi.org/10.1001/jama.1963.03060120024016>
15. Lawton MP, Brody EM. Assessment of older people: self-maintaining and instrumental activities of daily living. *Gerontologist*. 1969;9(3):179–86.
16. Pfeiffer E. A short portable mental status questionnaire for the assessment of organic brain deficit in elderly patients. *J Am Geriatr Soc*. 1975 Oct;23(10):433–41.
17. Parmelee PA, Thuras PD, Katz IR, Lawton MP. Validation of the Cumulative Illness Rating Scale in a geriatric residential population. *J Am Geriatr Soc*. 1995 Feb;43(2):130–7.
18. Rockwood K, Song X, MacKnight C, Bergman H, Hogan DB, McDowell I, et al. A global clinical measure of fitness and frailty in elderly people. *C Can Med Assoc J = J l'Association medicale Can*. 2005 Aug;173(5):489–95.
19. Volpicelli G, Elbarbary M, Blaivas M, Lichtenstein DA, Mathis G, Kirkpatrick AW, et al. International evidence-based recommendations for point-of-care lung ultrasound. In: *Intensive Care Medicine* [Internet]. Intensive Care Med; 2012 [cited 2020 Dec 9]. p. 577–91. Available from: <https://pubmed.ncbi.nlm.nih.gov/22392031/>
20. Bouhemad B, Mongodi S, Via G, Rouquette I. Ultrasound for “lung monitoring” of ventilated patients [Internet]. Vol. 122, *Anesthesiology*. Lippincott Williams and Wilkins; 2015 [cited 2020 Dec 3]. p. 437–47. Available from: <http://links.lww.com/ALN/B117>,
21. Volpicelli G, Gargani L, Perlini S, Spinelli S, Barbieri G, Lanotte A, et al. Lung ultrasound for the early diagnosis of COVID-19 pneumonia: an international multicenter study on behalf of the International Multicenter Study Group on LUS in COVID-19. *Intensive Care Med* [Internet]. 2021 [cited 2021 May 14];47:444–54. Available from: <https://doi.org/10.1007/s00134-021-06373-7>
22. Gargani L, Soliman-Aboumarie H, Volpicelli G, Corradi F, Pastore MC, Cameli M. Why, when, and how to use lung ultrasound during the COVID-19 pandemic: enthusiasm and caution. *Eur Hear J - Cardiovasc Imaging* [Internet]. 2020 Sep 1 [cited 2020 Dec 3];21(9):941–8. Available from: <https://academic.oup.com/ehjcardimaging/article/21/9/941/5855021>
23. Suleyman G, Fadel RA, Malette KM, Hammond C, Abdulla H, Entz A, et al. Clinical Characteristics and Morbidity Associated With Coronavirus Disease 2019 in a Series of Patients in Metropolitan Detroit. *JAMA Netw open*. 2020 Jun;3(6):e2012270.
24. McCarthy K, Hewitt J, Myint PK, Quinn T, Carter B. Frailty and mortality in patients with COVID-19 – Authors’ reply. *Lancet Public Heal* [Internet]. 2020 Nov 1;5(11):e582. Available from: [https://doi.org/10.1016/S2468-2667\(20\)30224-3](https://doi.org/10.1016/S2468-2667(20)30224-3)
25. Ciccarese F, Chiesa AM, Feletti F, Vizioli L, Pasquali M, Forti P, et al. The Senile Lung as a Possible Source of Pitfalls on Chest Ultrasonography and Computed Tomography. *Respiration* [Internet]. 2015 Jul 25 [cited 2020 Dec 3];90(1):56–62. Available from: <https://www.karger.com/Article/FullText/430994>
26. Sperandeo M, De Cata A, Molinaro F, Trovato FM, Catalano D, Simeone A, et al. Ultrasound signs of pulmonary fibrosis in systemic sclerosis as timely indicators for chest computed tomography. *Scand J Rheumatol* [Internet]. 2015 Sep 3 [cited 2020 Dec 3];44(5):389–98. Available from: <https://pubmed.ncbi.nlm.nih.gov/26099251/>
27. Bellan M, Soddu D, Balbo PE, Baricich A, Zeppegno P, Avanzi GC, et al. Respiratory and Psychophysical Sequelae Among Patients With COVID-19 Four Months After Hospital Discharge. *JAMA Netw open* [Internet]. 2021 Jan 4 [cited 2021 Apr 1];4(1):e2036142. Available from: <https://jamanetwork.com/>
28. Somani S, Richter F, Fuster V, De Freitas J, Naik N, Sigel K, et al. Characterization of Patients Who Return to Hospital Following Discharge from Hospitalization For COVID-19. *medRxiv : the preprint server for health sciences*. 2020.
29. George PM, Wells AU, Jenkins RG. Pulmonary fibrosis and COVID-19: the potential role for antifibrotic therapy. *Lancet Respir Med* [Internet]. 2020/05/15. 2020 Aug;8(8):807–15. Available from: <https://pubmed.ncbi.nlm.nih.gov/32422178>
30. Zhang C, Wu Z, Li J-W, Tan K, Yang W, Zhao H, et al. Discharge may not be the end of treatment: Pay attention to pulmonary fibrosis caused by severe COVID-19. *J Med Virol*. 2021 Mar;93(3):1378–86.
31. Alsahab M, Beishon L, Brown B, Burn E, Burton JK, Cox N, et al. Age and frailty are independently associated with increased COVID-19 mortality and increased care needs in survivors: results of an international multi-centre study. *Age Ageing* [Internet]. 2021 Feb 5 [cited 2021 May 15];50(3):617–30. Available from: <https://academic.oup.com/ageing/article/50/3/617/6128535>

Figures

Figure 1

Flow chart of the observational cohort study.

Figure 2

Lung ultrasound features of symptomatic and asymptomatic older patients with SARS-CoV2-infection. ** = p-value <0.01 *** = p-value <0.001 ns= not significant

Figure 3

Distribution of Lung Ultrasound Score in symptomatic and asymptomatic older patients with SARS-CoV-2 infection. *** = p-value <0.0001

T1 = Tertile 1 (LUS Score ≤ 0.5); T2= Tertile 2 (LUS Score 0.6 - 1.4); T3= Tertile 3 (LUS Score ≥ 1.5)

p= p value for trend

Figure 4

Three-months all-cause-mortality of older patients with SARS-CoV2 infection according to LUS score tertiles ** = p-value <0.01

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementalFigure.S1.tif](#)
- [SupplementalFigure2.tif](#)
- [SupplementalTable1.docx](#)