Nitrogen Fixation and Diazotrophic Community in Plastic-Eating Mealworms *Tenebrio Molitor* L.

Yu Yang (✉ yooyoung@bit.edu.cn)

Beijing Institute of Technology <https://orcid.org/0000-0002-4663-9753>

Lin Hu

Beijing Institute of Technology

Xiaoxi Li

Beijing Institute of Technology

Jialei Wang

Beijing Institute of Technology

Guishan Jin

Beijing Research Institute of Uranium Geology

Research Article

Keywords: nitrogen fixation, plastic waste, biodegradation, mealworm, Klebsiella

Posted Date: June 18th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-627936/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Microbial Ecology on January 21st, 2022.
See the published version at <https://doi.org/10.1007/s00248-021-01930-5>.

Abstract

Mealworms, the larvae of a coleopteran insect *Tenebrio molitor* L., are capable of eating, living on and degrading the non-hydrolyzable vinyl plastics as sole diet. However, vinyl plastics are carbon-rich but nitrogen-deficient. It remains puzzling how plastic-eating mealworms overcome the nutritional obstacle of nitrogen limitation. Here, we provide the evidence for nitrogen-fixation activity within plastic-eating mealworms. Acetylene reduction assays illustrate that the nitrogen-fixing activity ranges from 12.3 ± 0.7 to 32.9 ± 9.3 nmol ethylene·h⁻¹·gut⁻¹ and the corresponding fixed nitrogen equivalents of protein are estimated as 8.6 to 23.0 µg per day per mealworm. Nature nitrogen isotopic analyses of plastic-eating mealworms provide further evidence for the importance of nitrogen fixation as a new nitrogen source. Eliminating the gut microbial microbiota with antibiotics impairs the mealworm's ability to fix nitrogen from atmosphere, indicating the contribution of gut microbiota to nitrogen fixation. By using the traditional culture-dependent technique, PCR and RT-PCR of *nifH* gene, nitrogen-fixing bacteria diversity within the gut was detected and the genus *Klebsiella* was demonstrated to be an important nitrogen-fixing symbiont. These findings first build the relationship between the plastic degradation (carbon metabolism) and nitrogen fixation (nitrogen metabolism) within mealworms. Combined with previously reported plastic-degrading capability and nitrogen-fixing activity, mealworms may be potential candidates for up-recycling of plastic waste to produce protein sources.

Introduction

An increasing accumulation of non-hydrolyzable vinyl plastic wastes, such as polystyrene (PS), polyethylene (PE), polypropylene (PP), and polyvinyl chloride (PVC), has become a severe environmental and social concern (Ru et al., 2020; Yang et al., 2020). In 2015, mealworm, the larva of a coleopteran insect *Tenebrio molitor* L., was first reported to be able to eat PS foam (Styrofoam) as sole diet and lived as well as those fed with a normal food (wheat bran) over a period of one month (Yang et al., 2015a,b). Ingested Styrofoam was demonstrated to be efficiently degraded after passage through the larval gut within a retention time of less than 24 h, and further mineralized to CO₂ and incorporated into lipids (Yang et al., 2015a). Subsequently, the ability of mealworm to live on and degrade Styrofoam was ubiquitously confirmed by expanding assessments from 22 countries (Yang et al., 2018). Additionally, multiple recent studies showed that mealworm was also capable of surviving by eating other synthetic vinyl plastics, such as PE, PVC, PP, and Styrene-Butadiene rubber (SBR), as sole diet, and rapidly degrading these ingested vinyl plastics within the larval gut (Brandon et al., 2018; Aboelkheir et al., 2019; Peng et al., 2020; Yang et al., 2020). These findings have provided implications for future applications of mealworms in plastic waste management. However, there is a controversy that vinyl plastics of PS, PE, PVC, PP and SBR are carbon-rich but nitrogen-deficient diet, which could not meet the nitrogen requirements for the biosynthesis of proteins, nucleic acids and other metabolites within mealworms. Therefore, it is intriguing to know how mealworms overcome the nutritional obstacle of nitrogen limitation when they are fed with vinyl plastic as sole diet.

Many insects, like termite, shipworm, stag beetle and red turpentine beetle, intrinsically feed on nitrogen-deficient wood diet (Bar-Shmuel et al., 2020). These xylophagous insects have been extensively demonstrated to be capable of supplementing their diets by fixing nitrogen from atmosphere based on the acetylene reduction assays or nitrogen isotopic tracers (Breznak et al., 1973; Benemann, 1973; Carpenter and Culliney, 1975; Prestwich and Bentley, 1981; Pandey et al., 1992; Tayasu et al., 1994; Behar et al., 2005; Kuranouchi et al., 2006; Yamada et al., 2006; Morales-Jiménez et al., 2009; Ayayee et al., 2014; Sapountzis et al., 2016). The nitrogen-fixing activity of xylophagous insects could be abolished by feeding with antibiotics, indicating that gut bacteria may contribute to nitrogen fixation (Breznak et al., 1973; Benemann, 1973; Carpenter and Culliney, 1975). These findings promote a search for insect gut bacteria that are involved in nitrogen fixation (Nardi et al., 2002; Kneip et al., 2007; Orón Tamayo and Heil, 2015). By applying the traditional culture-dependent technique of growing the bacteria on a nitrogen-free medium, a few cultures of nitrogen-fixing bacteria, identified as the species from genera of *Azotobacter*, *Spirochaeta*, *Citrobacter*, *Enterobacter*, *Treponema*, *Klebsiella*, *Beijerinckia*, *Clostridium* and *Raoultella*, have been isolated from the guts of various nitrogen-fixing insects (Carpenter and Culliney, 1975; French et al., 1976; Potrikus and Breznak, 1977; Bridges, 1981; Waterbury et al., 1983; Lilburn et al., 2001; Risal and Ozawa, 2001; Gomathi et al., 2005; Behar et al., 2005; Doolittle et al., 2008; Morales-Jiménez et al., 2013). Nevertheless, the culture-dependent method may detect only a limited number of diazotrophic species because of the inability to culture most of environmental isolates. As the gene *nifH* encoding dinitrogenase reductase is genetically conserved (Zehr et al., 2003), the culture-independent molecular techniques based on the amplification of *nifH* gene has been used to detected an unexpected diversity of nitrogen-fixing bacteria within guts of these nitrogen-fixing insects (Ohkuma et al., 1996, 1999; Yamada et al., 2007; Du et al., 2012; Desai et al., 2012; Ayayee et al., 2014; Ceja-Navarro et al., 2014). However, the successful amplifications of *nifH* gene does not always indicate that the encoding dinitrogenase reductase is actually expressed by the corresponding bacteria, because the expressions of *nifH* genes are regulated at the transcriptional and post-transcriptional level (Zehr et al., 2003). Thus, detection of mRNA by a reverse transcription polymerase chain reaction (RT-PCR) is usually used to further support the expression of *nifH* genes and determine the actual activity of specific gut microbes in nitrogen fixation (Noda et al., 1999, 2002; Desai et al., 2012; Ayayee et al., 2014; Ceja-Navarro et al., 2014).

Like xylophagous insects, mealworm's gut also provides an ecological niche for the resident of a complex mixed microbial community (Yang et al., 2015b; Xia et al., 2020). The composition of gut bacterial community within mealworms from different sources has been investigated using both culture-dependent and culture-independent methods (Jung et al., 2014; Yang et al., 2015b; Wang et al., 2015; Li et al., 2016; Stoops et al., 2016; Xia et al., 2020; Urbanek et al., 2020; Przemieniecki et al., 2020). The culture-independent pyrosequencing-based analyses show that *Proteobacteria*, *Bacteroidetes*, *Firmicutes*, *Actinobacteria* and *Tenericutes* are the dominant phylum (Jung et al., 2014; Wang et al., 2015; Li et al., 2016; Stoops et al., 2016; Urbanek et al., 2020; Przemieniecki et al., 2020). The detected species belong to phylum *Proteobacteria* or *Firmicutes*, mainly *Citrobacter* spp., *Klebsiella* spp., *Enterobacter* spp., *Pantoea* spp., *Pseudomonas* spp., or *Clostridium* spp., are known potential nitrogen-fixing bacteria (Bar-Shmuel et al., 2020). These available data providing circumstantial evidence that nitrogen fixation may occur in

mealworms. However, the direct evidence for the nitrogen-fixing activity of vinyl plastic-eating mealworm and the involvement of gut nitrogen-fixing bacteria is still lacking.

Therefore, the specific aims of this study are to (i) determine whether the gut bacteria-mediated nitrogen fixation takes place within mealworms when they were fed with Styrofoam as sole diet by using the acetylene reduction assays and stable nitrogen isotope composition analyses, (ii) to uncover the diazotrophic diversity of the gut microbial symbiont by culture-dependent technique and culture-independent amplification of *nifH* genes, (iii) to illustrate whether the *nifH* genes encoding dinitrogenase reductase are actually expressed by gut nitrogen-fixing bacteria. The overall aim is to extend our present understanding that plastic-eating mealworms can adapt the nitrogen-deficient plastic diet by employing the nitrogen-fixing capacity of gut microbiota.

Materials And Methods

Collection and culture of mealworms

Mealworms (growth age at approximately 3 – 4 instars) were purchased from six different insect-breeding plants in northern (Beijing), eastern (Shanghai and Ningbo), western (Chengdu), southern (Guangzhou), and central (Luoyang) region of China, respectively. The distributions of body weight and size of the sampled mealworms are shown in SI Tables S1. The conventional diet for these sampled mealworms was bran.

Sampled mealworms from each source were divided into two groups. One group (denoted as Normal group, $n = 3$) were reared with Styrofoam blocks as a sole diet for 28 days. Other group (labeled as Antibiotic group, $n = 3$) were first fed with the antibiotic diet (60 mg gentamicin per 1 g bran) until the number of active gut bacteria became approximately zero. The number of active gut bacteria was detected by the series dilution method of plate counting on solid nitrogen-free plates (1.5% agar) (Burdman et al., 1998). These mealworms were subsequently fed with Styrofoam for 28 days. Both of “Normal” or “Antibiotic” groups were used for further investigations.

Acetylene reduction assays

Ten mealworms were randomly selected from each group and were quickly surface sterilized by immersion in 75% ethanol for 1 min and then rinsed 2 times with sterile phosphate saline buffer (PBS) prior to gut excision. Their guts were sterilely drawn out and pooled into a Hungate tube containing 5 mL of sterile PBS buffer. Then, Hungate tubes were stoppered with rubber plugs and the inside atmosphere was exchanged with argon gas (Ar). Acetylene was injected into the Hungate tubes to a final concentration of 20% (v/v) by replacement of an identical volume of argon gas. The Hungate tubes without guts but with acetylene gas were used as blank controls. The tubes for each group of mealworms and controls were prepared in triplicate. After 5 hours of incubation at 30 °C, to determine the reduction of acetylene to ethylene, 0.5 mL of gas sampled from the top of the tubes was analyzed by gas chromatography (GC, PANNA 91A, Changzhou, China) equipped with a flame ionization detector (FID)

and HP-PLOT Al_2O_3 S column (25 m $\times$ 0.32 mm $\times$ 8 μm , Agilent, USA). The operating temperature for inlet, column oven and FID was 150 $^\circ\text{C}$, 100 $^\circ\text{C}$ and 280 $^\circ\text{C}$, respectively. The gas flow rate for N_2 , H_2 and air was set as 20 $\text{mL}\cdot\text{min}^{-1}$, 40 $\text{mL}\cdot\text{min}^{-1}$ and 450 $\text{mL}\cdot\text{min}^{-1}$, respectively. Purified ethylene was used as the standard for the quantitative analysis. To estimate the amount of fixed nitrogen, the molar ratio of ethylene to nitrogen was defined as 4 : 1 and the average working time for nitrogen fixation by bacteria was assumed to be 16 h per day (Behar et al., 2005). The fixed nitrogen equivalent of protein was calculated based on a value of 16% nitrogen in protein (Behar et al., 2005).

Stable nitrogen isotope composition measurements

Ten mealworms were randomly selected from each group and lyophilized, ground to a homogeneous powder. The powder (about 1.0 mg) was packed in tin capsules and combusted to CO_2 and N_2 at 1000 $^\circ\text{C}$ in an on-line combustor. Subsequently, N_2 was separated in a He carrier stream on a Carbosieve G column and further introduced to an isotope ratio mass spectrometer (Finnigan MAT 253, Thermo Electron, Waltham, MA). Nitrogen isotope compositions were reported as $\delta^{15}\text{N}$ values relative to atmospheric nitrogen (a pure N_2 gas reference, 0.00‰). The system was calibrated using multiple runs of a powdered working standard (ammonium sulfate and sucrose mixture with $\delta^{15}\text{N}$ versus air of 1.33‰) and the detection method was periodically calibrated against international isotope standards (IAEA N1 and N3).

Isolation and characterization of nitrogen-fixing bacteria

Thirty mealworms from each group were randomly collected and quickly surface sterilized by immersion in 75 % ethanol for 1 min and then rinsed twice with sterile PBS. Their guts were drawn out and pooled in a 10 mL centrifuge tube containing 5 mL sterile PBS. The solution was roiled on a vortex mixer for 5 min in order to adequately release the microbial contents from the guts. After that, the clearing gut tissues were carefully picked out with a pipette, while the solution containing gut contents was used as gut microbial inoculum, which was 10-fold diluted and inoculated on the plates containing semi-solid nitrogen-free media (0.5% agar) (Burdman et al., 1998). These inoculated plates were incubated for 5 days at 30 $^\circ\text{C}$. Individual colonies were randomly picked and streaked repeatedly on Luria-Bertani (LB) agar to obtain pure cultures. Genomic DNA extraction, PCR amplification, sequencing and phylogenetic analysis of the 16S rRNA gene from these pure cultures were carried out according to the previously described method (Xia et al., 2020).

For acetylene reduction assays, pure cultures were inoculated in LB media and incubated in agitation for 24 h at 30 $^\circ\text{C}$. A well-known nitrogen-fixing bacterium *Azospirillum brasilense* was used as positive control (Pinto-Tomás et al., 2009). Bacterial pellets were washed 3 times with sterile PBS and re-suspended in 1 mL sterile PBS. One hundred microliter of this suspension were inoculated in 25 mL test tubes containing 15 mL of semi-solid nitrogen-free media. The test tubes stoppered with cotton plugs were incubated for 72 h at 30 $^\circ\text{C}$. Subsequently, the cotton plugs of test tubes were replaced with rubber plugs and 1 mL of acetylene was injected into the tubes. Ethylene production was measured periodically by GC as described above.

DNA extraction, PCR amplification, cloning and sequencing of *nifH* genes

Fifty of active superworms were randomly picked out from each group and were surface sterilized with the same method described above. Their guts were drawn out and pooled into a 50 mL centrifuge tube containing 25 mL sterile PBS on ice, respectively. After being roiled on a vortex mixer for 5 min, the microbial contents were released from the guts and the empty guts were carefully removed from the tube by using a pipette. The solution containing gut microbial contents were further centrifuged at 4 °C, 12,000 rpm for 5 min. The supernatants were discarded and the precipitates were used for the DNA extraction by using the TIANamp Stool DNA Kit (DP 328, TIANGEN, Beijing, China) according to the manufacturer's instructions. The average concentration of yielded genomic DNA was determined by NanoDrop One Microvolume UV-Vis Spectrophotometer (Thermo Scientific, USA). Genomic DNA extraction of the pure cultures were carried out according to the previously described method (Xia et al., 2020).

An internal fragment of the nitrogenase (*nifH*) gene (about 360 bp) was amplified with using each set of previously reported primer pairs (Table S2). The genomic DNA extract (1 µL, 300 ng·µL⁻¹), 1 µL of forward and 1µL of reverse primer were added to 11 µL of the PCR mixtures (2×M5 Taq HiFi PCR mix, Mei5 Biotechnology, Beijing). The final volume of PCR mixtures was 20 µL by adding double-distilled H₂O (ddH₂O). Negative controls without DNA were also included to test for contamination. PCR was carried out with one denaturation step of 8 min at 95°C, 35 cycles of 94°C for 0.5 min, 47°C for 0.5 min and 72°C for 1 min, and a final extension step of 7 min at 72°C. Amplification products were analyzed by electrophoresis in 1.5% (w/v) agarose gels in 1× TAE buffer. PCR products of the expected size were gel-purified using a GeneJET Gel Extraction Kit (Thermo Scientific, USA). PCR products were ligated into pJET1.2/blunt Cloning Vector and transformed into *Escherichia coli* XL-10-Gold cells using the CloneJET PCR Cloning Kit (Thermo Scientific, USA). Plasmid DNA was isolated from individual clones and purified using TIANprep Mini Plasmid Kit (DP103, Tiangen, Beijing, China). Nucleotide sequencing was done using ABI 3730XL genetic analyzer (Applied Biosystems, CA, USA).

RNA extraction and RT-PCR

Gut microbial contents were prepared with the same method as described above for DNA extraction. Total RNA was extracted from gut contents by using TRIzol™ Reagen (Invitrogen, USA). RNA samples were purified using RapidOut DNA Removal Kit (Thermo Scientific, USA) to remove DNA. Reverse transcription reactions were performed using GoScript™ Reverse Transcription System (Promega, USA). After reverse transcription, *nifH* genes were amplified in cDNA samples using PCR experimental conditions as described in the previous section. Each set of previously reported primer pairs was tested in the PCR reaction (Table S2). RNA samples without reverse transcription step were used as controls to check for residual DNA in the RNA preparations. RT-PCR and PCR negative controls were also included to test for reagent contamination. Generation of clone libraries and sequencing of RT-PCR amplified *nifH* products were performed by the same method described in the above section.

Phylogenetic analysis of *nifH* clones

All detected *nifH* gene nucleotide sequences were translated into amino acid sequences by using ExPASy bioinformatics resource tool (Gasteiger et al., 2003). Amino acid sequence with at least one amino acid change was considered as a unique operational taxonomic units (OTUs). The closest phylogenetic neighbours were obtained by aligning the representative amino acid sequence of each OTU against NCBI database using protein Basic Local Assignment Search Tool (BLASTp) (Altschul et al., 1990). Sequence alignment of all amino acid sequences was performed by using clustal x 2.0 software (Larking et al., 2007). Aligned sequences were used to reconstruct the maximum-likelihood (ML) phylogenetic trees using the program MEGA 6.0 (Felsenstein et al., 1981; Tamura et al., 2013), with bootstrap values based on 1000 replications.

Results

Nitrogen-fixing activity of Styrofoam-eating mealworm

We used acetylene reduction assay to examine the nitrogen-fixing activity of mealworm, because both acetylene and nitrogen are unsaturated compounds with covalent triple bonds and the conversion of acetylene to ethylene has been widely used as an indicator of nitrogen-fixing activity for both terrestrial and aquatic systems (Myrold, 2007). As shown in Fig. 1a, ethylene production from acetylene was detected by Styrofoam-eating mealworms from six different sources, with the rate from 12.3 ± 0.7 to 32.9 ± 9.3 nmol ethylene·h⁻¹·gut⁻¹. The acetylene-reducing (nitrogen-fixing) activity of Styrofoam-eating mealworm was similar to that of individual field-collected Mediterranean fruit fly (18.0 ± 1.5 nmol ethylene·h⁻¹·gut⁻¹) (Behar et al., 2005), but was one or two orders of magnitude higher than that of termite (1.25 ± 0.37 nmol acetylene·h⁻¹·gut⁻¹) and stag beetle (0.03 nmol acetylene·h⁻¹·gut⁻¹) (Breznak et al., 1973; Kuranouchi et al., 2006). When mealworms were fed with antibiotic diet (60 mg gentamicin per g bran), all detectable acetylene-reducing (nitrogen-fixing) activity was significantly weakened and decreased to the range of 8.7 ± 0.5 to 19.0 ± 0.4 nmol ethylene·h⁻¹·gut⁻¹ ($p < 0.01$, Fig. 1a), showing a correlation with the suppression of gut bacterial numbers detected by the serial dilution method of plate counting (SI Fig. S1). These results indicated that nitrogen fixation occurred in Styrofoam-eating mealworms and nitrogen-fixing bacteria were present within their guts. Additionally, we also determined the acetylene-reducing (nitrogen-fixing) activity of mealworms fed with the conventional diet bran (rich in carbohydrates and proteins). The bran-feeding mealworms had an acetylene-reducing (nitrogen-fixing) activity of 8.1 ± 2.1 to 12.2 ± 0.3 nmol ethylene·h⁻¹·gut⁻¹, which was significantly lower than that of Styrofoam-eating mealworms ($p < 0.01$, SI Table S1). This result agreed with the previously known phenomenon that the available nitrogen in the diet would repress the nitrogenase synthesis of nitrogen-fixing systems (Breznak et al., 1973).

According to the acetylene-reducing rate, the fixed nitrogen equivalents of protein were estimated as 8.6 to 23.0 µg per day per mealworm using the previous reported method (Behar et al., 2005). Investigation of natural stable nitrogen isotope compositions in organisms also can provide further evidence for the importance of nitrogen fixation as a source of new nitrogen, because the δ¹⁵N value of atmospheric N₂ is

0‰ and nitrogen-fixing organisms often have $\delta^{15}\text{N}$ close to 0‰ (Tayasu et al., 1994). Thus, we analyzed the $\delta^{15}\text{N}$ values of Styrofoam-eating mealworms, antibiotic-treated mealworms and bran-feeding mealworms that were sampled from six different source. As shown in Fig. 1b, all Styrofoam-eating mealworms had $\delta^{15}\text{N}$ values between 1.7‰ and 1.9‰, which were significantly lower than that of antibiotic-treated mealworms (2.3‰–2.4‰). Due to the suppression of gut microbiota within antibiotic-treated mealworms, low $\delta^{15}\text{N}$ values of Styrofoam-eating mealworms could be attributed to the presence of gut nitrogen-fixing bacteria that can fix nitrogen from atmosphere nitrogen and provide a substantial new nitrogen source for the mealworm's metabolism. Moreover, the $\delta^{15}\text{N}$ values of bran-feeding mealworms were also analyzed and found to be in the range of 1.9‰ to 2.1‰ (SI Table S1), which were higher than that of Styrofoam-eating mealworm. This result was consistent with the decreased acetylene-reducing (nitrogen-fixing) activity within bran-feeding mealworms in comparison to Styrofoam-eating mealworms.

Nitrogen-fixing bacterial cultures isolated from mealworm's gut

To assess the level of potential nitrogen-fixing bacteria in gut microbiota of Styrofoam-eating mealworms, gut contents from Styrofoam-eating mealworms were serially diluted and inoculated in a semi-solid nitrogen-free enrichment medium for the growth of nitrogen-fixing bacteria. The typical colonies of nitrogen-fixing bacteria came out on all plates inoculated with gut contents of Styrofoam-eating mealworms from six different sources (SI Fig. S1). The number of potential nitrogen-fixing bacteria was estimated at 5.5×10^4 CFU per gut. In contrast, the size of the total cultivable fraction on the a general LB medium, inoculated with the same dilution series, was estimated to be 5.0×10^6 CFU per gut (Yang et al., 2015b). Therefore, the level of functional nitrogen fixers was estimated at about 1.1% of the total bacterial population within the gut of Styrofoam-eating mealworm.

To further identify the specific gut microbes responsible to nitrogen fixation, we conducted extensive streaking-isolations using nitrogen-free medium. Colonies isolated from the nitrogen-free plates were identified by amplification and sequencing of the almost full length 16S rRNA gene. Phylogenetic analyses based on 16S rRNA (Fig. 2a) revealed that isolates belonged to the genera *Klebsiella*, *Mixta*, *Kluyvera* and *Citrobacter*. The abilities of isolates to reduce acetylene into ethylene were also determined. As shown in Fig. 2b, in comparison to a positive control of a known free-living diazotrophs *Azospirillum brasilense* ATCC 29145, isolates of the genus *Klebsiella* sp. showed a comparable nitrogen-fixing rate while isolates of other genera of *Mixta*, *Kluyvera* and *Citrobacter* had a relative lower rate than the *Klebsiella* isolates did. Given the noticeable nitrogen-fixation rates measured for the isolates of *Klebsiella*, *Mixta*, *Kluyvera* and *Citrobacter* and their prevalence in the guts of Styrofoam-eating mealworms from six different sources, we can therefore deduce that the gut species within the order of *Enterobacteriales* are the important nitrogen-fixing symbiont of Styrofoam-eating mealworms.

nifH genes diversity in Styrofoam-eating mealworm's gut

To further determine whether more uncultivable nitrogen-fixing bacteria resided in the Styrofoam-eating mealworm's gut, *nifH* genes were amplified from the DNA extracted from gut contents as well as the genomic DNA of nitrogen-fixing gut isolates. We have tested four previously reported primer pairs (SI Table S2) and found that only the primer pairs PolF-PolR could successfully produce the PCR products (about 360 bp) from all DNA samples of gut contents (Fig. 3a). With each of four tested primer pairs, the presence of *nifH* was only found in the nitrogen-fixing gut isolate from *Klebsiella* sp. but not *Mixta* sp., *Kluyvera* sp. and *Citrobacter* sp. (Fig. 3b). Nevertheless, these results revealed the presence of nitrogen-fixing bacteria within the gut microbiota of Styrofoam-eating mealworms.

In total, more than 100 clones from clone libraries of PCR products were picked and sequenced. However, several clones had similar DNA sequences and identical deduced amino acid sequences. The clones with identical amino acids sequence were classified into one OTU and only one sequence was taken as representative for each OTU. Finally, seven clones, representing different *nifH* amino acid sequences, were chosen for further study (SI Table S3). Phylogenetic analysis of the obtained *nifH* amino acid sequences with their closest neighbors was shown in Fig. 4. According to Zehr et al. (2003), *nifH* protein sequences can be classified into four major clusters (Clusters I–IV) in phylogeny. We found that all *nifH* clones from mealworm's gut fell into the previously described Cluster I, which consists of *nifH* genes from conventional eubacterial Mo-Fe operons and some *vnfH* (Zehr et al., 2003). Furthermore, cluster I could be divided into four subgroups, which consists of *nifH* sequences from the genus *Klebsiella*, *Pseudomonas*, unculture bacteria and *Azospirillum*, respectively. These results indicate that Styrofoam-eating mealworms have a unique *nifH* genes diversity, which is different from of the *nifH* genes within termite gut that generally belong to Clusters II or III compose of sequences from a diverse group of strict anaerobic microorganisms such as clostridia, sulfate reducers, spirochetes, and methanogens (Yamada et al., 2007; Du et al., 2012).

Expression of *nifH* genes in Styrofoam-eating mealworm's gut

To demonstrated whether the *nifH* genes encoding dinitrogenase reductase were actually expressed by gut nitrogen-fixing bacteria within Styrofoam-eating mealworms, transcribed *nifH* mRNA in the gut microbial communities was analyzed using RT-PCR. Amplifications were detected in the cDNA samples of gut contents from six different Styrofoam-eating mealworms but not in the control RNA sample of gut contents without the RT reaction step (Fig. 3c), indicating that the RNA preparation did not have any residual DNA after the DNAase treatment and *nifH* transcripts virtually presented in the total RNA pool extracted from Styrofoam-eating mealworms' guts. In comparison, no amplifications were observed in the cDNA samples of gut contents from antibiotic-treated mealworms and bran-feeding mealworms (SI Fig. S2), in line with their relative weaker acetylene-reducing (nitrogen-fixing) activities than Styrofoam-eating mealworms.

Clone libraries were constructed from *nifH* cDNA amplified by RT-PCR from total RNA. We recovered two unique representatives from the RT-PCR cDNA clone libraries. They clustered within group 1 of Cluster I and were exclusively *Klebsiella* phylotypes (Fig. 4 and SI Table S3). Given the high nitrogen-fixation rates

measured for the *Klebsiella* isolates (Fig. 2) and the apparent consistence of *Klebsiella nifH* genes in PCR and RT-PCR clone libraries (Figs. 3 and 4), it is likely that *Klebsiella*, the order of *Enterobacteriales*, is an important nitrogen-fixing symbiont of Styrofoam-eating mealworms.

Discussion

Since mealworm was first introduced as a plastic eater and digester in 2015, this coleopteran insect larva has been proven to survive by eating a series of non-hydrolyzable vinyl plastics, such as PS, PE, PP and PVC, as sole diet and effectively degrade them (Yang et al., 2015a; Brandon et al., 2018; Aboelkheir et al., 2019; Peng et al., 2020; Yang et al., 2020). In the perspective of nutrient elements content, the petroleum-derived vinyl plastics were carbon-rich but nitrogen-deficient diet. From a biology viewpoint, nitrogen was an indispensable element for biosynthesis of proteins, nucleic acids and other metabolites in insect (Bar-Shmuel et al., 2020). Hence, there was a pending question how mealworms can survive on nitrogen-deficient vinyl plastic as sole diet as well as that feed on normal nutritionally adequate diet (such as bran). Inspired by that the xylophagous insects were capable of supplementing their diets by fixing nitrogen from atmosphere (Bar-Shmuel et al., 2020), we attempted to reveal whether mealworms could also fix atmosphere nitrogen. Our results explicitly showed that Styrofoam-eating mealworm has a significant acetylene-reducing (nitrogen-fixing) activity (12.3 ± 0.7 to 32.9 ± 9.3 nmol ethylene·h⁻¹·gut⁻¹), which was one or two orders of magnitude higher than that of termite (1.25 ± 0.37 nmol acetylene·h⁻¹·gut⁻¹) and stag beetle (0.03 nmol acetylene·h⁻¹·gut⁻¹) (Breznak et al., 1973; Kuranouchi et al., 2006). This finding gives a reasonable interpretation on the question why mealworm can survive on nitrogen-deficient vinyl plastic as sole diet.

Furthermore, investigation of natural stable nitrogen isotope compositions in Styrofoam-eating mealworm has provided evidence for the incorporation of atmosphere nitrogen into the biomass. The fixed nitrogen equivalents of protein were estimated as 8.6 to 23.0 µg per day per mealworm based on the acetylene-reducing rate (Behar et al., 2005). Mealworms had a high protein content up to 50% and were widely used as food for pet and poultry breeding industry, and recently proposed as an edible insect for humans by Food and Agriculture Organization of the United Nations (FAO) (Li et al., 2016). Given that the ability of mealworms to degrade plastic and assimilated the degraded products of plastic as carbon source into biomass has been demonstrated, our finding of nitrogen-fixing capability will build the feasibility for mealworms to transform plastic waste into protein without the additional feed of nitrogen nutrient.

Notably, gut microbiota has been found to be responsible for the plastic degradation in our previous studies (Yang et al., 2015b) and nitrogen fixation within mealworm in this study. Hence, engineering a synthetic microbial consortium or cell factories, which can degrade plastic waste and fix atmosphere nitrogen at the same time, would be a possible way for up-recycling plastic waste to protein in vitro (Ru et al., 2020).

In conclusion, our work is the first demonstration for nitrogen-fixing activity within plastic-eating mealworms by using acetylene reduction assays and nature nitrogen isotopic analyses. Antibiotic suppression of gut microbial microbiota impaired the mealworm's ability to fix nitrogen from atmosphere, indicating the contribution of gut microbiota to nitrogen fixation. By applying the traditional culture-dependent technique, PCR and RT-PCR of *nifH* gene, the presence of nitrogen-fixing bacteria within the gut was demonstrated. These findings advance understandings of the role of gut microbiota in mealworms, and suggest that fixing atmosphere nitrogen is a way whereby plastic-eating mealworms overcome the nutritional obstacle of nitrogen limitation. Furthermore, it also indicates that mealworms may be applied for up-recycling of plastic waste to produce protein sources for food production and other commercial usage through the synergistic efforts of plastic-degrading and nitrogen-fixing activities of gut bacterial symbiont.

Declarations

Authors' contributions

HL, LX and WJ conducted acetylene reduction assays, isolation of gut nitrogen-fixing bacteria, and PCR and RT-PCR experiments. JG conducted bulk stable isotope measurements on mealworms. YY and HL analyzed data and wrote the manuscript. YY conceived and designed the research.

Funding

This work was supported by grants from National Natural Science Foundation of China (No. 51603004 and No. 31961133015), the Young Elite Scientist Sponsorship Program of the China Association of Science and Technology (No. 2017QNRC001) and the Beijing Institute of Technology Research Fund Program for Young Scholars (No. 3160011181804).

Availability of data and material

All studied strains of *Kluyvera* sp. BIT-1, *Citrobacter* sp. BIT-105, *Mixta* sp. BIT-56, *Klebsiella* sp. BIT-96 and were deposited in the public culture collection CGMCC or CICC with the collection number of CCTCC AB 2020371, CCTCC AB 2020194, CGMCC 1.17041, CGMCC 1.17846,

The GenBank accession numbers for 16S rRNA of strains *Kluyvera* sp. BIT-1, *Mixta* sp. BIT-56, *Klebsiella* sp. BIT-96 and *Citrobacter* sp. BIT-105 were MW422732, MW422733, MW422734 and MW422751, respectively. All detected *nifH* gene nucleotide sequences were deposited in GenBank with the accession numbers of MW442362 to MW442370.

Competing interests

The authors declare no competing interests.

Ethics approval

This article does not contain any studies with human participants or animals performed by any of the authors. All authors read and approved the manuscript.

References

1. Aboelkheir MG, Visconte LY, Oliveira GE, Toledo Filho RD, Souza FG (2019) The biodegradative effect of *Tenebrio molitor* Linnaeus larvae on vulcanized SBR and tire crumb. *Sci Total Environ* 649:1075–1082
2. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *J Mol Biol* 215:403–410
3. Ayayee P, Rosa C, Ferry JG, Felton G, Saunders M, Hoover K (2014) Gut microbes contribute to nitrogen provisioning in a wood-feeding cerambycid. *Environ Entomol* 43:903–912
4. Bar-Shmuel N, Behar A, Segoli M (2020) What do we know about biological nitrogen fixation in insects? Evidence and implications for the insect and the ecosystem. *Insect Sci* 27(3):392–403
5. Behar A, Yuval B, Jurkevitch E (2005) Enterobacteria-mediated nitrogen fixation in natural populations of the fruit fly *Ceratitis capitata*. *Mol Ecol* 14(9):2637–2643
6. Benemann JR (1973) Nitrogen fixation in termites. *Science* 181:164–165
7. Brandon AM, Gao SH, Tian R, Ning D, Yang SS, Zhou J, Wu WM, Criddle CS (2018) Biodegradation of polyethylene and plastic mixtures in mealworms (larvae of *Tenebrio molitor*) and effects on the gut microbiome. *Environ Sci Technol* 52:6526–6533
8. Breznak JA, Brill WJ, Mertins JW, Coppel HC (1973) Nitrogen fixation in termites. *Nature* 244:577–580
9. Bridges JR (1981) Nitrogen-fixing bacteria associated with bark beetles. *Microb Ecol* 7:131–137
10. Burdman S, Jurkevitch E, Schwartzburd B, Hampel M, Okon Y (1998) Aggregation in *Azospirillum brasilense*: effects of chemical and physical factors and involvement of extracellular components. *Microbiology* 144(7):1989–1999
11. Carpenter EJ, Culliney JL (1975) Nitrogen fixation in marine shipworms. *Science* 187:551–552
12. Ceja-Navarro JA, Nguyen NH, Karaoz U, Gross SR, Herman DJ, Andersen GL, Bruns TD, Pett-Ridge J, Blackwell M, Brodie EL (2014) Compartmentalized microbial composition, oxygen gradients and nitrogen fixation in the gut of *Odontotaenius disjunctus*. *ISME J* 8:6–18
13. Desai MS, Brune A (2012) Bacteroidales ectosymbionts of gut flagellates shape the nitrogen-fixing community in dry-wood termites. *ISME J* 6:1302–1313
14. Doolittle M, Raina A, Lax A, Boopathy R (2008) Presence of nitrogen fixing *Klebsiella pneumoniae* in the gut of the Formosan subterranean termite (*Coptotermes formosanus*). *Bioresour Technol* 99:3297–3300
15. Du X, Li X, Wang Y, Peng J, Hong H, Yang H (2012) Phylogenetic diversity of nitrogen fixation genes in the intestinal tract of *Reticulitermes chinensis* Snyder. *Curr Microbiol* 65:547–551

16. Felsenstein J (1981) Evolutionary trees from DNA sequences: a maximum likelihood approach. *J Mol Evol* 17:368–376
17. French JRJ, Turner GL, Bradbury JF (1976) Nitrogen fixation by bacteria from the hindgut of termites. *Microbiology* 95:202–206
18. Gasteiger E, Gattiker A, Hoogland C, Ivanyi I, Appel RD, Bairoch A (2003) ExPASy: the proteomics server for in-depth protein knowledge and analysis. *Nucleic Acids Res* 31:3784–3788
19. Gomathi V, Ramalakshmi A, Ramasamy K (2005) Isolation of nitrogen fixing bacteria from fungus termites. *Entomol Res* 35:75–78
20. Jung J, Heo A, Park YW, Kim YJ, Koh H, Park W (2014) Gut microbiota of *Tenebrio molitor* and their response to environmental change. *J Microbiol Biotechnol* 24(7):888–897
21. Kneip C, Lockhart P, Voss C, Maier UG (2007) Nitrogen fixation in eukaryotes—new models for symbiosis. *BMC Evol Biol* 7(1):55
22. Kuranouchi T, Nakamura T, Shimamura S, Kojima H, Goka K, Okabe K, Mochizuki A (2006) Nitrogen fixation in the stag beetle, *Dorcus* (*Macrodercus*) *rectus* (Motschulsky) (Col., Lucanidae). *J Appl Entomol* 130(9-10):471–472
23. Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, Valentin F, Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ, Higgins DG (2007) Clustal W and Clustal X version 2.0. *Bioinformatics* 23:2947–2948
24. Li L, Xie B, Dong C (2016) Can closed artificial ecosystem have an impact on insect microbial community? A case study of yellow mealworm (*Tenebrio molitor* L.). *Ecol Engrg* 86:183–189
25. Lilburn TG, Kim KS, Ostrom NE, Byzek KR, Leadbetter JR, Breznak JA (2001) Nitrogen fixation by symbiotic and free-living spirochetes. *Science* 292:2495–2498
26. Morales-Jiménez J, Zúñiga G, Villa-Tanaca L, Hernández-Rodríguez C (2009) Bacterial community and nitrogen fixation in the red turpentine beetle, *Dendroctonus valens* LeConte (Coleoptera: Curculionidae: Scolytinae). *Microb Ecol* 58(4):879–891
27. Morales-Jiménez J, Vera-Ponce de León A, García-Domínguez A, Martínez-Romero E, Zúñiga G, Hernández-Rodríguez C (2013) Nitrogen-fixing and uricolytic bacteria associated with the gut of *Dendroctonus rhizophagus* and *Dendroctonus valens* (Curculionidae: Scolytinae). *Microb Ecol* 66:200–210
28. Myrold DD (2007) Quantification of nitrogen transformations. In *Manual of Environmental Microbiology*. ASM 3:687–696
29. Nardi JB, Mackie RI, Dawson JO (2002) Could microbial symbionts of arthropod guts contribute significantly to nitrogen fixation in terrestrial ecosystems? *J Insect Physiol* 48(8):751–763
30. Noda S, Ohkuma M, Usami R, Horikoshi K, Kudo T (1999) Culture-independent characterization of a gene responsible for nitrogen fixation in the symbiotic microbial community in the gut of the termite *Neotermes koshunensis*. *Appl Environ Microbiol* 65:4935–4942

31. Noda S, Ohkuma M, Kudo T (2002) Nitrogen fixation genes expressed in the symbiotic microbial community in the gut of the termite *Coptotermes formosanus*. *Microbes Environ* 17:139–143
32. Ohkuma M, Noda S, Usami R, Horikoshi K, Kudo T (1996) Diversity of nitrogen fixation genes in the symbiotic intestinal microflora of the termite *Reticulitermes speratus*. *Appl Environ Microbiol* 62:2747–2752
33. Ohkuma M, Noda S, Kudo T (1999) Phylogenetic diversity of nitrogen fixation genes in the symbiotic microbial community in the gut of diverse termites. *Appl Environ Microbiol* 65:4926–4934
34. Orón Tamayo D, Heil M (2015) N fixation in insects: its potential contribution to N cycling in ecosystems and insect biomass. In de Bruijn, F. J. (Ed.), *Biological Nitrogen Fixation*. John Wiley & Sons, Inc, Hoboken, pp. 1141–1148
35. Pandey S, Waller DA, Gordon AS (1992) Variation in acetylene-reduction (nitrogen-fixation) rates in *Reticulitermes* spp. (Isoptera; Rhinotermitidae). *Va J Sci* 43(3):333–338
36. Peng BY, Chen Z, Chen J, Yu H, Zhou X, Criddle CS, Wu WM, Zhang Y (2020) Biodegradation of polyvinyl chloride (PVC) in *Tenebrio molitor* (Coleoptera: Tenebrionidae) larvae. *Environ Int* 145:106106
37. Pinto-Tomás AA, Anderson MA, Suen G, Stevenson DM, Chu FS, Cleland WW, Weimer PJ, Currie CR (2009) Symbiotic nitrogen fixation in the fungus gardens of leaf-cutter ants. *Science* 326:1120–1123
38. Potrikus CJ, Breznak JA (1977) Nitrogen-fixing *Enterobacter agglomerans* isolated from guts of wood-eating termites. *Appl Environ Microbiol* 33:392–399
39. Prestwich GD, Bentley BL (1981) Nitrogen fixation by intact colonies of the termite *Nasutitermes corniger*. *Oecologia* 49(2):249–251
40. Przemieniecki SW, Kosewska A, Ciesielski S, Kosewska O (2020) Changes in the gut microbiome and enzymatic profile of *Tenebrio molitor* larvae biodegrading cellulose, polyethylene and polystyrene waste. *Environ Pollut* 256:113265
41. Risal CP, Ozawa T (2002) Isolation and characterization of diazotrophs from the intestinal tract of an earthworm (*Pheretima vittata*). *Soil Sci Plant Nutr* 48:101–103
42. Ru JK, Huo YX, Yang Y (2020) Microbial degradation and valorization of plastic wastes. *Front Microbiol* 11:422
43. Sapountzis P, de Verges J, Rousk K, Cilliers M, Vorster BJ, Poulsen M (2016) Potential for nitrogen fixation in the fungus-growing termite symbiosis. *Front Microbiol* 7: 1993
44. Stoops J, Crauwels S, Waud M, Claes J, Lievens B, Van Campenhout L (2016) Microbial community assessment of mealworm larvae (*Tenebrio molitor*) and grasshoppers (*Locusta migratoria migratorioides*) sold for human consumption. *Food Microbiol* 53:122–127
45. Tamura K, Stecher G, Peterson D, Filipski A, Kumar S (2013) MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol* 30:2725–2729
46. Tayasu I, Sugimoto A, Wada E, Abe T (1994) Xylophagous termites depending on atmospheric nitrogen. *Naturwissenschaften* 81(5):229–231

47. Urbanek AK, Rybak J, Wróbel M, Leluk K, Mirończuk AM (2020) A comprehensive assessment of microbiome diversity in *Tenebrio molitor* fed with polystyrene waste. *Environ Pollut* 262:114281
48. Wang Y, Zhang Y (2015) Investigation of gut-associated bacteria in *Tenebrio molitor* (Coleoptera: Tenebrionidae) larvae using culture-dependent and DGGE methods. *Ann Entomol Soc Am* 108:941–949
49. Waterbury JB, Calloway CB, Turner RD (1983) A cellulolytic nitrogen-fixing bacterium cultured from the gland of Deshayes in shipworms (Bivalvia: Teredinidae). *Science* 221(4618):1401–1403
50. Xia M, Wang J, Huo YX, Yang Y (2020) *Mixta tenebrionis* sp. nov., isolated from the gut of the plastic-eating mealworm *Tenebrio molitor* L. *Int J Syst Evol Microbiol* 70:790–796
51. Yamada A, Inoue T, Noda S, Hongoh Y, Ohkuma M (2007) Evolutionary trend of phylogenetic diversity of nitrogen fixation genes in the gut community of wood-feeding termites. *Mol Ecol* 16:3768–3777
52. Yamada A, Inoue T, Wiwatwitaya D, Ohkuma M, Kudo T, Sugimoto A (2006) Nitrogen fixation by termites in tropical forests. *Thailand Ecosystems* 9(1):75–83
53. Yang SS, Ding MQ, He L, Zhang CH, Li QX, Xing DF, Wu WM (2020) Biodegradation of polypropylene by yellow mealworms (*Tenebrio molitor*) and superworms (*Zophobas atratus*) via gut-microbe-dependent depolymerization. *Sci Total Environ* 756:144087
54. Yang SS, Wu WM, Brandon AM, Fan HQ, Receveur JP, Li Y, Wang ZY, Fan R, McClellan RL, Gao SH, Ning D, Phillips DH, Peng BY, Wang H, Cai SY, Li P, Cai WW, Ding LY, Yang J, Zheng M, Ren J, Zhang YL, Gao J, Xing D, Ren NQ, Waymouth RM, Zhou J, Tao HC, Picard CJ, Benbow ME, Criddle CS (2018) Ubiquity of polystyrene digestion and biodegradation within yellow mealworms, larvae of *Tenebrio molitor* Linnaeus (Coleoptera: Tenebrionidae). *Chemosphere* 212:262–271
55. Yang Y, Wang J, Xia M (2020) Biodegradation and mineralization of polystyrene by plastic-eating superworms *Zophobas atratus*. *Sci Total Environ* 708:135233
56. Yang Y, Yang J, Wu WM, Zhao J, Song Y, Gao L, Yang R, Jiang L (2015) Biodegradation and mineralization of polystyrene by plastic-eating mealworms. 1. chemical and physical characterization and isotopic tests. *Environ Sci Technol* 49:12080–12086
57. Yang Y, Yang J, Wu WM, Zhao J, Song Y, Gao L, Yang R, Jiang L (2015) Biodegradation and mineralization of polystyrene by plastic-eating mealworms. 2. role of gut microorganisms. *Environ Sci Technol* 49:12087–12093
58. Zehr JP, Jenkins BD, Short SM, Steward GF (2003) Nitrogenase gene diversity and microbial community structure: a crosssystem comparison. *Environ Microbiol* 5:539–554

Figures

Figure 1

Evidence for nitrogen fixation in plastic-eating mealworms. (a) Nitrogen-fixation activity measured by acetylene reduction assays for normal Styrofoam-eating mealworms or antibiotic-treated mealworms from six different sources. (b) $\delta^{15}\text{N}$ values [N isotopic ratio, per mil (‰)] of normal Styrofoam-eating mealworms and antibiotic-treated mealworms from six different sources. All results are shown as mean $\pm$ standard deviation (SD) ($n = 3$ samples, 10 worms as a sample). The single asterisk and double asterisks indicate significant different with $p < 0.05$ and $p < 0.01$, respectively (t-test).

Figure 2

Nitrogen-fixing bacteria isolated from mealworm's gut. (a) The 16S rRNA gene-based maximum-likelihood phylogenetic tree of nitrogen-fixing bacteria isolated from the gut of plastic-eating mealworms. The 16S rRNA sequence of *Azospirillum brasilense* ATCC 29145 was used as outgroup. Scale bar indicates 2% estimated sequence divergence. Bootstrap support values ≥ 70 % are indicated on nodes. (b) Nitrogen-fixing activity measured by acetylene reduction assays for bacterial isolates obtained from the gut of mealworms. *Azospirillum brasilense* ATCC 29145, a known free-living nitrogen-fixing bacterium, was used as a positive control. Results are shown as mean $\pm$ standard deviation (SD) (n = 3).

Figure 3

PCR and RT-PCR amplification of *nifH* DNA and mRNA from the gut microbiota of plastic-eating mealworms. (a) PCR detections of *nifH* gene in the DNA extracts from the gut microbiota of plastic-eating mealworms from six different sources. (b) PCR products of *nifH* gene from the genomic DNA of potential nitrogen-fixing gut isolates. (c) PCR detections of reverse-transcribed *nifH* mRNA in the RNA extracts from the gut microbiota of plastic-eating mealworms from six different sources. The *nifH*-specific primers PolF-PolR was used in the PCR reaction. RNA preparations treated with DNase in the absence of RT did not show the presence of PCR products, indicating there no DNA contaminant in the RNA extracts.

Figure 4

nifH phylogenetic tree. Phylogenies were reconstructed from *nifH* amino acid sequences recovered from DNA (PCR), RNA (RT-PCR) and isolated cultures from the gut microbiota of plastic-eating mealworms. All gut phylotypes are indicated in red. The tree was constructed as described in the Materials and methods section.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SI.pdf](#)