

# Maternal Folic Acid Impacts DNA Methylation Profile in Male Rat Offspring Associated With Neurodevelopment and Learning/Memory Abilities

Xinyan Wang

Tianjin Medical University

Zhenshu Li

Tianjin Medical University

Yun Zhu

Tianjin Medical University

Jing Yan

Tianjin Medical University

Huan Liu

Tianjin Medical University

Guowei Huang

Tianjin Medical University

Wen Li (✉ [liwen828@tmu.edu.cn](mailto:liwen828@tmu.edu.cn))

Tianjin Medical University <https://orcid.org/0000-0002-9268-538X>

---

## Research

**Keywords:** folic acid, pregnancy, DNA methylation, learning/memory, synaptic plasticity

**Posted Date:** August 24th, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-62848/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published on January 11th, 2021. See the published version at <https://doi.org/10.1186/s12263-020-00681-1>.

## Abstract

**Background:** Periconceptional folic acid (FA) supplementation not only reduces the incidence of neural tube defects, but also improves cognitive performances in offspring. However, the genes or pathways that are epigenetically regulated by FA in neurodevelopment were rarely reported.

**Methods:** To elucidate the underlying mechanism, the effect of FA on the methylation profiles in brain tissue of male rat offspring was assessed by methylated DNA immunoprecipitation chip. Differentially methylated genes (DMGs) and gene network analysis were identified using DAVID and KEGG pathway analysis.

**Results:** Compared with the FA-N group, 1939 DMGs were identified in the FA-D group, and 1498 DMGs were identified in the FA-S group, among which 298 DMGs were overlapped. The pathways associated with neurodevelopment and learning/memory abilities were differentially methylated in response to maternal FA intake during pregnancy, and there were some identical and distinctive potential mechanisms under FA deficiency or supplemented conditions.

**Conclusions:** In conclusion, genes and pathways associated with neurodevelopment and learning/memory abilities were differentially methylated in male rat offspring in response to maternal FA deficiency or supplementation during pregnancy.

## Introduction

Emerging evidence has indicated that early life nutrition influences brain development, and exerts long-term consequences for memory abilities profoundly [1,2]. Periconceptional folic acid (FA) supplementation not only reduces the incidence of neural tube defects, but also improves cognitive performances in offspring [3,4]. Our previous studies demonstrated that maternal FA supplementation during pregnancy promoted offspring's neurogenesis and synaptogenesis at birth [5], stimulated sensory-motor reflex development in infancy [6,7], and exerted long-term beneficial effects on learning and memory abilities in adolescence and adult in rat offspring [6]. However, the genes or pathways that are epigenetically regulated by FA in neurodevelopment were rarely reported.

Through one-carbon metabolism, folate provides a labile source of methyl groups for numerous methylation reactions, including DNA methylation [8]. DNA methylation and histone modification are the most studied epigenetic modifications that are reprogrammed considerably in early embryonic development [9]. Epigenetic alterations are thought to module gene expression and silencing [10], and its dysregulation may contribute to numerous neurodevelopment disorders [11,12]. Maternal FA supplementation during pregnancy increased offspring's global DNA methylation level in rats [7], and influenced offspring's repeat element and imprinted gene methylation in human [13]. A meta-analysis included 1988 newborns from two European birth cohorts revealed that maternal plasma folate during pregnancy impacted epigenome-wide DNA methylation in cord blood, and some of the implicated genes had functional relevance to neurodevelopment [4]. Lambrot et al. [10] provided evidence in mice that

lifetime exposure to folate-deficiency may cause altered sperm epigenome and was associated with negative reproductive outcomes. Transient inhibition of DNA methylation by 5-Azacytidine treatment during development could induce neurodegeneration in neonatal mice, and long-lasting deficits in synaptic plasticity and memory abilities in adult mice [14]. Thereby DNA methylation modification may play critical roles in neurodevelopment and learning/memory ability in offspring.

In the present study, male offspring derived from dams fed with FA-deficient, FA-normal and FA-supplemented diets throughout pregnancy were sacrificed within 24 h after birth to detect DNA methylation profiles. We hypothesized that maternal FA intake during pregnancy might alter DNA methylation profiles established in utero that subsequently regulated neurodevelopment and long-term learning/memory abilities in male offspring.

## Methods

### Rats and dietary treatment

Three-month-old female Sprague-Dawley rats (Charles River Laboratories, China) were assigned randomly into three dietary groups: (1) FA-deficient diet (FA-D), (2) FA-normal diet (FA-N), and (3) FA-supplemented diet (FA-S). Dietary treatment began from 1 week before mating and throughout the end of pregnancy. Rats were housed in a specific pathogen-free facility under a controlled 12-h light/dark cycle and were provided with food and water ad libitum. All animal procedures were approved by the Tianjin Medical University Animal Ethics Committee (TMUaEC2015001).

Diet compositions were modified from AIN-93G diet (TestDiet, USA), which contains 2.1 mg FA/kg diet. The FA-D and FA-S diet contains 0.1 mg FA/kg diet and 3.5 mg FA/kg diet, respectively. The neonatal male offspring derived from 3 dams were sacrificed within 24 h after birth, and cerebral hemispheres were collected and stored at -80°C after liquid nitrogen flash-freezing for further methylated DNA immunoprecipitation (MeDIP) chip assay.

### MeDIP-Chip assay

Genomic DNA was extracted from brain tissue using a DNeasy Blood & Tissue Kit (Qiagen, USA). The purified DNA was quantified using nanodrop ND-1000. Then the genomic DNA was sonicated (Bioruptor sonicator, Diagenode, Belgian) to fragments between 200 and 1000 bp. The MeDIP was prepared as follows: 1 µg of sonicated genomic DNA fragments were denatured for 10 min at 94 °C, immunoprecipitated with mouse anti-5-methylcytosine antibody (1:400, Diagenode, Belgian) overnight at 4 °C, and incubated with anti-mouse IgG magnetic beads for 2 hours at 4 °C. After washing, the beads-DNA complex was resuspended in TE buffer with 0.25% SDS and 0.25 mg/mL proteinase K for 2 h at 65 °C. Lastly, the MeDIP DNA was purified using Qiagen MinElute columns (Qiagen, USA).

MeDIP DNA samples and Input samples were labeled with Cy5-9mer and Cy3-9mer primer, respectively, using a NimbleGen Dual-Color DNA Labeling Kit according to the manufacturer's instruction (Nimblegen

Systems, Inc., USA). Then the samples were hybridized to the Arraystar Rat RefSeq Promoter Array, which is designed to investigate the epigenetic modifications and transcription factor binding sites within 15,987 RefSeq Gene promoter regions totally covered by approximately 180,000 probes. Scanning was performed with the Agilent Scanner G2505C microarray scanner.

### **Identification of differentially methylated regions**

The  $\log_2$  (MeDIP/Input) getting from raw data was normalized using the Median-centering, quantile normalization and linear smoothing analysis by Bioconductor packages Ringo, limma, and MEDME. Then from the normalized  $\log_2$  (MeDIP/Input) data, a sliding-window peak-finding algorithm provided by NimbleScan v2.5 (Roche-NimbleGen) was applied to find the enriched peaks with specified parameters (sliding window width: 1500 bp; minimum probes per peak: 2; *P*-value minimum cut-off (-log10): 2), and peaks within 500 bp spacing were merged.

To compare the differentially enriched regions between two groups, the  $\log_2$  (MeDIP/Input) was averaged for each group (Experiment and Control), and the *M'* value for each probe was calculated as following:

$$M' = \text{Average}(\log_2 \text{MeDIP}_E/\text{Input}_E) - \text{Average}(\log_2 \text{MeDIP}_C/\text{Input}_C)$$

Then the NimbleScan sliding-window peak-finding algorithm was reran on these data to find the differential enrichment peaks (DEPs) between groups, filtered according to the following criteria:

- 1) At least one of the two groups had a median ( $\log_2$  MeDIP/Input)  $\geq 0.3$  and a median (*M*)  $> 0$ .
- 2) At least half of probes in a peak might have coefficient of variability  $\leq 0.8$  in both two groups.

### **Promoter classification**

Promoters were defined as the regions [-1300 bp, +500 bp] around the transcription start sites (TSS). Promoters were classified into 3 categories as follows:

- 1) High CpG-density promoter (HCP): promoters containing a 500-bp window with GC content  $\geq 55\%$ , and CpG observed to expected ratio (O/E)  $\geq 0.6$  within -700 bp to +200 bp around TSS.
- 2) Low CpG-density promoter (LCP): promoters that contained no 500 bp interval and with a CpG O/E  $\geq 0.4$ .
- 3) Intermediate CpG-density promoter (ICP): remaining promoters that were neither HCP nor LCP.

### **Functional analysis**

Genes containing DEPs in the promoter were considered as the differentially methylated genes (DMGs). Enrichments in biological process of gene ontology (GO) and pathways that have functional relevant to

neurodevelopment and learning/memory abilities were analyzed using DAVID Bioinformatics Resources 6.8 [15] and KEGG pathway analysis [16], respectively.

## Results

### **Maternal FA intake during pregnancy altered DNA methylation profiles in the brain tissue of male offspring**

To investigate the DNA methylation profiles altered by maternal FA intake during pregnancy, DEPs in promoter region between groups were identified using the NimbleScan v2.5 software (Roche-NimbleGen). Compared with the FA-N group, 1843 DEPs were identified in the FA-D group (744 hypermethylated and 1099 hypomethylated), and 1406 DEPs were identified in the FA-S group (738 hypermethylated and 668 hypomethylated).

Since one single peak can be assigned to more than one gene promoter, and one promoter may contain more than one peak, the peaks-to-promoters are not always one-to-one correspondence. The DEPs were further analyzed according to its assigned gene promoter. Compared FA-D group with FA-N group (FA-D/FA-N), there are 996, 438, and 433 DEPs located in HCP, ICP, and LCP, respectively. While there were 699, 392, and 337 DEPs located, representatively, in the HCP, ICP and LCP in the FA-S group compared with the FA-N group (FA-S/FA-N). Those DEPs constituted 1939 DMGs in FA-D/FA-N, and 1498 DMGs in FA-S/FA-N, among which 298 DMGs were overlapped.

Taken together, the results indicated that maternal FA intake during pregnancy altered DNA methylation profiles in brain tissue of neonatal male offspring, and DEPs localized mostly (approximately 50%) in HCP.

### **Maternal FA intake during pregnancy altered biological process involved in neurodevelopment and learning/memory abilities in the brain tissue of male offspring**

We identified some identical and distinctive DMGs altered in response to maternal FA deficiency and supplementation during pregnancy. For this reason, data of the maternal FA deficiency (FA-D) or supplementation group (FA-S) are compared with FA normal group (FA-N) separately. Go analysis revealed that DMGs in FA-D/FA-N were enriched in a wide variety of biological process that involved in neural tube development, neural crest cell migration, brain development, neuron differentiation, glia cell differentiation, synapse organization, chemical synaptic transmission, and learning/memory (Table 1). And DMGs in FA-S/FA-N were enriched in biological process that involved in neural tube development, brain development, neuron differentiation, synapse organization, myelination, chemical synaptic transmission, and memory (Table 2).

Although some biological process involved in neurodevelopment and learning/memory abilities are shared by different FA status, compared with FA-S/FA-N, more DMGs implicated in brain development and neuron differentiation were found in FA-D/FA-N. Additionally, DMGs in FA-D/FA-N distinctively

enriched in neural crest cell migration, glia cell differentiation; while DMGs in FA-S/FA-N distinctively enriched in myelination (Table 1, 2).

Taken together, the results indicated that gene implicated in neurodevelopment and learning/memory abilities were differentially methylated in response to maternal FA intake during pregnancy, and there were some identical and distinctive biological process altered under FA deficiency and supplemented conditions.

### **Maternal FA intake during pregnancy impacting pathways associated with neurodevelopment of male offspring**

To further discover the potential mechanisms whereby maternal FA intake during pregnancy impacts neurodevelopment through DNA methylation, KEGG pathway analysis was performed on DMGs to identify the enriched pathways associated with neurodevelopment in FA-D/FA-N and FA-S/FA-N. The results showed that DMGs altered by maternal FA deficiency were enriched in pathways including signaling pathways regulating pluripotency of stem cells, autophagy, tight junction, calcium signaling pathway, mammalian/mechanistic target of rapamycin (mTOR) signaling pathway, Notch signaling pathway, neurotrophin signaling pathway, and protein processing in endoplasmic reticulum (ER) (FA-D/FA-N in Table 3). While DMGs altered by maternal FA supplementation were enriched in pathways including signaling pathways regulating pluripotency of stem cells, autophagy, tight junction, calcium signaling pathway, mitogen-activated protein kinase (MAPK) signaling pathway, transforming growth factor-beta (TGF- $\beta$ ) signaling pathway, and axon guidance (FA-S/FA-N in Table 3).

The results showed that signaling pathways regulating pluripotency of stem cells, autophagy, tight junction, and calcium signaling pathway were overlapped between FA-D/FA-N and FA-S/FA-N, indicating the mechanisms that shared from the FA-deficient status to the FA-normal status and to the FA-supplemented status. Additionally, DMGs in FA-D/FA-N enriched distinctively in mTOR signaling pathway, Notch signaling pathway, neurotrophin signaling pathway, and protein processing in ER; while DMGs in FA-S/FA-N enriched distinctively in MAPK signaling pathway, TGF- $\beta$  signaling pathway and axon guidance (Table 3), indicating distinctive mechanisms of influencing neurodevelopment in male offspring under FA deficient or supplemented conditions.

Taken together, the results indicated that pathways associated with neurodevelopment were differentially methylated in response to maternal FA intake during pregnancy, and there were some identical and distinctive potential mechanisms impacting neurodevelopment under FA deficiency or supplemented conditions.

### **Maternal FA intake during pregnancy impacting pathways associated with learning/memory abilities of male offspring**

To illustrate the potential mechanisms of maternal FA during pregnancy impacting offspring's learning/memory abilities by DNA methylation, enriched pathways associated with synaptic plasticity and learning/memory abilities were identified using KEGG pathway analysis. The results showed that

DMGs altered by maternal FA deficiency were enriched in pathways including cGMP-PKG signaling pathway, gap junction, calcium signaling pathway, mTOR signaling pathway, Rap1 signaling pathway, Notch signaling pathway, endocytosis, and neurotrophin signaling pathway (FA-D/FA-N in Table 4). While DMGs altered by maternal FA supplementation were enriched in pathways including cGMP-PKG signaling pathway, gap junction, calcium signaling pathway, MAPK signaling pathway, ErbB signaling pathway, glutamatergic synapse, and neuroactive ligand-receptor interaction (FA-S/FA-N in Table 4).

The results showed that cGMP-PKG signaling pathway, gap junction and calcium signaling pathway were overlapped between FA-D/FA-N and FA-S/FA-N, indicating the mechanisms that shared from the FA-deficient status to the FA-normal status and to the FA-supplemented status. Additionally, DMGs in FA-D/FA-N enriched distinctively in mTOR signaling pathway, Rap1 signaling pathway, Notch signaling pathway, endocytosis and neurotrophin signaling pathway; while DMGs in FA-S/FA-N enriched distinctively in MAPK signaling pathway, ErbB signaling pathway, glutamatergic synapse, and neuroactive ligand-receptor interaction (Table 4), indicating the distinctive mechanisms of influencing learning/memory abilities in male offspring under maternal FA deficient or supplemented conditions.

Taken together, the results indicated that pathways associated with learning/memory abilities were differentially methylated in response to maternal FA intake during pregnancy, and there some identical and distinctive potential mechanisms impacting learning/memory abilities under FA deficiency or supplemented conditions.

## Discussion

In present study, the results showed that maternal FA intake during pregnancy altered DNA methylation profiles in brain tissue of neonatal male offspring, and pathways associated with neurodevelopment and learning/memory were differentially methylated. DMGs altered by maternal FA deficiency (FA-D/FA-N) were enriched in eight pathways associated with neurodevelopment and eight pathways associated with learning/memory abilities. While DMGs altered by maternal FA supplementation (FA-S/FA-N) were enriched in seven pathways associated with neurodevelopment and seven pathways associated with learning/memory abilities. There were some identical and distinctive potential mechanisms impacting neurodevelopment or learning/memory abilities under FA deficiency or supplemented conditions.

### Potential mechanisms of maternal FA intake during pregnancy impacting offspring's neurodevelopment through DNA methylation

DNA methyltransferases (DNMT) 3a-null mice showed impairments in postnatal neurogenesis [17], and DNMT3b knockdown in human embryonic stem cells altered the timing of neuronal differentiation and maturation [18], supporting vital roles of DNA methylation in neurodevelopment. Consistently, the results in present study illustrated that the DMGs altered by maternal FA deficiency and supplementation were both enriched in pathways of signaling pathways regulating pluripotency of stem cells, autophagy, tight junction, and calcium signaling pathway, all of which are associated with neurodevelopment. Signaling pathways regulating pluripotency of stem cells activates a core transcriptional network including Oct4,

Nanog and Sox2. These three transcription factors coordinate with their downstream target genes promotes self-renewal and pluripotency of stem cells, thereby impacting embryonic development and neurodevelopment. Autophagy, a self-degradative process that acts as a pro-survival or pro-death mechanism depending on different physiological or pathological conditions [19], is important for embryonic development, neural proliferation and differentiation during brain development [20,21]. Tight junctions are essential for establishing the selectively permeable barrier between neighboring cells, tight junctions formed between brain capillary endothelial cells help creating blood-brain barrier that act to protect the brain from harmful substances and maintain brain homeostasis, modified tight junction protein expression and disrupted blood-brain barrier structure are observed in cerebral pathologies [22,23]. Calcium signaling regulates nearly every aspect of neural development, including neural induction, neural rosettes formation, neural progenitor cells proliferation, neuronal migration and differentiation [24-26], specific synaptic connections development, and impairments in calcium-dependent network maturation are associated with neurodevelopment disorders [27].

Additionally, DMGs altered by maternal FA deficiency were enriched distinctively in mTOR signaling pathway, Notch signaling pathway, neurotrophin signaling pathway and protein processing in ER. The mTOR signaling plays vital roles in dendritic formation and axon guidance during normal brain development [28], and aberrant mTOR signaling has been implicated in neurodevelopment disorders that characterized by cognitive deficits, such as tuberous sclerosis and fragile X syndrome [29]. Notch signaling maintains the neural stem cells pool during brain development [30], regulates cortical cell-type differentiation in an orderly progression [31], and interacts with Reelin signaling regulating cortical neuronal migration [32]. Neurotrophins are growth factors consisting of nerve growth factor, brain derived neurotrophic factor, neurotrophin (NT) 3, and NT-4, that participated in a wide aspects of neurodevelopment, including neural survival, proliferation, differentiation, myelination, apoptosis, and axonal growth [33]. The ER is a subcellular organelle where proteins are folded with the help of luminal chaperones. Proteins that correctly folded are packaged into transport vesicles that shuttle to Golgi complex; while accumulation of unfolded or misfolded proteins in the ER lumen triggers ER stress and unfolded protein response that may induce aberrant neuronal differentiation and inhibition of dendrite outgrowth [34].

While DMGs altered by maternal FA supplementation were enriched distinctively in MAPK signaling pathway, TGF- $\beta$  signaling pathway and axon guidance. MAPKs have been involved in modulation of hippocampal development. Extracellular signal-regulated kinase (ERK) signaling regulates maintenance of neural progenitor cells in dentate gyrus [35]. ERK and p38 activities are dynamically regulated during postnatal development that may correlates with the occurrence of hippocampal programmed cell death and synaptogenesis [36]. TGF- $\beta$  signaling has been implicated in axonal formation and specification, and neuronal migration during brain development [37,38]. Axon guidance including axon outgrowth, axon repulsion and axon attraction, represent key stages in the formation of neuronal network.

## Potential mechanisms of maternal FA intake during pregnancy impacting offspring's long-term learning/memory abilities through DNA methylation

DNA methylation on cytosine residues in CpG sequences is a mitotically stable epigenetic modification that can produce long-term changes in gene expression [11,39], and transient inhibition of DNA methylation during development exerts long-term deleterious effects on synaptic plasticity and memory abilities in adult mice [14]. Consistently, the results in present study showed that the DMGs altered by maternal FA deficient and supplementation were both enriched in pathways of cGMP-PKG signaling pathway, gap junction and calcium signaling pathway. cGMP-PKG signaling mediates early memory consolidation and early-phase LTP [40], in addition, NO-cGMP-PKG signaling pathway regulates fear memory consolidation by promoting both presynaptic and postsynaptic alterations following fear conditioning at lateral amygdala synapses via NO-driven "retrograde signaling" [41]. Gap junctions in mammalian brain act to synchronize neuronal activity and connect glial cells participating in the regulation of brain metabolism and homeostasis, and suggested to contribute to fear learning and memory [42,43]. Postsynaptic calcium signaling is suggested to induce activity-dependent LTP in CA1 and CA2 hippocampal neurons; calcium/calmodulin may activates many plasticity-inducing pathways, such as CaMK, Ras/ERK, and PKA [44], and astroglial calcium signaling displays short-term plasticity and down-regulates synaptic transmission [45].

Additionally, DMGs altered by maternal FA deficiency were enriched distinctively in mTOR signaling pathway, Rap1 signaling pathway, Notch signaling pathway, endocytosis and neurotrophin signaling pathway. mTOR regulates protein translation in response to neuronal activity, thereby modulating synaptic plasticity and long-term memory formation, and has been implicated in fear memories acquisition/consolidation in the hippocampus/amygdala [46]. Small GTPase Rap1 signals synaptic glutamate receptor (GluR)-2/3 AMPARs removes by activating Rap1-p38MAPK signaling during LTD [47]. Notch signaling plays a critical role in brain development. Recently, emerging evidence has suggested that Notch signaling also plays an important role in synaptic plasticity and spatial memory formation [48,49]. Neurotrophins are firstly described as growth factors that support neuronal differentiation and survival. Later studies have indicated their roles in synapse formation and plasticity, including increasing neurotransmitter output, membrane excitability, and synaptic contacting size locally at synapse, and affecting gene expression and phenotypic enhancement distally at the neuronal soma [50].

While DMGs altered by maternal FA supplementation were enriched distinctively in MAPK signaling pathway, ErbB signaling pathway, Glutamatergic synapse, and Neuroactive ligand-receptor interaction. MAPK is activated after learning or LTP induction in hippocampus and amygdala, and pharmacological inhibition of MAPK activation impairs LTP, learning, and long-term memory [51]. ErbB signaling modulates hippocampal metabotropic GluRI-dependent LTD and object recognition memory [52], and pharmacological inhibition of ErbB signaling during adolescence impairs reference memory in adult mice [53]. In glutamatergic synapse, glutamate released from neuron presynaptic terminal acts on postsynaptic ionotropic or metabotropic GluRs to regulate synaptic plasticity, and afterwards, are removed from the synaptic cleft by EAATs located either on the presynaptic terminal, postsynaptic neuron, or neighboring glial cells. Signals resulting from ligand-receptor interactions trigger intracellular signaling that ultimately modulate gene expression, which could be the underlying basis of learning and memory [54].

# **Conclusion**

In conclusion, the present study found that maternal FA deficiency and supplementation during pregnancy altered DNA methylation profiles in brain tissue of male offspring, and those DMGs were associated with neurodevelopment and learning/memory abilities. These findings provide novel insights into the mechanism studies of maternal FA intake impacting offspring's neurodevelopment and learning/memory abilities with DNA methylation involved.

## **Abbreviations**

DEPs, differential enrichment peaks; DMGs: differentially methylated genes; DNMT, DNA methyltransferases; ER, endoplasmic reticulum; FA, folic acid; GluR, glutamate receptor; GO, gene ontology; HCP, high-CpG-density promoter; ICP, intermediate-CpG-density promoter; LCP, low-CpG-density promoter; LTD, long-term depression; LTP, long-term potentiation; MAPK, mitogen-activated protein kinase; MeDIP, methylated DNA immunoprecipitation; mTOR, mammalian/mechanistic target of rapamycin; TGF, transforming growth factor; TSSs, transcription start sites

## **Declarations**

### **Acknowledgment**

We would like to thank all the participants in the trial.

### **Author Contributions**

X.W. and W.L. conducted data curation; X.W. wrote the original draft; X.W. and Z.L. analyzed the data; X.W., Y.Z. and J.Y. contributed methodology; H.L. contributed software; G.H. and W.L. reviewed and edited the draft; G.H. and W.L. contributed funding acquisition. All authors have read and agreed to the published version of the manuscript.

### **Funding**

This work was supported by the National Natural Science Foundation of China (No. 81602849, 81472967, 81730091), Natural Science Foundation of Tianjin City (No. 19JCQNJC11700), and young elite scientist's sponsorship program by CAST (No. 2017QNRC001).

### **Availability of data and materials**

All of the data are available with reasonable request from the corresponding author.

### **Ethics approval and consent to participate**

All animal procedures were approved by the Tianjin Medical University Animal Ethics Committee (TMUaEC2015001).

### **Consent for publication**

Not applicable

### **Conflicts of interest**

The authors declare that they have no conflicts of interest.

### **Author details**

<sup>1</sup> Department of Nutrition and Food Science, School of Public Health, Tianjin Medical University, Tianjin 300070, China.

<sup>2</sup> Department of Epidemiology and Biostatistics, School of Public Health, Tianjin Medical University, Tianjin 300070, China.

<sup>3</sup> Department of Social Medicine and Health Administration, School of Public Health, Tianjin Medical University, Tianjin 300070, China.

<sup>4</sup> Tianjin Key Laboratory of Environment, Nutrition and Public Health, Center for International Collaborative Research on Environment, Nutrition and Public Health, Tianjin 300070, China

## **References**

1. Queiroz MP, Lima MDS, Barbosa MQ, de Melo M, Bertozzo C, de Oliveira MEG, Bessa RJB, Alves SPA, Souza MIA, Queiroga R, Soares JKB (2019) Effect of Conjugated Linoleic Acid on Memory and Reflex Maturation in Rats Treated During Early Life. *Frontiers in neuroscience* 13:370. doi:10.3389/fnins.2019.00370
2. Gould JM, Smith PJ, Airey CJ, Mort EJ, Airey LE, Warricker FDM, Pearson-Farr JE, Weston EC, Gould PJW, Semmence OG, Restall KL, Watts JA, McHugh PC, Smith SJ, Dewing JM, Fleming TP, Willaime-Morawek S (2018) Mouse maternal protein restriction during preimplantation alone permanently alters brain neuron proportion and adult short-term memory. *PNAS* 115 (31):E7398-e7407. doi:10.1073/pnas.1721876115
3. McGarel C, Pentieva K, Strain JJ, McNulty H (2015) Emerging roles for folate and related B-vitamins in brain health across the lifecycle. *The Proceedings of the Nutrition Society* 74 (1):46-55. doi:10.1017/S0029665114001554
4. Joubert BR, den Dekker HT, Felix JF, Bohlin J, Ligthart S, Beckett E, Tiemeier H (2016) Maternal plasma folate impacts differential DNA methylation in an epigenome-wide meta-analysis of newborns. *Nature communications* 7:10577. doi:10.1038/ncomms10577

5. Wang X, Li W, Li Z, Ma Y, Yan J, Wilson JX, Huang G (2018) Maternal Folic Acid Supplementation During Pregnancy Promotes Neurogenesis and Synaptogenesis in Neonatal Rat Offspring. *Cerebral cortex* (New York, NY : 1991). doi:10.1093/cercor/bhy207
6. Wang X, Li W, Li S, Yan J, Wilson JX, Huang G (2018) Maternal Folic Acid Supplementation During Pregnancy Improves Neurobehavioral Development in Rat Offspring. *Molecular neurobiology* 55 (3):2676-2684. doi:10.1007/s12035-017-0534-2
7. Li W, Li Z, Li S, Wang X, Wilson JX, Huang G (2018) Periconceptional Folic Acid Supplementation Benefit to Development of Early Sensory-Motor Function through Increase DNA Methylation in Rat Offspring. *Nutrients* 10 (3). doi:10.3390/nu10030292
8. Lan X, Field MS, Stover PJ (2018) Cell cycle regulation of folate-mediated one-carbon metabolism. 10 (6):e1426. doi:10.1002/wsbm.1426
9. Zhao P, Wang H, Wang H (2019) Essential roles of HDAC1 and 2 in lineage development and genome-wide DNA methylation during mouse preimplantation development. 1-17. doi:10.1080/15592294.2019.1669375
10. Lambrot R, Xu C, Saint-Phar S, Chountalos G, Cohen T, Paquet M, Suderman M, Hallett M, Kimmins S (2013) Low paternal dietary folate alters the mouse sperm epigenome and is associated with negative pregnancy outcomes. *Nature communications* 4:2889. doi:10.1038/ncomms3889
11. Mastrototaro G, Zaghi M, Sessa A (2017) Epigenetic Mistakes in Neurodevelopmental Disorders. 61 (4):590-602. doi:10.1007/s12031-017-0900-6
12. Lewis EM, Kroll KL (2018) Development and disease in a dish: the epigenetics of neurodevelopmental disorders. *Epigenomics* 10 (2):219-231. doi:10.2217/epi-2017-0113
13. Haggarty P, Hoad G, Campbell DM, Horgan GW, Piyathilake C, McNeill G (2013) Folate in pregnancy and imprinted gene and repeat element methylation in the offspring. *Am J Clin Nutr* 97 (1):94-99. doi:10.3945/ajcn.112.042572
14. Subbanna S, Nagre NN, Shivakumar M, Basavarajappa BS (2016) A single day of 5-azacytidine exposure during development induces neurodegeneration in neonatal mice and neurobehavioral deficits in adult mice. *Physiology & behavior* 167:16-27. doi:10.1016/j.physbeh.2016.08.036
15. Huang da W, Sherman BT, Lempicki RA (2009) Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nature protocols* 4 (1):44-57. doi:10.1038/nprot.2008.211
16. Kanehisa M, Goto S (2000) KEGG: kyoto encyclopedia of genes and genomes. *Nucleic acids research* 28 (1):27-30. doi:10.1093/nar/28.1.27
17. Wu H, Coskun V, Tao J, Xie W, Ge W, Yoshikawa K, Li E, Zhang Y, Sun YE (2010) Dnmt3a-dependent nonpromoter DNA methylation facilitates transcription of neurogenic genes. *Science* (New York, NY) 329 (5990):444-448. doi:10.1126/science.1190485
18. Martins-Taylor K, Schroeder DI, LaSalle JM, Lalonde M, Xu RH (2012) Role of DNMT3B in the regulation of early neural and neural crest specifiers. *Epigenetics* 7 (1):71-82. doi:10.4161/epi.7.1.18750

19. Fimia GM, Stoykova A, Romagnoli A, Giunta L, Di Bartolomeo S, Nardacci R, Corazzari M, Fuoco C, Ucar A, Schwartz P, Gruss P, Piacentini M, Chowdhury K, Cecconi F (2007) Ambra1 regulates autophagy and development of the nervous system. *Nature* 447 (7148):1121-1125. doi:10.1038/nature05925
20. Lei J, Calvo P, Vigh R, Burd I (2018) Journey to the Center of the Fetal Brain: Environmental Exposures and Autophagy. *Frontiers in cellular neuroscience* 12:118. doi:10.3389/fncel.2018.00118
21. Sotthibundhu A, Promjuntuek W, Liu M, Shen S, Noisa P (2018) Roles of autophagy in controlling stem cell identity: a perspective of self-renewal and differentiation. 374 (2):205-216. doi:10.1007/s00441-018-2829-7
22. Abbott NJ, Patabendige AA, Dolman DE, Yusof SR, Begley DJ (2010) Structure and function of the blood-brain barrier. *Neurobiology of disease* 37 (1):13-25. doi:10.1016/j.nbd.2009.07.030
23. Berndt P, Winkler L, Cording J, Breitkreuz-Korff O, Rex A, Dithmer S, Rausch V, Blasig R, Richter M, Sporbert A, Wolburg H, Blasig IE, Haseloff RF (2019) Tight junction proteins at the blood-brain barrier: far more than claudin-5. 76 (10):1987-2002. doi:10.1007/s00018-019-03030-7
24. Toth AB, Shum AK, Prakriya M (2016) Regulation of neurogenesis by calcium signaling. *Cell calcium* 59 (2-3):124-134. doi:10.1016/j.ceca.2016.02.011
25. Horigane SI, Ozawa Y, Yamada H, Takemoto-Kimura S (2019) Calcium signalling: a key regulator of neuronal migration. *Journal of biochemistry* 165 (5):401-409. doi:10.1093/jb/mvz012
26. Hribkova H, Grabiec M (2018) Calcium signaling mediates five types of cell morphological changes to form neural rosettes. 131 (3). doi:10.1242/jcs.206896
27. Lohmann C (2009) Calcium signaling and the development of specific neuronal connections. *Progress in brain research* 175:443-452. doi:10.1016/s0079-6123(09)17529-5
28. Takei N, Nawa H (2014) mTOR signaling and its roles in normal and abnormal brain development. *Frontiers in molecular neuroscience* 7:28. doi:10.3389/fnmol.2014.00028
29. Xu J, Kang E, Mintz CD (2018) Anesthetics disrupt brain development via actions on the mTOR pathway. *Communicative & integrative biology* 11 (2):1-4. doi:10.1080/19420889.2018.1451719
30. Byun SH, Kwon M, Lee SM, Noh H, Yoon K (2019) PACT increases mammalian embryonic neural stem cell properties by facilitating activation of the notch signaling pathway. *Biochemical and biophysical research communications* 513 (2):392-397. doi:10.1016/j.bbrc.2019.04.010
31. Wu ZQ, Li D, Huang Y, Chen XP, Huang W, Liu CF, Zhao HQ, Xu RX, Cheng M, Schachner M, Ma QH (2017) Caspr Controls the Temporal Specification of Neural Progenitor Cells through Notch Signaling in the Developing Mouse Cerebral Cortex. *Cerebral cortex (New York, NY : 1991)* 27 (2):1369-1385. doi:10.1093/cercor/bhv318
32. Hashimoto-Torii K, Torii M, Sarkisian MR, Bartley CM, Shen J, Radtke F, Gridley T, Sestan N, Rakic P (2008) Interaction between Reelin and Notch signaling regulates neuronal migration in the cerebral cortex. *Neuron* 60 (2):273-284. doi:10.1016/j.neuron.2008.09.026
33. Arevalo JC, Wu SH (2006) Neurotrophin signaling: many exciting surprises! *Cellular and molecular life sciences : CMLS* 63 (13):1523-1537. doi:10.1007/s00018-006-6010-1

34. Kawada K, Iekumo T, Saito R, Kaneko M, Mimori S, Nomura Y, Okuma Y (2014) Aberrant neuronal differentiation and inhibition of dendrite outgrowth resulting from endoplasmic reticulum stress. *Journal of neuroscience research* 92 (9):1122-1133. doi:10.1002/jnr.23389
35. Vithayathil J, Pucilowska J (2015) Dentate Gyrus Development Requires ERK Activity to Maintain Progenitor Population and MAPK Pathway Feedback Regulation. 35 (17):6836-6848. doi:10.1523/jneurosci.4196-14.2015
36. Costa AP, Lopes MW, Rieger DK, Barbosa SG, Goncalves FM, Xikota JC, Walz R, Leal RB (2016) Differential Activation of Mitogen-Activated Protein Kinases, ERK 1/2, p38(MAPK) and JNK p54/p46 During Postnatal Development of Rat Hippocampus. *Neurochemical research* 41 (5):1160-1169. doi:10.1007/s11064-015-1810-z
37. Zhang F, Yu J, Yang T, Xu D, Chi Z, Xia Y, Xu Z (2016) A Novel c-Jun N-terminal Kinase (JNK) Signaling Complex Involved in Neuronal Migration during Brain Development. *The Journal of biological chemistry* 291 (22):11466-11475. doi:10.1074/jbc.M116.716811
38. Yi JJ, Barnes AP, Hand R, Polleux F, Ehlers MD (2010) TGF-beta signaling specifies axons during brain development. *Cell* 142 (1):144-157. doi:10.1016/j.cell.2010.06.010
39. del Rosario MC, Ossowski V, Knowler WC, Bogardus C, Baier LJ, Hanson RL (2014) Potential epigenetic dysregulation of genes associated with MODY and type 2 diabetes in humans exposed to a diabetic intrauterine environment: an analysis of genome-wide DNA methylation. *Metabolism: clinical and experimental* 63 (5):654-660. doi:10.1016/j.metabol.2014.01.007
40. Bollen E, Puzzo D, Rutten K, Privitera L, De Vry J, Vanmierlo T, Kenis G, Palmeri A, D'Hooge R, Balschun D, Steinbusch HM, Blokland A, Prickaerts J (2014) Improved long-term memory via enhancing cGMP-PKG signaling requires cAMP-PKA signaling. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology* 39 (11):2497-2505. doi:10.1038/npp.2014.106
41. Ota KT, Monsey MS, Wu MS, Schafe GE (2010) Synaptic plasticity and NO-cGMP-PKG signaling regulate pre- and postsynaptic alterations at rat lateral amygdala synapses following fear conditioning. *PloS one* 5 (6):e11236. doi:10.1371/journal.pone.0011236
42. Juszczak GR, Swiergiel AH (2009) Properties of gap junction blockers and their behavioural, cognitive and electrophysiological effects: animal and human studies. *Progress in neuropsychopharmacology & biological psychiatry* 33 (2):181-198. doi:10.1016/j.pnpbp.2008.12.014
43. Bissiere S, Zelikowsky M, Ponnusamy R, Jacobs NS, Blair HT, Fanselow MS (2011) Electrical synapses control hippocampal contributions to fear learning and memory. *Science (New York, NY)* 331 (6013):87-91. doi:10.1126/science.1193785
44. Evans PR, Parra-Bueno P, Smirnov MS, Lustberg DJ (2018) RGS14 Restricts Plasticity in Hippocampal CA2 by Limiting Postsynaptic Calcium Signaling. 5 (3). doi:10.1523/eneuro.0353-17.2018
45. Sibille J, Zapata J, Teillon J, Rouach N (2015) Astroglial calcium signaling displays short-term plasticity and adjusts synaptic efficacy. *Frontiers in cellular neuroscience* 9:189.

doi:10.3389/fncel.2015.00189

46. Giovannini MG, Lana D, Pepeu G (2015) The integrated role of ACh, ERK and mTOR in the mechanisms of hippocampal inhibitory avoidance memory. *Neurobiology of learning and memory* 119:18-33. doi:10.1016/j.nlm.2014.12.014
47. Stornetta RL, Zhu JJ (2011) Ras and Rap signaling in synaptic plasticity and mental disorders. *The Neuroscientist : a review journal bringing neurobiology, neurology and psychiatry* 17 (1):54-78. doi:10.1177/1073858410365562
48. Sargin D, Botly LC, Higgs G, Marsolais A, Frankland PW, Egan SE, Josselyn SA (2013) Reprint of: disrupting Jagged1-Notch signaling impairs spatial memory formation in adult mice. *Neurobiology of learning and memory* 105:20-30. doi:10.1016/j.nlm.2013.07.001
49. Alberi L, Liu S, Wang Y, Badie R, Smith-Hicks C, Wu J, Pierfelice TJ, Abazyan B, Mattson MP, Kuhl D, Pletnikov M, Worley PF, Gaiano N (2011) Activity-induced Notch signaling in neurons requires Arc/Arg3.1 and is essential for synaptic plasticity in hippocampal networks. *Neuron* 69 (3):437-444. doi:10.1016/j.neuron.2011.01.004
50. Lim KC, Lim ST, Federoff HJ (2003) Neurotrophin secretory pathways and synaptic plasticity. *Neurobiology of aging* 24 (8):1135-1145. doi:10.1016/j.neurobiolaging.2003.09.003
51. Ryu HH, Lee YS (2016) Cell type-specific roles of RAS-MAPK signaling in learning and memory: Implications in neurodevelopmental disorders. *Neurobiology of learning and memory* 135:13-21. doi:10.1016/j.nlm.2016.06.006
52. Ledonne A, Mango D, Latagliata EC, Chiacchierini G, Nobili A, Nistico R, D'Amelio M, Puglisi-Allegra S, Mercuri NB (2018) Neuregulin 1/ErbB signalling modulates hippocampal mGluRI-dependent LTD and object recognition memory. *Pharmacological research* 130:12-24. doi:10.1016/j.phrs.2018.02.003
53. Golani I, Tadmor H, Buonanno A, Kremer I, Shamir A (2014) Disruption of the ErbB signaling in adolescence increases striatal dopamine levels and affects learning and hedonic-like behavior in the adult mouse. *European neuropsychopharmacology : the journal of the European College of Neuropsychopharmacology* 24 (11):1808-1818. doi:10.1016/j.euroneuro.2014.09.011
54. Hook M, Xu F, Terenina E, Zhao W, Starlard-Davenport A, Mormede P, Jones BC, Mulligan MK, Lu L (2019) Exploring the involvement of Tac2 in the mouse hippocampal stress response through gene networking. *Gene* 696:176-185. doi:10.1016/j.gene.2019.02.013

## Tables

Table 1 Gene implicated in neurodevelopment and learning/memory abilities that were differentially methylated in FA-D/FA-N

Go Term (BP)	DMGs associated
Dorsal/ventral neural tube patterning	5 genes: Gsc, Kif3a, Psen1, Pax7, Ptch1
Neural crest cell migration	14 genes: Ret, Efnb1, Pax6, Kitlg, Isl1, Sox8, Coro1c, Sema5a, Sema6c, Sema6d, Sema7a, Gbx2, Sema4b, Sema3b
Brain development, hippocampus development, orbitofrontal cortex development, cerebellum development	59 genes: Cast, Fgfr1, Pfkfb3, Stk36, Hmgcs1, Pax6, Phlpp2, Ptprs, Lmx1a, Ogdh, Cdk5, Tp73, Igf1r, Bbs2, Casp3, Atp2b4, Akap5, Dclk2, Lhx5, Reln, Pafah1b1, Rara, Eif2b4, Fgfr2, Sstr1, Glud1, Gbx2, Sdf4, Rere, Dmrt2, Acat2, Ctnnbp2, Ace, Tmem57, Cxcr4, Ndrg4, Sharpin, Synj2, Unc5c, Irs2, Col4a1, Ntf3, Gsx2, Mdga1, Gabra5, Ak2, Dpysl2, Ager, Pcdh19, Ddit4, Ptpn11, Notch1, Psen1, Hes5, Abcb1b, Snrbp, Pygo2, Ptch1, Cln5
Neuron differentiation, neuron projection development, positive regulation of neuron projection development, negative regulation of neuron projection development, axonogenesis, negative regulation of axonogenesis, axon guidance, negative regulation of axon extension involved in axon guidance, axon regeneration, negative regulation of axon regeneration, negative regulation of dendrite development	112 genes: Wnt10a, Ret, Wnt5b, Cebpb, Sox11, Neurog1, Fgfr1, Pld2, Ptprm, Ncdn, Vapa, Rb1, Itgb1, Ager, Cdk5, Pten, Uhmkl1, Igf1r, Lingo1, Wnt7b, App, Cd44, Sema7a, Camsap1, Map4, Runx3, Arsb, Ltk, Prkci, Katnb1, Ankrd1, Kidins220, Kif3c, Fam134c, Nme2, Ndrg4, Ntrk1, Bcl11a, Ntrk2, Adamts1, Neu1, Reln, Ngfr, Scarb2, Tmem30a, Bag5, Mag, Lrp1, Fkbp4, Adcy5, Cspg4, Lrig2, Pafah1b1, Paqr3, Fgfr2, Actb, Adcy1, Atl1, Stk11, Erbb3, Nkx2-8, Lrrn3, Lrtm2, Flot1, Pax6, Rtn4r, Ptpn11, Slitrk2, Notch1, Picalm, Lmx1a, Isl1, Gas7, Ddit4, Casp3, Wnt3, Kdm2a, Sema5a, Sema6c, Sema6d, Sema4b, Sema3b, Dhfr, Folr1, Jak2, Ctnna1, Eph4, Klk8, Fat3, Cyth2, Arf6, Bag1, Psen1, Hes5, Ppia, Acsl4, Runx2, Wnt7a, Lrp4, Mbp, Ntf3, Kif5a, Efna2, Efnb1, Dpysl2, Ttc8, Gbx2, Etv1, Unc5c, B3gnt2, Kif26a, Kif26b
Astrocyte differentiation, oligodendrocyte differentiation	11 genes: Notch1, S100b, Hes5, Pax6, Ptprj, Ntrk2, Bnip3, Olig1, Sox8, Cdk5, Nkx6-1
Synapse organization	7 genes: Sncg, Wnt7b, Klk8, Lrrtm1, Lmx1a, Wnt7a, Lrp4
Synaptic transmission, glutamatergic, positive regulation of synaptic transmission, glutamatergic	13 genes: Clcn3, P2rx1, Cnih2, Grin2d, Grin1, Unc13b, Cdk5, Egfr, Cckbr, Ntrk1, Ntrk2, Reln, Ngfr
Learning or memory, memory, associative learning	29 genes: Egfr, Casp3, Sts, Psen1, S100b, Dyx1c1, Prkar1b, Ntrk1, Grin1, Pafah1b1, Pten, Abca7, Klk8, Cebpb, Paip2, Th, Igf2, Lmx1a, Itpr3, Sorcs3, Adrb1, Plk2, Reln, Chrb2, Cacna1d, Gabra5, Neurod2, Nrgn, Cdk5

The MeDIP-Chip assay was performed on brain tissue from folic acid-deficient (FA-D) and folic acid-normal (FA-N) male offspring (n=3/group). Enriched biological processes (BP) of GO terms by DAVID were used to identify the differentially methylated genes (DMGs) associated with neurodevelopment and learning/memory abilities.

Table 2 Gene implicated in neurodevelopment and learning/memory abilities that were differentially methylated in FA-S/FA-N

Go Term (BP)	DMGs associated
Neural fold formation	3 genes: Nodal, Cfl1, Cecr2
Forebrain development, lateral ventricle development, hindbrain development	16 genes: Dlc1, App, Kdm2b, Dkk1, Htra2, Sstr1, Zic5, Apaf1, Neurog3, Hspa8, Smarca4, Aqp1, Tsku, Dbi, Hoxa1, Neurod1
Neuron projection development	16 genes: Stx3, Ptprm, Ncdn, Clmn, Itgb1, Pten, App, Npy, Cdnf, Sema7a, Camsap1, Shc1, Areg, Ikbkb, Ehd1, Runx3
Synapse maturation	3 genes: Erbb4, Shank1, Pten
Myelination	9 genes: Amigo1, Dhh, Plp1, Nab2, Nab1, Nfasc, Olig2, Afg3l2, Tgfb1
Positive regulation of synaptic transmission, GABAergic, synaptic vesicle docking, long-term synaptic potentiation, long term synaptic depression	15 genes: Erbb4, Tacr1, Tac1, Prkce, Car7, Stx4, Stx3, Syde1, Slc8a2, S100b, Plk2, Dbi, Pten, Shank2, Drd5
Memory	11 genes: Pak6, Slc8a2, Cebpb, Syt4, S100b, Plk2, Slc6a4, Pten, Cacna1d, Shank2, Neto1

The MeDIP-Chip assay was performed on brain tissue from folic acid-supplemented (FA-S) and folic acid-normal (FA-N) male offspring (n=3/group). Enriched BP of GO terms by DAVID were used to identify the DMGs associated with neurodevelopment and learning/memory abilities.

Table 3 Pathways associated with neurodevelopment that were differentially methylated in brain of males

Group	Pathway	No of DMGs (%) <sup>a</sup>	P-value
FA-D/FA-N	Signaling pathways regulating pluripotency of stem cells*	25/139 (17.99)	1.05E-03
	Autophagy – animal*	23/134 (17.16)	3.07E-03
	Tight junction*	25/170 (14.71)	1.54E-02
	Calcium signaling pathway*	27/188 (14.36)	1.63E-02
	mTOR signaling pathway	26/157 (16.56)	2.86E-03
	Notch signaling pathway	10/52 (19.23)	2.09E-02
FA-S/FA-N	Neurotrophin signaling pathway	19/124 (15.32)	2.17E-02
	Protein processing in endoplasmic reticulum	23/164 (14.02)	3.23E-02
	Signaling pathways regulating pluripotency of stem cells*	18/139 (12.95)	8.27E-03
	Autophagy – animal*	18/134 (13.43)	5.67E-03
	Tight junction*	21/170 (12.35)	7.82E-03
	Calcium signaling pathway*	27/188 (14.36)	2.73E-04
	MAPK signaling pathway	37/298 (12.42)	4.51E-04
	TGF-beta signaling pathway	13/87 (14.94)	7.30E-03
	Axon guidance	19/178 (10.67)	4.35E-02

<sup>a</sup> The ratio of No. of DMGs was calculated by the number of DMGs in a given pathway divided by the total number of genes in this pathway. \* The pathways overlapped between FA-D/FA-N and FA-S/FA-N. The P-value was calculated by Fisher's exact test.

Table 4 Pathways associated with learning/memory abilities that were differentially methylated in brain of males

Group	Pathway	No of DMGs (%) <sup>a</sup>	P-value
FA-D/FA-N	cGMP-PKG signaling pathway*	29/167 (17.37)	7.83E-04
	Gap junction*	15/88 (17.05)	1.61E-02
	Calcium signaling pathway*	27/188 (14.36)	1.63E-02
	mTOR signaling pathway	26/157 (16.56)	2.86E-03
	Rap1 signaling pathway	30/217 (13.82)	1.93E-02
	Notch signaling pathway	10/52 (19.23)	2.09E-02
	Endocytosis	35/263 (13.31)	2.09E-02
	Neurotrophin signaling pathway	19/124 (15.32)	2.17E-02
FA-S/FA-N	cGMP-PKG signaling pathway*	25/167 (14.97)	2.38E-04
	Gap junction*	14/88 (15.91)	3.07E-03
	Calcium signaling pathway*	27/188 (14.36)	2.73E-04
	MAPK signaling pathway	37/298 (12.42)	4.51E-04
	ErbB signaling pathway	15/89 (16.85)	1.22E-03
	Glutamatergic synapse	14/116 (12.07)	3.20E-02
	Neuroactive ligand-receptor interaction	29/291 (9.97)	3.44E-02

<sup>a</sup> The ratio of No. of DMGs was calculated by the number of DMGs in a given pathway divided by the total number of genes in this pathway. \* The pathways overlapped between FA-D/FA-N and FA-S/FA-N groups. The P-value was calculated by Fisher's exact test.