Identification of promoter elements in the *Dolichospermum circinale* AWQC131C saxitoxin gene cluster and the experimental analysis of their use for heterologous expression.

Paul Michael D'Agostino

Technische Universitat Munchen

Bakir Al-Sinawi

University of New South Wales

Rabia Mazmouz

University of New South Wales

Julia Muenchhoff

University of New South Wales

Brett Anthony Neilan

University of Newcastle

Michelle Moffitt (✉ m.moffitt@westernsydney.edu.au)

University of Western Sydney <https://orcid.org/0000-0001-9872-3175>

Research article

Keywords: cyanobacteria, *Dolichospermum circinale*, *Anabaena circinalis*, saxitoxin, paralytic shellfish toxins, regulation, promoter, transcription, heterologous expression

Posted Date: February 3rd, 2020

DOI: <https://doi.org/10.21203/rs.2.15635/v3>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Microbiology on February 17th, 2020. See the published version at <https://doi.org/10.1186/s12866-020-1720-3>.

Abstract

Abstract Background: *Dolichospermum circinale* is a filamentous bloom-forming cyanobacterium responsible for biosynthesis of the paralytic shellfish toxins (PST), including saxitoxin. PSTs are neurotoxins and in their purified form are important analytical standards for monitoring the quality of water and seafood and biomedical research tools for studying neuronal sodium channels. More recently, PSTs have been recognised for their utility as local anaesthetics. Characterisation of the transcriptional elements within the saxitoxin (sxt) biosynthetic gene cluster (BGC) is a first step towards accessing these molecules for biotechnology. **Results:** In *D. circinale* AWQC131C the sxt BGC is transcribed from two bidirectional promoter regions encoding five individual promoters. These promoters were identified experimentally using 5' RACE and their activity assessed via coupling to a lux reporter system in *E. coli* and *Synechocystis* sp. PCC 6803. Transcription of the predicted drug/metabolite transporter (DMT) encoded by sxtPER was found to initiate from two promoters, P sxtPER1 and P sxtPER2 . In *E. coli*, strong expression of lux from P sxtP , P sxtD and sxtPER1 was observed while expression from P orf24 and P sxtPER2 was remarkably weaker. In contrast, heterologous expression in *Synechocystis* sp. PCC6803 showed that expression of lux from P sxtP , P sxtPER , and P orf24 promoters was statistically higher compared to the non-promoter control, while P sxtD showed poor activity under the described conditions. **Conclusions:** Both of the heterologous hosts investigated in this study exhibited high expression levels from three of the five sxt promoters. These results indicate that the majority of the native sxt promoters appear active in different heterologous hosts, simplifying initial cloning efforts. Therefore, heterologous expression of the sxt BGC in either *E. coli* or *Synechocystis* could be a viable first option for producing PSTs for industrial or biomedical purposes.

Background

Saxitoxin (STX) is a neurotoxin produced by cyanobacteria and dinoflagellates and is a member of the broader group of alkaloids known as the paralytic shellfish toxins (PSTs) [1]. When high concentrations of PSTs are consumed by humans, acute poisoning can lead to death due to respiratory paralysis [2-4]. Therefore, PSTs are needed as analytical standards for the monitoring and protection of commercial seafood and freshwater reservoirs, as well as for use in biomedical research. While the PSTs pose a significant public health risk and economic burden on society during algal bloom events, their scientific and pharmaceutical potential is well known [3, 4]. Purified PSTs have been a critical tool for researchers investigating neuronal sodium channels, where the toxins specifically block site 1 of voltage-gated sodium channels [5]. Under controlled administration, PSTs are potent anaesthetics, particularly in combination with other local anaesthetics [6, 7]. Further attempts to utilise STX in clinical trials are hindered by its toxicity, but more recent approaches, such as generating liposomal formulations of STX, resulted in blockage of the sciatic nerves in rats with no myotoxic, cytotoxic or neurotoxic effects [8]. It has been postulated that the same delivery could provide effective localised treatment for severe joint pain [9]. Other PSTs, such as the gonyautoxins (GTXs), also have clinical potential and have been utilised for the treatment of anal fissures and chronic tension type headaches [10-12].

Obtaining significant quantities of purified PSTs for clinical research or water quality analysis is difficult. Chemical synthesis and biocatalytic synthesis of PSTs is complex, difficult to scale up and may not produce all relevant naturally occurring isoforms [13, 14]. The most common form of obtaining purified compounds involves extraction and isolation from dinoflagellate blooms, cyanobacterial cultures or contaminated shellfish coupled with synthetic conversion to additional PSTs [15-17]. Analytical calibration standards are commercially available from the National Research Council Canada (NRC), which are obtained from extractions of PST producing dinoflagellates or contaminated shellfish, and in some instances, semi-synthetic conversions of PSTs are required to obtain a broad array of analogues. Thus, the process is difficult, inefficient and costly [18]. These issues clearly highlight the need for an alternative and reliable method for the production and purification of commercial quantities of PSTs.

Heterologous expression of cyanobacterial biosynthetic gene clusters (BGCs) is not well established, in comparison to heterologous expression of Actinomycete BGCs. *Streptomyces* expression hosts successfully produced more than 90 Actinomycete NPs, most of which are heterologously expressed using native promoters [19]. On the other hand, only 10 cyanobacterial NPs have been successfully produced from heterologous systems using both native and heterologous promoters [20], highlighting the need to better understand the function of native promoters in heterologous hosts. *E. coli* is a suitable host for the heterologous expression of cyanobacterial pathways based on its fast growth rate as previously demonstrated [21, 22]. Initial studies used native promoters to produce the ribosomal peptides patellamide A and C, and the microviridins [23, 24]. Recently, there has been a focus on the heterologous expression of cyanobacterial natural product biosynthetic gene clusters (BGC) including the lyngbyatoxin (*ltx*) BGC in *E. coli* using the tetracycline-inducible *Ptet_O* promoter [21, 25]. The *ltx* BGC has been the focus of multiple heterologous expression studies due to its relatively small size. It has also been expressed in the cyanobacterium *Anabaena* sp. PCC 7120, and *E. coli* GB05-MtaA [26, 27]. While the native *ltx* promoters were active in *Anabaena* sp. PCC 7120 and drive the production of lyngbyatoxin A, the native promoters were not active in *E. coli*. The addition of *Anabaena* sp. PCC 7120 sigma factors to the *E. coli* host also failed to induce expression of lyngbyatoxin A, suggesting that the heterologous host was unable to recognise the cyanobacterial ribosome binding sites [28]. Subsequently, titres of lyngbyatoxin A close to the native producer were achieved when using the cyanobacteria *Anabaena* sp. PCC 7120 as a heterologous host [26]. This highlights that discrepancies between cyanobacterial promoter efficiencies in different host organisms remain poorly understood. Therefore, it is beneficial to test the activity of promoters in heterologous hosts and expression vectors using reporter systems prior to cloning complex biosynthetic pathways for biotechnological applications [29].

For the most part, cyanobacterial transcription machinery is similar to that found in *E. coli*, with the main difference being the widespread absence of the -35 hexamer in cyanobacteria, which is believed to be replaced by a transcription factor binding site to initiate transcription [30]. In *E. coli*, σ^{70} is able to recognise the majority of promoters while in cyanobacteria, a range of different sigma factors have been identified [31, 32].

The saxitoxin (*sxt*) BGC has been characterised in six cyanobacterial species from the order Nostocales and one from the order Oscillatoriales [33-37]. Each *sxt* BGC encodes a 'core' set of enzymes putatively responsible for STX biosynthesis, supplemented with 'tailoring' and 'auxiliary' genes that give rise to PST analogues or perform functions after PST biosynthesis. Information regarding the regulation of transcriptional elements of cyanobacterial secondary metabolite biosynthesis remains limited to the microcystin (*mcy*) and jamaicamide (*jam*) BGCs [38-43]. However, the regulation of PSTs at the molecular level, including the transcriptional elements of the *sxt* BGC remains largely unknown [44, 45].

Here, we identify the transcription units of the *sxt* BGC within the cyanobacterium *Dolichospermum circinale* AWQC131C which enabled the experimental isolation of five promoter regions. We then assessed the reliability of a *lux* reporter system to assay the activity of uncharacterised cyanobacterial promoters in the heterologous hosts *E. coli* and *Synechocystis* sp. PCC 6803 for the first time. Characterisation of these cyanobacterial promoters and determination of their activity in *E. coli* and *Synechocystis*, highlights the unpredictability of cyanobacterial promoters of natural product BGCs in heterologous hosts. This study is the first essential phase of understanding the cloning strategy expression of PST biosynthesis, identifying the need for promoter engineering or exchange in future experiments.

Results

Identification of transcriptional units within the *sxt* biosynthetic gene cluster

Reverse-transcriptase PCR revealed that the *sxt* BGC in *D. circinale* AWQC131C is transcribed as five transcriptional units from two bidirectional promoter regions (Figure 1; Figure S1). All five transcripts appear to be constitutively expressed under standard laboratory conditions, since *sxt* mRNA was detected across all time points. Operon 1, *sxtDV*EABC* (* indicates disrupted ORF of *sxtV* [34]), spans 7.3 kb, is transcribed in the reverse direction and encodes several proteins predicted to be involved early in PST biosynthesis. Operon 2, *sxtPQR*, spans 3.5 kb and is transcribed in the forward direction. The catalytic functions of SxtP, SxtQ and SxtR are unknown, but they are likely to be essential for PST biosynthesis as their presence and organisation is conserved amongst all reported *sxt* clusters. The third transcriptional unit is monocistronic and encodes SxtPER, a putative permease of the drug/metabolite transporter family of proteins and is transcribed from two promoters, as discussed further below. Operon 4, is transcribed in the forward direction and spans 12.8 kb. Operon 4 encodes a protein of unknown function, Orf24 that is conserved in most *sxt* clusters, followed by genes encoding twelve enzymes involved in PST biosynthesis, resulting in the polycistron *orf24sxtSTUNGHMIJKLO*.

The 3' ends of operons 1-4 were bioinformatically screened for putative Rho-dependent and Rho-independent transcriptional termination sites using the programs TransTerm and TranstermHP, respectively [46, 47]. Rho-independent transcription termination sites were identified in the non-coding regions of three out of four *sxt* mRNA transcripts (Table S1). Rho-dependent or Rho-independent termination sites were not identified in the sequence of mRNA encoding operon 1.

Transcription start sites and promoter regions of the *sxt* operons

The transcriptional start sites (TSSs) of each operon were experimentally identified *via* 5' rapid amplification of cDNA ends (5' RACE) (Table 1; Figure S2). The upstream region of each TSS was screened for a promoter sequence consistent with conserved binding sequences of group 1, 2 and 3 sigma factors [31]. All promoters identified in this study displayed sequence similarity to the consensus -10 hexamer (Pribnow box) of the prokaryotic RNA polymerase binding site, while there was sporadic presence of the -35 hexamer binding site (Table 1). These results suggest that the *sxt* promoters of *D. circinale* AWQC131C are activated by an RNA polymerase core enzyme in conjunction with a group 1 or group 2 sigma factor [31, 48]. For the -10 promoter sequences identified, a search was conducted for an extended -10 binding site and upstream (UP) element. The 5' untranslated region (UTR) of each operon was also bioinformatically screened for the presence of consensus ribosomal binding site (RBS) sequences, although previously reported bioinformatics surveys of cyanobacterial genomes were not able to identify the consensus RBS sequence in all genes [49, 50]. Based on the 5' RACE and bioinformatics data, the *D. circinale* AWQC131C *sxt* BGC includes of a total of five TSSs under standard culture conditions (Figure 2).

Operon 1 (*sxtDV*EABC*) contains a short 5' UTR of -32 bp upstream of the translation start site and a promoter (*PsxtD*) with high sequence similarity to the *E. coli* σ^{70} -10 and -35 hexamers. *PsxtP* initiates the transcription of operon 2, possesses a short 5' UTR spanning 34 bp and contains both -10 and -35 regions. The transcript initiated by *PsxtP* also displayed a likely RBS (AAGA) 6 nucleotides upstream of the *sxtP* translation start site. A conserved -35 sequence was also identified 21 bp upstream of the extended -10 sequence, resulting in an unusually long distance between the two hexamers. *Porf24* has a perfectly conserved -10 consensus sequence, including the extended -10 TGn motif (Table 1). The 5' UTR for *orf24* is 160 bp in length.

Unusually, transcription of the putative transporter, *sxtPER*, was initiated from two promoters, *PsxtPER1* and *PsxtPER2*. *PsxtPER1* is located 91 bp upstream of the annotated TSS of *sxtPER* (Figure 2) and contains a highly conserved -10 and -35 RNA polymerase binding site. *PsxtPER2* is located 94 bp downstream of the translational start site and contains a highly conserved -10 sequence, including the single nucleotide seen in extended -10 promoters as well as an RBS (AAAGAAG).

The activity of *sxt* promoters in *E. coli*

The five promoters identified in the *D. circinale* *sxt* cluster using 5' RACE, *PsxtP*, *PsxtD*, *PsxtPER1*, *PsxtPER2* and *Porf24*, were amplified by PCR and cloned into the *E. coli* expression vector, pET28b (Novagen), directly in front of a luciferase reporter (*lux*) operon (Figure S4). Expression of luciferase from each of these promoters was measured and compared with negative controls; *pET28-lux* harbouring a non-promoter region from within the *sxtO* gene and the pET28-*lux* plasmid with no added promoter. Unpaired t-tests showed that all promoters exhibited significant levels of expression (Table S2) when compared to the pET28-*lux* negative control. Under the described culture conditions, the heterologous

PsxtD, *PsxtP*, and *PsxtPER1* promoters mediated the highest levels of luciferase expression in *E. coli* (Table S3). There was a statistically significant difference ($p < 0.0001$) between the highest performing promoter *PsxtD* and all the other promoters, as well as the controls (*sxtO* and pET28-*lux*) (Table S4).

The promoter responsible for the transcription of *orf24* and the second promoter of *sxtPER*, *PsxtPER2* were weaker than the other promoters, but still significantly stronger than the controls (Figure 3A). The incorporation of both promoters into the *lux* expression constructs resulted in a 12-27-fold increase in luciferase expression over *sxtO-lux* (Figure 3B), and 810- 1770-fold increase in luciferase expression over the pET28-*lux* control. These results indicate that the promoters are active, albeit weaker than the other three promoters.

The activity of *sxt* promoters in *Synechocystis* sp. PCC 6803

Four *sxt* promoters were active in *Synechocystis* sp. PCC 6803 (Figure 4). Unpaired t-tests showed that expression of luciferase from the *PsxtP*, *PsxtPER1*, and *Porf24* were significantly different to expression in the control strain, while expression from *PsxtD* was not statistically different to the control strain ($P < 0.05$; Table S5).

The *sxtD* promoter regulates the transcription of operon 1 of the *D. circinale* *sxt* cluster, which carries the core biosynthetic genes, including the polyketide synthase-like enzyme, *sxtA*. Strains harbouring *PsxtD* had very low luciferase expression levels that were only 1.3-fold higher than expression levels in the promoter-less control strain (Table S6), and were statistically lower than the other 3 promoters (Table S7). The lack of statistically significant expression from *PsxtD* indicates the promoter as the only candidate for exchange for heterologous expression of PSTs in *Synechocystis* sp. PCC 6803. *PsxtP* and *Porf24* mediated consistent levels of luciferase expression per OD₇₃₀ throughout the experiment (Figure 4B) *PsxtPER1* mediated expression levels that were initially up to three-fold higher than *PsxtP*, however, the rate of expression decreased over the course of growth.

Discussion

Information regarding the transcriptional elements of cyanobacterial secondary metabolite biosynthesis is an essential first step in both understanding how these metabolites may be regulated within the native producers as well as harnessing these genes for future heterologous expression experiments. Here, we targeted the transcriptional units of the *D. circinale* AWQC131C *sxt* BGC and measured promoter activity within two possible future heterologous hosts, *E. coli* and *Synechocystis*.

The entire *D. circinale* AWQC131C *sxt* cluster was found to be transcribed on a total of five transcriptional units from two bidirectional promoter sites. The isolated promoters all contained a -10 Pribnow box, as expected, but sporadically contained other transcriptional elements including the -35 hexamer, UP element and RBS, which is commonly observed in cyanobacteria[30, 49, 50]. A further feature was the identification of a second promoter region initiating transcription of the proposed permease SxtPER. One of these included an intragenic promoter region, which results in the expression of a 258 aa truncated

isoform of SxtPER. While uncommon, the use of a second TSS to produce two protein isoforms has previously been reported [51]. For example, the bacterocin colicin V is secreted by a membrane transporter, where both the full length CvaA and shorter CvaA* proteins, translated from the same *cvaA* gene, are both required for optimal excretion [52]. Interestingly, BLASTp analysis revealed that a complete RhaT super domain of the drug/metabolite transporter family [53] is present within both isoforms of the protein. The larger encoded protein contains additional sequence at the N-terminus. While our previous proteomics experiments in *D. circinale* were unable to detect both isoforms of SxtPER protein under standard conditions, further experiments are required to determine if the two isoforms of the proteins are required for excretion of PST in *D. circinale*.

The expression of *PsxtP* is an interesting example of the promoter elements required for heterologous expression of cyanobacterial promoters in *E. coli*. *PsxtP* does not seem to have a discernible -35 binding region yet does have a RBS and promoted high expression levels in *E. coli*. Previous studies have shown that while the distance between the -10 and -35 sequences can effect transcription in cyanobacteria, the -35 hexamer is not always required [54-56]. Thus, the competing preferences between the TSS sequence and position, taken together with other elements of the promoter such as -10, and -35 sequences, transcription factors, the sequence length between the -10 and -35 regions, and the RBS, highlights the complexity of transcriptional regulation and shows the importance of experimental validation of promoter activation data to further improve bioinformatics databases.

It was found that the expression levels of *PsxtPER1* initiated at a high level but decreased over the course of growth. This indicates the majority of toxins could be exported from the cell early in culture and retained in the cell as the culture progresses. This would allow future research to optimise PST extraction at different culture stages, either from the cell free component or the cell mass. Alternatively, since *PsxtPER1* is active at the early growth stages within the heterologous host, it could be a target for repression to limit toxin export and therefore retaining toxin within the cell. This will increase the efficiency of toxin isolation from the cell biomass.

It is known that gene expression levels will have significant impact on the amount of PST molecule produced. Here, the promoters that regulate PST biosynthesis mediated lowered expression of luciferase in *Synechocystis* sp. PCC 6803 compared to *E. coli*. The significant decrease in luciferase expression by the cyanobacterial promoters in *Synechocystis* sp. PCC 6803 has previously been observed in studies of the zinc inducible promoter, *Psmt*, from *Synechococcus* sp. PCC7002. *Psmt* mediated higher levels of protein synthesis and therefore higher levels of ethylene production in *E. coli*, compared to *Synechocystis* sp. PCC 6803, which only produced residual levels [57]. Conversely, high expression levels impacted lyngbyatoxin biosynthesis in *E. coli*. Heterologous expression of lyngbyatoxin (*ltxA-D*) in *E. coli* was only successful when the strong T7 phage promoter was replaced with the weaker *Ptet_O* promoter [27]. *Ptet_O* has since been exploited for the heterologous expression of multiple cyanobacterial BGCs in *E. coli* [21, 22, 25]. Subsequent heterologous expression of lyngbyatoxin in *Anabaena* using non-native promoters was more successful than the *E. coli* system while expression from the native promoter did not occur in either case [23]. Together, these results suggest that native promoters are recognised differently in

heterologous hosts and that while successful transcription of cyanobacterial BGCs in heterologous hosts is important, other factors play a role in the efficiency of the host production of secondary metabolites. This study has identified each of the five native *sxt* promoters and established through the use of the lux reporter, which of those were not recognised in both *E. coli* and *Synechocystis*.

Conclusion

PSTs have a range of biomedical applications and thus heterologous expression of the *sxt* BGC should be explored as a potential tool for the characterisation, manipulation and sustainable production of these compounds. Heterologous expression of cyanobacterial natural product BGCs has had mixed success in the past and further characterisation of cyanobacterial promoters is required for successful expression of complex biosynthetic pathways such as the PST biosynthetic pathway. This study identified five putative *sxt* promoters in *D. circinale* AWQC131C and tested their activity in *E. coli* and *Synechocystis* sp. PCC 6803. In *E. coli*, *PsxtD*, *PsxtP* and *PsxtPER1* promoted high levels of luciferase expression while *Porf24* was significantly weaker. Further, if *Synechocystis* sp. PCC6803 is to be used as a host, the inactive *PsxtD* should be replaced by host-compatible promoters. Successfully manipulating the *sxt* BGC within a heterologous host at the transcription level is the first step to efficiently access the PSTs for a range of biotechnological applications.

Methods

Strains and culture conditions

D. circinale AWQC131C was maintained in Jaworski's medium (JM) [58] at 24°C ±1°C and illuminated with 11 μmol m⁻² s⁻¹ of photons on a 12:12 h light/dark cycle. *Synechocystis* sp. PCC 6803 was maintained in BG11 medium supplemented with 100 μg mL⁻¹ spectinomycin when required, at 30°C under constant illumination. Unless otherwise specified, *E. coli* strains (Table 2) were maintained in Luria broth or on agar plates supplemented with 100 μg mL⁻¹ ampicillin or 50 μg mL⁻¹ kanamycin and grown at 37°C.

Total RNA extraction, cDNA synthesis and transcriptional analysis

To extract high quality total RNA, cell pellets were snap-frozen in liquid nitrogen and ground into a fine powder with a mortar and pestle prior to extraction with the RNeasy Plant Mini kit (QIAGEN). Residual genomic (g) DNA was removed from total RNA samples using TURBO DNA-free™ DNase as described by the manufacturer (Ambion). Removal of contaminating gDNA was confirmed via PCR with the 27F/809R PCR primer set targeting the cyanobacterial 16S rRNA gene [60]. RNA quality was also checked via formaldehyde gel electrophoresis, while gDNA was checked by agarose gel electrophoresis.

The Superscript® III First Strand synthesis system (Invitrogen) was used to reverse transcribe 1 μg of total RNA primed with an antisense gene-specific primer (GSP). Transcriptional units were determined by PCR amplification in a 20 μL reaction mixture containing 2.5 mM MgCl₂, 1 × PCR buffer (Fisher Biotec, Geneworks), 10 pmol dNTPs (Austral Scientific), 10 pmol of GSP, 1 U of *Taq* polymerase (Fisher Biotec,

Geneworks) and sterile Milli-Q water. Thermal cycling was performed in a Bio-Rad 96-well iCycler (Bio-Rad) and began with an initial denaturation cycle of 95°C for 4 min, followed by 35 cycles of DNA denaturation at 95°C for 20 s and primer annealing at 55°C for 20 s. DNA strand extension was altered to 1 min for every 1 kb of amplified product. A final extension at 72°C for 7 min and a final holding temperature of 4°C completed the thermal cycling. Each reaction contained cDNA as the template and two primers (Table S8), which were designed to target an adjacent gene. Amplification was observed if the two adjacent genes were located on the same mRNA transcript. The positive control for each PCR contained gDNA. Two negative control reactions were performed by adding template from a cDNA synthesis reaction, the first omitting reverse transcriptase and the second reaction omitting a nucleic acid template.

Isolation of *D. circinale* AWQC131C *sxt* biosynthetic gene cluster transcription start sites (TSS) and promoters using 5' rapid amplification of cDNA ends (5'RACE)

To isolate the promoter of each transcriptional unit, TSSs were localised with the FirstChoice® RLM-RACE kit for 5' RACE (Ambion) with 10 µg total RNA as starting material. The 5' RACE adapter was ligated directly onto RNA followed by reverse transcription cDNA synthesis. First round PCR reactions were performed using a 5' outer adapter primer in conjunction with four reverse GSPs at approximately 50-100 bp intervals (Figure S3, Table S9). Reactions containing amplified products from the first round PCR became the template for second round nested PCR containing a 5' adapter inner primer in conjunction with the same four reverse primers. Amplicons of interest were analysed on a 2% (w/v) agarose gel and purified using a QIAquick spin gel extraction kit (QIAGEN). Purified PCR products were then cloned into the pGEM-T Easy vector (Promega) and sequenced using an ABI 3730 capillary sequencer at the Ramaciotti Centre for Genomics, UNSW.

Cloning and transformation

The TOPO TA cloning® kit (Invitrogen) and the pGEM®-T Easy Vector kit (Promega) were used for the cloning and transformation of *E. coli* (Table 2). Cloning with the TOPO TA cloning® kit involved setting up a ligation reaction containing 4 µL of PCR product, 1 µL of Invitrogen salt solution (1.2 M NaCl, 0.06 M MgCl₂) and 10 ng pCR®2.1-TOPO® plasmid DNA (Invitrogen). The ligation reaction was incubated for 20 min at room temperature and was then ready for transformation. The pGEM®-T Easy vector ligation reaction contained 1 × rapid ligation buffer (Promega), 50 ng of pGEM®-T Easy vector DNA (Promega), 3 Weiss U of T4 DNA ligase (Promega) and 3 µL of PCR product. The ligation reaction was left to incubate overnight at 4°C and was then ready for transformation. Positive transformants were selected by blue and white colony screening and the presence of a cloned insert was confirmed by colony PCR using either the primer sets M13F and M13R (pCR®2.1-TOPO) or T7F and M13R (pGEM-T Easy). Plasmids containing an insert were then sequenced.

Engineering the *sxt* promoter-luciferase reporter constructs for expression in *E. coli* DH5α

Five promoters (*PsxtD*, *PsxtP*, *PsxtPER1*, *Porf24*, *PsxtPER2*) and a non-promoter region within the *sxtO* open reading frame were cloned into the pET28b expression vector along with the luciferase reporter (*luxCDABE*) operon from *Photorhabdus luminescens* (Figure S4A). The luciferase operon (*luxCDABE*) was amplified from the pLuxNSII plasmid [59] via PCR (denaturation at 98°C for 3 min, followed by 30 cycles of denaturation at 98°C for 15 s, annealing step at 60°C for 20s, extension at 72°C for 30s/kb, and a final extension at 72°C for 10 min), and the pET28b backbone was also PCR amplified to remove the T7 promoter region. All primers were designed using the NEBuilder assembly tool (Table S10

). Double-stranded PCR fragments were amplified using the KAPA HiFi Hotstart DNA polymerase (KAPA biosystems). The pET28b backbone, *sxt* promoter (*Psxt*), and *lux* operon were assembled using the Gibson assembly master mix (NEB) [61], and incubated at 50°C, for 1 h. The reaction was transformed into chemically competent *E. coli* DH5α and positive colonies selected as above.

Engineering the *sxt* promoter-luciferase reporter constructs for expression in *Synechocystis* sp. PCC 6803

The *Psxt-lux* integration vector was engineered through classical restriction/ligation cloning using the restriction enzymes *NotI* and *KpnI* (NEB). The *Psxt-lux* fragments were amplified from the pET28b-P-*lux* vectors using the *lacI*-P-*lux*_NotI_F, and *lacI*-P-*lux*_KpnI_R primers (Table S10). Linear DNA fragments were digested, purified and ligated into the pSYN_6.3 vector (Figure S4B) using T4 DNA ligase at 22°C for 1 h, followed by transformation into *E. coli* DH5α and colony screening. Plasmid constructs were confirmed by terminal-end sequencing.

The integration of the *Psxt-lux* fragments into the *Synechocystis* sp. PCC 6803 genome (Figure S4C) was achieved via the natural competence of the host [62]. *Synechocystis* sp. PCC 6803 was grown at 30°C, with shaking at 100 rpm, under constant light until exponential phase, and used to inoculate a 50 mL BG11 medium to an initial OD₇₃₀ of 0.05. After ~4 days of photoautotrophic growth, the cells reached an OD₇₃₀ of 0.5, and were harvested by centrifugation at 2,750 *g* for 5 min. Cells were resuspended in 2 mL of fresh BG11 medium, divided into 0.5 mL aliquots (OD₇₃₀ of 2.5), combined with 10 µg of DNA and incubated at 30°C for 6 h. A sterile Immobilon Transfer membrane (Merk Millipore) was placed on each BG11 agar plate, overlaid with 200 µL of the transformation mixture and incubated for 12 h under constant illumination at 30°C. Membranes were transferred to BG11 agar plates containing 25 µg mL⁻¹ spectinomycin. The plates were incubated for a further two days at 30°C under constant illumination, before the membrane was transferred to plates containing 50 µg mL⁻¹ spectinomycin and incubated for a further 7-10 days until colonies become visible. Recombinant *Synechocystis* sp. PCC 6803 colonies were picked and streaked onto BG11 agar plates supplemented with 100 µg mL⁻¹ spectinomycin and subcultured a further three times to achieve integration of the cloned reporter fragment and full chromosomal segregation. Transformants were confirmed using the PhaCaF and PhaCbR PCR primers (Table S10).

Activity of *sxt* promoters in *E. coli* DH5α

Promoter-luciferase reporter constructs were transformed into *E. coli* DH5α and grown on M9 minimal medium supplemented with 50 µg mL⁻¹ kanamycin at 37°C for 24 h. Bioluminescence (RLU) and optical density measurements were measured at one-hour intervals until the OD₆₀₀ reached 0.8. A final measurement was taken at 24 h. The strength of each promoter was measured as the highest bioluminescence, normalised to OD₆₀₀. One-way ANOVA (Graphpad Prism 7) was used to calculate any statistical differences between the promoters. Unpaired t-tests were also used to determine the statistical differences between the strains and the control.

Activity of *sxt* promoters in *Synechocystis* sp. PCC 6803

Promoter-luciferase reporter constructs were transformed into *Synechocystis* sp. PCC 6803 strains, which were inoculated into BG11 medium supplemented with 100 µg mL⁻¹ of spectinomycin and grown at 30°C with shaking under constant illumination. Optical density and relative light units (RLU) were measured every 24 h for 400 h. Promoter strength was measured by determining the highest RLU per OD₇₃₀. One-way ANOVA (Graphpad Prism 7) was used to calculate any statistical differences between the promoters. Unpaired t-tests were also used to determine the statistical differences between the strains and the control.

Abbreviations

5' RACE – rapid amplification of cDNA ends

BGC – biosynthetic gene cluster

GSP – gene specific primer

PST – paralytic shellfish toxins

RBS – ribosomal binding site

TSS – transcriptional start site

UP – upstream

UTR – untranslated region

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and material

All data generated or analysed during this study are included in this published article and its supplementary information files. Sequences of plasmids used in this study are available at the end of the supplementary information file.

Competing interests

The authors declare that they have no competing interests

Funding

This research was funded by the Australian Research Council, grant number DP0880264. PMD and BA were supported by the Australian Postgraduate award. MM was supported by the Western Sydney University Research Development Strategic Initiative Scheme.

Authors' contributions

PMD, RM, JM, MCM and BAN designed the overall project. PMD performed experimental transcriptional analysis, identification of transcription start site by 5' RACE and bioinformatic analysis. BA and RM designed, cloned and performed promoter activity assays. PMD and BA wrote the manuscript while all authors edited and approved the manuscript.

Acknowledgements

The authors would like to thank Dr. Adam Jones and Dr. Daniela Ferreira for helpful experimental advice regarding cyanobacterial transcript analysis and identification of the transcription start site. The authors would also like to thank Anna Glöckle and Leanne Pearson for reading and editing the manuscript. The authors would like to thank Professor Carl Johnson for providing the pluxNS II plasmid carrying the luciferase operon, and Professor Dean Price for providing the *Synechocystis* 6803 strain.

References

1. Wiese M, D'Agostino PM, Mihali TK, Moffitt MC, Neilan BA: **Neurotoxic alkaloids: Saxitoxin and its analogs.** *Marine Drugs* 2010, **8**:2185-2211.
2. Etheridge SM: **Paralytic shellfish poisoning: Seafood safety and human health perspectives.** *Toxicon* 2010, **56**:108-122.
3. Lagos N: **Clinical applications of paralytic shellfish poisoning toxins.** In *Toxins and Biologically Active Compounds from Microalgae. Volume 2.* Edited by Rossini GP: CRC Press; 2014: 309–329.
4. Adams HJ, Blair MR, Jr., Takman BH: **The local anesthetic activity of saxitoxin alone and with vasoconstrictor and local anesthetic agents.** *Archives Internationales de Pharmacodynamie et de*

Therapie 1976, **224**:275-282.

5. Cestèle S, Catterall WA: **Molecular mechanisms of neurotoxin action on voltage-gated sodium channels.** *Biochimie* 2000, **82**:883-892.
6. Rodriguez-Navarro Alberto J, Lagos N, Lagos M, Braghetto I, Csendes A, Hamilton J, Figueroa C, Truan D, Garcia C, Rojas A, et al: **Neosaxitoxin as a local anesthetic preliminary observations from a first human trial.** *Anesthesiology* 2007, **106**:339-345.
7. Barnet CS, Tse JY, Kohane DS: **Site 1 sodium channel blockers prolong the duration of sciatic nerve blockade from tricyclic antidepressants.** *Pain* 2004, **110**:432-438.
8. Epstein-Barash H, Shichor I, Kwon AH, Hall S, Lawlor MW, Langer R, Kohane DS: **Prolonged duration local anesthesia with minimal toxicity.** *Proceedings of the National Academy of Sciences of the United States of America* 2009, **106**:7125-7130.
9. Chorny M, Levy RJ: **Site-specific analgesia with sustained release liposomes.** *Proceedings of the National Academy of Sciences of the United States of America* 2009, **106**:6891-6892.
10. Garrido R, Lagos N, Lattes K, Abedrapo M, Bocic G, Cuneo A, Chiong H, Jensen C, Azolas R, Henriquez A, Garcia C: **Gonyautoxin: New treatment for healing acute and chronic anal fissures.** *Diseases of the Colon & Rectum* 2005, **48**:335-343.
11. Garrido R, Lagos N, Lattes K, Azolas CG, Bocic G, Cuneo A, Chiong H, Jensen C, Henriquez AI, Fernandez C: **The gonyautoxin 2/3 epimers reduces anal tone when injected in the anal sphincter of healthy adults.** *Biological Research* 2004, **37**:395-403.
12. Lattes K, Venegas P, Lagos N, Lagos M, Pedraza L, Rodriguez-Navarro AJ, Garcia C: **Local infiltration of gonyautoxin is safe and effective in treatment of chronic tension-type headache.** *Neurological Research* 2009, **31**:228-233.
13. Walker JR, Merit JE, Thomas-Tran R, Tang DTY, Du Bois J: **Divergent Synthesis of Natural Derivatives of (+)-Saxitoxin Including 11-Saxitoxinethanoic Acid.** *Angewandte Chemie International Edition* 2019, **58**:1689-1693.
14. Chun SW, Narayan ARH: **Biocatalytic synthesis of α -amino ketones.** *Synlett : accounts and rapid communications in synthetic organic chemistry* 2019, **30**:1269-1274.
15. Watanabe R, Suzuki T, Oshima Y: **Preparation of calibration standards of N1-H paralytic shellfish toxin analogues by large-scale culture of cyanobacterium *Anabaena circinalis* (TA04).** *Marine drugs* 2011, **9**:466-477.
16. Quilliam M: **Certified reference materials for marine toxins.** 2006: 139-144.
17. Laycock MV, Thibault P, Ayer SW, Walter JA: **Isolation and purification procedures for the preparation of paralytic shellfish poisoning toxin standards.** *Natural Toxins* 1994, **2**:175-183.
18. Thomas K, Blay P, Burton IW, Cembella A, Craft C, Crain S, Hardstaff WR, Laycock M, LeBlanc D, LeBlanc P: **Certified reference materials for marine algal toxins.** In *Harmful Algae 2002: Florida Fish and Wildlife Conservation Commission and Intergovernmental Oceanographic Commission of UNESCO*. Edited by Steidinger KA, Landsberg JH, Tomas CR, Vargo GA2004: 216-218.

19. Nah H-J, Pyeon H-R, Kang S-H, Choi S-S, Kim E-S: **Cloning and Heterologous Expression of a Large-sized Natural Product Biosynthetic Gene Cluster in Streptomyces Species.** *Frontiers in microbiology* 2017, **8**:394-394.
20. Zhang JJ, Tang X, Moore BS: **Genetic platforms for heterologous expression of microbial natural products.** *Natural Product Reports* 2019, **36**:1313-1332.
21. Liu T, Mazmouz R, Ongley SE, Chau R, Pickford R, Woodhouse JN, Neilan BA: **Directing the Heterologous Production of Specific Cyanobacterial Toxin Variants.** *ACS Chemical Biology* 2017, **12**:2021-2029.
22. D'Agostino PM, Gulder TAM: **Direct pathway cloning combined with sequence- and ligation-independent cloning for fast biosynthetic gene cluster refactoring and heterologous expression.** *ACS Synthetic Biology* 2018, **7**:1702-1708.
23. Long PF, Dunlap WC, Battershill CN, Jaspars M: **Shotgun cloning and heterologous expression of the patellamide gene cluster as a strategy to achieving sustained metabolite production.** *ChemBioChem* 2005, **6**:1760-1765.
24. Ziemert N, Ishida K, Liaimer A, Hertweck C, Dittmann E: **Ribosomal synthesis of tricyclic depsipeptides in bloom-forming cyanobacteria.** *Angewandte Chemie International Edition* 2008, **47**:7756-7759.
25. Greunke C, Duell ER, D'Agostino PM, Glöckle A, Lamm K, Gulder TAM: **Direct pathway cloning (DiPaC) to unlock natural product biosynthetic potential.** *Metabolic Engineering* 2018, **47**:334-345.
26. Videau P, Wells KN, Singh AJ, Gerwick WH, Philmus B: **Assessment of *Anabaena* sp. strain PCC 7120 as a heterologous expression host for cyanobacterial natural products: Production of lyngbyatoxin A.** *ACS Synthetic Biology* 2016.
27. Ongley SE, Bian X, Zhang Y, Chau R, Gerwick WH, Müller R, Neilan BA: **High-titer heterologous production in *E. coli* of lyngbyatoxin, a protein kinase C activator from an uncultured marine cyanobacterium.** *ACS Chemical Biology* 2013, **8**:1888-1893.
28. Wells KN, Nelson D, Eiting JE, Videau P, Philmus B: **The influence of sigma factors and ribosomal recognition elements on heterologous expression of cyanobacterial gene clusters in Escherichia coli.** *FEMS Microbiology Letters* 2018, **365**.
29. Huang H-H, Camsund D, Lindblad P, Heidorn T: **Design and characterization of molecular tools for a synthetic biology approach towards developing cyanobacterial biotechnology.** *Nucleic Acids Research* 2010, **38**:2577-2593.
30. Curtis SE, Martin JA: **The transcription apparatus and the regulation of transcription initiation.** In *The Molecular Biology of Cyanobacteria. Volume 1*. Edited by Bryant DA. Dordrecht: Kluwer Academic Publishers; 1994: 613-639.[*Advances in Photosynthesis and Respiration*].
31. Imamura S, Asayama m: **Sigma factors for cyanobacterial transcription.** *Gene Regulation and Systems Biology* 2009, **3**:65-87.
32. Imamura S, Yoshihara S, Nakano S, Shiozaki N, Yamada A, Tanaka K, Takahashi H, Asayama M, Shirai M: **Purification, characterization, and gene expression of all sigma factors of RNA polymerase**

- in a cyanobacterium. *Journal of Molecular Biology* 2003, **325**:857-872.
33. Kellmann R, Mihali TK, Jeon YJ, Pickford R, Pomati F, Neilan BA: **Biosynthetic intermediate analysis and functional homology reveal a saxitoxin gene cluster in cyanobacteria.** *Applied and Environmental Microbiology* 2008, **74**:4044-4053.
  34. Mihali TK, Kellmann R, Neilan BA: **Characterisation of the paralytic shellfish toxin biosynthesis gene clusters in *Anabaena circinalis* AWQC131C and *Aphanizomenon* sp. NH-5.** *BMC Biochemistry* 2009, **10**:8.
  35. Stucken K, John U, Cembella A, Murillo AA, Soto-Liebe K, Fuentes-Valdés JJ, Friedel M, Plominsky AM, Vásquez M, Glöckner G: **The Smallest Known Genomes of Multicellular and Toxic Cyanobacteria: Comparison, Minimal Gene Sets for Linked Traits and the Evolutionary Implications.** *PLOS ONE* 2010, **5**:e9235.
  36. Mihali TK, Carmichael WW, Neilan BA: **A putative gene cluster from a *Lyngbya wollei* bloom that encodes paralytic shellfish toxin biosynthesis.** *PLoS ONE* 2011, **6**:e14657.
  37. Cullen A, D'Agostino PM, Mazmouz R, Pickford R, Wood SA, Neilan BA: **Insertions within the saxitoxin biosynthetic gene cluster result in differential toxin profiles.** *ACS Chemical Biology* 2018, **13**:3107-3114.
  38. Martin-Luna B, Sevilla E, Hernandez JA, Bes MT, Fillat MF, Peleato ML: **Fur from *Microcystis aeruginosa* binds *in vitro* promoter regions of the microcystin biosynthesis gene cluster.** *Phytochemistry* 2006, **67**:876-881.
  39. Ginn HP, Pearson LA, Neilan BA: **NtcA from *Microcystis aeruginosa* PCC 7806 is autoregulatory and binds to the microcystin promoter.** *Applied and Environmental Microbiology* 2010, **76**:4362-4368.
  40. Neilan BA, Pearson LA, Muenchhoff J, Moffitt MC, Dittmann E: **Environmental conditions that influence toxin biosynthesis in cyanobacteria.** *Environmental Microbiology* 2012, **15**:1239-1253.
  41. Kaebernick M, Neilan BA, Börner T, Dittmann E: **Light and the transcriptional response of the microcystin biosynthesis gene cluster.** *Applied and Environmental Microbiology* 2000, **66**:3387-3392.
  42. Kaebernick M, Dittmann E, Börner T, Neilan BA: **Multiple alternate transcripts direct the biosynthesis of microcystin, a cyanobacterial nonribosomal peptide.** *Applied and Environmental Microbiology* 2002, **68**:449-455.
  43. Jones A, Gerwick L, Gonzalez D, Dorrestein P, Gerwick W: **Transcriptional analysis of the jamaicamide gene cluster from the marine cyanobacterium *Lyngbya majuscula* and identification of possible regulatory proteins.** *BMC Microbiology* 2009, **9**:247.
  44. Ongley SE, Pengelly JJL, Neilan BA: **Elevated Na⁺ and pH influence the production and transport of saxitoxin in the cyanobacteria *Anabaena circinalis* AWQC131C and *Cylindrospermopsis raciborskii* T3.** *Environmental Microbiology* 2016, **18**:427-438.
  45. D'Agostino PM, Song X, Neilan BA, Moffitt MC: **Proteogenomics of a saxitoxin-producing and non-toxic strain of *Anabaena circinalis* (cyanobacteria) in response to extracellular NaCl and phosphate depletion.** *Environmental Microbiology* 2016, **18**:461-476.

46. Jacobs GH, Chen A, Stevens SG, Stockwell PA, Black MA, Tate WP, Brown CM: **Transterm: A database to aid the analysis of regulatory sequences in mRNAs.** *Nucleic Acids Research* 2009, **37**:D72-D76.
47. Kingsford C, Ayanbule K, Salzberg S: **Rapid, accurate, computational discovery of Rho-independent transcription terminators illuminates their relationship to DNA uptake.** *Genome Biology* 2007, **8**:R22.
48. Camsund D, Lindblad P: **Engineered transcriptional systems for cyanobacterial biotechnology.** *Frontiers in Bioengineering and Biotechnology* 2014, **2**.
49. Omotajo D, Tate T, Cho H, Choudhary M: **Distribution and diversity of ribosome binding sites in prokaryotic genomes.** *BMC genomics* 2015, **16**:604-604.
50. Mutsuda M, Sugiura M: **Translation initiation of cyanobacterial rbcS mRNAs requires the 38-kDa ribosomal protein S1 but not the shine-dalgarno sequence.** *Journal of Biological Chemistry* 2006, **281**:38314-38321.
51. Meydan S, Vázquez-Laslop N, Mankin AS: **Genes within genes in bacterial genomes.** *Microbiology Spectrum* 2018, **6**.
52. Hwang J, Manuvakhova M, Tai PC: **Characterization of in-frame proteins encoded by *cvaA*, an essential gene in the colicin V secretion system: CvaA* stabilizes CvaA to enhance secretion.** *Journal of Bacteriology* 1997, **179**:689-696.
53. Tate CG, Muir JA, Henderson PJ: **Mapping, cloning, expression, and sequencing of the *rhaT* gene, which encodes a novel L-rhamnose-H⁺ transport protein in *Salmonella typhimurium* and *Escherichia coli*.** *Journal of Biological Chemistry* 1992, **267**:6923-6932.
54. Dutheil J, Saenkham P, Sakr S, Leplat C, Ortega-Ramos M, Bottin H, Cournac L, Cassier-Chauvat C, Chauvat F: **The AbrB2 autorepressor, expressed from an atypical promoter, represses the hydrogenase operon to regulate hydrogen production in *Synechocystis* strain PCC6803.** *Journal of Bacteriology* 2012, **194**:5423-5433.
55. Subudhi S, Kurdrud P, Hongsthong A, Sirijuntarut M, Cheevadhanarak S, Tanticharoen M: **Isolation and functional characterization of *Spirulina* D6D gene promoter: Role of a putative GntR transcription factor in transcriptional regulation of D6D gene expression.** *Biochemical and Biophysical Research Communications* 2008, **365**:643-649.
56. Shibato J, Agrawal G, Kato H, Asayama M, Shirai M: **The 5'-upstream *cis*-acting sequences of a cyanobacterial *psbA* gene: Analysis of their roles in basal, light-dependent and circadian transcription.** *Molecular Genetics and Genomics* 2002, **267**:684-694.
57. Guerrero F, Carbonell V, Cossu M, Correddu D, Jones PR: **Ethylene synthesis and regulated expression of recombinant protein in *Synechocystis* sp. PCC 6803.** *PLOS ONE* 2012, **7**:e50470.
58. Thompson AS, Rhodes JC, Pettman I: *Culture collection of algae and protozoa: Catalogue of strains. Natural Environment Research Council.* 5 edn. Ambleside, United Kingdom: Freshwater Biological Association; 1988.
59. Woelfle MA, Xu Y, Qin X, Johnson CH: **Circadian rhythms of superhelical status of DNA in cyanobacteria.** *Proceedings of the National Academy of Sciences* 2007, **104**:18819-18824.

60. Jungblut A, Hawes I, Mountfort D, Hitzfeld B, Dietrich DR, Burns BP, Neilan BA: **Diversity within cyanobacterial mat communities in variable salinity meltwater ponds of McMurdo Ice Shelf, Antarctica.** *Environmental Microbiology* 2005, **7**:519-529.
61. Gibson DG, Young L, Chuang R-Y, Venter JC, Hutchison CA, Smith HO: **Enzymatic assembly of DNA molecules up to several hundred kilobases.** *Nat Meth* 2009, **6**:343-345.
62. Eaton-Rye JJ: **Construction of gene interruptions and gene deletions in the cyanobacterium *Synechocystis* sp. strain PCC 6803.** In *Photosynthesis Research Protocols*. Edited by Carpentier R. Totowa, NJ: Humana Press; 2011: 295-312.

Tables

Table 1: Characteristics of promoter regions in the *sxt* biosynthetic gene cluster of *D. circinale* AWQC131C.

Promoter	Promoter sequence 5' à 3'	Position ^Δ
<i>PsxtD</i>	tgtc TTGTGG ....(14bp)....GAg TATACT tgactagt A	-32
<i>PsxtP</i>	gtat CTATCA ....(12bp)....GTg TATACT agtcaagt A	-34
<i>PsxtPER1</i>	ttcc TTGCAA ....(15bp)....AGt TACAAT tacatg A	-91
<i>PsxtPER2</i>	tgag ATGACA ....(21bp)....CGa TATATT ttgggt G	+94
<i>Porf24</i>	aaaa TTTCCT ....(15bp)....TGc TATAAT gaaatc T	-160
<i>E. coli</i> σ ⁷⁰ consensus	.... TTGACA ....(14bp)....TGn TATAAT .....N	

The -10 and -35 hexamers are capitalised and conserved nucleotides are in bold print. N indicates transcriptional start site (TSS). ^Δ Position of promoter relative to TSS.

Table 2: Strains and plasmids

Organism	Purpose	Ref.
DH5 α	Expression host of <i>lux</i> reporter assay	Promega
TOP 10	Transformation and propagation of plasmids	Invitrogen
<i>D. circinale</i> AWQC131C	Organism harbouring native <i>sxt</i> cluster	[34]
<i>Synechocystis</i> sp. PCC6803	Expression host of <i>lux</i> reporter assay	
<i>Synechocystis</i> SXL7	$\Delta phbC :: PsxtD-luxCDABE$ - Spec ^R	This study
<i>Synechocystis</i> SXL8	$\Delta phbC :: PsxtP-luxCDABE$ - Spec ^R	This study
<i>Synechocystis</i> SXL9	$\Delta phbC :: PsxtPER1-luxCDABE$ - Spec ^R	This study
<i>Synechocystis</i> SXL10	$\Delta phbC :: Porf24-luxCDABE$ - Spec ^R	This study
<i>Synechocystis</i> SXL11	$\Delta phbC :: luxCDABE$ - Spec ^R	This study
Plasmids	Purpose	Ref.
pCR ^{TM} 2.1 TOPO	Cloning and sequencing of 5' RACE TSS products	Invitrogen
pGEM T-Easy	Cloning and sequencing of 5' RACE TSS products	Promega
pET28b	Cloning and expression of <i>sxt</i> promoters	Novagen
pSYN_6.3	Cloning and transformation of <i>sxt</i> promoters in <i>Synechocystis</i> sp. PCC 6803	Unpublished
pLUX NS II	<i>luxCDABE</i>	[59]
pSXL1	pET28b:: <i>PsxtD-luxCDABE</i>	This study
pSXL2	pET28b:: <i>PsxtP-luxCDABE</i>	This study
pSXL3	pET28b:: <i>PsxtPER1-luxCDABE</i>	This study
pSXL4	pET28b:: <i>Porf24-luxCDABE</i>	This study
pSXL5	pET28b:: <i>PsxtPER2-luxCDABE</i>	This study
pSXL6	pET28- <i>sxtO-luxCDABE</i>	This study
pET28- <i>lux</i>	pET28- <i>luxCDABE</i>	This study
pSXL7	pSYN_6.3:: <i>PsxtD-luxCDABE</i>	This study
pSXL8	pSYN_6.3:: <i>PsxtP-luxCDABE</i>	This study
pSXL9	pSYN_6.3:: <i>PsxtPER1-luxCDABE</i>	This study
pSXL10	pSYN_6.3:: <i>Porf24-luxCDABE</i>	This study
pSXL11	pSYN_6.3:: <i>luxCDABE</i>	This study

Figures

Figure 1

Transcriptional analysis of the *D. circinale* AWQC131C *sxt* cluster. Genes required for PST biosynthesis are transcribed by a minimum of four mRNA transcripts from two bi-directional promoter regions encoding five promoters; *PsxtD* (operon 1), *PsxtP* (operon 2), *PsxtPER1* (TU 3), *PsxtPER2* (TU 3) and *Porf24* (operon 4). Direction of transcription is indicated by black arrows.

A

TCTTAATAGGATTAATATTGTATCTATCATTTTGTATCTATCAATTGGTAA
 CCTTAGTGTATACTAGTCAAGTATACTCTTATTAACAAATTTCCACAAGA
 CAACTCATGAAAAGGCTCAGTTCGTTAGTTCTAGCAGCACTCTTGTTTGT

SxtD
 PsxtD +1
 PsxtP +1
 -35
 RBS
 -10
 -10
 -35
 SxtP

B

GTAAAAACCACAGCAGCCATTGTAAAGCGAACTGCATTAGCACTTCTTT
 AGAGATGTTAGTGATTAAACCTTTAGTAATTGGCAAGATTGAGCTATAGA
 CTAGAGTTGCCAGAATAATCAACAGCACCCAAATATATCGGTTATTT
 TCTGGGGCAGATGTGTTCATCTCAACCTTATCCAAGCTATTTTGGAGTTC
 TGTTAAATCTTGGTTTACCATCTCAACTTAAAAATTCCTTAGGTCTGGGC
 GTAGTGCTATAATGAAATCTTTGGCTGTACGGTATATCGCTAATAACGCG
 AATAATGTGAGTCATGTAATTGTAACTCATAAATCCCTTGAATTGCAAGG
 AAATAAAACAGATGCTTACATCTGAATAAAAACCAAGCAATGTTTATAC

SxtPER2
 RBS
 PsxtPER2 +1
 -35
 -10
 SxtPER
 -10
 Porf24 +1
 -35
 PsxtPER +1
 -10
 -35
 Orf24

Figure 2

Sequence analysis of the five promoters present within the *D. circinale* AWQC131C sxt biosynthetic gene cluster. Individual promoters include, PsxtD (green), PsxtP (red), PsxtPER2 (light blue), PsxtPER1 (dark blue) and Porf24 (yellow). Transcriptional elements including promoters, TSS, translation start site, and RBS are shown for the five sxt promoters.

Figure 3

Heterologous expression from of luciferase from sxt promoters in *E. coli* DH5α. (A) Luciferase expression was normalised to the optical density, OD₆₀₀. The activity of five promoters was tested: PsxtD, PsxtP, PsxtPER1, Porf24, and PsxtPER2. An intergenic sxtO sequence was used as the non-promoter control. The highest RLU/OD for the *E. coli* DH5α strains expressing luxCDABE. (*) Denotes statistically significant differences ($p < 0.0001$) between the *E. coli* DH5α strains and the pET28-lux control in unpaired t-tests. (B) Expression relative to the sxtO-lux control. The mean normalised luciferase expression for each promoter was divided by the mean normalised expression of sxtO. The three strongest promoters (PsxtD, PsxtP, PsxtPER) exhibit between 1000-9500 fold higher levels of luciferase expression over the sxtO control. The Porf24 and PsxtPER2 promoters had only a 12 and 27-fold increase in expression, respectively.

Figure 4

Heterologous expression of luciferase from sxt promoters in *Synechocystis* sp. PCC 6803. (A) Luciferase expression from the four main sxt promoters in *Synechocystis* sp. PCC 6803. The highest RLU/OD for the *Synechocystis* sp. PCC 6803 strains expressing luxCDABE. (*) Denotes statistically significant differences ($p < 0.05$) between the *Synechocystis* sp. PCC 6803 strains and the *Synechocystis* sp. PCC 6803-lux control in unpaired t-tests. (B) Normalised luciferase expression over time in *Synechocystis* PCC6803. RLU/OD₇₃₀ over 400 h of growth. PsxtD normalised expression was low. PsxtP and Porf24 had consistent levels of luciferase expression. PsxtPER showed high initial expression that declined over growth to expression reach levels similar to PsxtP and Porf24.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [200129SupportinginfofinalPD.pdf](#)