Extracellular Vesicles of Mesenchymal Stromal Cells Can be Taken up by Microglial Cells and Partially Prevent The Stimulation Induced by β -Amyloid

Dorota Kaniowska (✉ dorota.kaniowska@izi-extern.fraunhofer.de)

Fraunhofer-Institut für Zelltherapie und Immunologie: Fraunhofer-Institut für Zelltherapie und Immunologie IZI <https://orcid.org/0000-0001-5128-2165>

Kerstin Wenk

Institute for Clinical Immunology, University of Leipzig

Phil Rademacher

Institute for Clinical Immunology, University of Leipzig

Ronald Weiß

Institute for Clinical Immunology, University of Leipzig

Claire Fabian

Fraunhofer-Institut für Zelltherapie und Immunologie IZI: Fraunhofer-Institut für Zelltherapie und Immunologie IZI

Isabell Schulz

Fraunhofer-Institut für Zelltherapie und Immunologie IZI: Fraunhofer-Institut für Zelltherapie und Immunologie IZI

Max Guthardt

Fraunhofer-Institut für Zelltherapie und Immunologie: Fraunhofer-Institut für Zelltherapie und Immunologie IZI

Franziska Lange

Fraunhofer-Institut für Zelltherapie und Immunologie IZI: Fraunhofer-Institut für Zelltherapie und Immunologie IZI

Sebastian Greiser

Fraunhofer-Institut für Zelltherapie und Immunologie IZI: Fraunhofer-Institut für Zelltherapie und Immunologie IZI

Matthias Schmidt

UFZ Centre for Environmental Research Leipzig-Halle: Helmholtz-Zentrum für Umweltforschung UFZ

Ulf-Dietrich Braumann

Fraunhofer-Institut für Zelltherapie und Immunologie IZI: Fraunhofer-Institut für Zelltherapie und Immunologie IZI

Frank Emmrich

Fraunhofer-Institut für Zelltherapie und Immunologie IZI: Fraunhofer-Institut für Zelltherapie und Immunologie IZI

Ulrike Köhl

Fraunhofer-Institut für Zelltherapie und Immunologie: Fraunhofer-Institut für Zelltherapie und Immunologie IZI

Yarúa Jaimes

Fraunhofer-Institut für Zelltherapie und Immunologie IZI: Fraunhofer-Institut für Zelltherapie und Immunologie IZI

Research Article

Keywords: neuroinflammation, Amyloid beta, Alzheimer Disease, microglia, mesenchymal stromal/stem cells, extracellular vesicles

Posted Date: June 28th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-631345/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Version of Record: A version of this preprint was published at Stem Cell Reviews and Reports on January 26th, 2022. See the published version at <https://doi.org/10.1007/s12015-021-10261-4>.

Abstract

Mesenchymal stromal/stem cells (MSCs) have great capacity for immune regulation. MSCs provide protective paracrine effects, which are partially exerted by extracellular vesicles (EVs). It has been reported that MSCs-derived EVs (MSC-EVs) contain soluble factors, such as cytokines, chemokines, growth factors and even microRNAs, which confer them similar anti-inflammatory and regenerative effects to MSCs. Moreover, MSCs modulate microglia activation through a dual mechanism of action that relies both on cell contact and secreted factors. Microglia cells are the central nervous system immune cells and the main mediators of the inflammation leading to neurodegenerative disorders.

Here, we investigated whether MSC-EVs affect the activation of microglia cells by β -amyloid aggregates. We show that the presence of MSC-EVs can prevent the upregulation of pro-inflammatory mediators such as tumor necrosis factor (TNF)- α and nitric oxide (NO). Both are up-regulated in neurodegenerative diseases representing chronic inflammation, as in Alzheimer's disease. We demonstrate that MSC-EVs are internalized by the microglia cells. Further, our study supports the use of MSC-EVs as a promising therapeutic tool to treat neuroinflammatory diseases.

Significance Statement

It has been reported that mesenchymal stromal/stem cells and MSC-derived small extracellular vesicles have therapeutic effects in the treatment of various degenerative and inflammatory diseases. Extracellular vesicles are loaded with proteins, lipids and RNA and act as intercellular communication mediators. Here we show that extracellular vesicles can be taken up by murine microglial cells. In addition, they partially reduce the activation of microglial cells against β -amyloid aggregates. This inhibition of microglia activation may present an effective strategy for the control/therapy of neurodegenerative diseases such as Alzheimer's disease.

Introduction

Neuroinflammation is a major contributor to Alzheimer's disease (AD) progression. Brain tissue of patients with AD exhibits chronic inflammation, which is attributed to the activation of microglia cells. The primary role of microglia in the healthy brain is to survey the nervous tissue environment. Despite that, microglia are very reactive cells, their activation results in the overproduction of pro-inflammatory mediators and neuro-toxic cytokines [1, 15, 24]. Additionally, activated microglia are commonly associated with amyloid – β (A β) containing plaques [29] which can directly stimulate microglia cells to acquire a pro-inflammatory phenotype [35]. In this study, we used A β aggregates as stimulus to induce inflammatory responses in the mouse microglia cell line BV-2 as model system since the production of pro-inflammatory mediators by microglia is significantly elevated after A β aggregates treatment [13]. Several studies indicated that inhibition of microglial activation provides an effective strategy for the control/therapy of many injuries and diseases, including stroke, multiple sclerosis, neurodegenerative diseases, along with brain trauma [4].

In the last decade, mesenchymal stromal/stem cells (MSCs) have received special interest in the treatment of inflammatory medical conditions, including AD [5, 26, 42]. MSCs support their niches *in vivo* by nurturing and promoting the proliferation and differentiation of surrounding cells. Moreover, when transplanted, these cells migrate to sites of inflammation and injury promoting immunomodulation and tissue repair [44]. An easily accessible MSC source with excellent proliferation and differentiation capacity [49, 50], low immunogenicity [28, 31], and ability for immunomodulation [9, 21, 25, 33, 46] are adipose stromal/stem cells (ASCs). Injection of ASCs into the cerebral cortex of AD mice led to recovery of spatial learning and memory [27], and intravenous injection of ASCs mitigate dementia in AD mice [5, 19]. The supernatant of ASC cultures is enriched with extracellular vesicles (EVs) which mimic their immune regulatory and regenerative action.

EVs are a heterogeneous population loaded with proteins, lipids and RNA bound by a phospholipid bilayer and functioning as intercellular communication mediators [7]. EVs of MSCs (MSC-EVs) have been reported to display therapeutic effects on the treatment of various degenerative and inflammatory diseases [2, 12, 20, 23, 43]. We recently showed that mouse MSC-EVs possess the ability to prevent microglia activation through the modulation of inflammatory cytokines [17]. In order to evaluate different aspects of MSC-EVs in neuroinflammation, we investigated the uptake of human ASC-EVs by BV-2 cells and their capacity to modulate the pro-inflammatory responses of BV-2 cells stimulated with A β aggregates.

Materials And Methods

Culture of human ASCs from adipose tissue

ASCs were cultured in Dulbecco's modified Eagle's medium, 1 g/L D-glucose and GlutaMAX (DMEM-Low Glucose [LG], Life Technologies) supplemented with 20% FBS (Life Technologies). Medium was changed every 3–4 days. For subculture, cells were passaged weekly with 0.05% trypsin/ 4 mM EDTA (Life Technologies) and seeded at $3\text{--}4.5 \times 10^3$ cells per cm² in T175 flasks.

Flow cytometry

Surface marker expression was analyzed by flow cytometry. ASCs were used from 4 donors. Cells were incubated at 4°C for 15 minutes in the dark with the following antibodies: CD45 (1:50, REA747), CD73 (1:50, REA804), CD90 (1:50, REA897), CD105 (1:50, REA794) as well as the isotype control (1:50, REA293) (all from Miltenyi Biotec, Germany). Flow cytometry measurement was done on a MACSQuant10 (Miltenyi Biotec) and data was analyzed using FlowLogic (Miltenyi Biotec). A minimum of 10,000 events was analyzed for each sample.

C57BL/6 Mouse Mesenchymal Stem Cells

Mouse bone-marrow MSCs were purchased from Cyagen US Inc. (Santa Clara, USA). MSCs were cultured in MEM-alpha medium with GlutaMAX, supplemented with 15% FBS (Life Technologies), 1% penicillin-

streptomycin (Life Technologies) and 2 ng/ml recombinant murine basic fibroblast growth factor (rMu bFGF, Peprotech), with medium changes every three days. The company provides the identity of the cells. Mouse MSCs from Cyagen are positive for CD34, CD44, Sca-1 and negative for CD117. Also, they present multipotential to differentiate towards osteogenic, chondrogenic and adipogenic lineages.

Extracellular vesicles isolation

MSCs culture-supernatant containing the EVs was collected every 72 h starting at passage 2 up until passage 8, at an optical cell confluency higher than 90%. The EVs were isolated using differential (ultra)-centrifugation steps as previously described [17] (Supplementary Fig. 1). Briefly, cell culture supernatant was centrifuged at 500 x g for 10 min to remove cells and cell debris (Hettich centrifuge, MA). To deplete large vesicles, samples were centrifuged again at 10,000 x g during 30 min and the final concentration of EVs was achieved by centrifuging the supernatant at 70,000 x g during 90 min (Thermo Scientific High-Speed Centrifuge, Germany). All centrifugation steps were performed at 4°C. The supernatant was discarded via aspiration. EV pellet was resuspended in 0.9% NaCl or BV-2 cell culture medium and stored at -80°C, depending on the application. Quantification of EVs was performed by flow cytometry or NanoSight nanoparticle tracking analysis (NTA).

Extracellular vesicles quantification using flow cytometry

The first quantification of EVs was done by FACS Canto II flow cytometer (BD Biosciences) with beads (0.2 µm and 1 µm) as standard. Based on this quantification, 8 EVs per BV-2 cell were used for the initial experiments. Since quantification by flow cytometry can only be done between 0.2 µm and 1 µm, later quantification was done using NTA.

Extracellular vesicles quantification and size evaluation using NTA

To evaluate the size distribution and quantity of the EVs in more detail, the samples were evaluated using nanoparticle tracking analysis (NTA, NanoSight LM10, Malvern Panalytical, Software "NTA 3.0"). All samples were diluted 40 folds in 0.9% NaCl to a final volume of 1 ml. The ideal measurement concentrations were found by pre-testing the optimal particle per frame value (20–100 particles/frame). Following settings were set according to the manufacturer's software manual (NanoSight NS300 User Manual, MAN0541-01-EN-00, 2017): camera (type sCMOS) level was increased until 8 so that all particles were distinctly visible, slide shutter 350, slide gain 250, and temperature 25°C. The software NTA 3.0 was used for the analysis of three videos of 60 s recorded per sample, with a detection threshold of 5.

Scanning helium-ion microscopy examination of ASC-EVs

EVs isolated from the culture medium of ASCs were morphologically evaluated using scanning helium-ion microscopy. The ASC-EVs were fixed in 2.5% glutaraldehyde in PBS, pH 7.4 and incubated at 4°C at least for 24 h. Later, the vesicles were slowly filtered onto a ceramic filter with a pore size of 20 nm (GE Healthcare, Sweden) supported by a mixed cellulose ester (MCE) membrane filter with 0.45 µm pore size in a manual filtration unit (Sartorius). Still in the filtration unit, the ASC-EVs on the filter were washed with

500 µl of PBS. Subsequently, the filter was removed and dehydrated in a graded ethanol-series starting with 30% ethanol in a dilution series up to absolute ethanol. Then the filter was critical point dried (CPD) using a Leica CPD300 critical point drier (Leica Microsystems, Germany) in order to make them vacuum-compatible. During 20 CPD-cycles the ethanol was exchanged by super-critical CO₂, which afterwards evaporated. After the drying process, a part of the ceramic filter was glued onto an aluminum-specimen stub (Agar Scientific, UK) using air-drying conductive silver epoxy (Acheson 1415 Plano, Germany). Then, the samples were sputter-coated with a conductive layer of gold-palladium (9:1) of 30 nm thickness using a tabletop sputtering device Leica EM SCD 500. The micrographs were acquired with a Zeiss Orion NanoFab HIM (Carl Zeiss Microscopy, MA). He⁺ ions with an ion-landing energy of 25 kV and a beam current of about 0.7 pA allowed for high surface sensitivity and minimum damaging of the specimen. Secondary electrons were detected using an Everhart-Thornley-type secondary electron detector for imaging, with a pixel dwell time of 0.5 µs. Noise reduction was achieved by scanning and averaging each line for 64 times.

Multiplex analysis of ASC-EV surface markers

The presence of vesicle surface markers in ASC-EVs was analyzed using the MACSPlex exosome, human, Kit (Miltenyi Biotec) following the manufacturer's instructions. Briefly, 50 µl of the antibody-coated capture beads targeting thirty-seven different membrane antigens were incubated during 16 h with 1.5x10⁹ ASC-EVs resuspended in 120 µl of 0.9% NaCl. Later, an antibody cocktail targeting the tetraspanins CD9, CD81 and CD63 was added and incubated for 1 h. The flow cytometry analyses were done using the FACS Canto II (BD Biosciences) and the FACS Diva software (BD Biosciences). The calibration of the flow cytometer was set as recommended by the kit manufacturer using the setup beads (Singlets) provided. The values of median fluorescence intensity below the provided corresponding control antibodies were considered as threshold and as negative. Data are normalized to the mean values of the median fluorescence intensities of the tetraspanins (CD9, CD63 and CD81). The samples were evaluated in three independent runs.

BV-2 stimulation with β-amyloid aggregates

The immunomodulatory capacity of human ASC-EVs or mouse MSC-EVs in the prevention of the microglia cells activation towards stimulation with Aβ aggregates was tested *in vitro*. BV-2 cells were seeded at 2.1x10⁵ cells/cm² in a 24-well plate and incubated 24 h in DMEM-HG with GlutaMAX and 2% FBS (Life technologies). Microglia cells were primed using 1 µg/ml of lipopolysaccharides (LPS, Sigma-Aldrich) for 3 h. Later, the cells were washed with cell culture media, prior stimulation with 10 µM of Aβ aggregates for 6 h (for evaluation of gene expression) or 24 h (for evaluation of cytokine secretion and nitric oxide (NO) production) at 37°C and 8% of CO₂. Unstimulated cells were used as negative control (NC). Eight human ASC-EVs or murine MSC-EVs per cell (flow cytometry quantification) were added to BV-2 cells in the respective wells. As control for the effect of the vesicles on BV-2 cells, the cells were incubated with the vesicles after priming and with no further stimulation.

Live cell imaging of EV/microglia co-culture

Immediately after adding the EVs stained with PKH26 Red Fluorescent Cell Linker or CellTracker™ Deep Red dye to BV-2 cells, live cell imaging was performed using an inverse microscope ZEISS Axio Observer.Z1 (Carl Zeiss AG, Germany) equipped with AxioCam MR Rev3 camera (image size of 1388 x 1040 pixel, bit depth of 12), Plan-Apochromat 63x/1.40 oil objective and live cell incubation setup consisting of Heating Unit XL at temperature of 37°C and CO₂ Module S at 8% CO₂ (PeCon GmbH, Germany). Measurements were done in z-stacks with Brightfield and Fluorescence exposure time of each 300 ms, filter set no. 50 (Carl Zeiss AG) for Cy5, excitation filter at 625–655 nm, emission filter at 665–715 nm and dichroic mirror at 660 nm for EVs with CellTracker™ Deep Red dye as well as brightfield exposure time of 80 ms, fluorescence exposure time of 220 ms, filter set no. 43 (Carl Zeiss AG) for DsRed, excitation filter at 538–562 nm, emission filter at 570–640 nm and dichroic mirror at 570 nm for EVs with PKH26 Red Fluorescent Cell Linker dye.

Statistical Analysis

Each experiment was performed at least three independent times. The data represents the mean ± SEM calculated from all assays. Statistical significance was calculated using the Mann Whitney U-Test or Student's *t*-test as indicated and shown with asterisks (**p* < 0.05, ***p* < 0.01, ****p* < 0.001). All analyses were done with the Graph Pad Prism v.6.02 (San Diego, CA, USA) or SigmaPlot 14.0 (Systat Software) software.

Results

Human ASCs from donors of different ages maintain MSC characteristics

The adipose-derived stromal vascular fraction (SVF) is a heterogeneous population of cells including MSC-like cells, endothelial progenitor cells, and hematopoietic cells. ASCs were isolated from adipose tissue obtained from eyelid and breast (Fig. 1A) and analyzed according to the position statement of the International Society for Cellular Therapy [10, 45]. The ASC yield was much greater for donors aged less than 40 years compared to older donors; the yield gradually fell as the donor age rose (data not shown). The human ASCs have grown *in vitro* as typical spindle-like structures adherent on tissue culture plastic without obvious changes in shape, size or density (Fig. 1A). The surface protein expression of human ASCs at early passage was examined by flow cytometry (Fig. 1B). ASCs were positive for stem cell markers CD73, CD90, and CD105. Conversely, isolated ASCs had a low or no expression of the hematopoietic marker CD45. There was no difference between groups regarding cell morphology (data not shown). ASCs were also evaluated in terms of their multilineage capacity to exhibit chondrogenic, osteogenic and adipogenic differentiation (Fig. 1C). Three-lineage differentiation assay was performed to test the multipotency of human ASCs. After two-week adipogenic differentiation, massive amounts of lucent lipid droplets of a high refractivity could be visualized in the cytoplasm as observed under a microscope, and the color red was developed after using Oil Red O. The osteogenically-differentiated cells in the monolayer culture were stained by Fast red and proteoglycan-rich cartilaginous matrix produced

during chondrogenic differentiation were visualized by Alcian blue. These results indicated that we obtained human ASCs with high purity and multipotent ability.

Screening the ASC-EVs immunomodulatory capacity

Previously we demonstrated the immunomodulatory capacity of mouse MSC-EVs towards LPS stimulated BV-2 cells [17]. In order to assess whether human ASC-EVs act as BV-2 immune modulators in a comparable way to mouse MSC-EVs, we used the same LPS stimulation assay of BV-2 cells and measured the TNF- α secretion using ELISA. EVs from the cell culture supernatants were isolated via differential (ultra)-centrifugation [17] (Supplementary Fig. 1). The capacity of human ASC-EVs to prevent TNF- α secretion by LPS stimulated BV-2 cells was highly variable among all donors (Fig. 2A; $n = 12$). However, the half of the tested donors ($n = 6$) showed no effect on the secretion of TNF- α , regardless of the concentration, whereas the other half ($n = 6$) showed a strong immunomodulatory potential (Fig. 2B and 2C). We selected the immunomodulatory donors for EV characterization and one donor for testing the effect on BV-2 cells after stimulation with A β aggregates (Fig. 2D).

Characterization of ASC-derived EVs

Human ASC-derived EVs were characterized according to the minimal information for studies of EVs [39] using helium ion microscopy for structural evaluation and visualization, NanoSight for quantification and size evaluation, and MACSPlex assays to evaluate the present surface antigens. Although EV biogenesis, cargo, and function might widely differ, EVs remain mainly classified according to their size unless specific markers for subcellular origin are established. Using helium ion microscopy, the ASC-EVs were visualized as spherical structures with sizes ranging from small EVs (~ 30 nm) to medium/large EVs (~ 400 nm), confirming herewith the heterogeneity of the EV population (Fig. 3A). The NanoSight evaluation also showed that the human ASC-EVs had dimensions ranging from 30 to 400 nm with a mean of about 157 nm and a particle concentration of 3.1×10^8 particles/ml of original cell culture supernatant (Fig. 3B). In addition, using flow cytometry, we analyzed the surface expression of various molecules on human ASC-EVs (Fig. 3C). We detected 11 different molecules variably expressed; in particular, the tetraspanins and genuine EV markers CD9, CD63 and CD81. Moreover, EVs released from human ASCs were positive for the MSC markers: CD29, CD44, CD105 and CD146 and negative for CD45 (data not shown). Besides, a higher expression of MCSP and ROR1, cancer markers, with strong median fluorescence intensities (MFI) were found.

Stimulation of BV-2 cells with A β aggregates

Neuroinflammation in AD is associated with an increase of the number of microglia, accompanied by phenotypic changes leading to an enhanced cytokine and chemokine release. A β aggregates stimulate microglia cells to produce pro-inflammatory signals, which are neurotoxic mediators and might propagate an inflammatory cycle in the central nervous system (CNS) [35]. In order to establish an *in vitro* model for microglia stimulation by A β aggregates, we used a widely reported protocol for A β (1–42) aggregation during 86 h at 37°C in PBS [22, 38]. Resting BV-2 cells did not show to be responsive towards A β aggregates without prior stimulation (data not shown). Hence, BV-2 cells were primed during 3 h with LPS

to induce microglia activation. An overview of the experimental procedure is shown in Supplementary Fig. 2.

Activated BV-2 cells were stimulated with different concentrations of A β aggregates (1 μ M, 2.5 μ M, 5 μ M, 10 μ M and 20 μ M) and responses were assessed by measuring TNF- α secretion (Supplementary Fig. 3A) and nitric oxide (NO) release (Supplementary Fig. 3B). A clear upregulation of TNF- α (Fig. 4A) and NO (Fig. 4B) was reached at 10 μ M of A β aggregates, hence that was the concentration selected for all further stimulation experiments.

Suppression of pro-inflammatory molecules by EVs in BV-2 cells stimulated with A β aggregates

We recently showed that EVs derived from mouse MSCs attenuate the production of inflammatory molecules induced by LPS stimulation of BV-2 cells [17]. In accordance with these investigations, we examined the influence of human ASC-EVs and mouse MSC-EVs on cytokine production by A β -stimulated mouse BV-2 cells to assess whether EVs of different species performed the same function on mouse BV-2 cells.

Therefore, BV-2 cells were co-cultured with human ASC-EVs or mouse MSC-EVs in the presence and absence of A β aggregates. The concentrations of EVs was the same used in our previous study, where we also tested the immunomodulatory potential of mouse MSC-EV on microglia cells [17]. The mRNA expressions of TNF- α and PTGS2 were evaluated 6 h after stimulation, while secretions of TNF- α and NO were analyzed after 24 h. As shown in Fig. 5A and B the stimulation of BV-2 cells with A β aggregates induced an upregulation of TNF- α and PTGS2 transcripts, compared to the controls. Importantly, the presence of human ASC-EVs or mouse MSC-EVs significantly prevented the upregulation of these pro-inflammatory molecules. Addition of human ASC-EVs or mouse MSC-EVs to only LPS primed cells did not have any significant effect on the transcription of TNF- α and PTGS-2 (Fig. 5A and B).

However, A β -stimulated BV-2 cells showed a two times higher concentration of TNF- α and NO in the cell culture supernatant compared to non-stimulated cells (data not shown). A significant EV effect on A β -stimulated BV-2 cells were observed on TNF- α and NO protein levels induced by mouse MSC-EVs. The human ASC-EVs had no effect on TNF- α and NO secretion of A β -stimulated mouse BV-2 cells. (Fig. 5C and D)

The β -amyloid receptor CD36 is affected by human and mouse MSC-derived EVs

The cell surface receptor CD36 is a class B scavenger receptor, expressed on microglia cells in normal brains and brains from AD patients. CD36 mediates macrophages and microglia responses to A β aggregates, playing a major role in the pro-inflammatory events associated with AD [11]. Approximately 65% of BV-2 cells express CD36 after priming (Fig. 6A). We observed that the CD36 expression decreases to approximately 51% after stimulation with A β aggregates during 24 h, which has been described as a

common process, due to the internalization of the molecules during phagocytosis [8, 36]. Interestingly, BV-2 cells stimulated with A β aggregates in the presence of human ASC-EVs or mouse MSC-EVs show a significant increase of CD36, compared to cells stimulated in absence of the EVs (64%, $p = 0.0286$ and 72%, $p = 0.0095$, respectively).

MSC-derived EVs prevent the upregulation of CD206 by BV-2 after β -amyloid stimulation

CD206 is a C-type lectin and mannose receptor usually used as a surface marker for M2 macrophages [6, 48]. We observed that CD206 is expressed by 7% of the LPS primed BV-2 cells (NC) (Fig. 6B). While, after stimulation with A β aggregates, the number of CD206 expressing cells gained up to 25%. Interestingly, co-culture with human ASC-EVs or mouse MSC-EVs significantly prevented the upregulation triggered by A β stimulation (10%, $p = 0.0152$ and 9%, $p = 0.0087$ respectively). The expression of the cell surface molecules CD45, CD11b, CD86 and CD80 was also evaluated and remained unaffected after BV-2 stimulation with A β aggregates (data not shown).

Human ASC-EVs and mouse MSC-EVs do not affect the uptake of latex beads by BV-2 cells

The internalization of the A β complexes occurs via phagocytosis [13]. Given the observation that internalization of CD36 was prevented in presence of human ASC-EVs and mouse MSC-EVs (Fig. 6A), we evaluated the general phagocytic capacity of microglia cells in the presence of EVs. Therefore, we established a phagocytosis assay based on flow cytometry that measures the bulk phase uptake of latex beads to examine the phagocytic response. First, we incubated BV-2 cells with latex beads and found that almost 67% of all BV-2 cells took up at least one bead. LPS priming of the cells slightly suppressed the phagocytosis of latex beads. However, no statistically significant difference was found in beads phagocytosis between LPS untreated and treated cells (Supplementary Fig. 4). Stimulation of BV-2 cells with A β aggregates prevented the latex bead uptake. However, the presence of human ASC-EVs or mouse MSC-EVs did not show any influence on the capacity of BV-2 cells to take up latex beads.

ASC-EVs are internalized by microglia cells

EVs are expected to have different mechanisms for interacting with the cells. In order to deliver a signal, EVs need to fuse with the membrane of the target cells, either with the plasma membrane or after endocytic uptake with the endosomal membrane [30]. We demonstrated that human ASC-EVs influence the inflammatory responses of stimulated microglia cells (Fig. 5). However, the mechanism of vesicle-cell interaction is still an open question. In order to understand how human ASC-EVs might deliver a signal to BV-2 cells, two different EV labelling protocols were used: (i) the vesicle content stained with CellTracker™ Deep Red dye and (ii) the vesicle membrane stained with PKH26 Red Fluorescent Cell Linker. It was

observed utilizing live cell imaging that the vesicles only internalize in a subset of BV-2 cells for both protocols.

Moreover, the content of the vesicles (Fig. 7A), if released, and the vesicle membrane (Fig. 7B) were spread anywhere inside of a BV-2 cell. Based on z-stacks obtained during live cell imaging, and after subsequent 3D image post processing and 3D visualization applying a combination of both surface and volume rendering using Wolfram Mathematica 12 (Wolfram Research Inc., Champaign, IL, USA), we could determine that the vesicle content was dissolved and the membranes of the ASC-EVs seemed to be remaining centrally inside the cytoplasm of the BV-2 cells. This indicates that the mechanism of interaction involves the uptake of the complete vesicle and not only the fusion of the EVs with the cell membrane.

Furthermore, the EV-membranes were exchanged via cell junctions between cells in culture over a timescale of 5 days (Supplementary video 1). Hence, we demonstrate that ASC-EVs are internalized by BV-2 cells and that the vesicle content is released inside the cells.

Discussion

Neuroinflammation is a major problem in neurodegenerative diseases as Alzheimer disease [40]. The microglia cells, as the CNS immune cells, are in charge of the immune surveillance, and become easily activated towards various stimuli. However, the functionality of microglia cells is important for the homeostasis maintenance. Hence, a tight balance between responsive but not activated microglia cells is essential in order to regulate the chronic inflammation leading to neurodegeneration. In a previous study, we had demonstrated that mouse MSC-EVs represent powerful modulators of mouse microglia responses towards LPS stimulation [17]. Here, we show that MSC-EVs can act also as immune modulators of microglia responses after stimulation with A β aggregates.

Human microglia cells can be obtained from human brains post mortem only. Therefore, most of the research about the inflammatory responses of microglia cells has been gathered using murine cells, *in vitro* and/or *in vivo*. Hence, assessing new therapeutic approaches for neurodegenerative diseases involving microglia cells represents an enormous challenge. To develop *in vitro* models that mimic the inflammatory processes occurring in the diseased CNS is of major importance for initial screening of possible therapeutic agents. The mouse cell line BV-2 has been repetitively used as an alternative model system for primary microglia cultures or for animal experiments evaluating brain inflammation [16]. Our goal was to assess the immunomodulatory capacity of MSC-EVs towards microglia cells. In this study, we demonstrated that EVs derived from human ASCs, exert effects comparable to those from mouse MSCs on modulating the inflammatory responses of microglia cells on mRNA level. Furthermore, our analysis of EVs derived from human ASCs confirmed the concurrent presence of tetraspanins as EV marker and of MSC-derived marker. Importantly, the $\alpha 5$ integrin subunit of the fibronectin receptor (CD49e) was detected on the vesicles. The presence of low levels from this marker on MSCs has been associated with osteoblast differentiation in various studies [18, 32]. In AD, the aggregates that form A β

plaques are closely associated with activated microglia cells. These cells produce molecules as TNF- α , IL-1 β , IL-6 and NO which are known to promote neuroinflammation [41]. The A β peptides are metabolism products usually formed by 36 to 43 amino acids. There are several A β forms found in the brain, but toxicity to the cells has been demonstrated by the A β peptides 1–42 [34]. *In vitro* models to assess the inflammatory responses occurring in the AD brain represent the first approach to evaluate possible therapies. Here, we established an *in vitro* model for microglia stimulation with A β 1–42 aggregates [22, 38] in which we can induce the main responses reported in the AD brain for activated microglia cells. Our model allows assessing the production of pro-inflammatory molecules as TNF- α , IL-6 and NO, which are prevalent in chronic neuroinflammation diseases [3]. Interestingly, we did not observe an upregulation of IL-1 β expression in microglia cells after stimulation with A β 1–42 aggregates (data not shown). Moreover, following our described protocol [17], BV-2 cells require previous priming in order to induce inflammatory responses strong enough for assessing the molecular methods available at our laboratory.

The EVs by themselves do not trigger any inflammatory stimuli; however, their specific content affects the microenvironment. An immunosuppressive capacity of MSC-EVs has been reported previously by Wang et al., showing that human MSC-EVs prevent life-threatening acute Graft Versus Host Disease by modulating immune responses [43]. Subsequently, Sun et al. observed that human MSC-EVs promote functional recovery in spinal cord injury via attenuating inflammation [37]. We could show that both human ASC-EVs and mouse MSC-EVs significantly down-regulated the transcripts of pro-inflammatory mediators as TNF- α and PTGS2. However, only BV-2 cells stimulated with A β in presence of mouse MSC-EVs showed a significant lower secretion of TNF- α and NO, indicating less inflammatory and more neuroprotective effects. The observation of neuroprotective effects from MSC-EVs mediated by the prevention of NO secretion was also previously reported [14]. The group reported that MSC-EVs present high catalase activity, which might provide the protective action against neuronal oxidative stress. The human ASC-EVs had no significant effect on secreted TNF- α nor on NO, which may be explained by the species barrier between human ACS-EVs and mouse BV-2 cells. However, we were able to show that EVs can have the potential to modify the inflammatory response of immune cells by using the mouse-derived MSC-EVs on mouse microglia BV-2 cells.

The percentage of BV-2 cells expressing the surface A β receptor CD36 decreased after stimulation with A β aggregates. This reduction was described to be associated with the A β internalization [11]. Interestingly, the presence of human ASC-EVs and mouse MSC-EVs prevented this reduction, which might imply a prevention of A β internalization. Due to this observation, we hypothesized that the MSC-EVs could affect the phagocytic capacity of microglia cells. However, we did not observe any significant influence in the capacity of BV-2 cells to internalize latex beads. Further it could be important, to evaluate whether the MSC-EVs directly affect the A β internalization. In contrast, upregulation of CD36 and CD206 expression in response to A β aggregates was significantly prevented in presence of human and mouse MSC-EVs. The mannose receptor CD206 is commonly associated with the anti-inflammatory activation of macrophages [6, 48], and increased in the late stage of lesions developed by ischemia patients [47]. Taking together, we observed that the autologous setting, using mouse MSC-EVs showed a slightly

stronger effect at preventing pro-inflammatory responses of BV-2 cells towards A β than using human ASC-EVs.

The interaction mechanism of EVs with the target cell might highly vary [30]. Here, we could demonstrate using live cell microscopy that the human ASC-EVs are internalized by a random subset of BV-2 cells, and the EV-content is distributed in the cytoplasm. The EV-membranes were spread towards neighbored BV-2 cells via tubuli, which suggests that the signals delivered by the EVs might reach more than one cell. Hence, from a therapeutic perspective it should be considered that even the local application of MSC-EVs could provide broader systemic effects. As we showed previously, in BV-2 cells, MSC-EVs prevent the MAPK phosphorylation signaling cascade which occurs after LPS stimulation, therewith leading to lower transcription of genes associated with inflammation [17]. MSC-EVs transport a broad number of bioactive molecules; however, the specific EV delivered molecules that are involved in preventing the inflammatory responses from microglia cells, remain to be elucidated.

Conclusion

In conclusion, our findings demonstrate that EVs derived from human ASC and mouse MSCs modulate the activation of BV-2 microglia cells and can prevent the pro-inflammatory response. Therefore, ASC-EVs might be a promising tool for new therapeutic approaches targeting chronic inflammation leading to neurodegeneration, including AD. Moreover, we have shown that human ASC-EVs are internalized by microglia cells, and that the content is distributed to neighboring cells. Thus, EVs have the potential to amplify signals which might be used in therapeutic approach. Furthermore, MSCs are very attractive in regenerative medicine due to their potential in tissue protection and regeneration. Thus, our findings provide hints that the mechanism of tissue protection may also be carried out by MSC derived EVs.

Declarations

ACKNOWLEDGEMENTS

We would like to thank Dr. med. Markus Tränkle from Klinik am Rosenthal, Leipzig and all those involved in donating samples for biomedical research. The authors thank Alessa Webers and Dr. Hannah Scheiblich, from the Henneka Lab at the German Center for Neurodegenerative Diseases, for their valuable help to establish the protocol for stimulation of BV-2 cells using A β aggregates. We thank Dr. Alexander Ewe, from the Rudolf-Boehm-Institute for Pharmacology and Toxicology at the Leipzig University, for the introduction to the NanoSight. The authors are also grateful for using the analytical facilities, in particular the scanning helium-ion microscope, at ProVIS - Centre for Chemical Microscopy at UFZ Leipzig, which is supported by European Regional Development Funds (EFRE – Europe funds Saxony) and the Helmholtz Association. Finally, we would like to thank Katrin Arnold and Annett Friedrich-Stöckigt who have shared their knowledge and experience in the MSC field. The authors also thank Eric Possardt for his support during the live cell imaging of the MSC-EVs and Dr. Thomas Grunwald for his advices and discussions. This study was supported by the High-Performance Center for Chemical

and Biosystems, and co-financed with tax funds from Ministry of Saxony (Grant number 100312141). YJ was supported by the TALENTA speed up program (Confirmation date 19.12.2017) from the Fraunhofer organization and by the Fraunhofer Cluster of Excellence for Immune-mediated Diseases CIMD (Project number 601241). PR was supported by a doctoral scholarship from the medical faculty of Leipzig (Project number 990101-113).

Author contributions

Dorota Kaniowska: Conception and design, Collection and/or assembly of data, Data analysis and interpretation, Manuscript writing

Kerstin Wenk: Collection and/or assembly of data, Data analysis and interpretation, Manuscript writing

Phil Rademacher: Collection and/or assembly of data, Data analysis and interpretation

Ronald Weiss: Collection and/or assembly of data, Data analysis and interpretation

Claire Fabian: Collection and/or assembly of data, Data analysis and interpretation

Isabell Schulz: Collection and/or assembly of data, Data analysis and interpretation

Max Guthardt: Collection and/or assembly of data, Data analysis and interpretation

Franziska Lange: Administrative support, Collection and/or assembly of data, Data analysis and interpretation

Sebastian Greiser: Collection and/or assembly of data, Data analysis and interpretation

Matthias Schmidt: Collection and/or assembly of data, Data analysis and interpretation

Ulf-Dietrich Braumann: Data analysis and interpretation

Frank Emmrich: Administrative support, Financial support

Ulrike Koehl: Administrative support, Financial support

Yarúa Jaimes: Conception and design, Collection and/or assembly of data, Data analysis and interpretation, Manuscript writing

All authors contributed to the study conception and design and have read and approved the final manuscript.

Funding

This study was supported by the High-Performance Center for Chemical and Biosystems, and co-financed with tax funds from Ministry of Saxony (Grant number 100312141). YJ was supported by the TALENTA

speed up program (Confirmation date 19.12.2017) from the Fraunhofer organization and by the Fraunhofer Cluster of Excellence for Immune-mediated Diseases CIMD (Project number 601241). PR was supported by a doctoral scholarship from the medical faculty of Leipzig (Project number 990101-113).

Ethical Approval

Approval to perform the study was obtained from the Ethical Commission (291-16-ek) of the University of Leipzig and after written informed consent from all the donors.

Competing interests

We have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Consent to Publish

Authors are responsible for correctness of the statements provided in the manuscript.

Consent to participate

Informed consent was obtained from all individual participants included in the study.

Availability of data and materials

The datasets generated and analyzed during the current study are available from the corresponding author on reasonable request.

References

1. Boche, D., Perry, V. H., & Nicoll, J. A. R. (2013). Review: activation patterns of microglia and their identification in the human brain. *Neuropathology and applied neurobiology*, 39(1), 3–18. doi:10.1111/nan.12011
2. Bonventre, J. V., & Yang, L. (2011). Cellular pathophysiology of ischemic acute kidney injury. *The Journal of clinical investigation*, 121(11), 4210–4221. doi:10.1172/JCI45161
3. Brennan, F. M., Maini, R. N., & Feldmann, M. (1995). Cytokine expression in chronic inflammatory disease. *British medical bulletin*, 51(2), 368–384. doi:10.1093/oxfordjournals.bmb.a072967
4. Carniglia, L., Ramírez, D., Durand, D., Saba, J., Turati, J., Caruso, C., et al. (2017). Neuropeptides and Microglial Activation in Inflammation, Pain, and Neurodegenerative Diseases. *Mediators of inflammation*, 2017, 5048616. doi:10.1155/2017/5048616.
5. Chang, K. A., Kim, H. J., Joo, Y., Ha, S., & Suh, Y. H. (2014). The therapeutic effects of human adipose-derived stem cells in Alzheimer's disease mouse models. *Neuro-degenerative diseases*, 13(2–3), 99–102. doi:10.1159/000355261

6. Cherry, J. D., Olschowka, J. A., & O'Banion, M. K. (2014). Neuroinflammation and M2 microglia: the good, the bad, and the inflamed. *Journal of Neuroinflammation*, 11, 98. doi:10.1186/1742-2094-11-98
7. Cocucci, E., & Meldolesi, J. (2015). Ectosomes and exosomes: shedding the confusion between extracellular vesicles. *Trends in cell biology*, 25(6), 364–372. doi:10.1016/j.tcb.2015.01.004
8. Coraci, I. S., Husemann, J., Berman, J. W., Hulette, C., Dufour, J. H., Campanella, G. K., et al. (2002). CD36, a Class B Scavenger Receptor, Is Expressed on Microglia in Alzheimer's Disease Brains and Can Mediate Production of Reactive Oxygen Species in Response to β -Amyloid Fibrils. *The American Journal of Pathology*, 160(1), 101–112
9. Cui, L., Yin, S., Liu, W., Li, N., Zhang, W., & Cao, Y. (2007). Expanded adipose-derived stem cells suppress mixed lymphocyte reaction by secretion of prostaglandin E2. *Tissue engineering*, 13(6), 1185–1195. doi:10.1089/ten.2006.0315
10. Dominici, M., Le Blanc, K., Mueller, I., Slaper-Cortenbach, I., Marini, F., Krause, D., et al. (2006). Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society for Cellular Therapy position statement. *Cytotherapy*, 8(4), 315–317. doi:10.1080/14653240600855905
11. El Khoury, J. B., Moore, K. J., Means, T. K., Leung, J., Terada, K., Toft, M., et al. (2003). CD36 mediates the innate host response to beta-amyloid. *The Journal of experimental medicine*, 197(12), 1657–1666. doi:10.1084/jem.20021546
12. Farzamfar, S., Hasanpour, A., Nazeri, N., Razavi, H., Salehi, M., Shafei, S., et al. (2019). Extracellular micro/nanovesicles rescue kidney from ischemia-reperfusion injury. *Journal of cellular physiology*, 234(8), 12290–12300. doi:10.1002/jcp.27998
13. Gabandé-Rodríguez, E., Keane, L., & Capasso, M. (2020). Microglial phagocytosis in aging and Alzheimer's disease. *Journal of neuroscience research*, 98(2), 284–298. doi:10.1002/jnr.24419
14. de Godoy, M. A., Saraiva, L. M., de Carvalho, L. R. P., Vasconcelos-Dos-Santos, A., Beiral, H. J. V., Ramos, A. B., et al. (2018). Mesenchymal stem cells and cell-derived extracellular vesicles protect hippocampal neurons from oxidative stress and synapse damage induced by amyloid- β oligomers. *The Journal of biological chemistry*, 293(6), 1957–1975. doi:10.1074/jbc.M117.807180
15. Graeber, M. B., Li, W., & Rodriguez, M. L. (2011). Role of microglia in CNS inflammation. *FEBS letters*, 585(23), 3798–3805. doi:10.1016/j.febslet.2011.08.033
16. Henn, A., Lund, S., Hedtjärn, M., Schrattenholz, A., Pörzgen, P., & Leist, M. (2009). The suitability of BV2 cells as alternative model system for primary microglia cultures or for animal experiments examining brain inflammation. *ALTEX*, 26(2), 83–94. doi:10.14573/altex.2009.2.83
17. Jaimes, Y., Naaldijk, Y., Wenk, K., Leovsky, C., & Emmrich, F. (2017). Mesenchymal Stem Cell-Derived Microvesicles Modulate Lipopolysaccharides-Induced Inflammatory Responses to Microglia Cells. *Stem cells (Dayton, Ohio)*, 35(3), 812–823. doi:10.1002/stem.2541
18. Keselowsky, B. G., Wang, L., Schwartz, Z., Garcia, A. J., & Boyan, B. D. (2007). Integrin α (5) controls osteoblastic proliferation and differentiation responses to titanium substrates presenting

- different roughness characteristics in a roughness independent manner. *Journal of biomedical materials research. Part A*, 80(3), 700–710. doi:10.1002/jbm.a.30898
19. Kim, S., Chang, K. A., Kim, J., Park, H. G., Ra, J. C., Kim, H. S., et al. (2012). The preventive and therapeutic effects of intravenous human adipose-derived stem cells in Alzheimer's disease mice. *PloS one*, 7(9), e45757. doi:10.1371/journal.pone.0045757
 20. Kordelas, L., Rebmann, V., Ludwig, A. K., Radtke, S., Ruesing, J., Doeppner, T. R., et al. (2014). MSC-derived exosomes: a novel tool to treat therapy-refractory graft-versus-host disease. *Leukemia*, 28(4), 970–973. doi:10.1038/leu.2014.41
 21. Kronsteiner, B., Wolbank, S., Peterbauer, A., Hackl, C., Redl, H., van Griensven, M., et al. (2011). Human mesenchymal stem cells from adipose tissue and amnion influence T-cells depending on stimulation method and presence of other immune cells. *Stem cells and development*, 20(12), 2115–2126. doi:10.1089/scd.2011.0031
 22. Kummer, M. P., Hermes, M., Delekarte, A., Hammerschmidt, T., Kumar, S., Terwel, D., et al. (2011). Nitration of tyrosine 10 critically enhances amyloid β aggregation and plaque formation. *Neuron*, 71(5), 833–844. doi:10.1016/j.neuron.2011.07.001
 23. Lai, P., Chen, X., Guo, L., Wang, Y., Liu, X., Liu, Y., et al. (2018). A potent immunomodulatory role of exosomes derived from mesenchymal stromal cells in preventing cGVHD. *Journal of hematology & oncology*, 11(1), 135. doi:10.1186/s13045-018-0680-7
 24. Lee, D. Y., Oh, Y. J., & Jin, B. K. (2005). Thrombin-activated microglia contribute to death of dopaminergic neurons in rat mesencephalic cultures: dual roles of mitogen-activated protein kinase signaling pathways. *Glia*, 51(2), 98–110. doi:10.1002/glia.20190
 25. Lee, J. M., Jung, J., Lee, H. J., Jeong, S. J., Cho, K. J., Hwang, S. G., et al. (2012). Comparison of immunomodulatory effects of placenta mesenchymal stem cells with bone marrow and adipose mesenchymal stem cells. *International immunopharmacology*, 13(2), 219–224. doi:10.1016/j.intimp.2012.03.024
 26. Liew, L. C., Katsuda, T., Gailhouste, L., Nakagama, H., & Ochiya, T. (2017). Mesenchymal stem cell-derived extracellular vesicles: a glimmer of hope in treating Alzheimer's disease. *International immunology*, 29(1), 11–19. doi:10.1093/intimm/dxx002
 27. Ma, T., Gong, K., Ao, Q., Yan, Y., Song, B., Huang, H., et al. (2013). Intracerebral transplantation of adipose-derived mesenchymal stem cells alternatively activates microglia and ameliorates neuropathological deficits in Alzheimer's disease mice. *Cell transplantation*, 22(Suppl 1), S113–26. doi:10.3727/096368913X672181
 28. McIntosh, K., Zvonic, S., Garrett, S., Mitchell, J. B., Floyd, Z. E., Hammill, L., et al. (2006). The immunogenicity of human adipose-derived cells: temporal changes in vitro. *Stem cells (Dayton, Ohio)*, 24(5), 1246–1253. doi:10.1634/stemcells.2005-0235
 29. Meyer-Luehmann, M., Spire-Jones, T. L., Prada, C., Garcia-Alloza, M., de Calignon, A., Rozkalne, A., et al. (2008). Rapid appearance and local toxicity of amyloid- β plaques in a mouse model of Alzheimer's disease. *Nature*, 451(7179), 720–724. doi:10.1038/nature06616

30. Mulcahy, L. A., Pink, R. C., & Carter, D. R. F. (2014). Routes and mechanisms of extracellular vesicle uptake. *Journal of extracellular vesicles*. 3, doi:10.3402/jev.v3.24641
31. Niemeyer, P., Kornacker, M., Mehlhorn, A., Seckinger, A., Vohrer, J., Schmal, H., et al. (2007). Comparison of immunological properties of bone marrow stromal cells and adipose tissue-derived stem cells before and after osteogenic differentiation in vitro. *Tissue engineering*, 13(1), 111–121. doi:10.1089/ten.2006.0114
32. Olivares-Navarrete, R., Raz, P., Zhao, G., Chen, J., Wieland, M., Cochran, D. L., et al. (2008). Integrin alpha2beta1 plays a critical role in osteoblast response to micron-scale surface structure and surface energy of titanium substrates. *Proceedings of the National Academy of Sciences of the United States of America*, 105(41), 15767–15772. doi:10.1073/pnas.0805420105
33. Puissant, B., Barreau, C., Bourin, P., Clavel, C., Corre, J., Bousquet, C., et al. (2005). Immunomodulatory effect of human adipose tissue-derived adult stem cells: comparison with bone marrow mesenchymal stem cells. *British journal of haematology*, 129(1), 118–129. doi:10.1111/j.1365-2141.2005.05409.x
34. Querfurth, H. W., & LaFerla, F. M. (2010). Alzheimer's disease. *The New England journal of medicine*, 362(4), 329–344. doi:10.1056/NEJMr0909142
35. Rojanathammanee, L., Floden, A. M., Manocha, G. D., & Combs, C. K. (2015). Attenuation of microglial activation in a mouse model of Alzheimer's disease via NFAT inhibition. *Journal of Neuroinflammation*. 12, doi:10.1186/s12974-015-0255-2
36. Stewart, C. R., Stuart, L. M., Wilkinson, K., van Gils, J. M., Deng, J., Halle, A., et al. (2009). CD36 ligands promote sterile inflammation through assembly of a Toll-like receptor 4 and 6 heterodimer. *Nature immunology*, 11(2), 155–161. doi:10.1038/ni.1836
37. Sun, G., Li, G., Li, D., Huang, W., Zhang, R., Zhang, H., et al. (2018). hucMSC derived exosomes promote functional recovery in spinal cord injury mice via attenuating inflammation. *Materials science & engineering. C, Materials for biological applications*, 89, 194–204. doi:10.1016/j.msec.2018.04.006
38. Teplow, D. B. (2006). Preparation of Amyloid β -Protein for Structural and Functional Studies. *Amyloid, Prions, and Other Protein Aggregates, Part C (Vol (413 vol., pp. 20–33). Methods in Enzymology)* Elsevier
39. Théry, C., Witwer, K. W., Aikawa, E., Alcaraz, M. J., Anderson, J. D., Andriantsitohaina, R., et al. (2018). a position statement of the International Society for Extracellular Vesicles and update of the MISEV2014 guidelines. *Journal of extracellular vesicles*, 7(1), 1535750. doi:10.1080/20013078.2018.1535750.Minimal information for studies of extracellular vesicles 2018 (MISEV2018
40. Turgeman, G. (2015). The therapeutic potential of mesenchymal stem cells in Alzheimer's disease: converging mechanisms. *Neural regeneration research*, 10(5), 698–699. doi:10.4103/1673-5374.156953

41. Walker, D. G., & Lue, L. F. (2005). Investigations with cultured human microglia on pathogenic mechanisms of Alzheimer's disease and other neurodegenerative diseases. *Journal of neuroscience research*, 81(3), 412–425. doi:10.1002/jnr.20484
42. Wang, F., Jia, Y., Liu, J., Zhai, J., Cao, N., Yue, W., et al. (2017). Dental pulp stem cells promote regeneration of damaged neuron cells on the cellular model of Alzheimer's disease. *Cell biology international*, 41(6), 639–650. doi:10.1002/cbin.10767
43. Wang, L., Gu, Z., Zhao, X., Yang, N., Wang, F., Deng, A., et al. (2016). Extracellular Vesicles Released from Human Umbilical Cord-Derived Mesenchymal Stromal Cells Prevent Life-Threatening Acute Graft-Versus-Host Disease in a Mouse Model of Allogeneic Hematopoietic Stem Cell Transplantation. *Stem cells and development*, 25(24), 1874–1883. doi:10.1089/scd.2016.0107
44. Wang, N., Li, Q., Zhang, L., Lin, H., Hu, J., Li, D., et al. (2012). Mesenchymal stem cells attenuate peritoneal injury through secretion of TSG-6. *PloS one*, 7(8), e43768. doi:10.1371/journal.pone.0043768
45. Wuchter, P., Bieback, K., Schrezenmeier, H., Bornhäuser, M., Müller, L. P., Bönig, H., et al. (2015). Standardization of Good Manufacturing Practice-compliant production of bone marrow-derived human mesenchymal stromal cells for immunotherapeutic applications. *Cytotherapy*, 17(2), 128–139. doi:10.1016/j.jcyt.2014.04.002
46. Yoo, K. H., Jang, I. K., Lee, M. W., Kim, H. E., Yang, M. S., Eom, Y., et al. (2009). Comparison of immunomodulatory properties of mesenchymal stem cells derived from adult human tissues. *Cellular immunology*, 259(2), 150–156. doi:10.1016/j.cellimm.2009.06.010
47. Zrzavy, T., Hametner, S., Wimmer, I., Butovsky, O., Weiner, H. L., & Lassmann, H. (2017). Loss of 'homeostatic' microglia and patterns of their activation in active multiple sclerosis. *Brain: a journal of neurology*, 140(7), 1900–1913. doi:10.1093/brain/awx113
48. Zrzavy, T., Machado-Santos, J., Christine, S., Baumgartner, C., Weiner, H. L., Butovsky, O., et al. (2018). Dominant role of microglial and macrophage innate immune responses in human ischemic infarcts. *Brain pathology (Zurich, Switzerland)*, 28(6), 791–805. doi:10.1111/bpa.12583
49. Zuk, P. A., Zhu, M., Ashjian, P., de Ugarte, D. A., Huang, J. I., Mizuno, H., et al. (2002). Human adipose tissue is a source of multipotent stem cells. *Molecular biology of the cell*, 13(12), 4279–4295. doi:10.1091/mbc.e02-02-0105
50. Zuk, P. A., Zhu, M., Mizuno, H., Huang, J., Futrell, J. W., Katz, A. J., et al. (2001). Multilineage cells from human adipose tissue: implications for cell-based therapies. *Tissue engineering*, 7(2), 211–228. doi:10.1089/107632701300062859

Figure 7

Figure 7 is a video that can be found in the supplementary file section

Figures

Figure 1

Isolation and characterization of human adipose derived cells (ASCs). (A) Human subcutaneous adipose tissues (2-10 gr) from different age groups and donors were digested with collagenase IV. After centrifugation, the stromal-vascular fraction was isolated and transferred to flasks completely filled with culture medium. The ASCs were cultured in 20% FBS DMEM-LG with medium changes every 3–4 days. The cell surface expression of the common ASC markers was evaluated by flow cytometry of 4

immunomodulatory donors (B), showing that the cells did not express hematopoietic lineage markers, such as CD45, and were positive for CD73, CD90 and CD105 n=4. (C) Cytochemical staining of differentiated and undifferentiated ASCs. Fast Red staining after osteogenic differentiation show a calcium deposition. Alcian blue staining after chondrogenic differentiation show marked deposition of glycosaminoglycans in the matrix. Oil red O stain after induction of adipogenic differentiation show cytoplasmic neutral triglyceride droplets. Scale bars = 100 μ m.

Figure 2

Immunomodulatory screening of extracellular vesicles from adipose stromal/stem cells (ASCs) of different donors. EVs from the ASC supernatant were isolated via differential (ultra)-centrifugation. Resting BV-2 cells were primed for 3 h with 1 μ g/ml LPS before addition of 8 EVs/cell (quantification by flow cytometry) for 24 h. The secretion of TNF- α was evaluated by ELISA in the cell-culture supernatant as an activation read-out for the LPS-stimulated microglia cells. (A) The TNF- α levels from all donors (n=12) were compared. The donors were separated into (B) non-immunomodulatory (n=6) and (C) immunomodulatory donors (n=6). Statistics were calculated with the Mann Whitney U-Test. NC =

negative control (D) Further characterization and functional analysis of EVs were carried out for the immunomodulatory EVs as shown in the scheme.

Figure 3

Characterization of EVs isolated from adipose stromal/stem cells (ASCs) culture supernatant. EVs from the ASC supernatant were isolated via differential (ultra)-centrifugation. (A) One representative micrograph of EVs isolated from ASCs taken by scanning helium-ion microscopy. (B) One representative graph of ASC-EV size evaluation using NanoSight nanoparticle tracking (NTA) analysis. ASC-EVs isolated from cell culture supernatant show a mean size of 157 nm. (C) MACSPlex analysis of the ASC-EV surface markers.

Figure 4

Activation of BV-2 cells with A β aggregates. Resting BV-2 cells were primed for 3 h with 1 μ g/ml LPS before stimulation with 10 μ M of A β aggregates for 24 h. The cell culture supernatants were finally analyzed for secretion of (A) TNF- α by ELISA and (B) NO using a Griess reagent. Data represent means $\pm$ SEM (n=3). Statistics were calculated with the Student's t-test.

Figure 5

The human ASC-EVs and mouse MSC-EVs treatment partially prevents the upregulation of pro-inflammatory molecules in BV-2 cells after $A\beta$ stimulation. EVs from the human ASC and mouse MSC supernatants were isolated via differential (ultra)-centrifugation. Resting BV-2 cells were primed for 3 h with 1 μ g/ml LPS before stimulation with 10 μ M of $A\beta$ aggregates in the presence and absence of 8 EVs/cell (flow cytometry quantification). Gene transcription of (A) TNF_{α} and (B) $PTGS2$ were evaluated by RT-PCR 6 h after stimulation. After 24 h of stimulation, secretion of the pro-inflammatory molecules (C) TNF_{α} and (D) NO to the cell culture supernatant were evaluated using ELISA and Griess reagent

reactions, respectively. Data represent means $\pm$ SEM (n=3). Statistics were calculated with the Mann Whitney U-Test (A and B) or with the Student's t-test (C and D).

Figure 6

The human ASC-EVs and mouse MSC-EVs affect the expression of CD36 and CD206 of BV-2 cells after A β stimulation. EVs from the human ASC and mouse MSC supernatants were isolated via differential (ultra)-centrifugation. Resting BV-2 cells were primed for 3 h with 1 μ g/ml LPS before stimulation with 10 μ M of A β aggregates in the presence and absence of $6.5 \times 10^4 \pm 1.5 \times 10^4$ EVs/cell (NTA quantification). After 24 h of stimulation surface expressions of (A) CD36 and (B) CD206 on BV-2 cells were evaluated by flow cytometry. Data represent means $\pm$ SEM (n = 3). Statistics were calculated with the Mann Whitney U-Test.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Graphicalabstract.tif](#)
- [210328SupplemantarydatacfDKRW.docx](#)
- [Fig.7video.avi](#)