Nasopharyngeal Metastasis From Colorectal Cancer: A Case Report

Ye-Huan Liu

First Affiliated Hospital of Wenzhou Medical University

Bei-Bei Lin

Wenzhou Medical University First Affiliated Hospital

Shi-Xu Lv (✉ lvshixu@foxmail.com)

The First Affiliated Hospital of Wenzhou Medical University

Case report

Keywords: nasopharyngeal metastasis, colorectal cancer

Posted Date: October 23rd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-63557/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Nasopharyngeal metastasis from colorectal cancer has never been reported in the past.

Case presentation: In this paper, we establish a case report of a 79-year-old man suffering from adenocarcinoma of the rectum with distant metastases to the liver, lung, and nasopharynx. Over the past 7 years, he received radical surgery for rectal cancer (miles surgery), chemotherapy, hepatectomy, and pneumonectomy.

Conclusions: We equally discuss the histopathological and clinical aspects associated with this condition.

Background

Metastases from colorectal cancer can occur either by lymphatic or hematogenous spread, with the most common sites involved being the lung and liver. Atypical sites of metastases from colorectal cancer involves organs such as the spleen, thyroid gland, spermatic cord, and skeletal muscles[1-4]. Even though metastases to these sites might occur as a feature of end-stage disease, metastasis to the nasopharynx from colorectal cancer had never been observed. For the very first time, we hereby report a case of nasopharyngeal metastasis from colorectal cancer 7 years after radical surgery for rectal cancer (miles surgery). Furthermore, we also perform a concise review of the histopathological and clinical aspects of this rare entity, and ultimately evaluate adequate diagnostic and therapeutic options based on our observation of the case.

Case Presentation

In October 2012, a 72-year-old man was admitted to our hospital with a chief complain of passing bloody stools for 2 months without seeking any medical attention. The patient denied any abdominal pain and tenesmus. His body was very healthy before. The patient also recorded a weight lost of 2 kg within the past 6 months of presenting at our hospital. A colonoscopy revealed a budding tumor in the rectum that easily bled. A histopathological analysis of the biopsy showed chronic inflammation of the rectal mucosa with high-grade intraepithelial neoplasia, features consistent with highly differentiated carcinoma. A computed tomography (CT) scan of the chest, abdomen, and pelvis, and brain magnetic resonance imaging (MRI) depicted no abnormalities. He underwent radical surgery for rectal cancer (miles surgery) with a subsequent histopathological finding demonstrating an ulcerative type of moderately-poorly differentiated colorectal adenocarcinoma (size 3*3cm), infiltrating all the layers of the intestinal wall and the adipose tissue outside the serous membrane. Four regional lymph nodes were resected and none showed metastases (pT4N0M0, Stage II). Immunohistochemical analysis displayed Bcl-2(-) P53(+) k-ras(-) (Figure1). The status of MSI proteins and BRAF was not verified. There was no family history of colorectal cancer or malignancy, or familial predisposition for colorectal cancer in the patient. The operation was smooth and the postoperative recovery was good. Two months later, he received a

chemotherapeutic regimen including oxaliplatin and capecitabine. However, the oxaliplatin injection was discontinued due to a severe allergic reaction elicited at the first dose and capecitabine was stopped following the grade III hand-foot syndrome (HFS) after four cycles.

During the 7 years following surgical resection, he was on regular follow-up every 3 months with serological and radiological monitoring for disease recurrence. Unfortunately, the patient underwent three operations during this period due to distant metastasis to the liver and lung. In June 2014, he received a hepatectomy after a CT Scan of the abdomen demonstrated a single low density lesion in the right hepatic lobe (Figure2). The patient subsequently underwent two endoscopic pulmonary wedge resections of isolated metastatic pulmonary lesions from colorectal cancer detected by a chest CT Scan in November 2016 and February 2017 (Figure3A,B,4A,B). Meanwhile, blood tumor marker examinations were always within a normal range.

In August 2019, the 79-year-old man presented at his regular follow-up visit with headaches, diplopia, ptosis and right sided hearing loss which had been present for one week. There was no enlargement of the cervical lymph nodes during the physical examination. Serum NSE and CYFRA21-1 were elevated with values of 17.9 ng/ml [0-15] and 17.7 ng/ml [0-13] respectively. Brain MRI displayed multiple lacunar infarctions, along with a right bony mass in the skull base and clivus involving the parapharyngeal space, cephalic longus muscle, and cavernous sinus was observed, for which metastases or nasopharyngeal primary malignancies were considered as differentials (Figure5). Further examination by electronic nasopharyngoscopy revealed a large number of new cauliflower-like masses in the posterior roof wall of the nasopharynx and the right pharyngeal recess which appeared necrotic macroscopically (Figure 6). A biopsy was performed and the subsequent histopathological exam demonstrated adenocarcinoma with extensive necrosis which was consistent with intestinal adenocarcinoma metastasis to the nasopharynx based on immunohistochemical results and clinical history. Immunohistochemical analysis displayed CDX-2(+) CK20(-) NapsinA(-) TTF-1(focal+) Villin(+) (Figure7). Local nasopharyngeal radiotherapy was therefore commenced while the gene detection illustrated a CODE600 V600E mutation, *kras* [0-0], *nras* [0-0]. (DT:6000cGy/30f, 200cGy/f). Afterwards, the patient refused targeted therapy of cetuximab. Post radiotherapy, the patient developed side effects, including pharyngoxerosis, dysgeusia (bitter taste), odynophagia and so on. A chest CT scan exhibited small nodules suffused in both lungs, with the biggest nodule (size 2.5*1.8CM) located proximal to the the right oblique fissure (Figure8). Multiple enlarged lymph nodes were present in both pulmonary hila and the mediastinum, with local pleural thickening, all suggesting disease recurrences. Despite the fact that chemotherapy was the most appropriate treatment option for metastatic colon cancer, the patient did not consent, principally because of intolerance to previous chemotherapeutic regimens and the fear of severe allergic reactions. After radiotherapy, the patient's quality of life significantly declined mainly due to nasosinusitis, tympanitis, hearing loss, pharyngoxerosis, hypogeusia, and restriction of mouth opening. During follow-up post palliative radiotherapy, the patient deteriorated with progressive dyspnoea secondary to mass-effects of the metastatic deposit in the nasopharynx. He died 7 months post commencement of palliative therapy such as pain relief for end-stage disease in March 2020.

Discussion

Colorectal cancer, commonly known as bowel cancer, comprises cancerous growth in the colon, rectum, and depending on the definition criteria used can also include those found in the anus. Bowel cancer is classified according to the tumor node metastasis (TNM) staging system into stages I, II, III, and IV. Cancer that metastasizes to distal sites (stage-IV) is usually non-curable. The regional lymph nodes, liver, and lungs are the most common sites of metastasis associated with colorectal cancer. Infrequent sites of metastasis, including the spleen, thyroid gland, spermatic cord, and skeletal muscle have also been recorded[1-4]. Although metastases from colorectal cancer to these unusual sites might occur with the status of widespread disease, nasopharyngeal metastasis from colorectal cancer had never been reported prior to this article. To our knowledge, this is the first case of nasopharyngeal metastasis from primary colorectal cancer to be reported in the literature. At present, there are four mechanisms of metastasis linked with colorectal cancer which consist of lymphatic spread, direct extension, hematogenous spread, and planting spread. Theoretically, based on the analysis of the lymphatic reflux principle and anatomical location of the nasopharynx, we speculate that hematogenous spread of the primary colorectal cancer represents the most likely mechanism of metastasis in this case.

Nasopharyngeal carcinoma often develops at the roof of the nasopharyngeal posterior wall, generally with a cauliflower-like shape which is consistent with our report. Primary cancer is common in nasopharyngeal malignancies, thus making it difficult to determine whether this lesion is metastatic or not. In this regard, attention should be paid to the patient's other symptoms, signs, and medical history which are essential for diagnosis [4-5]. The main clinical manifestations of nasopharyngeal carcinoma include neck mass (40%), retracted blood snot (18.7%), and ear disorders (17.0%). The rate of lymph node metastasis was 82.3% at the time of diagnosis[6]. End-stage patients may present with ocular symptoms due to the nasopharyngeal carcinoma invading the external ocular muscles and/or oculomotor nerve[7]. Likewise, nasopharyngeal metastatic carcinoma can also cause the aforementioned symptoms. Luckily, histopathological and immunohistochemical examinations enable us to identify the tumor origin of the same tissue class. In our case, the caudate homologous transcription factor 2(CDX-2) testing positive indicates that the nasopharyngeal neoplasm is a metastasis of colorectal adenocarcinoma to the nasopharynx. CDX-2 is a tumor suppressor gene, and related studies have proven that it is closely related to the development and differentiation of normal intestinal epithelial cells into cancer[8-9].

As everyone knows, common treatment options for colorectal carcinoma consist of surgery, chemotherapy, radiotherapy and/or targeted therapy. Radical surgery is the main treatment, but postoperative recurrence and metastasis seriously affect the survival of patients. Adjuvant chemotherapy as the primary treatment after surgery can reduce the risk of recurrence and metastasis. However, it is still controversial about the patients of stage Ⅲ colorectal cancer whether to accept the auxiliary therapy when the primary tumors and metastases have received local R0 resections[10-11]. On the other side, according to the findings of the current study, chemotherapy did not produce a favorable effect on elderly patients with nasopharyngeal malignancies. On the other hand, radiotherapy is the most commonly used and effective treatment modality for head and neck tumors. Except for some severe acute reactions of

radiotherapy, both elderly and young patients are generally able to tolerate radiotherapy [12-14]. Over the past 40 years, with the advancement of technology and pharmacotherapy, more effective therapies for cancer have been developed under integrated clinical management. We conclude that these multiple relapses could be explained by a lack of adjuvant chemotherapy. Our case of atypical colorectal metastasis reminds us of the fact that the risk posed by hematogenous spread increases as the disease progresses.

In conclusion, nasopharyngeal metastases from colorectal cancer are extremely rare clinical entities. To the best of our knowledge, this is the first case reporting this occurrence, which not only enriches the database of this rare clinical entity but also reminds clinicians to be aware that metastatic carcinoma should be a top differential diagnosis when head and facial symptoms appear during the treatment of metastatic colorectal cancer or in patients with a past history of colorectal cancer. In these cases, nasopharyngoscopy and brain MRI should be performed promptly. These head and facial symptoms cannot be ignored, since they may indicate rapid disease progression which correlates with poor prognosis. Strict monitoring of patients with colorectal cancer after primary treatment would lead to the early diagnosis of such metastases and give patients more opportunities of treatment for a better prognosis.

Abbreviations

Computed tomography (CT)

Magnetic resonance imaging (MRI)

Hand-foot syndrome (HFS)

Tumor node metastasis (TNM)

Caudate homologous transcription factor 2 (CDX-2)

Declarations

Ethics approval and consent to participate

Written informed consent was obtained from the patient and legal guardian for publication of this research. The study was approved by the ethics committee of the first affiliated hospital of Wenzhou Medical University (381-06).

Consent for publication section

Written informed consent was obtained from the patient and legal guardian for publication of this research.

Availability of data and materials

Ture.

Competing interests

The authors declare that they have no competing interest.

Funding

None.

Acknowledgments

Not applicable.

Authors' contributions

LYH drafted and designed the manuscript, and also collected the patient's data and clinical records.

LBB assisted with the collection of the patient's data and clinical records. LSX helped to edit and revise the manuscript. All the authors have read and approved the final manuscript.

Authors' information (optional)

Ye-Huan Liu¹ Master Email:1259932365@qq.com

Bei-Bei Lin^{1*} Master Email: LBB0711@163.com

Shi-Xu Lv² Master Email:lvshixu@foxmail.com

¹Department of General medicine, The First Affiliated Hospital of Wenzhou Medical University, Wenzhou, Zhejiang, People's Republic of China

²Department of Thyroid & Breast Surgery, The First Affiliated Hospital of Wenzhou Medical University, Wenzhou, Zhejiang, People's Republic of China

References

1. Bigot P, Goodman C, Hamy A, et al. Isolated splenic metastasis from colorectal cancer: report of a case. J Gastrointest Surg 2008; 12: 981-982
2. Choi PW, Kim CN, Kim HS, et al. Skeletal Muscle Metastasis from Colorectal Cancer: Report of a Case. J Korean Soc Coloproctol 2008; 24: 492-496

3. Matsumoto G, Ise H, Inoue H, et al. Metastatic colon carcinoma found within an inguinal hernia sac: report of a case. *Surg Today* 2000; 30: 74-77
4. Phillips JS, Lishman S, Jani P. Colonic carcinoma metastasis to the thyroid: a case of skip metastasis. *J Laryngol Otol* 2005; 119: 834-836
5. Ariaschamorro B, Galeasanaya F, Salinassánchez J, et al. Maxillofacial metastasis of genitourinary origin. A report of 3 cases and review of the literature[J]. *Medicina Oral Patologia Oral Y Cirugia Bucal*, 2012, 17(2):e242-e245.
6. Yi junlin, gao li, huang xiaodong, et al. Clinical manifestations and prognosis of nasopharyngeal carcinoma [J].*Journal of Chinese academy of medical sciences*, 2006,28 (3) : 315.
7. Jiang bin, wang zhenchang, xianjunfang. MRI features of nasopharyngeal carcinoma first diagnosed with ophthalmic symptoms [J].*Chinese journal of radiation oncology*, 2010,44 (10) : 1045.
8. Mao Sheng-dong. The Expression of CDX2 and CK in Colorectal Cancer Tissue and its Clinical Significance [J]. *Journal of Pharmaceutical Biotechnology*, 2019, 026(001):37-40.
9. Zhang J, Feng X Y, Lin Y. Analysis of risk factors for invasion and metastasis of gastric cancer [J]. *Journal of Shanxi Medical University*, 2017, 048(008):836-839.
10. GAO P, HUANG XZ, SONG YX, et al. The Impact of timing of adjuvant chemotherapy on survival in stage Ⅲ colon cancer: a population based study [J]. *BMC Cancer*, 2018,18 (1) : 234.
11. LIU Q, LUO D, AN H, et al. Survival benefit of adjuvant chemotherapy for patients with poorly differentiated stage Ⅲ A colon cancer [J]. *J Cancer*, 2019,10 (5) : 1209-1215.
12. Mitsuhashi N, Hayakawa K, Yamakawa M, et al. Cancer in patients aged over 90 years or older: radiation therapy [J] . *Radiology*, 1999, 211(3) : 829- 833.
13. Pignon T, Horiot J C, Van den Bogaert W, et al. No age limit for radical radiotherapy in head and neck tumours [J] . *Eur J Cancer*, 1996, 32A(12) : 2075- 2081.
14. Zeng Qi, Guo Xiang ning-wei li, et al. Clinical Characteristics and Prognosis of Aged Nasopharyngeal Carcinoma Patients: A Report of 313 Cases [J]. *Cancer (English edition)*, 2008, 027 (003) : 289-294.

Figures

Figure 1

Tall malignant columnar cells lining large irregular glands (hematoxylin-eosin staining, magnification×100). Histopathological exam revealed an ulcerative type of moderately - poorly differentiated adenocarcinoma. Immunohistochemical analysis displayed BcL-2(-) P53(+) k-ras(-).

Figure 2

Abdominal CT scan demonstrating, in the horizontal view, a single low density lesion (arrow) in the right hepatic lobe (size 3.8*2.8cm).

A

B

Figure 3

A. Chest CT scan demonstrating, in the horizontal view, a isolated nodule of 9mm in diameter (arrow) in the lobe of the left lung. B. Adenocarcinoma spread to the left lung parenchyma (hematoxylin-eosin staining, magnification $\times 100$). Histopathological exam revealed tubular adenocarcinoma with a small amount of necrosis. Immunohistochemical analysis displayed CDX-2(+++) CK20(-) CK7(-) TTF-1(-) Villin(++).

A

B

Figure 4

A.Chest CT scan demonstrating, in horizontal view, a isolated 11mm in diameter nodule (arrow) in the upper lobe of the right lung. B.Adenocarcinoma spread to the right lung parenchyma (hematoxylin-eosin staining, magnification×100). Histopathological exam revealed of moderately differentiated tubular adenocarcinoma. Immunohistochemical analysis displayed CDX-2(+) CEA(+) CK20(+) CK7(-) TTF-1(-) Villin(+).

Figure 5

Brain magnetic resonance imaging demonstrating, in the horizontal view, a right bone mass (arrow) in the skull base and clivus invading the parapharyngeal space, cephalic longus muscle, and cavernous sinus (T1W, FLAIR).

Figure 6

Electronic nasopharyngoscopy illustrating a large number of new cauliflower-like masses in the roof of the posterior wall of the nasopharynx and right pharyngeal recess, with a brittle texture and easy to bleed during biopsy.

Figure 7

Adenocarcinoma spread to the nasopharyngeal parenchyma (hematoxylin-eosin staining, magnification×100).The histopathological exam revealed adenocarcinoma with extensive necrosis. Immunohistochemical analysis displayed CDX-2(+) CK20(-) NapsinA(-) TTF-1(focal +) Villin(+).

Figure 8

Chest CT scan demonstrating, in the horizontal view, small nodules suffused in both lungs, with the largest one (arrow) proximal to the right oblique fissure (size 2.5*1.8CM).