Deep spatial profiling of COVID19 brains reveals neuroinflammation by compartmentalized local immune cell interactions and targets for intervention

Bertram Bengsch (✉ bertram.bensch@uniklinik-freiburg.de)

University Medical Center Freiburg <https://orcid.org/0000-0003-2552-740X>

Marius Schwabenland

Institute of Neuropathology, Faculty of Medicine, University of Freiburg <https://orcid.org/0000-0003-2205-5427>

Henrike Salié

University Medical Center Freiburg

Jovan Tanevski

Heidelberg University Hospital <https://orcid.org/0000-0001-7177-1003>

Saskia Killmer

Department of Medicine II, Freiburg Freiburg University Medical Center, Faculty of Medicine, University of Freiburg, Freiburg, Germany

Jakob Matschke

University Medical Center Hamburg-Eppendorf

Klaus Püschel

University Medical Center Hamburg-Eppendorf

Henrik Mei

German Rheumatism Research Center Berlin (DRFZ), a Leibniz Institute, Berlin, Germany

Tobias Boettler

Department of Medicine II (Gastroenterology, Hepatology, Endocrinology and Infectious Diseases), Freiburg University Medical Center, Faculty of Medicine, University of Freiburg, Freiburg, Germany
<https://orcid.org/0000-0002-1195-055X>

Christoph Neumann-Haefelin

Department of Medicine II (Gastroenterology, Hepatology, Endocrinology and Infectious Diseases), Freiburg University Medical Center, Faculty of Medicine, University of Freiburg, Freiburg, Germany

Maike Hofmann

University Hospital Freiburg <https://orcid.org/0000-0001-8410-8833>

Julio Saez-Rodriguez

Heidelberg University Hospital

Klaus-Peter Knobeloch

University of Freiburg <https://orcid.org/0000-0003-1587-1944>

Thomas Blank

University of Freiburg

Robert Thimme

Department of Medicine II, Freiburg Freiburg University Medical Center, Faculty of Medicine, University of Freiburg, Freiburg, Germany <https://orcid.org/0000-0003-1417-4135>

Markus Glatzel

University Medical Center Hamburg-Eppendorf <https://orcid.org/0000-0002-7720-8817>

Marco Prinz (✉ marco.prinz@uniklinik-freiburg.de)

University of Freiburg <https://orcid.org/0000-0002-0349-1955>

Research Article

Keywords: SARS-CoV-2, COVID-19, encephalopathy, brain autopsy, microglia, T cells, high-dimensional imaging, neuroinflammation, neurological deficits, machine learning, single-cell analysis, mass cytometry

Posted Date: August 28th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-63687/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Deep spatial profiling of COVID19 brains reveals neuroinflammation by compartmentalized local immune cell interactions and targets for intervention

Marius Schwabenland^{1*}, Henrike Salié^{2*}, Jovan Tanevski³, Saskia Killmer², Jakob Matschke⁴, Klaus Püschel⁵, Henrik E. Mei⁶, Tobias Boettler², Christoph Neumann-Haefelin², Maike Hofmann², Julio Saez-Rodriguez³, Klaus-Peter Knobeloch¹, Thomas Blank¹, Robert Thimme², Markus Glatzel⁴, Marco Prinz^{1,7,8#} & Bertram Bengsch^{2,8#}

* equal contribution first authors # equal contribution last authors

¹Institute of Neuropathology, Faculty of Medicine, University of Freiburg, Freiburg, Germany

²Faculty of Medicine, Department of Medicine II, Gastroenterology, Hepatology, Endocrinology, and Infectious Disease, University Medical Center Freiburg, Freiburg, Germany

³ Institute for Computational Biomedicine, Faculty of Medicine, Heidelberg University and Heidelberg University Hospital, Heidelberg, Germany

⁴Institute of Neuropathology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany

⁵Institute of Legal Medicine, University Medical Center Hamburg-Eppendorf, Hamburg, Germany

⁶German Rheumatism Research Center Berlin (DRFZ), a Leibniz Institute, Berlin, Germany

⁷Center for Basics in NeuroModulation (NeuroModulBasics), Faculty of Medicine, University of Freiburg, Freiburg, Germany

⁸Signaling Research Centers BLOSS and CIBSS, University of Freiburg, Freiburg, Germany

Correspondence to:

Marco Prinz

Institute of Neuropathology

University of Freiburg

Breisacher Str. 64

D-79106 Freiburg, Germany

Phone: +49-761-270-51050

E-mail: marco.prinz@uniklinik-freiburg.de

or Bertram Bengsch

University Medical Center Freiburg

Department of Medicine II

Hugstetter Str. 55

D-79106 Freiburg, Germany

Phone: +49-761-270-32870

E-mail: Bertram.Bensch@uniklinik-freiburg.de

ABSTRACT

COVID-19 causes neurological symptoms that can be potentially life-threatening in up to 67 % of the patients. The underlying pathophysiological mechanisms of COVID-19 associated encephalopathy, the involved immune cells, their spatial distribution and their cellular interactions during disease remain largely unclear. In this study, we performed a 38-biomarker imaging mass cytometry analysis of the brain stem from 25 patients and additional controls to understand the local immune response during SARS-CoV-2 infection at a spatially resolved, high-dimensional single-cell level. Importantly, utilizing an unbiased image segmentation and cell classification pipeline, we observed a significant immune activation in the central nervous system (CNS) and identified novel context-specific CD8 T cell and microglial clusters. Spatially resolved single-cell analysis identified distinct phenotypes of T cells and microglial clusters, their presence in specific anatomical regions and their cellular interactions. Our analysis further highlights microglial nodules and perivascular immune cell clusters as key sites of the local immune response. It also demonstrates that disease-associated neuroinflammation is associated with severe axonal damage as a structural basis for the neurologic deficits. Finally, we identified compartment- and cluster-specific immune checkpoints that can be used for future therapeutic interventions.

KEYWORDS

SARS-CoV-2, COVID-19, encephalopathy, brain autopsy, microglia, T cells, high-dimensional imaging, neuroinflammation, neurological deficits, machine learning, single-cell analysis, mass cytometry

INTRODUCTION

COVID-19 due to the SARS-CoV-2 pandemic is emerging as a multifaceted disease with multi-organ complications. One major complication are neurological symptoms that can occur during acute infection in up to 67 % of the patients but also persist or even emerge after viral elimination and are summarized as COVID-19 encephalopathy¹⁻³. Typically, COVID-19 encephalopathy manifests with anosmia but symptoms range from gait disorders, breathing difficulties to coma suggesting involvement of several brain regions such as the olfactory bulb but also infratentorial regions like the medulla⁴. Thromboembolic events including stroke were reported to be a major complication of COVID-19, but are unlikely to explain COVID-19 neurological symptoms in the majority of cases. Of note, SARS-CoV-2 has been detected in brain samples⁵ and immune activation in COVID-19 brains is debated based on differing observations from a limited number of small case series⁶⁻¹⁰, raising the possibility that immune-mediated mechanisms may drive COVID-19 encephalopathy. However, comprehensive data with detailed dissection of immune cell populations on a large collection of brains from COVID-19 patients is currently not yet available. Previous studies on peripheral immune functions in COVID-19 patients have uncovered patterns of immune activation and deviation in the peripheral blood and identified distinct immunotypes with differences in CD4 and CD8 activation, effector memory CD8 T cell differentiation and differential induction of B cell responses that are linked to different disease severity^{11,12}. In particular, an immunotype with strong CD4 T cell activation, and expansion of plasmablasts was linked to highest NIH COVID severity scales^{11,12}. However, it is currently unclear whether the immune responses assessed in the peripheral blood reflect the immune responses in the tissue and which immune populations are involved in mediating organ pathology. Thus, in order to understand the cellular, immunological and anatomical basis of COVID-19 encephalopathy, we performed a highly multiplexed spatial analysis of the adaptive and innate immune system in brain sections from 25 COVID-19 patients that succumbed to the disease. We

utilized imaging mass cytometry (IMC) that allowed the simultaneous detection of several key immune populations, including novel disease-linked clusters of CD4 and CD8 T cells, NK cells, B cells, and numerous myeloid and brain-specific innate immune populations, their cellular interactions and importantly, their compartmentalization in distinct anatomical regions of the brain stem. In addition, we performed a thorough molecular profiling of immune checkpoint receptor expression together with immune activation and functional analyses. Spatial mapping of the disease-associated altered brain immune response using imaging mass cytometry revealed significant infiltration of distinct immune cell clusters with a prominent role for an activated CD8 T cell cluster with concomitant expansion of microgliosis-related cell clusters in distinct anatomic niches in the majority of patients. Our results indicate that during COVID-19 encephalopathy, the adaptive as well as the CNS endogenous innate immune system are concomitantly activated, and highlight disease-specific immune cell clusters that physically interact in distinct CNS compartments. Compartment-enriched immune cell clusters revealed distinct immune marker profiles that could be used for therapeutic interventions. Furthermore, the presence of immune cell clusters correlates with the extent of axonal degeneration as a morphological correlate of neuronal damage. Our data provide a foundation for novel therapeutic avenues aimed at reducing neuroinflammation in COVID-19 patients.

RESULTS

Highly multiplexed imaging of the brain microarchitecture by imaging mass cytometry indicates strong immune activation in COVID-19 patients

To understand the immune response in COVID-19-associated neuropathology, we performed highly multiplexed high-dimensional imaging using imaging mass cytometry (IMC) together with immunohistochemistry to identify the spatial brain immune profile in brain stem sections from patients that succumbed to COVID-19 and compared it to control patients that died of non-infectious and non-neurological reasons (**Figure 1A**). Classical immunohistochemistry for the cytotoxic T-cell marker CD8 and the myeloid cell marker Iba1 revealed significant CD8 T cell infiltration and an elevated number of Iba1 parenchymal cells with typical microglial morphology (such as round to spindle-shaped somata and a distinct arborization pattern) indicating a robust microgliosis (**Figure 1B**). Microgliosis occurred either in a diffuse manner or in clusters that are referred to as microglia nodules. The observed CD8 immunoreactivity in 17 out of 25 patients and pronounced microglia activation in 20 out of 25 patients suggested that classical neuroinflammation¹³ is a histopathological hallmark of COVID-19-related neuropathology.

However, since classical immunohistochemistry allows analysis of only a few markers, limiting insight into the fine composition of cellular phenotypes and immune cell subsets, we designed and validated a 38-marker IMC approach to comprehensively profile key brain immune and parenchymal cell populations. Tissue sections were immunolabeled with metal isotope coupled antibodies. Spatial expression data was obtained after spot-by-spot laser ablation of the tissue at 1 μ m resolution and the subsequent mass spectrometry analysis of metal labels. Resolution of IMC data was comparable to conventional immunohistochemistry (**Figure 1B**) but the high-parametric information provided by this technique allowed deeper insights into the cellular composition in specific anatomical compartments, such as areas with dense microglial accumulation known as microglial nodules, that are absent in healthy brain tissue^{13,14}. In a next

step, expression of markers for canonical lymphoid and myeloid immune populations and parenchymal brain cells was assessed together with markers of immune activation, function and regulation (**Figure 1C**). Parallel manual count analysis indicated a significant immune infiltration of the brain stem from COVID-19 patients by CD8 T cells and (albeit to a lower degree) CD4 T cells and some B cells (**Figure 1D**). Importantly, this adaptive immune infiltration by immune cells that are largely absent in control brains occurred in the presence of significantly increased numbers of Iba1⁺ microglial cells as well as CD163⁺ perivascular macrophages (**Figure 1D**). Interestingly, we also observed a reduction in CD3⁺GzmB⁺ NK cells in diseased patients, indicative of a ~50% depletion of brain NK cells that has been described in the periphery of COVID-19 patients and is implicated in a more severe clinical course of COVID-19^{15,16}. Given the profound immune activation in COVID-19 brains, we wondered whether this robust immune activation pattern in the brain stem was linked to tissue destruction and neuronal damage. We thus quantified amyloid precursor protein (APP) deposits in the brains of COVID-19 patients and controls, respectively. Notably, significantly higher numbers of APP-positive deposits were observed in COVID-19 patients (**Figure 1E**). This damage occurred in the absence of an apparent tissue necrosis, suggesting a specific and likely directed immune-mediated pathology.

Together, these analyses reveal strong signs of neuroinflammation with significant immune infiltration, activation of the central nervous system (CNS) endogenous innate immune system and accompanying increased neuronal damage in the brains of patients with COVID-19.

Identification of distinct brain immune cell clusters during COVID-19-associated neuropathology

To understand the cellular phenotypes in the CNS during SARS-CoV-2 infection in greater detail, we first utilized supervised machine learning to generate cell masks that allowed image segmentation using a reference pipeline adapted for better resolution of microglial cells¹⁷ (**Figure**

2A). We then extracted high-dimensional single-cell level data and clustered using Phenograph algorithm to identify the cellular composition in different patients (**Figure 2A**). Phenograph identifies cellular clusters on a graph constructed based on the high-dimensional phenotypic similarities of cells in the dataset before detecting discrete communities on this graph. It was chosen based on its robustness to identify even rare but phenotypically distinct populations¹⁸. This clustering strategy identified 31 distinct cellular clusters in the IMC analysis based on the high-dimensional segmented single cell data (**Figure 2A**). Most abundant clusters displayed molecular expression signatures of stromal (clusters c1, c4, c6, c8, c14, c21, c28), neuronal (c2, c20) and astrocytic (c3, c5, c18) cells, in agreement with the dominance of these cell types in the CNS (**Figure 2B, Supplementary Figure 1A**). Other identified clusters included cells with typical expression of endothelial cell markers (e.g. CD34, c9), or pericytes (c19). Importantly, this analysis also clearly identified multiple distinct immune clusters, such as several microglial clusters with varying expression of Iba1 and markers of immune activation and lysosomal function such as HLA-DR and CD68 in clusters c7, c11, c12, c13, c15, c16, as well as different subsets of myeloid immune cells i.e. CD163⁺CD204⁺ perivascular macrophages (c27) and a related microglial cluster (c11). Lymphoid clusters detected included B cells, CD8 and CD4 T and NK cell subsets (clusters c23, c22, c29, c26, respectively) (**Figure 2B**). To understand changes in the immune landscape in COVID-19-linked neuroinflammation, we next visualized the immune cell clusters on a brain immune map generated via *t*-SNE and analyzed their abundance in patients and controls. This approach revealed major changes in the brain immune cell composition in the brains of COVID-19-affected patients, that were qualitatively and quantitatively dominated by activated microglial clusters (e.g., c7, c11, c13, c15, c16) and adaptive immune cell infiltration (c22, c23, c29) (**Figures 2C-E**). In contrast, only few immune clusters were present in control brains (e.g. c12, c26) and reduced in COVID-19 patients, suggesting a role in immune homeostasis for these microglial and NK cell populations. Taken together, our analyses identified

distinct adaptive and innate immune cell clusters in the brain during SARS-CoV-2 infection that were not present in control individuals.

The cellular interactome points to immune cell-crosstalk at the perivascular compartment during COVID-19 neuroinflammation

We next wondered which cellular clusters were differentially enriched or depleted in the CNS of COVID-19 patients across the cohort. As expected based on the analysis above, comparison of high-dimensional clusters indicated a significant enrichment of CD8 T cell cluster c22, the CD4 T cluster c29, and perivascular macrophage cluster c27 across COVID-19 patients (**Figure 3A**). Interestingly, these data also revealed that COVID-19-associated neuroinflammation is linked to significant expansion of an astrocytic (c3) and pericyte cluster (c19) (**Figure 3A**). Moreover, SARS-CoV-2 receptor ACE2 expression was found on some brain vessels, suggesting the possibility of viral antigens presented in the brain vasculature (**Supplementary Figure 1B**). Therefore, it is tempting to speculate that COVID-19 immune pathology might be linked to uniform vascular activation.

To understand co-regulated cell dynamics during COVID-19 encephalopathy in an unbiased manner, we next performed a linear regression analysis between all clusters to determine highly correlated immune infiltration patterns. This analysis identified modules of highly inter-correlated immune cell clusters, including one that was composed of the clusters c3, c7, c22, c26, c27 and c29, thus involving the major COVID-19-associated cellular phenotypes identified above (**Figure 3B**). As such, these patterns pointed to a close link between the vasculature-linked c27 and T cell clusters c22 and c29, indicating T cell infiltration. We therefore asked if CD8 T cells were also spatially linked to the perivascular compartment. Indeed, more than half of CD8 T cells in cluster c22 were found in significantly closer proximity to vascular collagen-I in COVID-19 samples

compared to a random distribution (**Figure 3C**). Neighborhood analysis further confirmed a close spatial interaction between c22 and vascular cell clusters, including endothelial cell cluster c9, pericyte cluster c19 and perivascular macrophage cluster c27 (**Figure 3D**). Importantly, the inflammatory spatial interaction pattern was pronounced in COVID-19 patients but not in controls, in which cluster c22 only displayed significant interactions with cells from the same cluster or homeostatic NK cell cluster c26 (**Figure 3E, Supplementary Figure 1C**). In sum, these data demonstrate a clear link between CD8 T cell infiltration and vascular activation during COVID-19-related encephalitis.

Activation of CD8 T cell cluster c22 in COVID-19 brains

The strong enrichment of c22 representing CD8 T cells observed in COVID-19 brains pointed to a central role of this immune cell population. To understand their activation and function, we tested this cluster for the expression of markers informative about T cell differentiation, transcriptional programming, activation/exhaustion and function. We observed a strong enrichment of cluster c22 cells expressing CD45RO indicative of non-naïve T cells (**Figure 3F**). Cells in c22 also expressed immune checkpoints PD-1 and Tim-3 as well as transcription factors Tox and Eomes, a phenotype compatible with highly activated effector cells with a possible induction of features of T cell exhaustion¹⁹. Of note, these cells also expressed granzyme B, indicating cytotoxic effector T cell function (**Figure 3F**). In sum, the strong enrichment of CD8 T cells with significant activation involving several immune checkpoints in this immune cluster implicates that this immune cell population is a central mediator of neuroinflammation during SARS-CoV-2 infection.

Immune activation in brains of COVID-19 patients occurs in distinct anatomic niches

Our previous analysis indicated a strong perivascular immune infiltration linked to CD8 T cluster c22. However, importantly, we also noted that not all cells in this cluster were associated with the perivascular space (**Figure 3C**). Indeed, we also observed additional COVID-19-specific neighborhood relationships of CD8 T cluster c22 with microglial clusters c7, c11 and c15 outside the perivascular region, in particular with microglial cluster c7 (**Figure 3E**), suggesting additional anatomic niches linked to specific microglial immune clusters in the brain parenchyma.

To systematically dissect these anatomic regions, we generated automated maps based on collagen-I and Iba1 expression that informed about the cellular distance to the vasculature and identified microglial hotspots (**Figures 4A,B**). Importantly, these analyses highlighted microglial nodules with an accumulation of densely packed microglial cells as an additional major site of infiltration by CD8 T cell cluster c22 (**Figure 4C**). Presence of microglial nodules was observed in the medulla of 28 % of COVID-19 patients (7 out of 25) and in none of the control cases. Microglia nodules differed from more scattered parenchymal microgliosis observed in the majority of COVID-19 patients (76%, 19 out of 25) and in 20 % in the control cases (**Figure 4C**). These data highlighted the perivascular and microglial nodule compartment as key sites of COVID-19 immune responses. As shown exemplarily in **Figure 4D**, however, some additional CD8 T cell cluster c22 cells could also be identified scattered in the parenchyma. Thus, to dissect the role of distinct anatomic compartments as places of local immune responses, we performed a detailed cluster composition analysis in the perivascular, juxtavascular, parenchymal and microglial nodule compartments. Clearly, we identified major differences in the cluster composition of these compartments (**Figure 4D**). Most changes in cluster composition were found in microglial nodules that were enriched in microglial clusters c7, c15 and CD8 T cell cluster c22 (**Figure 4D,E**). Interestingly, in microglial nodules, only microglial clusters c7 and c15, but not microglial clusters c11, c13, c16 were enriched, indicating context-specific roles for these microglial populations (**Figures 4D,E**). In contrast, the perivascular compartment displayed enrichment of endothelial

cell cluster c9, pericyte cluster c19 and perivascular macrophage cluster c27, with some enrichment of CD8 T cell cluster c22, NK cell cluster c26 and stromal cluster c28 (**Figures 4D,E**). Only subtle differences in cluster composition were found between the juxtavascular and perivascular compartment, but their composition clearly differed from the nodule and perivascular compartment (**Figure 4D**). These data illustrate differential immune activation consisting of defined cell clusters at distinct anatomical regions in COVID-19-associated brains and identified the key adaptive and innate immune subsets in each compartment.

Differential immune activation at distinct anatomical sites

As shown above, activated CD8 T cells in cluster c22 were significantly enriched in the CNS of COVID-19 patients. However, the presence of heterogeneous immune cell subsets and their different activation states at distinct anatomical niches suggested varying immunological roles for these compartments. We next tested whether these anatomic compartments had an impact on CD8 T cell activation and differentiation represented by changes in cluster c22. Interestingly, the different anatomical regions were indeed linked to differences in T cell activation observed in c22. The diffuse parenchymal compartment displayed the lowest levels of PD-1, CD39, Tim-3 and Eomes in c22, while high expression of PD-1, CD39 and particularly Tim-3 was observed in cluster c22 cells in the perivascular compartment (**Figure 5A**). These expression patterns differed from c22 cells in microglial nodules, that displayed high co-expression of PD-1 and CD39, but not Tim-3, however, the c22 cells in microglial nodules also featured higher induction of Eomes and HLA-DR, indicating a more profound T-cell activation and a possible bias towards exhaustion. Of note, while quantitative differences between the juxtavascular compartment and diffuse parenchyma had not been observed based on cluster composition analysis (**Figure 4D**), qualitative differences were apparent for CD8 T cell cluster c22, with elevated PD-1 expression in the juxtavascular space (**Figure 5A**), suggesting a closer qualitative relationship of the juxtavascular c22 cluster

cells to their perivascular counterparts. Together, these results indicate higher activation of CD8 T cells in c22 associated with the vasculature and microglial nodules but also differences in the expression of specific immune regulatory molecules between these anatomic sites.

Pervasive inflammatory effect of microglial nodules

Differential T cell activation patterns can be influenced by antigens or the inflammatory microenvironment. Since immune infiltration was found throughout our cohort of COVID-19 patients (microgliosis: 80 %), but microglial nodules were observed in only 28 % of COVID-19 cases, we wondered if the presence of microglial nodules might orchestrate more severe neuroinflammation. This analysis was influenced by our observation of a higher antigen-presentation capacity of CD8 T cell cluster c22 expressing HLA-DR, but also higher HLA-DR expression by CD4 T cell cluster c29 and microglial cluster c7 in the nodules, indicating strong inflammation (**Figure 5B**). We thus analyzed the immune activation patterns of key immune cells in the different compartments depending on presence or absence of microglial nodules (**Figure 5C, Supplementary Figure 1E**). Of note, analysis of CD8 T cells cluster c22 revealed higher PD-1 expression in the parenchyma and higher PD-1 and CD39 co-expression in the juxtavascular and parenchymal compartment in patients with microglial nodules. Together, these data clearly illustrate a pervasive pro-inflammatory effect linked to the presence of microglial nodules across all brain compartments. We next wondered if a similar immune compartmentalization also affected other immune clusters. Indeed, as shown in **Figures 5B,D** and **Supplementary Figure 1F** we observed a similar induction of HLA-DR on CD4 T cell cluster c29 or microglial cluster c7 and c15 in microglial nodules. Similar to the CD8 T cell c22, these changes were associated with a differential up-regulation of additional co-regulatory immune checkpoints, such as PD-L1, CD39 or CD33 (**Figure 5D, Supplementary Figure 1F**). Notably, co-expression of immune checkpoints PD-1 and PD-L1 on adjacent immune cells suggests active modulation of brain immunity through

263 the PD-1:PD-L1 pathway (**Supplementary Figure 1G**). Fitting to these observations, patients
264 with microglial nodules tended to have higher numbers of APP deposits (**Supplementary Figure**
265 **1H**), suggesting a link between immune activation and axonal damage. These results
266 demonstrate significant differences in the anatomic immune regulation in COVID-19 affected
267 brains and point towards a specific role for microglial nodules and CD8 T cell-microglia crosstalk
268 in orchestrating the brain immune responses. Expression of multiple immune checkpoints by
269 these immune populations indicates potential for therapeutic manipulation.

DISCUSSION

Here, by performing highly multiplexed spatial analysis of the cellular composition and immunological phenotype of CNS tissue from patients that died with COVID-19, we identified a profound immune activation dominated by specific CD8 T cell clusters affecting (1) the vasculature and (2) significant CD8 T cell - microglial crosstalk in the parenchyma that frequently manifests in the formation of microglial nodules. Remarkably, the degree of neuroinflammation was variable but major changes in microglial immune activation were observed in about 80 % of patients. Importantly, this immune infiltration was linked to axonal damage, highlighting significant severe neuroinflammation in COVID-19 patients that concomitantly affects neuronal populations in the brain stem region, which might explain the high prevalence of transitory and long-lasting neurological comorbidities in SARS-CoV-2 infection. These results underline the need for therapeutic interventions aimed at maintaining neuronal homeostasis in SARS-CoV-2 infection.

Key findings of our study are the broad immune activation and anatomic compartmentalization of the altered brain immune responses in COVID-19 patients. Briefly, perivascular immune activation and infiltration by distinct CD8 and CD4 T cell clusters and expansion of disease-associated clusters of perivascular macrophages, pericytes and microglial cells was a dominant feature found in the majority (80%) of patients. Importantly, our analysis also revealed compartmentalized changes in brain immunity since the vascular immune response pattern differed quantitatively and qualitatively from the parenchymal and microglial nodule immune response. Indeed, presence of a moderate or severe degree of microgliosis and microglial nodules were observed in 48% and 28% of patients, and were linked to CD8 T cell infiltration and high activation. Of note, these pathologic features of microgliosis and microglial nodule are linked to increasing and pervasive immune activation. In particular, cellular analysis in microglial nodules revealed high HLA-DR expression and significant T cell-microglial crosstalk that occurred together with significantly

higher immune activation in the rest of the tissue in these patients, including the perivascular compartment. This broad immune activation was linked to significant axonal damage observed in the patient cohort with trends to more axonal damage in patients with microglial nodules, which provides a mechanistic basis for COVID-19 encephalopathy.

The origin and onset of perivascular inflammation, microglial activation and expansion and their relationship to each other are currently unclear. One possible explanation for the perivascular immune activation is the expression of the SARS-CoV-2 receptor ACE2 found in endothelial cells. SARS-CoV-2 virus has been detected in COVID-19 brains, but anatomic fine-mapping of infection sites is lacking⁵. ACE2-mediated virus uptake might lead to local immune activation by adaptive immune responses reacting to viral antigen expressed in the vascular compartment. Alternatively, dysregulated systemic cytokine responses reported in many COVID-19 patients might contribute to the perivascular phenotype²⁰. A possible sequence of events compatible with our data could be that perivascular inflammation due to antigen recognition breaches the blood-brain barrier and permits adaptive immune cell infiltration into the brain parenchyma. However, another potential mechanism could be linked to the uptake of SARS-CoV-2 via olfactory nerves described in mice²¹ and fitting to the anosmia reported in the majority of patients. Uptake via the olfactory nerves could lead to viral antigen presented in the brain parenchyma in the absence of a compromised blood-brain barrier. This mechanism might cause primary activation of brain resident immunity, including microglial clusters, initially independent of adaptive immunity. While temporal or mechanistic relationships could not be established in the cross-sectional post-mortem cohort analyzed here, our work highlights detailed anatomic immune profiling as an important disease feature required to understand COVID-19 encephalopathy.

In our study, we further tested for the expression of multiple targetable immune checkpoints, against which therapeutic agents have been tested in clinical trials. These targets include PD-1, PD-L1, CD38, CD39, Tim-3 and TIGIT. The expression of multiple immune checkpoints, including

320 PD-1, on CD8 and CD4 T cells suggests that these cells can be modulated through these
321 pathways. In particular, the PD-1/PD-L1 pathway is of interest, as high expression of PD-1 on
322 CD8 T cell cluster cells and PD-L1 on closely interacting microglial clusters was observed.
323 However, it remains unclear if boosting adaptive immunity by checkpoint blockade antibodies may
324 benefit disease outcomes, as even higher T cell activation would be predicted. Indeed, the up-
325 regulation of these immune checkpoints could potentially explain the limited necrosis observed in
326 the tissues analyzed, despite the presence of CD8 effector T cells armed with lytic granula. It is
327 therefore possible that brief interruption of inhibitory signaling might boost the T cell immune
328 response to eliminate viral antigen in the brain. Nevertheless, such a scenario would require the
329 boosted immune response to eliminate viral antigen quickly before causing significant and
330 widespread immune pathology. There is precedent that such an approach can be effective, as
331 illustrated by the successes in treating JC-virus-associated progressive multifocal
332 leukoencephalopathy with PD-1 blockade approaches²². Additional investigations of the role of
333 SARS-CoV-2 neurotropism and immunoregulation will be required to better delineate the
334 relationship between virus, antigen, neuroinflammation and immunomodulation.

335 Microglial cells fulfill many functions in brain immunity and homeostasis, and adapt quickly to
336 different challenges and disease settings. Recently, significant microglial heterogeneity during
337 development and has been highlighted by single cell expression techniques based on RNA
338 expression profiling in murine models ²³. Here, in order to understand the broad and
339 heterogeneous nature of the human brain immune responses during SARS-CoV-2 infection, we
340 developed an integrated IMC method that allowed the quantification of 38 immune markers and
341 included key microglial markers together with markers against major innate and adaptive immune
342 cell populations to determine the spatial brain immune response with unprecedented detail.
343 Cellular segmentation revealed significant heterogeneity in brain-specific immune populations in
344 controls and COVID-19 patients, which identified several subsets of microglial cell populations

and homeostatic innate immunity clusters. Importantly the heterogeneity and composition of the 7 newly identified microglial clusters in our study was significantly changed in COVID-19 brains towards more activated microglial clusters. In particular, two microglial clusters (clusters c7 and c15) emerged that were significantly linked to the formation of microglial nodules, i.e. densely packed areas with high microglial content that can be approximated using a novel microglial nodule index. These microglial clusters displayed high immune activation and expression of microglial functional markers. Importantly, in microglial nodules, they interacted closely with CD8 and CD4 T cells. This interaction seemed to have a pervasive effect on the global brain immune response as significantly higher immune activation was observed in distant anatomic compartments in patients with microglial nodules and suggests that the microglial nodules serve as orchestrating immune hubs not unlike tertiary lymphoid structures in other organs. Of note, the broad immune infiltration in COVID-19 brains was linked to an expense of a likely homeostatic microglial cluster with high expression of CD68, which appeared to interact with a GzmB⁺ CD3⁻ NK-cell-like immune population. While homeostatic microglial-NK cell relationships have been suggested by some studies^{24,25}, evidence for a role of such spatially interacting clusters in human disease was lacking. Moreover, it remains to be clarified if the reduction of microglial homeostatic clusters and emergence of activated inflammatory microglial clusters can be explained by a potential activation and expansion of these microglial clusters. Nevertheless, while the immune profiling in this study focused on the pro-inflammatory changes associated with immune infiltration in COVID-19 brains, the observed dysbalancing of homeostatic brain immunity during COVID-19 suggests that strategies fostering homeostatic immune cluster interactions could also represent interesting therapeutic approaches to ameliorate immune activation in the brain.

The broad neuroinflammation observed by deep spatial immune analysis in this study points to a profound CNS manifestation of COVID-19. In our large prospectively collected post-mortem cohort, 80 % of patients displayed significant microglial alterations, 68% of patients had notable

370 parenchymal CD8 infiltration and 28 % of patients had the most severe phenotype with profound
371 microglial nodule formation. Moreover, axonal degeneration was typically found in the medulla of
372 COVID-19 patients, but this neuropathology occurred in the absence of broad necrosis. The
373 immunological and clinically relevant alterations observed in our work fit to the high rate of
374 neurologic symptoms during COVID-19⁴. However, these results were unexpected in light of
375 earlier assessments that mainly reported hypoxic changes and thromboembolic events and did
376 not report significant encephalitis⁹. Of note, only 16 % of patients in our cohort had
377 thromboembolic events that also were localized to distant brain regions, suggesting that the
378 profound neuroinflammation observed in the brain stem and the thromboembolic events are
379 unrelated. The high prevalence of neuroinflammation during COVID-19 observed in our work
380 highlights the need for better strategies targeting or preventing this inflammatory encephalopathy.

381 Our study clearly underlines the relevance of high-parametric deeply resolved spatial analysis of
382 brain immunity that is able to comprehensively dissect brain immune populations, such as distinct
383 microglial clusters and multiple brain compartments in order to understand the immunological and
384 pathophysiological changes during COVID-19. The identification of disease-linked CD8 T cell-
385 microglial cross-talk that occurred in specific anatomic compartments and between defined
386 cellular clusters that involve multiple immune checkpoints highlights possible cellular and
387 molecular targets for therapeutic intervention.

ACKNOWLEDGMENTS

This project was supported by grants from the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) (SFB 992, SFB1160, SFB/TRR167, SFB/TRR179, German Excellency strategy CIBSS - EXC-2189– Project ID390939984) and special research funds from the Ministry for Science, Research and Art of Baden-Wuerttemberg dedicated to “COVID-19 research” and “Neuroinflammation”. B.B. was further supported by DFG grant BE-5496/5-1 and M.P. was further supported by the Sobek Foundation, the Ernst-Jung Foundation, the Reinhart-Koselleck-Grant and Gottfried Wilhelm Leibniz-Prize. H.E.M. was supported by DFG ME-3644/5-1. Graphical visualizations including the graphical abstract were created with BioRender.com. We thank E. Barleon, M. Ditter and S. Hirner for excellent technical assistance.

METHODS

Ethics approvals: The analyses were performed with the approval of the Institutional Review Boards (Ethics Committee of the Albert-Ludwigs-University, Freiburg: 322/20, 10008/09; Ethics Committee of the Hamburg Chamber of Physicians: WF-051/20, PV7311). The study was performed in agreement with the principles expressed in the Declaration of Helsinki (2013).

Specimen collection: Formaldehyde-fixed paraffin-embedded sections of the upper medulla oblongata of 25 COVID-19 patients that had been tested positive for SARS-CoV-2 were obtained from autopsies that underwent routine neuropathological workup performed at the Institute of Legal Medicine of the University Medical-Center of Hamburg-Eppendorf. Five aged-matched control tissues of the same location were obtained from autopsies performed at the Department of Neuropathology, Freiburg prior to the pandemic and selected due to the causes of death of the patients being unrelated to neurological or infectious diseases.

Clinical data: Patient clinical data is summarized in **Table 1**.

Immunohistochemistry: For initial deparaffinization, slides were incubated at 80°C for one hour, then deparaffinized in Xylene and in EnVision FLEX Target Retrieval Solution High pH cooking buffer for 20 minutes (DAKO, cat. # K8000). Endogenous tissue peroxidase was blocked in 3% hydrogen peroxidase for 10 minutes (Merck Millipore, cat. no 1.08597.1000). Samples were blocked with 10% normal goat serum (NGS, BioScience, cat. # 0060-01) plus 1% Triton X 100 (Roth, cat. # 3051.4) in TRIS buffer (EnVision Flex Wash Buffer, Dako, cat. # DM831) for one hour. Double-immunolabelling for Iba1 and CD8 was performed sequentially: the sections were first incubated with Iba1 antibody (abcam, ab178846, 1:1000) in 10% NGS 1%, Triton X 100 in TRIS buffer overnight at +4°C. After washing three times with TRIS buffer, sections were incubated with a secondary antibody (goat anti-rabbit, Southern BioScience, 1:300, cat. # 4050-08) in 10% normal goat serum, 1% Triton X 100 in TRIS buffer for 45 minutes at RT. Three more washing steps were performed before incubating the sections with 0.1% streptavidin peroxidase (BioScience, cat. # 7100-05) in TRIS buffer for 45 minutes at RT. Specimens were rinsed three times with TRIS buffer. Next, slides were incubated with DAB solution: 1 drop EnVision Flex DAB Chromogen (Dako, cat. # DM827) per 1 ml EnVision Flex Substrate Buffer (Dako, cat. # DM823). The incubation time was determined by continuous

monitoring of the staining progress using a light microscope. Thereafter, the procedure was repeated for the immunohistochemical reaction for CD8 using a goat anti-mouse secondary antibody (Southern BioScience, 1:200, cat. # 1031-08) and 0.1% streptavidin-AP (BioScience, cat. # 7100-04). Permanent Red (Monosan Permanent AP-Red Kit, Monosan, cat. # MON-APP185) served as chromogen. Finally, the slides were counterstained with Gill's Hematoxylin solution (1.05175.0500, MerckMillipore). Coverslips were mounted with Kaisers Glycerin-Gelatine (Roth, cat. # 6474.1). Images were taken with a 20x objective (Keyence, BioRevo). The immunohistochemical reactions for APP (Merck, cat. # MAB348) and ACE2 (Novus Biologicals cat. # NBP2-80035) were performed using the EnVision FLEX Target Retrieval Solution Low pH cooking buffer and the EnVision FLEX system (DAKO, # K8000). Slides were counterstained with Gill's Hematoxylin solution (1.05175.0500, MerckMillipore). Coverslips were mounted with xylene-based Vitro-Clud mounting medium (R. Langenbrinck, cat. # 04-0001). APP-positive deposits were counted manually using a light microscope (Olympus, BX41). Representative pictures were acquired with a 20x and 40x objective (Leica, DFC450).

IMC antibody panel: An IMC panel was designed including markers for the adaptive and innate immune system as well as brain-specific cell populations (Table 2). Metal-labeled antibodies were either obtained pre-conjugated (Fluidigm) or labeled in-house by conjugating purified antibodies to lanthanide metals using the Maxpar X8 antibody labelling kit (Fluidigm) according to the manufacturer's instructions. 89-Yttrium (III) nitrate tetrahydrate (Sigma Aldrich, cat. # 217239-10G) and 157-Gadolinium (III) chloride (Trace Sciences Int.) were diluted in L-buffer to a 1M stock solution and further diluted to a 50 μ M working solution for subsequent antibody labelling with the Maxpar X8 labelling kit. Metal-conjugated antibodies were titrated and validated on brain, liver and tonsil tissue.

Sample preparation and staining for IMC: Tissue sections were baked at 60°C for one hour, dewaxed in Xylene twice for 15 minutes and rehydrated in descending grades of ethanol (100% - 100% - 95% - 80%) for five minutes each, then rinsed in TBS (pH 7.6) for 10 minutes. Epitope retrieval was performed in a decloaking chamber (Biocare Medical) with EnVision FLEX Target Retrieval Solution High pH for 30 minutes at 95°C. Slides were cooled down to room temperature for 20 minutes inside the buffer and 10 minutes in TBS. Tissue sections were encircled with a PAP pen (ImmEdge, Vector laboratories, cat. # H-

452 4000) and blocked for 45 minutes at room temperature using SuperBlock (TBS) Blocking Buffer
453 (ThermoFisher Scientific, cat. # 37581). Afterwards, the sections were stained with a mix of metal-labeled
454 primary antibodies diluted in TBS with 0.05% BSA and incubated at 4°C overnight. Slides were rinsed in
455 TBS-T twice (TBS supplemented with 0.2% Tween-20) and twice in TBS for 5 minutes each. Tissue
456 sections were then stained with 1:2000 Iridium Cell-ID intercalator (500 µM, Fluidigm cat. # 201192B) in
457 TBS for 30 minutes at room temperature. Slides were rinsed three times for 5 minutes in TBS, dipped in
458 ddH₂O for 5 seconds and air-dried. Slides were stored at room temperature until image acquisition.

459 *Image acquisition:* One 1500 µm² image per sample was acquired using a Hyperion Imaging System
460 (Fluidigm). Briefly, tuning of the instrument was performed according to the manufacturer's instructions.
461 Regions of interest were determined using an Iba1 and CD8a double immunohistochemistry staining. Laser
462 power was set to three and tissue sections were laser-ablated spot-by-spot at 200 Hz resulting in a pixel
463 size/resolution of 1 µm². Preprocessing of the raw data was conducted using the CyTOF software v7.0
464 (Fluidigm). Image acquisition control was performed using MCD Viewer v1.0.560.6 (Fluidigm).

465 *Data processing:* MCD files containing the raw data were converted into tiff files and segmented into single
466 cells using an unbiased, supervised analysis pipeline adapted from
467 <https://github.com/BodenmillerGroup/lmcSegmentationPipeline> and processed as reported previously¹⁷.
468 Briefly, nucleated cells were segmented using a combination of CellProfiler v3.1.9 and Ilastik v1.3.3. Pixels
469 were classified into nuclei, cytoplasm and background using Ilastik. A probability map for the three
470 classifications was generated and used to create a cell mask with CellProfiler. For combination with a
471 subsequent Iba1 mask, Iba1 signal was classified as background. A second cell mask was then created in
472 CellProfiler using Iba1 expression of the images. Both masks were combined in CellProfiler using the "AND"
473 function in ImageMath.

474 Data folders containing tiff images of the 38 markers, seven background channels and the combined cell
475 mask were loaded into histoCAT v1.76 (<https://github.com/BodenmillerGroup/histoCAT>) and mean marker
476 intensity of pixels and spatial features of segmented cells were calculated.

477 *Manual cell count:* CD3⁺CD8⁺, CD3⁺CD4⁺, CD3⁺GzmB⁺, CD20⁺, parenchymal Iba1⁺ microglia cells and
478 CD163⁺ perivascular macrophages were visualized in MCD viewer and counted manually by two
479 independent blinded scientists.

480 *Data transformation and normalization:* The presented data was normalized to 99th percentile for
481 Phenograph clustering (in histoCAT) Fig. 2A und 2B. For opt-SNE analysis presented in Fig. 2A, 2C and
482 2E, heatmap visualization and gating in Omiq.ai (Santa Clara, CA), all channels used for further analysis
483 were arcsinh transformed with a cofactor of 0.2. In heatmaps (Fig. 2B, 3D and 3E) z-scored cluster means
484 are depicted.

485 *High-dimensional analysis:* Phenograph clustering algorithm implemented in histoCAT (v1.76) was used to
486 identify even rare clusters of cells with distinct marker expressions using the following settings: Nearest
487 neighbors = 15, random seed = 2 and clustering was performed based on data from channels containing
488 lineage and activation markers: CD103, CD163, CD204, CD20, CD33, CD34, CD39, CD3, CD45RO, CD45,
489 CD4, CD68, CD8a, Collagen I, Eomes, GFAP, GzmB, HLA-DR, Iba1, Ki-67, NeuN, PD-1, PD-L1, Tim-3.
490 Segmentation events with an area of < 20 pixels were excluded. Single-cell cluster data was integrated with
491 expression data and exported for further analysis to FlowJo (TreeStar, Inc.) and Omiq (Omiq, Inc.). Analysis
492 was performed on Phenograph clusters containing more than 100 cells in the total dataset consisting of
493 total n = 96,882 cells.

494 *t-SNE analysis:* Visualization of the global single-cell landscape was performed using Opt-SNE in Omiq
495 based on channels (CD103, CD163, CD204, CD20, CD33, CD34, CD39, CD3, CD45RO, CD45, CD4,
496 CD68, CD8a, Collagen I, Eomes, GFAP, GzmB, HLA-DR, Iba1, Ki-67, NeuN, PD-1, PD-L1, Tim-3), random
497 seed 3800, max iterations 1000, opt-SNE end 5000, perplexity 30, theta 0.5 and verbosity 25.
498 Subsequently, Phenograph clusters identified as immune cells were visualized using opt-SNE calculated
499 on immune cell markers (CD103, CD163, CD204, CD20, CD33, CD39, CD3, CD45RO, CD45, CD4, CD68,
500 CD8a, Eomes, GzmB, HLA-DR, Iba1, Ki-67, PD-1, PD-L1, Tim-3, TOX), random seed 691, max iterations
501 1000, opt-SNE end 5000, perplexity 30, theta 0.5 and verbosity 25.

502 *Heatmaps:* Signal intensity of marker expression in Phenograph clusters was determined in Omiq and
503 visualized in a heatmap in R studio v4.0.2 Pheatmap package.

504 *Cluster compositions (stacked bar graph)*: Clusters were gated on patients and counts were exported from
505 Omiq. The cluster composition between patient groups was visualized using the ggplot2 stacked bar graph
506 function in R studio.

507 *Correlation heatmaps*: Linear regression analysis of Phenograph cluster frequencies in COVID-19 samples
508 was performed pairwise across all Phenograph clusters (JMP 12.2.0, SAS Institute, Inc). P values for
509 individual cluster pairs were determined and summarized in a cluster matrix. P values were then visualized
510 as a heatmap using R v4.0.2 Pheatmap package.

511 *Neighborhood analysis*: Spatial cellular interaction patterns were tested by neighborhood analysis as
512 implemented in histoCAT, v1.76. Briefly, it was examined per image whether a cell of a certain Phenograph
513 cluster was located significantly ($p = 0.05$) more frequently in close proximity (4 pixel) to a cell of another
514 cell cluster than expected by a random distribution of cells. The results for each cluster pair were
515 summarized and visualized in a heatmap using R v4.0.2 Pheatmap package.

516 *Vascular proximity map*: FIJI, a distribution of ImageJ, v1.52p, was used for the calculation of the proximity
517 to the closest Collagen-I-positive structure. Background and noise were reduced by excluding extremely
518 weak signal in the 169Tm-Collagen-I imaging mass cytometry channel (threshold 1.65) and the median
519 filter (radius 0.5 pixels). The values for all 2.25 million pixels per image were obtained and plotted with
520 ggplot2 function in R studio.

521 *Microglia nodule index map*: The calculation of the microglia nodule index map was performed in a similar
522 way to the vascular proximity map. The starting point was the signal in the 141Pr-Iba1 channel. The 141Pr-
523 Iba1-positive pixels were measured in a radius of 15 μm and put in relation to the total area. The values
524 were calculated for all 2.25 million pixels in the image. The map was again visualized using ggplot2 in R
525 studio.

526 *Microglia nodule and microgliosis quantification and pie chart*: The composition of the cohort was visualized
527 based on the criteria of microgliosis (moderate microgliosis with > 80 microglia cells per mm^2 and severe
528 microgliosis > 160 cells per mm^2) and the presence or absence of microglia nodules.

529 *CNS compartments:* Distances to vasculature inferred by distance to nearest collagen-I expressing pixel
530 and microglia nodule index were used to identify 4 compartments: Microglia nodules (nodule index ≥ 0.5),
531 perivascular (nodule index < 0.5 , distance to collagen-I $< 5 \mu\text{m}$), juxtavascular (nodule index < 0.5 , distance
532 to collagen-I $\geq 5 \mu\text{m}$ and $\leq 20 \mu\text{m}$), parenchymal (nodule index < 0.5 , distance to collagen-I $> 20 \mu\text{m}$).

533 Statistical analysis: Statistical analyses were performed using GraphPad Prism 8 (GraphPad Software,
534 Inc.). Differences in cell counts of clusters and cell populations as well as APP density between COVID-19
535 and control patients in Figures 1D, 1E, 3A and 3F were analyzed by t-tests with Welch's correction.
536 Frequencies of cell populations between different CNS compartments in figures 4E and 4F were analyzed
537 by Mann-Whitney-U tests. If not indicated otherwise in figure legends, throughout the manuscript * indicates
538 $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$, **** $p < 0.0001$.

FIGURE LEGENDS

Figure 1: Highly multiplexed imaging mass cytometry analysis of COVID-19 brains.

A: Graphical illustration of the experimental workflow. Medulla oblongata tissue slices from autopsies of COVID-19 patients (n = 25) and controls (n = 5) underwent parallel conventional immunohistochemistry and high-dimensional imaging mass cytometry (IMC) with 38 markers.

B: Exemplary immunohistochemistry (left) and corresponding IMC visualization (right) of the upper medulla tissue of a COVID-19 patient. The area of interest is magnified in the right upper corner. Scale bars indicate 100 μ m and 50 μ m, respectively. Left: Double-chromogenic immunohistochemical reaction for Iba1 (brown) and CD8 (pink), counterstained with haematoxylin (blue). Arrows indicate CD8a⁺ T cells. Right: Image overlay visualization of Iba1 (red), CD8 (green), collagen-I (light blue), CD163 (yellow) and histone H3 (blue) IMC data out of the 38-marker dataset.

C: Visualization of the indicated marker expression data from channels shown exemplarily for one area of interest as indicated in B). The scale bar represents 50 μ m.

D: Manual counts of microglial cells, perivascular macrophages, CD4, CD8 T cells, B cells and NK cells was performed on the IMC dataset and compared between controls (blue) and COVID-19 patients (green). Data are shown as bar graphs $\pm$ SEM.

E: Robust axonal damage in COVID-19 brains: immunohistochemical reaction for APP (brown) in the upper medulla of control and COVID-19 patients. Counterstaining with hematoxylin (blue). Scale bars indicate 100 μ m and 10 μ m in the inserts, respectively. The quantification of APP-positive deposits in the whole section is shown next to the histological images. Data are shown as bar graphs $\pm$ SEM.

Significant differences were determined by unpaired *t* tests with Welch's correction (D,E). In D all controls were negative for manual B cell counts, therefore a one sample *t* test was used here.

Figure 2: Molecular and cellular census of neuroinflammation during SARS-CoV-2 infection.

A: Segmentation of the highly multiplexed IMC images into cellular masks representing single cells was performed on all patients and controls using supervised machine learning. Single-cell data was extracted and cells were clustered using Phenograph and visualized on a *t*-SNE map.

B: Heatmap displaying mean marker expression (shown as z-score) of the Phenograph clusters. Clusters were annotated using the expression and spatial distribution of the cells.

C: The neuroimmune landscape of SARS-CoV-2 infected and non-infected patients was visualized by *t*-SNE based on the immunological clusters determined in B.

D: The immune cell cluster composition was assessed between the two groups. Stacked bar charts indicate mean counts per group.

E: The brain immune map from C is plotted according to cellular origin: cells from COVID-19 patients are depicted in blue; cells from controls are shown in orange.

Figure 3: Profiling of the CNS immune response during COVID-19 indicates a localized and orchestrated adaptive immune infiltration.

578 A: High-dimensional annotated cells were compared across patients and controls. Typical cell clusters with
579 significantly different abundances are displayed. Means are presented as bar graphs, error bars are
580 indicating $\pm$ SEM.

581 B: Pairwise linear regression analysis between all cluster frequencies in the brains of COVID-19 patients
582 was conducted and p-values of individual cluster pairs were visualized in a heatmap.

583 C: Violin plot indicating the distance of CD8⁺ cells to the nearest vessel (collagen-I-positive) structure in
584 COVID-19 patients. A random distribution is shown for comparison. The 25th, 50th and 75th percentiles are
585 depicted. The CD8 T cell distribution is further visualized by an overlaid scattered dot plot, showing all CD8⁺
586 cells and the corresponding distance to the closest vessel structure in 25 COVID-19 patients. Each dot
587 represents one cell.

588 D: Spatial interactions between cell clusters were assessed in COVID-19 patients by permutation based
589 neighborhood analysis and displayed as hierarchically clustered heatmap of the neighborhood interaction
590 (fraction of significant images) ranging from -1 (avoidance) to +1 (preferential interaction). Rows indicate
591 the neighborhood of a cell phenotype of interest. Columns indicate the enrichment or depletion of a cell in
592 other neighborhoods.

593 E: Cluster 22 spatial neighborhood interactions were determined in COVID-19 and control patients. Rows
594 indicate the neighborhood of a cell phenotype of interest.

595 F: CD8 T-cell activation in the brains of COVID-19 patients. Cluster 22 cells were tested for expression of
596 immune markers involved in T-cell activation, differentiation, exhaustion and function. Absolute cell counts
597 were compared between patient groups. Data are presented as boxplots showing the median, IQR and
598 upper and lower whiskers and dots represent samples.

599 Significant differences were determined by unpaired *t* tests with Welch's correction (A, F).

600

601 **Figure 4: Spatial compartmentalization of immune cell clusters to distinct anatomical**
602 **niches in the brain of COVID-19 patients.**

603 A: Left: Representative IMC image showing Iba1 (red) and CD8a (green) expression. Middle: Microglial
604 distribution was assessed using the accumulation of Iba1 signal across a 15 μ m and visualized in a
605 microglia nodule index map. Right: magnified inserts of the area of interest indicated in the left and middle
606 panel. Scale bars indicate 100 μ m and 50 μ m, respectively.

607 B: Vascular distance was estimated as distance to collagen-I. A proximity map displaying the distance of
608 each pixel to the next Collagen-I-positive structure was generated and overlaid with collagen-I expression
609 signal. Close proximity is indicated by increasing darkness. Collagen-I expression is indicated in blue.

610 C: Patient sections were assessed for presence of microgliosis or microglial nodules and visualized in a pie
611 chart. There was a significant difference in the prevalence of microglial alterations (microgliosis and/or
612 nodules) between the two groups.

613 D: Cluster composition of anatomical compartments. Images show examples of anatomical niches in one
614 patient sample. Image overlay visualization of Iba1 (red), CD8a (green), collagen I (light blue), CD163
615 (yellow) and histone H3 (blue). Scale bar indicates 50 μ m. Arrows and asterisks point to CD8 T cells located
616 inside a vessel (perivascular), close to a vessel (juxtavascular), infiltrating into the parenchyma
617 (parenchymal) and inside a microglial nodule (nodule compartment), respectively. Cluster composition of
618 the four compartments was evaluated for all COVID-19 patients and displayed as stacked bar graphs of
619 cluster frequencies per anatomical compartment.

620 E: Enrichment of individual clusters in anatomic compartments was assessed by calculating and visualizing
621 the enrichment index (mean cluster frequency in given compartment / mean frequency in the total COVID-
622 19 cohort).

623 Unpaired *t* test with Welch's correction was used in B.

624 **Figure 5: Assessment of immune cell activation in different anatomical compartments**
625 **suggests pervasive inflammatory effect of microglial nodules on distant T-cell activation**
626 **and immune checkpoint expression.**

627 A: Analysis of CD8 T cells in different CNS compartments. Frequencies of PD-1⁺, CD39⁺, PD-1⁺ CD39⁺,
628 Tim3⁺ and Eomes⁺ and HLA-DR⁺ cells of cluster 22 were compared across perivascular, juxtavascular,
629 parenchymal and nodule compartments. Boxplots show the median with IQR and upper and lower whiskers.

630 B: HLA-DR expression of CD8 T cells, CD4 T cells and activated microglial cells in different CNS
631 compartments. Left: Spatial heatmap displaying HLA-DR signal intensities in segmented cells on the image.
632 Scale bar indicates 100 μ m. Right: Frequencies of HLA-DR⁺ cells of cluster c22 (CD8 T cells), c29 (CD4 T
633 cells), c7 (activated microglia) and c15 (activated microglia) compared across perivascular, juxtavascular,
634 parenchymal and nodule compartments. Boxplots show the median with IQR and upper and lower whiskers.

635 C: CD8 T-cell activation in separate anatomical regions. Patients were grouped based on the presence or
636 absence of microglial nodules in the image and the frequencies of PD-1⁺, CD39⁺, PD-1⁺CD39⁺, Eomes⁺
637 and HLA-DR⁺ of cluster c22 were compared between those patients in the perivascular, juxtavascular and
638 parenchymal compartment to reveal a pervasive effect of microglial nodules on the activation of CD8 T
639 cells. Boxplots show the median with IQR and upper and lower whiskers.

640 D: Compartment-dependent expression of activation and immunoregulatory molecules by activated
641 microglial cells. Frequencies of CD68⁺, CD33⁺, CD39⁺, Tim-3⁺ and PD-L1⁺ cells of cluster c7 were
642 compared across the four different CNS compartments. Boxplots show the median with IQR and upper and
643 lower whiskers.

644 E: Exemplary image showing PD-1 (blue) and PD-L1 (red) expression on a spatial heatmap of segmented
645 cells.

646 Mann-Whitney-U non-parametric t-tests were used in A-D.

SUPPLEMENTARY DATA

Supplementary figure 1:

A: Bar graph visualizing cell counts of Phenograph clusters. Clusters are enumerated based on abundance of a given cluster in the combined analysis of all tissue samples.

B: Immunohistochemical reaction for ACE2 (brown) in the upper medulla of a control patient. Counterstaining with haematoxylin (blue). An ACE2-positive structure, histopathologically corresponding to a vessel, is indicated with an arrow. The scale bar represents 50 μm .

C: Cluster 22 spatial neighborhood interactions were determined in COVID-19 patients and controls. Spatial interactions of single c22 cells in other neighborhoods are shown.

D: Image overlay visualization of Iba1 (red), CD8a (green), collagen I (light blue), CD163 (yellow) and histone H3 (blue). Scale bar indicates 100 μm . Boxes indicate exemplary images of anatomical compartments shown in Figure 4D. Region 1: perivascular compartment, region 2: juxtavascular compartment, region 3: parenchyma, region 4: nodule compartment.

E: Frequency of Tim-3⁺ cells of cluster c22 compared between COVID-19 patients without microglial nodules (orange) and with microglial nodules (red) in the perivascular (left), juxtavascular (middle) and parenchymal (right) compartment respectively. Boxplots show the median with IQR and upper and lower whiskers.

F: Compartment-dependent expression of activation and immunoregulatory molecules by activated microglial cells. Frequencies of CD68⁺, CD33⁺, CD39⁺, Tim-3⁺ and PD-L1⁺ cells of cluster c15 were compared across the four different CNS compartments. Boxplots show the median with IQR and upper and lower whiskers.

G: PD-1 and PD-L1 signal intensities in segmented cells on the image.

H: Axonal damage in controls and COVID-19 patients: data from Figure 1E is depicted according to the presence or absence of microglia nodules. Data are shown as scattered dot plot. Error bars indicate SEM. Each dot represents one patient.

Mann-Whitney-U non-parametric t-tests were used in E and F.

SUPPLEMENTARY TABLES

Table 1: Cohort and clinical data

Table showing patient group (COVID-19 and control), sex, age, place of death (ER: Emergency room, hosp.: Hospital ward, ICU: Intensive care unit, NH: Nursing home), post-mortem interval (PMI), cause of death (pe: pulmonary artery embolisms, scd: sudden cardiac death), comorbidities (CI: cardiac insufficiency, COPD: chronic obstructive pulmonary disease, DM: diabetes mellitus, IHD: ischemic heart disease, RI: renal insufficiency, Tx: Transplantation) and details of the brain autopsy for each patient.

Table 2: IMC brain adaptive and innate immune system panel

Table listing antibody clones used for IMC staining with information on metal label, company of which the antibody was obtained, catalog number and dilution at which the antibody was used.

REFERENCES

- 685 1 Mao, L. *et al.* Neurologic Manifestations of Hospitalized Patients With Coronavirus
686 Disease 2019 in Wuhan, China. *JAMA Neurol*, doi:10.1001/jamaneurol.2020.1127 (2020).
- 687 2 Helms, J. *et al.* Neurologic Features in Severe SARS-CoV-2 Infection. *N Engl J Med* **382**,
688 2268-2270, doi:10.1056/NEJMc2008597 (2020).
- 689 3 Varatharaj, A. *et al.* Neurological and neuropsychiatric complications of COVID-19 in 153
690 patients: a UK-wide surveillance study. *Lancet Psychiatry*, doi:10.1016/S2215-
691 0366(20)30287-X (2020).
- 692 4 Ellul, M. A. *et al.* Neurological associations of COVID-19. *Lancet Neurol*,
693 doi:10.1016/S1474-4422(20)30221-0 (2020).
- 694 5 Puelles, V. G. *et al.* Multiorgan and Renal Tropism of SARS-CoV-2. *N Engl J Med* **383**,
695 590-592, doi:10.1056/NEJMc2011400 (2020).
- 696 6 Kantonen, J. *et al.* Neuropathologic features of four autopsied COVID-19 patients. *Brain*
697 *Pathol*, doi:10.1111/bpa.12889 (2020).
- 698 7 Reichard, R. R. *et al.* Neuropathology of COVID-19: a spectrum of vascular and acute
699 disseminated encephalomyelitis (ADEM)-like pathology. *Acta Neuropathol* **140**, 1-6,
700 doi:10.1007/s00401-020-02166-2 (2020).
- 701 8 Schaller, T. *et al.* Postmortem Examination of Patients With COVID-19. *JAMA*,
702 doi:10.1001/jama.2020.8907 (2020).
- 703 9 Solomon, I. H. *et al.* Neuropathological Features of Covid-19. *N Engl J Med*,
704 doi:10.1056/NEJMc2019373 (2020).
- 705 10 von Weyhern, C. H., Kaufmann, I., Neff, F. & Kremer, M. Early evidence of pronounced
706 brain involvement in fatal COVID-19 outcomes. *Lancet* **395**, e109, doi:10.1016/S0140-
707 6736(20)31282-4 (2020).
- 708 11 Mathew, D. *et al.* Deep immune profiling of COVID-19 patients reveals distinct
709 immunotypes with therapeutic implications. *Science*, doi:10.1126/science.abc8511
710 (2020).
- 711 12 Kuri-Cervantes, L. *et al.* Comprehensive mapping of immune perturbations associated
712 with severe COVID-19. *Sci Immunol* **5**, doi:10.1126/sciimmunol.abd7114 (2020).
- 713 13 Prinz, M. & Priller, J. The role of peripheral immune cells in the CNS in steady state and
714 disease. *Nat Neurosci* **20**, 136-144, doi:10.1038/nn.4475 (2017).
- 715 14 Tay, T. L. *et al.* A new fate mapping system reveals context-dependent random or clonal
716 expansion of microglia. *Nat Neurosci* **20**, 793-803, doi:10.1038/nn.4547 (2017).
- 717 15 Market, M. *et al.* Flattening the COVID-19 Curve With Natural Killer Cell Based
718 Immunotherapies. *Front Immunol* **11**, 1512, doi:10.3389/fimmu.2020.01512 (2020).
- 719 16 Zheng, M. *et al.* Functional exhaustion of antiviral lymphocytes in COVID-19 patients. *Cell*
720 *Mol Immunol* **17**, 533-535, doi:10.1038/s41423-020-0402-2 (2020).
- 721 17 Schapiro, D. *et al.* histoCAT: analysis of cell phenotypes and interactions in multiplex
722 image cytometry data. *Nat Methods* **14**, 873-876, doi:10.1038/nmeth.4391 (2017).
- 723 18 Levine, J. H. *et al.* Data-Driven Phenotypic Dissection of AML Reveals Progenitor-like
724 Cells that Correlate with Prognosis. *Cell* **162**, 184-197, doi:10.1016/j.cell.2015.05.047
725 (2015).
- 726 19 Bengsch, B. *et al.* Epigenomic-Guided Mass Cytometry Profiling Reveals Disease-
727 Specific Features of Exhausted CD8 T Cells. *Immunity* **48**, 1029-1045 e1025,
728 doi:10.1016/j.immuni.2018.04.026 (2018).
- 729 20 Sinha, P., Matthay, M. A. & Calfee, C. S. Is a "Cytokine Storm" Relevant to COVID-19?
730 *JAMA Intern Med*, doi:10.1001/jamainternmed.2020.3313 (2020).
- 731 21 Netland, J., Meyerholz, D. K., Moore, S., Cassell, M. & Perlman, S. Severe acute
732 respiratory syndrome coronavirus infection causes neuronal death in the absence of

733 encephalitis in mice transgenic for human ACE2. *J Virol* **82**, 7264-7275,
734 doi:10.1128/JVI.00737-08 (2008).

735 22 Cortese, I. *et al.* Pembrolizumab Treatment for Progressive Multifocal
736 Leukoencephalopathy. *N Engl J Med* **380**, 1597-1605, doi:10.1056/NEJMoa1815039
737 (2019).

738 23 Masuda, T. *et al.* Spatial and temporal heterogeneity of mouse and human microglia at
739 single-cell resolution. *Nature* **566**, 388-392, doi:10.1038/s41586-019-0924-x (2019).

740 24 Lunemann, A. *et al.* Human NK cells kill resting but not activated microglia via NKG2D-
741 and NKp46-mediated recognition. *J Immunol* **181**, 6170-6177,
742 doi:10.4049/jimmunol.181.9.6170 (2008).

743 25 Hao, J. *et al.* Central nervous system (CNS)-resident natural killer cells suppress Th17
744 responses and CNS autoimmune pathology. *J Exp Med* **207**, 1907-1921,
745 doi:10.1084/jem.20092749 (2010).

Fig. Graphical Abstract

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Supplementary Fig. 1

ID	Group	Sex	Age	POD	PMI	CauseOfDeath	Comorbidities	Brainweight[g]	Brain edema	Brain atrophy	Macroscopic findings
1	COVID-19	m	75	NH	3	sudden cardiac death	Parkinson's disease	1110	1	0	none
2	COVID-19	f	87	NH	0	pneumonia	COPD, dementia, IHD, RI	1215	0	1	none
3	COVID-19	f	85	hosp.	0	pneumonia	atrial fibrillation, CI, IHD, myelofibrosis, RI	1240	0	1	Fresh infarction in territory of ACP
4	COVID-19	f	86	NH	0	pneumonia	COPD, dementia, IHD	1250	0	1	Fresh infarction in territory of ACP
5	COVID-19	m	90	NH	2	pneumonia	atrial fibrillation, dementia, DM, history of stroke	1015	0	2	Old infarctions in territory of ACP
6	COVID-19	m	90	hosp.	3	emphysema with respiratory decompensation	CI, COPD	1440	0	1	none
7	COVID-19	m	77	hosp.	2	pneumonia	aortic aneurysm, atrial flutter, cardiac hypertrophy, emphysema, RI	1590	2	0	none
8	COVID-19	m	76	ICU	3	pe, respiratory tract infection	CI, COPD	1460	1	0	none
9	COVID-19	m	70	hosp.	1	pneumonia (aspiration)	CI, COPD, IHD, Parkinson's disease	1430	1	0	none
10	COVID-19	m	93	hosp.	3	pneumonia	DM, hypertension	1400	1	0	none
11	COVID-19	m	66	ER	2	pneumonia	DM, IHD	1450	1	0	none
12	COVID-19	f	54	hosp.	1	pneumonia	trisomy 21, epilepsy	950	0	3	Grey matter heterotopia
13	COVID-19	f	86	NH	2	sepsis, pneumonia	emphysema, epilepsy, hypoxic brain damage, IHD, RI	1210	0	1	none
14	COVID-19	f	85	home	1	pneumonia	CI, COPD	1180	0	1	none
15	COVID-19	f	70	ICU	3	pneumonia	CI	1150	0	1	none
16	COVID-19	f	75	ICU	4	pneumonia	cardiac arrhythmia, IHD	1210	0	1	none
17	COVID-19	m	93	hosp.	2	pneumonia	atrial fibrillation, CI, IHD, obstructive sleep apnoea syndrome	1000	0	2	Old cerebellar infarction
18	COVID-19	f	82	hosp.	4	purulent bronchitis	COPD, history of pulmonary embolism, RI	1080	0	2	none
19	COVID-19	m	63	ICU	1	pe, pneumonia	CI	1435	1	0	Fresh infarction in territory of ACA
20	COVID-19	m	84	hosp.	5	pneumonia, septic encephalopathy	DM, history of stroke, hypertension, IHD, ulcerative colitis	1350	1	0	none
21	COVID-19	m	71	ICU	2	pe, pneumonia	CI, DM, lung granuloma	1665	2	0	none
22	COVID-19	m	59	hosp.	12	pneumonia	cardiomyopathy	1575	2	0	none
23	COVID-19	m	85	ICU	2	pneumonia	COPD, history of aortic valve replacement, hypertension, IHD	1400	1	0	none
24	COVID-19	f	75	home	2	pulmonary artery embolism	hypertension, IHD	1095	0	2	none
25	COVID-19	f	76	hosp.	2	pneumonia	breast cancer, hypertension	1180	0	1	none
26	control	f	82	hosp.	3	aspiration	hypertension	1380	0	0	none
27	control	m	81	hosp.	2	respiratory failure / invasive lung carcinoma	lung squamous cell carcinoma, DVT, pulmonary embolisms, CI	1450	0	0	none
28	control	f	88	hosp.	1	cardiac failure with cardiomegaly	tension pneumothorax, aortic valve replacement	1260	0	0	none
29	control	m	87	hosp.	3	hemorrhagic shock / GI bleeding	atrial fibrillation, anticoagulation	1110	0	0	none
30	control	f	78	hosp.	2	ambilateral fulminant pulmonary embolism	hypertension	1300	0	0	none

m: male, f: female

n/a: not available

POD (place of death): ER: Emergency room, hosp.: Hospital ward, ICU: Intensive care unit, NH: Nursing home

cause of death: pe: pulmonary artery embolisms, scd: sudden cardiac death

PMI: postmortem interval (days)

co-morbidities: CI: cardiac insufficiency, COPD: chronic obstructive pulmonary disease, DM: diabetes mellitus, IHD: ischemic heart disease, RI: renal insufficiency, Tx: Transplantation

Brain edema, brain atrophy, arteriosclerosis: 0: none, 1: basal, 2: moderate, 3: severe

Macroscopic findings: ACP: Arteria cerebri posterior, ACA: Arteria cerebri anterior, ACM: Arteria cerebri media

Metal	Target	Clone	Vendor	Catalog #	Dilution
89Y	CD45	D9M8I	CST	13917BF	1:50
115In	HLA-DR	TAL-1B5	abcam	ab176408	1:400
141Pr	Iba1	EPR16588	abcam	ab178846	1:200
142Nd	CD56	EPR2566	abcam	ab214436	1:100
143Nd	GFAP	EPR1034Y	abcam	ab218309	1:100
144Nd	CD69	EPR21814	abcam	ab234512	1:100
146Nd	CD39	EPR20627	abcam	ab236038	1:50
147Sm	CD163	EDhu-1	fluidigm	3147021D	1:200
148Nd	NeuN	1B7	BioLegend	834501	1:400
149Sm	CD103	EPR22590-27	abcam	ab2454201	1:50
150Nd	PD-L1	E1L3N	CST	13684S	1:50
151Eu	ACE-2	AC18F	Novus	NBP2-80035	1:50
152Sm	TCF-1	C63D9	CST	2203BF	1:100
153Eu	TOX	NAN448B	abcam	ab237009	1:100
154Sm	Tim-3	D5D5R	fluidigm	3154024D	1:800
155Gd	FoxP3	236A/E7	Thermo fisher	14-4777-82	1:100
156Gd	CD4	EPR6855	fluidigm	3156033D	1:400
157Gd	TIGIT	4A12	Enzo	ENZ-ABS698-0100	1:100
158Gd	IRF8	7G11A45	BioLegend	656501	1:100
159Tb	CD68	KP1	fluidigm	3159035D	1:2000
160Gd	T-bet	4B10	fluidigm	3160010B	1:100
161Dy	CD20	H1	fluidigm	3161029D	1:800
162Dy	CD8a	C8/144B	fluidigm	3162034D	1:800
163Dy	Eomes	WD1928	CST	14-4877-82	1:100
164Dy	Arginase-1	D4E3M	fluidigm	316027D	1:100
165Ho	PD-1	D4W2J	CST	86163BF	1:100
166Er	CD204	J5HTR3	invitrogen	14-9054-82	1:400
167Er	GranzymeB	EPR20129-217	fluidigm	3167021D	1:1600
168Er	Ki-67	B56	fluidigm	3168022D	1:100
169Tm	Collagen I	Poly	fluidigm	3169023D	1:800
170Er	CD3	Poly	fluidigm	3170019D	1:200
171Yb	Histone H3	D1H2	fluidigm	3171022D	1:3000
172Yb	CD38	EPR4106	abcam	ab176886	1:100
173Yb	CD45RO	UCHL1	fluidigm	3173016D	1:800
174Yb	CD33	SP266	abcam	ab238784	1:100
175Lu	CD34	EP373Y	abcam	ab198395	1:400
176Lu	Myelin-CNPase	SMI 91	BioLegend	836404	1:200

Figures

Figure 1

Highly multiplexed imaging mass cytometry analysis of COVID-19 brains. A: Graphical illustration of the experimental workflow. Medulla oblongata tissue slices from autopsies of COVID-19 patients (n = 25) and controls (n = 5) underwent parallel conventional immunohistochemistry and high-dimensional imaging

mass cytometry (IMC) with 38 markers. B: Exemplary immunohistochemistry (left) and corresponding IMC visualization (right) of the upper medulla tissue of a COVID-19 patient. The area of interest is magnified in the right upper corner. Scale bars indicate 100 μm and 50 μm , respectively. Left: Double-chromogenic immunohistochemical reaction for Iba1 (brown) and CD8 (pink), counterstained with haematoxylin (blue). Arrows indicate CD8a+ T cells. Right: Image overlay visualization of Iba1 (red), CD8 (green), collagen-I (light blue), CD163 (yellow) and histone H3 (blue) IMC data out of the 38-marker dataset. C: Visualization of the indicated marker expression data from channels shown exemplarily for one area of interest as indicated in B). The scale bar represents 50 μm . D: Manual counts of microglial cells, perivascular macrophages, CD4, CD8 T cells, B cells and NK cells was performed on the IMC dataset and compared between controls (blue) and COVID-19 patients (green). Data are shown as bar graphs $\pm$ SEM. E: Robust axonal damage in COVID-19 brains: immunohistochemical reaction for APP (brown) in the upper medulla of control and COVID-19 patients. Counterstaining with hematoxylin (blue). Scale bars indicate 100 μm and 10 μm in the inserts, respectively. The quantification of APP-positive deposits in the whole section is shown next to the histological images. Data are shown as bar graphs $\pm$ SEM. Significant differences were determined by unpaired t tests with Welch's correction (D,E). In D all controls were negative for manual B cell counts, therefore a one sample t test was used here.

Figure 2

Molecular and cellular census of neuroinflammation during SARS-CoV-2 infection. A: Segmentation of the highly multiplexed IMC images into cellular masks representing single cells was performed on all patients and controls using supervised machine learning. Single-cell data was extracted and cells were clustered using Phenograph and visualized on a t-SNE map. B: Heatmap displaying mean marker expression (shown as z-score) of the Phenograph clusters. Clusters were annotated using the expression

and spatial distribution of the cells. C: The neuroimmune landscape of SARS-CoV-2 infected and non-infected patients was visualized by t-SNE based on the immunological clusters determined in B. D: The immune cell cluster composition was assessed between the two groups. Stacked bar charts indicate mean counts per group. E: The brain immune map from C is plotted according to cellular origin: cells from COVID-19 patients are depicted in blue; cells from controls are shown in orange.

Fig. 3

Figure 3

Profiling of the CNS immune response during COVID-19 indicates a localized and orchestrated adaptive immune infiltration. A: High-dimensional annotated cells were compared across patients and controls. Typical cell clusters with significantly different abundances are displayed. Means are presented as bar graphs, error bars are indicating $\pm$ SEM. B: Pairwise linear regression analysis between all cluster frequencies in the brains of COVID-19 patients was conducted and p-values of individual cluster pairs were visualized in a heatmap. C: Violin plot indicating the distance of CD8+ cells to the nearest vessel (collagen-I-positive) structure in COVID-19 patients. A random distribution is shown for comparison. The 25th, 50th and 75th percentiles are depicted. The CD8 T cell distribution is further visualized by an overlaid scattered dot plot, showing all CD8+ cells and the corresponding distance to the closest vessel structure in 25 COVID-19 patients. Each dot represents one cell. D: Spatial interactions between cell clusters were assessed in COVID-19 patients by permutation based neighborhood analysis and displayed as hierarchically clustered heatmap of the neighborhood interaction (fraction of significant images) ranging from -1 (avoidance) to +1 (preferential interaction). Rows indicate the neighborhood of a cell phenotype of interest. Columns indicate the enrichment or depletion of a cell in other neighborhoods. E: Cluster 22 spatial neighborhood interactions were determined in COVID-19 and control patients. Rows indicate the neighborhood of a cell phenotype of interest. F: CD8 T-cell activation in the brains of COVID-19 patients. Cluster 22 cells were tested for expression of immune markers involved in T-cell activation, differentiation, exhaustion and function. Absolute cell counts were compared between patient groups. Data are presented as boxplots showing the median, IQR and upper and lower whiskers and dots represent samples. Significant differences were determined by unpaired t tests with Welch's correction (A, F).

Figure 4

Spatial compartmentalization of immune cell clusters to distinct anatomical niches in the brain of COVID-19 patients. A: Left: Representative IMC image showing Iba1 (red) and CD8a (green) expression. Middle: Microglial distribution was assessed using the accumulation of Iba1 signal across a 15 μ m and visualized in a microglia nodule index map. Right: magnified inserts of the area of interest indicated in the left and middle panel. Scale bars indicate 100 μ m and 50 μ m, respectively. B: Vascular distance was estimated as distance to collagen-I. A proximity map displaying the distance of each pixel to the next

Collagen-I-positive structure was generated and overlaid with collagen-I expression signal. Close proximity is indicated by increasing darkness. Collagen-I expression is indicated in blue. C: Patient sections were assessed for presence of microgliosis or microglial nodules and visualized in a pie chart. There was a significant difference in the prevalence of microglial alterations (microgliosis and/or nodules) between the two groups. D: Cluster composition of anatomical compartments. Images show examples of anatomical niches in one patient sample. Image overlay visualization of Iba1 (red), CD8a (green), collagen I (light blue), CD163 (yellow) and histone H3 (blue). Scale bar indicates 50 μ m. Arrows and asterisks point to CD8 T cells located inside a vessel (perivascular), close to a vessel (juxtavascular), infiltrating into the parenchyma (parenchymal) and inside a microglial nodule (nodule compartment), respectively. Cluster composition of the four compartments was evaluated for all COVID-19 patients and displayed as stacked bar graphs of cluster frequencies per anatomical compartment. E: Enrichment of individual clusters in anatomic compartments was assessed by calculating and visualizing the enrichment index (mean cluster frequency in given compartment / mean frequency in the total COVID-19 cohort). Unpaired t test with Welch's correction was used in B.

Figure 5

Assessment of immune cell activation in different anatomical compartments suggests pervasive inflammatory effect of microglial nodules on distant T-cell activation and immune checkpoint expression. A: Analysis of CD8 T cells in different CNS compartments. Frequencies of PD-1+, CD39+, PD-1+ CD39+, Tim3+ and Eomes+ and HLA-DR+ cells of cluster 22 were compared across perivascular, juxtavascular, parenchymal and nodule compartments. Boxplots show the median with IQR and upper and lower

whiskers. B: HLA-DR expression of CD8 T cells, CD4 T cells and activated microglial cells in different CNS compartments. Left: Spatial heatmap displaying HLA-DR signal intensities in segmented cells on the image. Scale bar indicates 100 μ m. Right: Frequencies of HLA-DR+ cells of cluster c22 (CD8 T cells), c29 (CD4 T cells), c7 (activated microglia) and c15 (activated microglia) compared across perivascular, juxtavascular, parenchymal and nodule compartments. Boxplots show the median with IQR and upper and lower whiskers. C: CD8 T-cell activation in separate anatomical regions. Patients were grouped based on the presence or absence of microglial nodules in the image and the frequencies of PD-1+, CD39+, PD-1+CD39+, Eomes+ and HLA-DR+ of cluster c22 were compared between those patients in the perivascular, juxtavascular and parenchymal compartment to reveal a pervasive effect of microglial nodules on the activation of CD8 T cells. Boxplots show the median with IQR and upper and lower whiskers. D: Compartment-dependent expression of activation and immunoregulatory molecules by activated microglial cells. Frequencies of CD68+, CD33+, CD39+, Tim-3+ and PD-L1+ cells of cluster c7 were compared across the four different CNS compartments. Boxplots show the median with IQR and upper and lower whiskers. E: Exemplary image showing PD-1 (blue) and PD-L1 (red) expression on a spatial heatmap of segmented cells. Mann-Whitney-U non-parametric t-tests were used in A-D.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [GraphicalAbstract.png](#)
- [SupplementaryData.pdf](#)