Scabies, Incomplete Lupus Erythematosus and Henoch-Schonlein Purpura: A Case Report

Yang Fang Wu

First Affiliated Hospital of Anhui Medical University

Jing Jing Wang

Jinling Hospital

Hui Hui Liu

First Affiliated Hospital of Anhui Medical University

Wei Xia Chen

First Affiliated Hospital of Anhui Medical University

Peng Hu (✉ hupeng28@aliyun.com)

First Affiliated Hospital of Anhui Medical University <https://orcid.org/0000-0002-2144-9806>

Case report

Keywords: Antinuclear antibody, Henoch-Schonlein purpura, immunoglobulin A, proteinuria, scabies

Posted Date: September 18th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-63971/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background Henoch-Schonlein purpura (HSP) is a common autoimmune vasculitis in childhood. Although the detailed etiology of HSP remains unknown, several triggers, especially for infectious agents, have been proved to be associated with HSP onset.

Case presentation: In the present report, we describe an unusual patient who suffered from HSP and incomplete lupus erythematosus (ILE) on day 8 after scabies, mainly based on nonthrombocytopenic purpura, abdominal pain, medium proteinuria, positive antinuclear antibody (ANA) and skin scrapings. A course of pulse methylprednisolone (10mg/kg/day) was administered for 6 days, followed by oral prednisone (1mg/kg/day). On day 18, the purpuric rashes almost disappeared, whereas medium proteinuria still existed. In this circumstance, a renal biopsy was performed with the informed consent of the parents. According to the criteria proposed by the International Study of Kidney Disease in Children, this patient was classified as class IIIb. The immunofluorescence microscopy revealed granular deposits of IgA, IgM and C3 in the glomerular mesangium. His proteinuria gradually decreased after treatment of corticosteroids plus mycophenolate mofetil (20mg/kg/day).

Conclusions: However, it should be noted that corticosteroids and immunosuppressors may increase the relapse risk of scabies, therefore this patient should be followed up carefully after discharging from our department.

Introduction

Henoch-Schonlein purpura (HSP) is an immunoglobulin A (IgA)-mediated, autoimmune, hypersensitivity vasculitis in childhood, mainly characterized by non-thrombocytopenic purpura, abdominal pain, arthritis and renal involvement. An epidemiological survey in Taiwan, China from 1997 to 2012 demonstrated that a total of 1002 HSP children were recruited with an annual incidence of 9.61 per 100,000 children younger than 18 years. The occurrence of HSP peaks between 7 and 9 years of age, and male children are more often affected than female children [1, 2]. Currently, the detailed etiology of HSP remains unknown. Several triggers, including infectious agents, foods, drugs, vaccinations and insect bites, have been reported to be associated with HSP onset. Recently, our research panel retrospectively reviewed the clinical records of 1200 HSP children, and found that streptococcus was the most frequent infectious trigger and identified in 205 cases (17.08%), followed by helicobacter pylori, mycoplasma, parainfluenza, respiratory syncytial virus, tubercle bacillus and toxoplasma gondii [2]. Scabies is a skin hypersensitivity reaction caused by infestation with the mite, *Sarcoptes scabiei* var. *hominis*, and its prevalence decreases in worldwide as the improvement of environmental hygiene [3]. In the present report, we encountered an unusual patient who suffered from HSP and incomplete lupus erythematosus (ILE) on day 8 after scabies.

Case Report

An 11-year-old boy was first referred to the Department of Dermatology in November 2018, with a history of nocturnal and generalized pruritus, itchy rashes on his upper trunk. Physical examination revealed multiple small(< 6 mm) papules and raised burrows on his upper trunk. By a light microscope under low power, a translucent mite around 0.2 to 0.5 mm long with four pairs of brown legs was seen on the direct examination of skin scrapings (Fig. 1a). He was diagnosed with scabies and treated with whole-body applications of 5% permethrin lotion and oral cetirizine (10 mg/day). On day 8, the papules and small burrows were significantly resolved, whereas edema and palpable purpuric skin lesions were found on his lower limbs (Fig. 1b), combined with transient abdominal pain. Under the circumstances, he was brought to our department for a further consultation. On admission, his blood pressure was 108/80 mmHg, heart rate was 65 beats/min, axillary temperature was 36.8 °C and respiratory rate was 18 breaths/min. Except for the characteristic purpura and periumbilical tenderness, the cardio-pulmonary examination revealed unremarkable. His initial blood counts were as follows: white blood cells 6.68×10^9 /l, red blood cells 5.01×10^{12} /l, hemoglobin 154 g/l, and platelets 221×10^9 /l. Other blood chemistry findings were erythrocyte sedimentation rate (ESR) 10 mm/h, C-reactive protein (CRP) 0.80 mg/l, creatinine (Cr) 63 μ mol/l, serum ferritin 20.61 μ g/l, complements 3 (C3) 1.12 g/l, complements 4(C4) 0.24 g/l and immunoglobulin A (IgA) 2.88 g/l. The autoimmune serologic workup came back positive for antinuclear antibody (ANA); negative for anti-double stranded DNA antibody (anti-dsDNA), anti-Smith antibody (anti-SM) and anti neutrophil cytoplasmic antibody (ANCA). Antibody tests for human immunodeficiency virus (HIV), hepatitis B virus (HBV), hepatitis C virus (HCV), Epstein-Barr virus (EBV), mycoplasma (MP) and tubercle bacillus (TB) were also not reactive. Urinalysis showed red blood cell: 1+, protein: 2 + with 24 h urine protein of 0.90 g. A routine stool test showed weak positive faecal occult blood but no erythrocytes. The electrocardiogram, abdominal and renal ultrasonography had no significant abnormalities. According to the above clinical and laboratory findings, he was diagnosed as having HSP and ILE. Taking into consideration medium proteinuria, a course of pulse methylprednisolone (10 mg/kg/day) was administered for 6 days, followed by oral prednisone (1 mg/kg/day). On day 18, the purpuric rashes almost disappeared, whereas medium proteinuria still existed. A renal biopsy was performed and the histopathological examination of the biopsy specimen showed 8 glomeruli with one evidence of global sclerosis, crescent formation was observed in 5/8 glomeruli. Mild mesangial proliferation and expansion were also seen in glomeruli. Moreover, mild expansion was noted in the interstitium, in association with tubular atrophy (Fig. 1c). The immunofluorescence microscopy revealed granular deposits of IgA, IgM and C3 in the glomerular mesangium (Fig. 1d). Based on these findings, he was diagnosed as having HSP nephritis (International Study of Kidney Disease in Children category: Class IIIb). The patient was treated with high-dosage intravenous methylprednisolone(30 mg/kg/day) followed by prednisone(1 mg/kg/day) combined with oral mycophenolate mofetil(20 mg/kg/day). The pulse therapy was performed three times in total and his proteinuria gradually decreased. The study protocol was approved by the Research Ethics Commission of the First Affiliated Hospital of Anhui Medical University.

Discussion And Conclusions

In this report, we encountered an unusual case who suffered from HSP on day 8 after scabies. As there is no definitive test for HSP diagnosis, the 2008 European League Against Rheumatism criteria are still the most commonly used guideline for HSP diagnosis [4]. On this basis, our patient fulfilled 3 criteria (nonthrombocytopenic purpura, abdominal pain and renal involvement), and thus the diagnosis of HSP was made undoubtedly after scabies. To the best of our knowledge, very few previous studies have documented that cutaneous vasculitis can be secondary to or coexisting with scabies [5]. Furthermore, Liu *et al.* [6] tracked 4481 scabies patients for a 7-year period and found that scabies lead to a 5.44-fold increased risk for hypersensitivity vasculitis, a 4.91-fold increased risk for dermatomyositis, a 2.89-fold increased risk for polyarteritis nodosa, a 2.73-fold increased risk for systemic lupus erythematosus respectively, compared with the healthy population. According to the current limited evidence, we speculated that the potential mechanisms of scabies-associated HSP may be attributed to the host response to antigens on *Sarcoptes scabiei* var. *Hominis*, such as immune-complex mediated injury, cytotoxic T-cell-mediated immune responses and autoimmune reactions. Further studies are warranted to clarify their shared etiologies and relationships.

As for the present case, another unique feature was the positive ANA, whereas the other diagnostic criteria for systemic lupus erythematosus (SLE) were absent. In fact, HSP and SLE share several commonalities such as their own characteristic rashes and a multi-system involvement. In our previous report, we encountered an unusual female patient who suffered from HSP at 3 months after diagnosis of SLE. Distinct from the present case, her autoimmune serologic workup indicated positive anti-dsDNA/anti-SM and low C3/C4 besides positive ANA [7]. ILE patients had at least one but less than four of the American College of Rheumatology(ACR) classification criteria of SLE and did not present distinctive clinical features or meet classification criteria of other connective tissue diseases [8]. Although the majority of ILE patients may never reach SLE classification, about 10%-50% of ILE patients are at high risk of progressing to complete SLE within 5 years [9]. In southern Sweden, Hallengren *et al.* [10] followed up 28 ILE patients for a 10-year period and found that 16 of 28 patients developed complete SLE, and malar rash and anti-cardiolipin antibodies were the potent predictors of complete SLE; in addition, positive Wasserman and ANA, thrombocytopenia, arthritis and early involvement of multiple organs could also serve as the candidates of risk factors for development of complete SLE. Based on the experiences from adult ILE, antimalarial drugs may play a beneficial role in limiting organ damages, delaying complete SLE onset, slowing the accrual of autoantibodies and reducing the need for drugs with greater toxicity, but which has not yet been formally tested in longitudinal studies and even in pediatric patients [11].

During the period of hospitalization, our patient suffered from acute kidney injury and had persistent proteinuria. The histopathological examination of the renal biopsy specimen showed that the pathological classification was class IIIb. As we all know, the degree of proteinuria has been proven to be not only a sign but an accelerator of nephropathies. A recent single-center study from Shanghai Children's Hospital showed that the pathological classification in 62.97% of HSP patients with nephrotic proteinuria were above the ISKDC class III [12]. Currently, the main treatments of HSP nephritis encompass corticosteroids alone or plus immunosuppressors. Feng *et al.* [12] administered methylprednisolone and

mycophenolate mofetil to 54 HSP patients with nephrotic proteinuria, and found that 68.52% of patients had complete remission and no patient progressed to end-stage renal disease after a follow-up of 6 months to 5 years. In the present report, it should be noted that corticosteroids and immunosuppressors may increase the relapse risk of scabies, therefore this patient should be followed up carefully after discharging from our department.

Abbreviations

American College of Rheumatology (ACR); anti-double stranded DNA antibody (anti-dsDNA); anti neutrophil cytoplasmic antibody (ANCA); antinuclear antibody (ANA); anti-Smith antibody (anti-SM); complements 3(C3); complements 4 (C4); C-reactive protein (CRP); creatinine (Cr); Epstein-Barr virus (EBV); erythrocyte sedimentation rate (ESR); hepatitis B virus (HBV); hepatitis C virus (HCV); Henoch-Schonlein purpura (HSP); human immunodeficiency virus (HIV); immunoglobulin A (IgA); incomplete lupus erythematosus (ILE); International Study of Kidney Disease in Children (ISKDC); mycoplasma (MP); systemic lupus erythematosus (SLE); tubercle bacillus (TB).

Declarations

Ethical approval and consent to participate

The study protocol was approved by the Research Ethics Commission of the First Affiliated Hospital of Anhui Medical University. Permission to carry out the study and access patient records was sought from the respective hospital administrators.

Consent for publication

Written informed consent was obtained from the patient's parents for publication of this case report and accompanying images.

Availability of data and materials

The datasets generated and/or analysed during current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

No.

Authors' contributions

Y.F.W. and J.J.W., study concept and design, manuscript development, writing; H.H.L., study support, data acquisition; W.X.C., study support, drafting/revising the image; P.H., critical revision of manuscript for intellectual content, drafting/revising the image, writing, and funding. All authors have read and agreed to the published version of the manuscript.

Acknowledgments

The authors would like to thank all those who lent their hands in the course of writing this paper. The authors would also like to thank Prof. Bo Hu and Dr. Wei Wei for their helpful comments.

References

1. Lei WT, Tsai PL, Chu SH, Kao YH, Lin CY, Fang LC, et al. Incidence and risk factors for recurrent Henoch-Schonlein purpura in children from a 16-year nationwide database. *Pediatr Rheumatol Online J*. 2018; 16: 25.
2. Xu Y, Wang JJ, Liu FF, Wu Y, Wu YF, Samadli S, et al. Predisposing factors of childhood Henoch-Schonlein purpura in Anhui province, China. *J Investig Med*. 2019; 67: 771-778.
3. Engelman D, Steer AC. Control Strategies for Scabies. *Trop Med Infect Dis*. 2018; 3: pii: E98.
4. Ozen S, Pistorio A, Iusan SM, Bakkaloglu A, Herlin T, Brik R, et al. EULAR/PRINTO/PRES criteria for Henoch-Schonlein purpura, childhood polyarteritis nodosa, childhood Wegener granulomatosis and childhood Takayasu arteritis: Ankara 2008. Part II: Final classification criteria. *Ann Rheum Dis*. 2010; 69: 798-806.
5. Clevy C, Brajon D, Combes E, Benzaquen M, Dales JP, Koeppl MC, et al. Scabietic vasculitis: Report of 2 cases. *Ann Dermatol Venereol*. 2017; 144: 349-355.
6. Liu JM, Chiu FH, Lin CY, Chang FW, Hsu RJ. Incidence of autoimmune diseases in patients with scabies: a nationwide population-based study in Taiwan. *Rheumatol Int*. 2017; 37: 1125-1134.
7. Hu P, Huang BY, Zhang DD, Jiang GM, Liu SY, Xia X, et al. Henoch-Schonlein purpura in a pediatric patient with lupus. *Arch Med Sci*. 2017; 13: 689-690.
8. Vilá LM, Mayor AM, Valentín AH, García-Soberal M, Vilá S. Clinical outcome and predictors of disease evolution in patients with incomplete lupus erythematosus. *Lupus*. 2000; 9: 110-115.
9. Laustrop H, Voss A, Green A, Junker P. Occurrence of systemic lupus erythematosus in a Danish community: an 8-year prospective study. *Scand J Rheumatol*. 2009; 38: 128-132.
10. Ståhl Hallengren C, Nived O, Sturfelt G. Outcome of incomplete systemic lupus erythematosus after 10 years. *Lupus*. 2004; 13: 85-88.
11. Aberle T, Bourn RL, Munroe ME, Chen H, Roberts VC, Guthridge JM, et al. Clinical and Serologic Features in Patients With Incomplete Lupus Classification Versus Systemic Lupus Erythematosus Patients and Controls. *Arthritis Care Res (Hoboken)*. 2017; 69: 1780-1788.
12. Feng D, Huang WY, Hao S, Niu XL, Wang P, Wu Y, et al. A single-center analysis of Henoch-Schonlein purpura nephritis with nephrotic proteinuria in children. *Pediatr Rheumatol Online J*. 2017; 15: 15.

Figures

Figure 1

Skin scrapings, manifestations, renal pathology and immunofluorescence. (a) Skin scrapings revealed translucent mite around 0.2 to 0.5 mm long with four pairs of brown legs. (b) Palpable purpura symmetrically distributed in the lower limbs. (c) Histopathological findings of the renal biopsy specimens showed 8 glomeruli with one evidence of global sclerosis, crescent formation was observed in 5/8 glomeruli. Mild mesangial proliferation and expansion were also seen in glomeruli. Moreover, mild expansion was noted in the interstitium, in association with tubular atrophy. The patient was diagnosed as having HSP nephritis (International Study of Kidney Disease in Children category: Class IIIb). (d) The immunofluorescence microscopy revealed granular deposits of IgA, IgM and C3 in the glomerular mesangium.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [CAREChecklist.pdf](#)
- [CAREChecklist.pdf](#)