

# Recent invasion and eradication of two members of the *Euwallacea fornicatus* species complex (Coleoptera: Curculionidae: Scolytinae) from tropical greenhouses in Europe

Hannes Schuler (✉ [hannes.schuler@unibz.it](mailto:hannes.schuler@unibz.it))

Free University of Bozen-Bolzano <https://orcid.org/0000-0001-8307-9831>

Radosław Witkowski

Poznan University of Life Sciences: Uniwersytet Przyrodniczy w Poznaniu

Bart van de Vossenberg

Dutch National Plant Protection Organization

Björn Hoppe

JKI Braunschweig Messeweg: Julius Kuhn-Institut Standort Braunschweig-Messeweg

Moritz Mittelbach

Senatsverwaltung für Umwelt- und Klimaschutz

Tibor Bukovinszki

Dutch National Plant Protection Organization

Stefan Schwembacher

Autonomous Province of Bolzano: Provincia Autonoma di Bolzano

Bas van de Meulengraaf

Dutch National Plant Protection Organization

Uwe Lange

Thüringer Landesamt für Landwirtschaft und Ländlichen Raum

Sabine Rode

Thüringer Landesamt für Landwirtschaft und Ländlichen Raum

Alessandro Andriolo

Autonomous Province of Bolzano: Provincia Autonoma di Bolzano

Marta Belka

Poznan University of Life Sciences: Uniwersytet Przyrodniczy w Poznaniu

Andrzej Mazur

Poznan University of Life Sciences: Uniwersytet Przyrodniczy w Poznaniu

Andrea Battisti

University of Padova: Universita degli Studi di Padova

## Research Article

**Keywords:** Ambrosia beetle, Euwallacea fornicatus, Euwallacea perbrevis, invasive species, Polyphagous shot hole borer, Tea shot hole borer

**Posted Date:** July 12th, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-640781/v2>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

1   **Recent invasion and eradication of two members of the *Euwallacea fornicatus* species complex (Coleoptera:  
2   Curculionidae: Scolytinae) from tropical greenhouses in Europe**

4   Hannes Schuler<sup>1,2\*</sup>, Radosław Witkowski<sup>3</sup>, Bart van de Vossenberg<sup>4</sup>, Björn Hoppe<sup>5</sup>, Moritz Mittelbach<sup>6</sup>, Tibor  
5   Bukovinszki<sup>4</sup>, Stefan Schwembacher<sup>7</sup>, Bas van de Meulengraaf<sup>4</sup>, Uwe Lange<sup>8</sup>, Sabine Rode<sup>8</sup>, Alessandro Andriolo<sup>9</sup>,  
6   Marta Bełka<sup>3</sup>, Andrzej Mazur<sup>3</sup>, Andrea Battisti<sup>10</sup>

7

8   <sup>1</sup> Faculty of Science and Technology, Free University of Bozen-Bolzano, Italy; ORCID: 0000-0001-8307-9831

9   <sup>2</sup> Competence Centre for Plant Health, Free University of Bozen-Bolzano, Italy

10   <sup>3</sup> Faculty of Forestry and Wood Technology, Poznań University of Life Sciences, Poland

11   <sup>4</sup> Dutch National Plant Protection Organization, National Reference Centre Wageningen, Netherlands

12   <sup>5</sup> Julius Kühn Institute (JKI) - Federal Research Centre for Cultivated Plants, Institute for National and International  
13   Plant Health, Braunschweig, Germany; ORCID: 0000-0001-8689-4911

14   <sup>6</sup> Senatsverwaltung für Umwelt, Verkehr und Klimaschutz, Berlin, Germany

15   <sup>7</sup> Plant Protection Service of the Autonomous Province of Bozen-Bolzano, Italy

16   <sup>8</sup> Thüringer Landesamt für Landwirtschaft und Ländlichen Raum, Referat 23 Pflanzenschutz und Saatgut, Jena,  
17   Germany

18   <sup>9</sup> Forest Service of Bozen-Bolzano, Italy

19   <sup>10</sup> DAFNAE, University of Padova, Italy; ORCID: 0000-0002-2497-3064

20

21   \* Corresponding author: Hannes Schuler, hannes.schuler@unibz.it

22

23   **Abstract**

24   Ambrosia beetles of the *Euwallacea fornicatus* species complex are emerging tree pests with a broad host range  
25   including important agricultural crops. Native to Southeast Asia, these species were introduced into various countries,  
26   where they cause considerable damage to many tree species. Here we report several outbreaks of *E. fornicatus* s.l. in  
27   Europe. The first individuals were found in 2017 in a palm house of a botanical garden in Poznań (Poland) whereas in  
28   2020 an outbreak was detected in a tropical greenhouse in Merano (Italy). In 2021, two additional outbreaks were  
29   detected in two greenhouses in Germany, in Erfurt and Berlin. For the latter cases it was possible to trace back the  
30   invasion to a distributor of exotic plants in the Netherlands where several infected plants were detected. Molecular  
31   analysis show that individuals from Poland and Italy are genetically identical but belong to a different mitochondrial  
32   clade than individuals in Germany which are identical to most individuals of two greenhouses in the Netherlands.  
33   Moreover, in the two greenhouses in the Netherlands we found beetles that belong to another haplotype of *E. fornicatus*  
34   and two haplotypes of *E. perbrevis*, a species in the *E. fornicatus* complex, which has not been previously intercepted in  
35   Europe. Our study provides novel insights into the invasion history of *E. fornicatus* and the first eradication measures in  
36   Europe. Considering the high potential of introduction and establishment of *Euwallacea* ambrosia beetles, particular  
37   attention should be paid to monitor the presence of these pests in greenhouses across Europe.

38

39   **Keywords**

40   Ambrosia beetle, *Euwallacea fornicatus*, *Euwallacea perbrevis*, invasive species, Polyphagous shot hole borer, Tea shot  
41   hole borer

42 **Declarations**

43 **Funding:** Internal funds of the Free University of Bozen-Bolzano to H.S., Forest Service of Bozen-Bolzano, Italy and  
44 European Union's Horizon 2020 Program for Research and Innovation 'HOMED' [grant no. 771271] to A.B.

45 **Conflicts of interest:** The authors declare that they have no conflicts of interest.

46 **Availability of data and material:** Sequence data will be deposited into the Genbank. The datasets generated and/or  
47 analyzed during the current study are available from the corresponding author on reasonable request.

48 **Ethics approval:** This article does not contain any studies with human participants or animals (vertebrates) performed  
49 by any of the authors.

50 **Consent to participate:** Not applicable

51 **Consent for publication:** Not applicable

52 **Author's contribution:** AB, HS conceived and designed research. AA, AM, BvdV, BH, BvdM, HS, MB, MM, SS, SR,  
53 RW, UL, TB, conducted experiments. HS, BvdV performed data analysis. HS and AB wrote the manuscript with  
54 contribution of all authors.

55 **Acknowledgments:** The authors would like to thank the administration of the Poznań Palm House for sharing materials  
56 and giving full access to the site. Erika Corretto, Patrizia Elena Vannucchi and Wietse den Hartog for technical  
57 assistance and Zvi Mendel and Stanley Freeman for helpful comments and suggestions.

58

59 **Introduction**

60 Members of the *Euwallacea fornicatus* species complex (Coleoptera: Curculionidae: Scolytinae) are severe agricultural  
61 pests in most parts of the world (Beaver 1976; Li et al. 2016; Ge et al. 2017). They are one of a few examples of  
62 ambrosia beetles that are able to infest healthy plants (Gomez et al. 2019). The *E. fornicatus* species complex has a  
63 broad host range of 412 plant species in 75 families (Gomez et al. 2019) including important agricultural crops such as  
64 tea (*Camellia sinensis*), mango (*Mangifera indica*) and avocado (*Persea americana*) (Eskalen et al. 2013; Carrillo et al.  
65 2016); Danthanarayana 1968; Yamaguchi et al. 2006). They are associated with at least three symbiotic fungi *Fusarium*  
66 *euwallaceae*, *Graphium euwallaceae* and *Paracremonium pembeum* (syn. *Acremonium pembeum*) (Carrillo et al. 2016;  
67 Freeman et al. 2016; Lynch et al. 2016). Some of these vectored fungi are plant pathogens causing avocado dieback in  
68 California (USA) and Israel (Eskalen et al. 2012, 2013; Freeman et al. 2013).

69

70 *Euwallacea fornicatus* s.l. is native to Southeast Asia with confirmed records in China, Japan, Malaysia, the  
71 Philippines, Taiwan, Sri Lanka, Thailand, and Vietnam (CABI 2021, Stouthamer et al. 2017). Outside its native range,  
72 *E. fornicatus* s.l. invaded various countries in North and Central America (USA, Panama, Costa Rica), Africa (South  
73 Africa, Réunion), Oceania (Australia, Papua New Guinea, Fiji and other countries), and Israel (Kirkendall and  
74 Ødegaard 2007; Cooperband et al. 2016, CABI 2021). In Europe, *E. fornicatus* s.l. is considered a quarantine pest and is  
75 included in the Annex II Part A within the group "Scolytidae spp. (non-European)" (Commission Implementing  
76 Regulation EU 2019/2072). Its explicit listing in Annex II is anticipated by the end of 2021. Its wide host range makes  
77 *E. fornicatus* s.l. a serious threat for agriculture but also forestry, ornamental plants, and botanical gardens.

78

79 Several outbreaks were recently discovered in tropical greenhouses of botanical gardens in Europe. In 2017, specimens  
80 of *E. fornicatus* were found on a sacred fig (*Ficus religiosa*) tree in a palm house in Poznań, Poland (EPPO 2019),  
81 whereas various tropical plants have been attacked by *E. fornicatus* in a greenhouse in Merano, Italy (EPPO 2020).  
82 Recently, two other cases have been reported in Germany in two tropical greenhouses in Thuringia (EPPO 2021a) and

83 Berlin (EPPO 2021b) and in a greenhouse in the Netherlands (EPPO 2021c). Although tracing information of  
84 consignments has been used to establish the origin of infested plants, analysis of molecular data may provide direct  
85 insights into the invasion history of *Euwallacea* and thus into the dynamics of introduction events across the different  
86 locations. Here we perform a comparative genetic analysis of insects from different locations in Europe and attempt to  
87 trace back the introduction pathway of the beetles. Moreover, we report the subsequent eradication measures. Our study  
88 provides new insights into the introduction and the subsequent steps to eradicate these invasive insect pest species.  
89

## 90 Materials and methods

### 91 Detection and identification

92 A sacred fig tree (*Ficus religiosa*) in the Poznań Palm House (Poland) showed branch dieback and visible frass, typical  
93 symptoms for infestation with ambrosia beetles in March 2017. An approximately 1 m long branch of the tree was cut  
94 and placed in a breeding container. Emerging beetles were collected for taxonomic identification. In April 2020, several  
95 trees in the tropical greenhouse of the Gardens of Trauttmansdorff Castle in Merano (Italy), showed boreholes and  
96 ejection of wooden debris. Emerging insects were directly collected from infested plants, from the walls of the  
97 greenhouse and dissected from trap logs deployed in the greenhouse. In addition, two plants, one *Mangifera indica* and  
98 one *Tectona grandis*, showed symptoms of a bark beetle infection in a tropical greenhouse in Erfurt (Germany) in  
99 January 2021 and two months later several plants in a greenhouse in Berlin (Germany) showed resin flows on stems and  
100 boreholes with small protruding tubes of compacted sawdust. Emerging beetles were collected and stored in ethanol. In  
101 the Netherlands, several plants were found to be infested at two plant nurseries (NL1 and NL2) and where beetles were  
102 extracted.

103

104 Identification of the insect material was carried out using taxonomic keys available in the literature (Smith et al. 2020;  
105 Wood and Bright 1992; Rabaglia et al. 2006; Wood 2007; Smith et al. 2019) by the Poznań University of Life Sciences  
106 in Poland, by the Laboratory for Virology and Diagnostics of the Research Centre Laimburg in Italy and by the federal  
107 plant protection agencies in Thuringia and Berlin and the National reference laboratory of the Julius-Kühn Institute  
108 (JKI) in Braunschweig in Germany and by the National Reference Centre of the NPPO in the Netherlands.

109

### 110 DNA extraction and sequencing

111 A total of 32 individuals were analyzed including three individuals from Poland, 9 individuals from Italy, three  
112 individuals from Erfurt and 17 from Berlin. Individuals were genotyped using the primers Lep-R1 (Hebert et al. 2004).  
113 Moreover, in the Netherlands, in a total of 32 beetle specimens from the Dutch outbreak locations (18 beetle specimens  
114 from NL1 and 14 beetle specimens from NL2) were included in the analysis. Details about the extraction protocol, PCR  
115 conditions and sequencing can be found in supplementary document S1.

116

### 117 Cluster analysis

118 Relationship of the obtained sequences was determined using 83 partial mitochondrial COI sequences obtained from  
119 NCBI GenBank and the 64 partial COI sequences generated in this study. Sequences were aligned using the MAFFT  
120 aligner (Katoh et al. 2002; Katoh and Standley 2013) incorporated in Geneious Prime v2021.1.1 (BioMatters, New  
121 Zealand). Terminal positions in the alignment that were not covered by all sequences in the dataset were masked  
122 resulting in a 564 bp alignment used for clustering analysis. A maximum likelihood tree was constructed with FastTree  
123 (Price et al. 2010) using the generalized time-reversible (GTR) model and 1,000 bootstraps to determine confidence

124 levels of internal nodes. The *cox1* sequence of *Euwallacea andamanensis* isolate PR13-238 (KU727039) was used to  
125 root the *E. fornicatus* s.l. sequences.

126

127 *Eradication and surveillance*

128 In Poland, immediately after the outbreak was noticed, actions limiting its spread were undertaken. The whole sacred  
129 fig tree was covered with Storanet® BASF Agrar (insect-proof net impregnated with alpha-cypermethrin) and the tree  
130 was subsequently removed and burnt under controlled conditions. The soil around the roots was excavated and the  
131 surrounding area was treated with the fungicide Topsin M (Thiophanate-Methyl, 100 ppm concentration). Beetle  
132 surveillance with 4 triangle barrier traps IBL-2 (baited with 98% ethanol) started immediately and continued for a year  
133 after removal of the infested tree. Furthermore, plant health conditions in the palm house were monitored by visual  
134 observation of symptoms.

135

136 After the first detection of *E. fornicatus* in Italy, an intense monitoring has started inside the greenhouse and in the  
137 surrounding area. Due to the advanced outbreak, all the plants in the greenhouse, including their roots, were removed  
138 and destroyed under official control in June 2021. The greenhouse was also subjected to solarization for a period of  
139 three months. Simultaneously, three sticky traps baited with quercivorol and alpha-copaene according to (Kendra et al.  
140 2019) and several trap logs of *Acer negundo* (diameter 4-10 cm, length 30-60 cm) were placed inside the greenhouse to  
141 verify the possible presence of adults. In the area external to the greenhouse, two traps were deployed at 500 and 1000  
142 m from the tropical greenhouse in each cardinal direction, for a total of eight traps. All traps were weekly checked.  
143 Visual inspections from the ground, ladders and hoisting platforms were implemented in- and outside of the infested  
144 greenhouse. Known host plants of the beetle (*Acer* spp., *Citrus* spp., *Platanus* spp., etc.) were monitored for boreholes  
145 especially after favorable weather conditions. Additionally, weakening tree parts or branches were inspected for  
146 symptoms.

147

148 After the introduction in Germany, monitoring inside and outside the respective greenhouse have been initiated, using  
149 the setup described above, consisting of trap logs containing quercivorol pheromones. In the plant nurseries of  
150 Netherlands, ethanol and acetic acid-baited traps were placed in all greenhouse compartments for one week (18 traps at  
151 NL1, 6 traps at NL2). Later sticky traps baited with quercivorol and alpha-copaene were placed in the greenhouse  
152 compartments at eight positions in both greenhouses.

153

154 **Results**

155 *Detection and eradication*

156 In Poland, the sacred fig sample yielded over 1,000 specimens of *E. fornicatus*. No beetles were caught by the traps for  
157 one year after the first detection. Plants, occurring in the same pavilion as the infested sacred fig and plants in  
158 neighboring pavilions, which were subjected to regular inspections did not show symptoms of ambrosia beetles and  
159 fungal infection.

160

161 In Italy, a total of 28 trees of 21 different species (*Annona muricata*, *Artocarpus heterophyllus*, *Averrhoa carambola*,  
162 *Bixa orellana*, *Bulnesia arborea*, *Cananga odorata*, *Clausena lansium*, *Crescentia cujete*, *Debregeasia edulis*,  
163 *Dimocarpus longan*, *Ficus altissima*, *Ficus* sp., *Justicia* sp., *Kigelia africana*, *Melicoccus bijugatus*, *Mangolia*  
164 *champaca*, *Millettia brandisiana*, *Persea americana*, *Terminalia catappa*, *Terminalia buceras*, *Theobroma cacao*)

165 showed boreholes and ejection of wooden debris. Intensity of infestation, diameter of infested trees and distribution of  
166 the boreholes on the plant differ strongly between species and individual plants. *Annona muricata* and *Bixa orellana*  
167 were the most heavily infested plants, showing more than five holes per dm<sup>2</sup>. The highest density of boreholes was  
168 observed next to fresh and older cut branches and on thinner parts in the crown/upper stem. The presence of boreholes  
169 was also observed in twigs less than 2 cm of diameter of *Dimocarpus longan* and *Justicia spp.* trees. No beetles were  
170 caught in the traps deployed outside of the tropical greenhouse and visual inspection of susceptible host trees did not  
171 result in finding of symptoms of beetle attack.

172

173 The two recent findings in Germany revealed that in Erfurt just two plants, one *Mangifera indica* and one *Tectona*  
174 *grandis*, were attacked by *E. fornicatus*, whereas in Berlin 136 shrubs and trees of *Clusia rosea*, *Heteropanax sp.*, *Ficus*  
175 *sp.*, and *Mangifera indica* showed symptoms. All infested plants were destroyed and surveillance started in both  
176 localities.

177

178 In the Netherlands, at both locations all infested plants were destroyed. The sale and transport of all woody plants and  
179 palms (Arecaceae) in the two greenhouses in the Netherlands was put on hold followed by intensive visual inspection of  
180 all consignments by the phytosanitary authorities.

181

#### 182 *Reconstruction of the invasion*

183 Trace-back investigations revealed that the *F. religiosa* tree infected with *E. fornicatus* in Poland was imported in  
184 November 2016 from the Netherlands. The infection has already been detected five months later and subsequent  
185 phytosanitary measures impeded the dispersion to other trees. The advanced outbreak in Italy with 21 different species  
186 attacked at the time of detection, hindered a detailed reconstruction of the invasion. The greenhouse in Italy was  
187 established in 2014 and most plants were purchased between 2013 and 2014. No symptoms of beetle attack were  
188 reported in the first years following the establishment. Additionally, one *T. cacao* plant was replaced in 2018, which  
189 was the same that two years later was discovered to be attacked by *E. fornicatus*. Both infested plants in Erfurt, and  
190 most of the infested plants in Berlin were acquired and imported from a distributor of exotic plants in the Netherlands in  
191 2020.

192

193 Following a trace-back of the notification in the Erfurt case, we were able to link these infestations to a Dutch  
194 greenhouse for commercial sale of tropical plants (NL1). A follow up survey resulted in several trees that were attacked  
195 by ambrosia beetles. Samples were collected from a *Ficus microcarpa* individual that contained specimens of the  
196 *Euwallacea fornicatus s.l.* in March 2021. Additionally, beetles were collected from *Bauhinia x blakeana*, *Ficus*  
197 *microcarpa* 'Panda', *Ficus sp.*, *Ficus maclellandii* 'Alii' and *Ficus* 'Amstel King' whereas one individual was collected  
198 from a trap. A following survey in another greenhouse associated with the company (NL2) resulted in the discovery of  
199 several additional symptomatic trees. At location NL2, individuals were collected from *F. benjamina* 'Exotica', *F.*  
200 *foliolae* and *Ficus lyrata*. A total of 32 individuals from the two populations in the Netherlands (18 from NL1, 14 from  
201 NL2) were subsequently genotyped and included in our cluster analysis.

202

#### 203 *Molecular identification*

204 All individuals from Poland and Italy were genetically identical (Genbank acc. Number TBA) but different from the  
205 individuals found in Germany and the Netherlands. The haplotype described in Poland and Italy clustered with a

206 haplotype of the PSHB clade which according to Smith et al. (2019) is *E. fornicate*s. However, none of the haplotypes  
207 described in *E. fornicate*s populations in their native and invasive range were identical to the haplotype described in  
208 these two localities. The cluster analysis showed that the prevalent COI haplotype in Europe was most related to a  
209 haplotype present in Vietnam (Stouthamer et al. 2017; Fig. 1). In contrast all individuals from Germany and the  
210 Netherlands share the same haplotype (Genbank acc. Number TBA) which is related to a population previously  
211 genotyped in Taiwan (Stouthamer et al. 2017). Moreover, two individuals from the Netherlands (NL2) belonged to a  
212 PSHB haplotype that did not match any of the haplotypes uncovered at the other locations.(Stouthamer et al. 2017)  
213

214 Sequence analysis of 32 individuals from the Netherlands revealed four haplotypes. Thirteen individuals of location  
215 NL1 clustered with *E. fornicate*s in the PSHB clade 3A (Stouthamer et al. 2017) whereas five individuals belonged to  
216 two haplotypes of the TSHBa clade (Genbank acc. Number TBA) which according to Smith et al. (2019) is now  
217 considered *Euwallacea perbrevis*. All sequenced individuals from NL2 belonged to two mitochondrial haplotypes that  
218 clustered with *E. fornicate*s in the PSHB clade 3A (Fig. 1). The CO1 sequences of 25 of the PSHB clade 3A individuals  
219 from both locations in the Netherlands were identical to those from the Erfurt and Berlin interceptions, but not to those  
220 from Italy and Poland interceptions (Fig. 1).

## 221     **Discussion**

222 Botanical gardens and greenhouses are hotspots for invasive species (Wang et al. 2015). The deliberate import of alien  
223 plants might lead to the invasion of non-native plants (Hulme 2011) but might also lead to the introduction of associated  
224 insects (Scott-Brown et al. 2018). Although introduced insects might suffer from the climatic conditions in the new  
225 environment, greenhouses play an important role in helping them to adapt to the new environment, eventually aiding  
226 the establishment and invasion of non-native insect species (Wang et al. 2015). Here we show an additional example of  
227 a recent introduction of two non-native insect pests in botanical gardens of Europe.  
228

229 Ambrosia beetles of the *Euwallacea fornicate*s species complex are invasive species introduced into various continents  
230 but had not been detected in Europe. Here we describe the recent outbreaks of *E. fornicate*s in four tropical greenhouses  
231 in Poland, Italy, and Germany. We reconstructed the invasion history and its invasion route. Molecular characterization  
232 of 12 individuals from Poland and Italy showed that all individuals share the same mitochondrial haplotype, suggesting  
233 that they were likely introduced from the same source population. The haplotype belongs to the Polyphagous shot hole  
234 borer clade (Stouthamer et al. 2017) and most related to a haplotype found in Vietnam (Stouthamer et al. 2017; Fig. 1).  
235 Since the haplotype was not described elsewhere, we could not determine the exact source population. In contrast, the  
236 populations in Germany and the Netherlands were related to a different haplotype which has been previously described  
237 in Taiwan (Stouthamer et al. 2017). This highlights that the cases from Germany likely resulted from an introduction  
238 via one of the greenhouses of the commercial nursery in the Netherlands whereas the cases from Poland and Italy  
239 resulted from an independent introduction event, likely from the same source population. The additional detection of *E.*  
240 *perbrevis* in a greenhouse in the Netherlands highlights a third introduction event of a species which has not been found  
241 elsewhere in Europe.  
242

243 The outbreaks in the different localities were in different epidemiological phases: *E. fornicate*s in Poland was present in  
244 a single sacred fig tree (*Ficus religiosa*) and not found in any other tree species in the same greenhouse, in Italy beetles  
245 have been detected already in 28 different plants belonging to 21 species. It seems likely that the *E. fornicate*s was

247 introduced to Poland with an infested *F. religiosa* tree in 2016 and eradicated a few months later in the initial phase of  
248 its establishment. In contrast, the outbreak in Italy was more advanced when it was discovered in 2020. The most  
249 plausible explanation is that the ambrosia beetle was introduced with the *T. cacao* plant purchased in 2018 and  
250 subsequently attacked other plant species in the greenhouse where the outbreak was detected two years later. In Erfurt  
251 both infected trees of *Mangifera indica* and *Tectona grandis*, as well as most of the 136 infected trees in Berlin were  
252 imported from the Netherlands. In both cases, the presence of the beetle was detected a few months after their  
253 introduction.

254  
255 Greenhouses might act as a springboard for non-native species if they are able to adapt and disperse to novel  
256 environments (Wang et al. 2015). Although *E. fornicatus* might not be able to survive the outdoor conditions prevailing  
257 in Poland, Germany and the Netherlands, Mediterranean conditions in Italy might allow an establishment of the beetles  
258 in this area. Moreover, all affected greenhouses are surrounded by several native and non-native trees and shrubs,  
259 known as potential hosts for *E. fornicatus*.

260  
261 The occurrence of *E. fornicatus* in the greenhouse of a retailer in the Netherlands and especially the description of *E.*  
262 *perbrevis*, which has not been described elsewhere in Europe highlights the need of more efficient examinations of  
263 imported exotic plants prior to re-sale. While eradication may be relatively simple in a greenhouse environment, it  
264 becomes much more problematic once a population is established outdoors. Therefore, we argue that surveillance  
265 should be also intensified in tropical greenhouses in order to reduce the likelihood of an establishment in nature.

## 266 Figures

267 **Fig. 1** Phylogenetic tree based on partial (564 bp) mitochondrial *cox1* gene sequences representing the relation of the  
268 individuals found in Europe (in bold and italics) with haplotypes described in other studies.

## 269 References

- 270  
271  
272 Beaver RA (1976) The biology of Samoan bark and ambrosia beetles (Coleoptera, Scolytidae and Platypodidae). B  
273 Entomol Res 65:531–548. <https://doi.org/10.1017/s0007485300006210>
- 274 CABI (2021) *Euwallacea fornicatus* distribution map <https://www.cabi.org/isc/datasheet/18360453> [accessed on 15  
275 April 2021].
- 276 Carrillo D, Cruz LF, Kendra PE, et al (2016) Distribution, pest status and fungal associates of *Euwallacea nr. fornicatus*  
277 in Florida Avocado Groves. Insects 7:55. <https://doi.org/10.3390/insects7040055>
- 278 Cooperband MF, Stouthamer R, Carrillo D, et al (2016) Biology of two members of the *Euwallacea fornicatus* species  
279 complex (Coleoptera: Curculionidae: Scolytinae), recently invasive in the U.S.A., reared on an ambrosia beetle  
280 artificial diet. Agric For Entomol 18:223–237. <https://doi.org/10.1111/afe.12155>
- 281 Danthanarayana W (1968) The distribution and host-range of the shot-hole borer (*Xyleborus fornicatus* Eichh.) of tea.  
282 Tea Q. 39: 61-69.
- 283 EPPO (2019) First finding of *Euwallacea fornicatus* in Poland, EPPO Reporting Service no. 02 – 2019
- 284 EPPO (2020) First report of *Euwallacea fornicatus* in Italy, EPPO Reporting Service no. 05 - 2020
- 285 EPPO (2021a) First report of *Euwallacea fornicatus* in Germany, EPPO Reporting Service no. 02 - 2021

- 286 EPPO (2021b) New finding of *Euwallacea fornicatus* in Germany, EPPO Reporting Service no. 03 - 2021
- 287 EPPO (2021c) First report of *Euwallacea fornicatus* sensu lato and cf. *Cryphalus* sp. in the Netherlands, EPPO  
288 Reporting Service no. 04 - 2021
- 289 Eskalen A, Gonzalez A, Wang DH, et al (2012) First Report of a *Fusarium* sp. and its vector Tea shot hole borer  
290 (*Euwallacea fornicatus*) causing fusarium dieback on avocado in California. Plant Dis 96:1070–1070.  
291 <https://doi.org/10.1094/pdis-03-12-0276-pdn>
- 292 Eskalen A, Stouthamer R, Lynch SC, et al (2013) Host range of fusarium dieback and its ambrosia beetle (Coleoptera:  
293 Scolytinae) vector in Southern California. Plant Dis 97:938–951. <https://doi.org/10.1094/pdis-11-12-1026-re>
- 294 Freeman S, Sharon M, Dori-Bachash M, et al (2016) Symbiotic association of three fungal species throughout the life  
295 cycle of the ambrosia beetle *Euwallacea nr. fornicatus*. Symbiosis 68:115–128. <https://doi.org/10.1007/s13199-015-0356-9>
- 297 Freeman S, Sharon M, Maymon M, et al (2013) *Fusarium euwallaceae* sp. nov.—a symbiotic fungus of *Euwallacea*  
298 sp., an invasive ambrosia beetle in Israel and California. Mycologia 105:1595–1606. <https://doi.org/10.3852/13-066>
- 299 Ge X, Jiang C, Chen L, et al (2017) Predicting the potential distribution in China of *Euwallacea fornicatus* (Eichhoff)  
300 under current and future climate conditions. Sci Rep 7:906. <https://doi.org/10.1038/s41598-017-01014-w>
- 301 Gomez DF, Lin W, Gao L, Li Y (2019) New host plant records for the *Euwallacea fornicatus* (Eichhoff) species  
302 complex (Coleoptera: Curculionidae: Scolytinae) across its natural and introduced distribution. Journal of Asia-  
303 Pacific Entomology 22:338–340. <https://doi.org/10.1016/j.aspen.2019.01.013>
- 304 Hebert PDN, Penton EH, Burns JM, et al (2004) Ten species in one: DNA barcoding reveals cryptic species in the  
305 neotropical skipper butterfly *Astraptes fulgerator*. P Natl Acad Sci Usa 101:14812–14817.  
306 <https://doi.org/10.1073/pnas.0406166101>
- 307 Hulme PE (2011) Addressing the threat to biodiversity from botanic gardens. Trends Ecol Evol 26:168–174.  
308 <https://doi.org/10.1016/j.tree.2011.01.005>
- 309 Katoh K, Misawa K, Kuma K, Miyata T (2002) MAFFT: a novel method for rapid multiple sequence alignment based  
310 on fast Fourier transform. Nucleic Acids Res 30:3059–3066. <https://doi.org/10.1093/nar/gkf436>
- 311 Katoh K, Standley DM (2013) MAFFT Multiple Sequence Alignment Software Version 7: Improvements in  
312 performance and usability. Mol Biol Evol 30:772–780. <https://doi.org/10.1093/molbev/mst010>
- 313 Kendra PE, Montgomery WS, Narvaez TI, Carrillo D (2019) Comparison of trap designs for detection of *Euwallacea*  
314 *nr. fornicatus* and other Scolytinae (Coleoptera: Curculionidae) that vector fungal pathogens of avocado trees in  
315 Florida. J Econ Entomol 113:980–987. <https://doi.org/10.1093/jee/toz311>
- 316 Kirkendall LR, Ødegaard F (2007) Ongoing invasions of old-growth tropical forests: establishment of three incestuous  
317 beetle species in southern Central America (Curculionidae: Scolytinae). Zootaxa 1588:53–62.  
318 <https://doi.org/10.11646/zootaxa.1588.1.3>
- 319 Li Y, Gu X, Kasson MT, et al (2016) Distribution, host records, and symbiotic fungi of *Euwallacea fornicatus*  
320 (Coleoptera: Curculionidae: Scolytinae) in China. Fla Entomol 99:801–804. <https://doi.org/10.1653/024.099.0441>
- 321 Lynch SC, Twizeyimana M, Mayorquin JS, et al (2016) Identification, pathogenicity and abundance of *Paracremonium*  
322 *pembeum* sp. nov. and *Graphium euwallaceae* sp. nov.—two newly discovered mycangial associates of the  
323 polyphagous shot hole borer (*Euwallacea* sp.) in California. Mycologia 108:313–329. <https://doi.org/10.3852/15-063>
- 325 Price MN, Dehal PS, Arkin AP (2010) FastTree 2 – Approximately maximum-likelihood trees for large alignments.  
326 Plos One 5:e9490. <https://doi.org/10.1371/journal.pone.0009490>

- 327 Rabaglia RJ, Dole SA, Cognato AI (2006) Review of American Xyleborina (Coleoptera: Curculionidae: Scolytinae)  
328 occurring North of Mexico, with an illustrated key. Ann Entomol Soc Am 99:1034–1056.  
329 [https://doi.org/10.1603/0013-8746\(2006\)99](https://doi.org/10.1603/0013-8746(2006)99)
- 330 Scott-Brown AS, Hodgetts J, Hall J, et al (2018) Potential role of botanic garden collections in predicting hosts at risk  
331 globally from invasive pests: a case study using *Scirtothrips dorsalis*. J Pest Sci 91:601–611.  
332 <https://doi.org/10.1007/s10340-017-0916-2>
- 333 Smith SM, Gomez DF, Beaver RA, et al (2019) Reassessment of the species in the *Euwallacea fornicatus* (Coleoptera:  
334 Curculionidae: Scolytinae) complex after the rediscovery of the “lost” type specimen. Insects 10:261.  
335 <https://doi.org/10.3390/insects10090261>
- 336 Stouthamer R, Rugman-Jones P, Thu PQ, et al (2017) Tracing the origin of a cryptic invader: phylogeography of the  
337 *Euwallacea fornicatus* (Coleoptera: Curculionidae: Scolytinae) species complex. Agric For Entomol 19:366–375.  
338 <https://doi.org/10.1111/afe.12215>
- 339 Wang C, Zhang X, Pan X, et al (2015) Greenhouses: hotspots in the invasive network for alien species. Biodivers  
340 Conserv 24:1825–1829. <https://doi.org/10.1007/s10531-015-0876-x>
- 341 Wood SL, Bright DE (1992) A catalog of Scolytidae and Platypodidae (Coleoptera), Part 2: Taxonomic index volume  
342 A. Great basin naturalist memoirs, No 13, A catalog of Scolytidae and Platypodidae (Coleoptera), Part 2:  
343 Taxonomic Index Volume A
- 344 Wood S. L. 2007., Bark and ambrosia beetles of South America (Coleoptera: Scolytidae) (Monte L. Bean Life Science  
345 Museum, Brigham Young University, Provo, 2007), 900 p.
- 346 Yamaguchi T, Iwamoto J, Goto H, Nojima H, Omatsu N, Torigoe H, Yasuda K, Setokuchi O, Hayashikawa S (2006)  
347 Insect pests of the mango plant, *Mangifera indica*, on the Amami islands, Japan. Kyushu Plant Protection Research  
348 52: 60-65.

# Figures




Figure 1

Phylogenetic tree based on partial (564 bp) mitochondrial cox1 gene sequences representing the relation of the individuals found in Europe (in bold and italics) with haplotypes described in other studies.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Suppldocument1.pdf](#)