Invasive liver abscess caused by hypervirulent *Klebsiella pneumoniae*: case report and literature review

Xiao-Guang Xiao

First Affiliated Hospital of Dalian Medical University

Zhao-Hong Mu

Dalian City Second People's Hospital

Qing-Dong Li

First Affiliated Hospital of Dalian Medical University

Yong-Li Zhang (✉ zylicu@163.com)

First Affiliated Hospital of Dalian Medical University <https://orcid.org/0000-0002-4263-8382>

Case Report

Keywords: *Klebsiella pneumoniae*, endogenous endophthalmitis, *Klebsiella pneumoniae* invasive liver abscess syndrome

Posted Date: August 28th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-64114/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: *Klebsiella pneumoniae* is a common pathogenic bacteria that cause infectious diseases, *Klebsiella pneumoniae* invasive liver abscess syndrome is a rare disease, which often caused by hypervirulent *Klebsiella pneumoniae*(hvKP), the hvKP patients with sepsis may have life-threatening multiple organ dysfunction and visual sequelae.

Case presentation: We report a case of *Klebsiella pneumoniae* invasive liver abscess syndrome caused by hvKP, the patient was treated with long-term antibiotic treatment, along with controlling blood sugar with Insulin, and the patient was eventually discharged from the hospital. Case analysis and literature review were performed to improve the understanding of invasive liver abscess caused by hvKP.

Conclusions Patients with hvKP invasive liver abscess are in relatively severe conditions and often have poor visual prognosis. Appropriate antimicrobial drugs should be administered to prevent complications and improve prognosis.

Background

Klebsiella pneumoniae is a well-known Gram-negative pathogen that causing various clinical infection patterns, such as pneumonia, bacteremia, urinary tract infection, endophthalmitis, as well as liver abscess and meningitis. *Klebsiella pneumoniae* has emerged as the predominant cause of pyogenic liver abscess in Asia, and also as a primary pathogen of pyogenic liver abscess in China^[1]. *Klebsiella pneumoniae* is divided into classic *Klebsiella pneumoniae* and hypervirulent *Klebsiella pneumoniae* (hvKP) according to the differences of characteristics in virulence.

The Hypervirulen phenotype of *Klebsiella pneumoniae* is associated with the development of a distinctive invasive syndrome^[2], Hypervirulen strains of *Klebsiella pneumoniae* often show hypermucoviscosity, are determined as hvKP, hvKP is the main pathogen of liver abscess^[3]. Multiple abscesses caused by hvKP bacteremia is also called *Klebsiella pneumoniae* invasive liver abscess syndrome, it is a rare clinical condition characterized by primary liver abscess associated with other organ abscesses, including eyes, lungs and central nervous system^[4], and it is more frequently in patients with hypertension and diabetes mellitus^[5]. The syndrome has been mentioned in many publications for many years^[6, 7], it occurs worldwide, and poor prognosis is common.

We herein report a case of *Klebsiella pneumoniae* invasive liver abscess syndrome caused by hvKP. We discuss clinical characteristics and made literature review.

Case Presentation

A 78-year-old woman presented to the intensive care unit(ICU) with a 6-day history of fever and a 1-day history of unconsciousness. His medical history included type 2 diabetes mellitus and hypertension. On physical examination, the patient was somnolence, and her right eye was swollen with purulent

secretions (Fig. 1a). Laboratory studies revealed a white blood cell count of 25,110 per cubic millimetre (reference range 4,000 to 10,000). Findings on computed tomography of eye, abdomen, and lung included the presence of uneven vitreous density of the right eye and exudation around the right eyeball (Fig. 1b), low-density areas in the right lobe of the liver, maximal diameter of 4.65 cm (Fig. 2a,2b), a small cavity and ground-glass opacities in the lung upper right lobe, and solitary nodule shadow in the lung lower right lobe close to the pleura (Fig. 3 [arrow]). Contrast-enhanced magnetic resonance imaging of the brain revealed ring-enhancing lesion (0.5 cm in diameter) located in the left forehead (Fig. 4 [arrow]). Based on the clinical data, we diagnosed her with invasive liver abscess syndrome accompanied by pneumoniae, endogenous endophthalmitis and brain abscess. Empiric treatment with intravenous Meropenem was started in the ICU. We chose not to perform percutaneous or surgical drainage of the liver abscess, as the abscess was not liquefied and systemic response to antibiotic therapy was favourable.

Blood culture grew *Klebsiella pneumoniae*, and the string test was performed, the test was positive, and a diagnosis of hvKP bloodstream infection was made, and *Klebsiella pneumoniae* invasive liver abscess syndrome was also diagnosed, examination showed that the hvKP sensitive to most antibiotics, including Meropenem, the patient was continually treated with Meropenem, along with controlling blood sugar with Insulin. Besides the right eyeball, abnormal images disappeared after the patient was treated with a 30-day course of antibiotic. The patient survived but had right eye visual sequelae. At three months follow-up after discharge from the hospital, the patient was doing well except for the loss of vision in the right eye.

Discussion

Klebsiella pneumoniae is one of the common pathogens in nosocomial and community-acquired infections, and it is the primary pathogen of liver abscess. *Klebsiella pneumoniae* invasive liver abscess syndrome caused by hvKP is a devastating infection that is uncommon in western countries, while increasingly prevalent in Asian countries^[8, 9]. The most striking aspect of hvKP strains is the ability to cause severe infections in otherwise healthy patients^[10], hvKP typically presents as a community-acquired infection causing a liver abscess. Strains of hvKP often associated with *rmpA* and *magA* genes and confer a mucoid phenotype. The string test is commonly used to determine the hypermucoviscous phenotype, and we will define the string test positive when a bacteriologic inoculation loop can generate a viscous string > 5 mm in length by stretching bacterial colonies on an agar plate^[11].

The distinctive feature of community-acquired hvKP is good susceptibility to most available antibiotics, but some studies reported that healthcare-associated hvKP showed different results of antimicrobial susceptibility^[11, 12]. The emergence of hypervirulent strains resistant to antimicrobial drugs will bring more challenges in clinical diagnosis and treatment. A rapid microbiological test to confirm the pathogen as early as possible is crucial for treatment. The case we reported was characterized by community-acquired hvKP infection, and the hvKP was sensitive to most antibiotics, the infection was quickly controlled after using Meropenem. However, some patients require a prolonged course of antimicrobial

treatment which has evolved over the months from inpatient treatment to outpatient parenteral antibiotic therapy.

Klebsiella pneumoniae invasive liver abscess syndrome affects multiple organs, in addition to liver abscesses, hvKP metastasize to distant sites, including lungs, brain, eyes, kidneys, spleen, bone marrow^[13, 14]. The symptoms of *Klebsiella pneumoniae* invasive liver abscess syndrome are nonspecific and can include fever, chill, abdominal pain, nausea, vomiting, dizziness and vision loss, but may also be driven by the location of the metastatic infection. Of these, sight-threatening endophthalmitis has been commonly reported, and *Klebsiella pneumoniae* endogenous endophthalmitis typically has poor visual prognosis, it may cause irreversible visual loss if not treated properly and on time^[15, 16, 17]. Early diagnosis and prompt treatment may salvage useful vision in some eyes.

The patients of *Klebsiella pneumoniae* invasive liver abscess syndrome need to be admitted to the ICU for monitoring and treatment, septic shock and multiple organ failure were risk factors for mortality^[18]. Therefore, high awareness, early and aggressive treatment of patients diagnosed with *Klebsiella pneumoniae* invasive liver abscess syndrome is critical in saving the lives of patients. It is also essential to guide clinical management from laboratory testing to optimize patient outcomes.

Abbreviations

hvKP : hypervirulent *Klebsiella pneumoniae*; ICU : intensive care unit

Declarations

Acknowledgements

Not applicable.

Ethical approval and Consent to participate

Not applicable.

Consent for the publication

Informed consent for the publication of all images was obtained from the patient.

Availability of supporting data

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Funding source

This work was supported by the Natural Science Foundation of Liaoning Province (20180551029).

Authors' contributions

XGX and ZHM contributed equally to this work and should be considered co-first authors. XGX, QDL and YLZ participated in the writing of the article, and YLZ edited the final draft. ZHM provided the clinical data and participated in the writing of the article. All authors had read and approved the final manuscript.

Authors' information

¹Xiao-Guang Xiao: Clinical Laboratory, The First Affiliated Hospital of Dalian Medical University, Zhongshan Road 222, DaLian, 116011, China,

²Zhao-Hong Mu: Department of Critical Care Medicine, Dalian Second People's Hospital, Hongji Street29, Xigang District, Dalian, 116014, China

³Qing-Dong Li: Department of Critical Care Medicine, The First Affiliated Hospital of Dalian Medical University, Zhongshan Road 222, DaLian, 116011, China

^{3*} Yong-Li Zhang: Department of Critical Care Medicine, The First Affiliated Hospital of Dalian Medical University, Zhongshan Road 222, DaLian, 116011, China

* corresponding author: Yong-Li Zhang Email: zylicu@163.com Phone: +86 18098875877

* Correspondence: zylicu@163.com Department of Critical Care Medicine, The First Affiliated Hospital of Dalian Medical University, Zhongshan Road 222, DaLian, 116011, China

References

1. Qian Y, Wong CC, Lai S, Chen H, He X, Sun L, et al. A retrospective study of pyogenic liver abscess focusing on *Klebsiella pneumoniae* as a primary pathogen in China from 1994 to 2015. *Scientific Reports*. 2016;6:38587.
2. Lee HC, Chuang YC, Yu WL, Ko WC. Clinical implications of hypermucoviscosity phenotype in *Klebsiella pneumoniae* isolates: association with invasive syndrome in patients with community-acquired bacteraemia. *J Intern Med*. 2006;59(6):606-14.
3. Zhang YW, Zhao CJ, Wang Q, Wang XJ, Chen HB, Li H, Zhang FF, Li SG, Wang RB, Wang H. High prevalence of hypervirulent *Klebsiella pneumoniae* infection in China: geographic distribution, clinical characteristics, and antimicrobial resistance. *Antimicrobial Agents and Chemotherapy*. 2016;60(10): 6115-20.
4. Van Keer J, Van Keer K, Van Calster J, Derdelinckx I. More Than Meets the Eye: *Klebsiella pneumoniae* Invasive Liver Abscess Syndrome Presenting with Endophthalmitis. *J Emerg Med*.

- 2017;52(6): e221-3.
5. Guo Y, Wang S, Zhan L, Yu F. Microbiological and Clinical Characteristics of Hypermucoviscous *Klebsiella pneumoniae* Isolates Associated with Invasive Infections in China. *Front Cell Infect Microbiol.* 2017;7:24.
 6. Wang B, Zhang P, Li Y, Wang Y. *Klebsiella pneumoniae*-induced multiple invasive abscesses: A case report and literature review. *Medicine (Baltimore).* 2019;98(39): e17362.
 7. Bradley ME, Scoular SK. Metastatic *Klebsiella pneumoniae* Invasive Liver Abscess Syndrome in Denver, Colorado. *J Pharm Pract.* 2019;10:1-5.
 8. Struve C, Roe CC, Stegger M, Krogfelt KA. Mapping the Evolution of Hypervirulent *Klebsiella pneumoniae*. *MBio.* 2015;6(4): e00630.
 9. Siu LK, Yeh KM, Lin JC, Fung CP, Chang FY. *Klebsiella pneumoniae* liver abscess: a new invasive syndrome. *The Lancet Infectious Diseases.* 2012;12(11):881-7.
 10. Martin RM, Bachman MA. Colonization, Infection, and the Accessory Genome of *Klebsiella pneumoniae*. *Front Cell Infect Microbiol.* 2018;8:4.
 11. Hyun M, Lee JY, Ryu SY, Kim HA. Antibiotic Resistance and Clinical Presentation of Health Care-Associated Hypervirulent *Klebsiella pneumoniae* Infection in Korea. *Microb Drug Resist.* 2019;25(8):1204-9.
 12. Chew KL, Lin RTP, Teo JWP. *Klebsiella pneumoniae* in Singapore: Hypervirulent Infections and the Carbapenemase Threat. *Front Cell Infect Microbiol.* 2017;7:515.
 13. Sturm E, Tai A, Lin B, Pollard J. Bilateral osteomyelitis and liver abscess caused by hypervirulent *Klebsiella pneumoniae*-a rare clinical manifestation (case report). *BMC Infect Dis.* 2018;18(1):380.
 14. Setsu T, Tsuchiya A, Yamagiwa S, Terai S. Invasive Liver Abscess Syndrome Caused by *Klebsiella pneumoniae*. *Intern Med.* 2017;56(22):3121-2.
 15. Chen YH, Li YH, Lin YJ, Chen YP, Wang NK, Chao AN, et al. Prognostic Factors and Visual Outcomes of Pyogenic Liver Abscess-Related Endogenous *Klebsiella pneumoniae* Endophthalmitis: A 20-year retrospective review. *Scientific Reports.* 2019;9(1):1071.
 16. Li YH, Chen YH, Chen KJ, Chen TL. Infectious Sources, Prognostic Factors, and Visual Outcomes of Endogenous *Klebsiella pneumoniae* Ophthalmol Retina. 2018;2(8): 771-8.
 17. Chen SC, Lee YY, Chen YH, Sheu SJ. *Klebsiella pneumoniae* Infection Leads to a Poor Visual Outcome in Endogenous Endophthalmitis: A 12-year Experience in Southern Taiwan. *Ocul Immunol Inflamm.* 2017; 25(6):870-7.
 18. Qian Y, Wong CC, Lai SC, Si JM. *Klebsiella pneumoniae* invasive liver abscess syndrome with purulent meningitis and septic shock: A case from mainland China. *World J Gastroenterol.* 2016;22(9):2861-6.

Figures

Figure 1

The appearance and CT of the right eye. a. The patient right eye was swollen with purulent secretions; b. The vitreous density of the right eye was uneven and exudation around the right eyeball.

Figure 2

Low-density areas in the right lobe of the liver. a. The maximal diameter of Low-density areas was 4.65cm; b. The flake low-density shadow in the right lobe of the liver.

Figure 3

The patient’s CT image in the lungs. a. A small cavity in the lung upper right lobe; b and c. Ground-glass opacities in the lung upper right lobe; d. Solitary nodule shadow in the lung lower right lobe close to the pleura.

Figure 4

Contrast-enhanced magnetic resonance imaging of the brain revealed ring-enhancing lesion (0.5cm in diameter) located in the left forehead. a. The image on the sagittal plane. b. The image on the coronal plane.