

# Knockdown of nrf2 exacerbates TNF- $\alpha$ -induced proliferation and invasion of rheumatoid arthritis fibroblast-like synoviocytes through activating JNK pathway

**Yu Du**

Department of Rheumatology & Immunology, Shanghai Jiao Tong University Affiliated Sixth People's Hospital

**Qian Wang**

Department of Rheumatology & Immunology, Shanghai Jiao Tong University Affiliated Sixth People's Hospital

**Na Tian**

Department of Rheumatology & Immunology, Shanghai Jiao Tong University Affiliated Sixth People's Hospital

**Meng Lu**

Department of Physiology and Pathophysiology, School of Basic Medical Sciences, Fudan University

**Xian-Long Zhang**

Department of Orthopedic Surgery, Shanghai Jiao Tong University Affiliated Sixth People's Hospital

**Sheng-Ming Dai** (✉ [shengmingdai@163.com](mailto:shengmingdai@163.com))

Shanghai Jiao Tong University Affiliated Sixth People's Hospital <https://orcid.org/0000-0001-8876-4205>

---

## Research article

**Keywords:** Rheumatoid arthritis, nuclear factor erythroid 2-related factor 2, reactive oxygen species, invasion, matrix metalloproteinase

**Posted Date:** August 28th, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-64396/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

## Background.

Fibroblast-like synoviocytes (FLS) in the synovial tissue of rheumatoid arthritis (RA) exhibit over-proliferative and aggressive phenotypes, which participate in the pathophysiology of RA. In RA, little is known about the non-antioxidant effect of nuclear factor erythroid 2-related factor 2 (nrf2), master regulator of redox homeostasis. In this study, we explored the expression and upstream regulatory factors of nrf2, and revealed its functions in modulating the proliferation and invasion in RA-FLS.

## Methods.

FLS were isolated from RA and osteoarthritis patients. Expression of nrf2 in the synovial tissues and FLS was analyzed by immunohistochemistry, real-time PCR, western blot, and immunofluorescence. Cell proliferation was examined by Cell Counting Kit-8, and cell invasion was tested by transwell assay. Phosphorylation of JNK was determined by Western blot.

## Results.

Nrf2 expression in the RA synovial tissues was upregulated. TNF- $\alpha$  promoted expression and nuclear translocation of nrf2 in RA-FLS, and increased the intracellular reactive oxygen species (ROS) level. Nrf2 nuclear translocation was blocked by ROS inhibitor N-acetylcysteine. Both knockdown of nrf2 by siRNA and inhibition of nrf2 by ML385 significantly promoted the TNF- $\alpha$ -induced proliferation and invasion of RA-FLS. Activation of nrf2 by sulforaphane (SFN) profoundly inhibited the TNF- $\alpha$ -induced proliferation and invasion of RA-FLS. Knockdown of nrf2 also enhanced the TNF- $\alpha$ -induced matrix metalloproteinases (MMPs) expression and phosphorylation of JNK in RA-FLS. Proliferation and invasion of RA-FLS incubated with TNF- $\alpha$  and nrf2 siRNA was inhibited by pre-treatment with JNK inhibitor SP600125.

## Conclusion.

Taken together, nrf2 is over-expressed in synovial tissues of RA patients, which may be induced by TNF- $\alpha$  and ROS levels. Increased nrf2 may suppress TNF- $\alpha$ -induced proliferation, invasion, and MMPs expression in RA-FLS through inhibiting JNK activation, indicating that nrf2 plays a protective role to relieve the severity of synovitis in RA.

## Background

Rheumatoid arthritis (RA) is an autoimmune disease that primarily affects the joints, characterized by synovial inflammation and hyperplasia, autoantibody production, as well as cartilage and bone destruction, which would ultimately lead to irreversible functional loss(1). Although major progress has

been made in the treatment of RA, there are still concerns about incomplete responses and toxicity. Further elucidation of RA pathogenesis and development of effective therapies remain an urgent need.

Synovium comprises an intimal lining and a sublining layer, where locates two major cell types: fibroblast-like synoviocytes (FLS) and macrophage-like synoviocytes. Normal synovium has a thin intimal lining consisting of two to three layers of cells; however, the intimal lining in RA becomes inflamed, hyperplastic with multiple layers of cells and expansion in both cell types, as well as invasive of cartilage and bone(2, 3). RA-FLS exhibits an aggressive tumor-like phenotype that increases the invasiveness into the extracellular matrix (ECM) and further aggravates the joint damage (4). However, the mechanisms underlying the abnormal growth and invasion of RA-FLS are not fully understood.

Oxidative stress occurs in RA due to an imbalance in the generation and removal of reactive oxygen species (ROS), and consequently leads to inflammation, abnormal cellular behavior, and tissue damage(5). Nuclear factor erythroid 2-related factor 2 (nrf2) as a transcriptional factor and a master regulator of antioxidant response could be activated by ROS, and has been considered as a therapeutic target for RA(6). Under basal conditions, nrf2 binds to the inhibitory Kelch-like ECH-associated protein 1 (Keap1) in cytoplasm, which targets nrf2 to proteasomal degradation(7). When being activated by ROS or electrophilic compounds such as sulforaphane (SFN), nrf2 translocates into the nucleus after release from Keap1 and subsequently regulates expression of targeted genes(8–10).

The transactivation of multiple genes involved in antioxidant response, immune process, and metabolism by nrf2 has been well recognized(9). It was originally considered to exert protection against oxidative stress and inflammation in a variety of diseases including RA. Nrf2 deficiency aggravated disease severity and inflammatory condition in RA animal models(11, 12). Chemicals that directly activate nrf2 showed anti-inflammatory and immunoregulatory effects(13, 14). Nrf2 also mediated the therapeutic effects of many drugs such as licochalcone A and resveratrol in RA(15–17). Nevertheless, recent studies noted that nrf2 could also play a role in the cancer progression, promoting proliferation and invasion in glioblastoma and hepatocellular carcinoma(18–20). These remind us that nrf2 might as well modulate RA disease severity through other mechanisms such as controlling cell growth and invasion. Upregulated nrf2 has been found in the FLS of RA patients, but how would nrf2 modulate the proliferation and invasion of RA-FLS remains unknown(11).

Here, we confirmed the higher expression of nrf2 in the RA synovium, and revealed that TNF- $\alpha$  contributed to the activation of nrf2 by inducing cellular ROS in RA-FLS. More importantly, we identified knockdown of nrf2 as an enhancer of proliferation, invasion, and the expression of the matrix metalloproteinases (MMPs) of RA-FLS by activating JNK pathway.

## Methods

### Tissue samples

Synovial tissues from RA and osteoarthritis (OA) patients were obtained during the total knee arthroplasty. Patients were diagnosed according to the 2010 ACR/EULAR classification criteria for RA, or 1986 revised ACR classification criteria for OA(21, 22).

### **Isolation and culture of human FLS**

Synovial tissues were used to isolate primary FLS. The fat, fibrous membranes, and cartilage fragments were removed. The synovial tissue was minced into fragments and digested with 1 mg/mL type II collagenase (Merck Millipore, USA) dissolved in [Dulbecco's Modified Eagle Medium](#) (DMEM) at 37°C overnight. After that, the digested mixture was pipette through a 70 µm mesh cell filter (Merck Millipore, USA) into a 50 mL tube, and centrifuged at 250 g for 10 min. The supernatant was removed and the cell pellet was washed with DMEM for 3 times. Finally, the cell pellet was re-suspended with DMEM supplemented with 10% fetal bovine serum (FBS) and 100 U/mL [penicillin-streptomycin](#) (Gibco, USA). The cells were cultured in 10 cm culture dishes at a density of  $2.5 \times 10^4$  cells/mm<sup>2</sup> in a humidified incubator (Thermo Fisher Scientific, USA) containing 95% air and 5% CO<sub>2</sub> at 37°C and sub-cultured upon confluence. Passage 3-8 were used for experiments.

### **Immunohistochemistry and immunocytochemistry**

Synovial tissues were fixed in 4% paraformaldehyde (PFA) for 48 h and embedded in paraffin. Serial sections (5 µm) were made for immunohistochemistry and hematoxylin and eosin (HE) staining. For immunohistochemistry, synovial tissue sections were deparaffinized with xylene and rehydrated with gradient ethanol. Antigen retrieval was conducted with citrate buffer at 95°C for 8 min. For immunocytochemistry, cells were fixed with 4% PFA for 10 min, and permeabilized with 0.25% Triton X-100 in phosphate buffered saline (PBS) for 10 min. Tissue sections or cells were blocked with 10% donkey serum (Jackson ImmunoResearch Labs, USA) in tris buffered saline (TBS) for 1 h at room temperature, and incubated with anti-nrf2 antibodies (Abcam, UK, 1: 500) or rabbit IgG (Merck Millipore, USA, 1:500) at 4°C overnight. Then, they were washed with TBS (0.025% Triton X-100) for 2 × 5 min. Next, tissue sections were incubated with horseradish peroxidase (HRP) conjugated donkey anti-rabbit IgG (Jackson ImmunoResearch Labs, USA, 1:1000) at room temperature for 1 h, and stained with [diaminobenzidine](#) (DAB, Beyotime Biotechnology, China) to visualize the staining. Cells were incubated with Alexa Fluor 488-conjugated donkey anti-rabbit IgG secondary antibody (Abcam, UK, 1: 200) in darkness for 30 minutes at room temperature. Afterwards, cells were mounted with mounting media containing DAPI (4',6-diamidino-2-phenylindole) (Beyotime Biotechnology, China). Digital images were captured from synovial tissue sections through an optical microscope (Olympus, Japan) and from cells with a fluorescence microscope (Olympus, Japan).

### **Western blot**

Total protein of cultured RA-FLS was extracted using RIPA lysis buffer (Beyotime Biotechnology, China) on ice. Nuclear protein was extracted with Nuclear and Cytoplasmic Protein Extraction Kit (Beyotime Biotechnology, China) according to the manufacturer's instructions. The lysates were added by 5 × Gel

Sample Loading Buffer (New Cell & molecular Biotech, China) and heated for 8 minutes at 100°C. The denatured samples were separated by 10% sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and transferred to polyvinylidene fluoride (PVDF) membranes (Merck Millipore, USA). After blocking with 5% bovine serum albumin (BSA) in TBS-Tween, the membranes were incubated overnight at 4°C with primary antibodies against nrf2 (Abcam, UK, 1: 500),  $\beta$ -actin (Affinity, China, 1:5000), Histone-3 (Cell Signaling Technology, USA, 1:1000), phospho-ERK (Cell Signaling Technology, USA, 1:1000), ERK (Cell Signaling Technology, USA, 1:1000),  $\beta$ -Tubulin (Affinity, China, 1:5000), phospho-JNK (Cell Signaling Technology, USA, 1:1000), JNK (Cell Signaling Technology, USA, 1:1000), and GAPDH (Affinity, China, 1:5000). After washing with TBS-Tween, membranes were incubated with HRP-conjugated goat anti-mouse (Affinity, China, 1:5000) or anti-rabbit (Affinity, China, 1:5000) antibodies, protein bands were visualized using NcmECL Ultra enhanced chemiluminescent detection reagents (New Cell & molecular Biotech, China).

### RNA extraction and real-time quantitative PCR

Total RNA of cultured RA-FLS was extracted by Trizol reagents (Thermo Fisher Scientific, USA) and reverse-transcribed into cDNA using the ReverTra Ace qPCR RT Master Mix with gDNA Remover kit according to the manufacturer's instructions (Toyobo, Japan). Real-time PCR analysis was conducted using the SYBR Green Realtime PCR Master Mix kit (Toyobo, Japan) on the [Quant Studios 7 Flex real-time PCR system](#) (Applied Biosystems, USA). The primer sequences are listed in Table 1.

**Table 1. Primer sequences for real-time PCR.**

Gene symbol	Forward primer (5'-3')	Reverse primer (5'-3')
nrf2	TCAGCGACGGAAAGAGTATGA	CCACTGGTTTCTGACTGGATGT
MMP1	TCAGGGGAGATCATCGGGAC	GTCCAAGAGAATGGCCGAGT
MMP2	GATACCCCTTTGACGGTAAGGA	CCTTCTCCCAAGGTCCATAGC
MMP3	ACTGGAGATTTGATGAGAAGAGAA	TGGGTCAAACCTCAAACGTGA
MMP9	AGACCTGGGCAGATTCCAAAC	CGGCAAGTCTTCCGAGTAGT
MMP14	CGAGGTGCCCTATGCCTAC	CTCGGCAGAGTCAAAGTGG
$\beta$ -actin	GGACCTGACTGACTACCTCAT	CGTAGCACAGCTTCTCCTTAAT

### siRNA transfection

RA-FLS were seeded in six-well plates and reached 80% confluence at the day of transfection. Cells were transfected with 70 nM nrf2 siRNA or scramble siRNA (Ribobio, China) by Lipofectamine RNAiMAX

transfection reagent (Invitrogen, USA). Briefly, 7  $\mu\text{L}$  of nrf2 siRNA (20  $\mu\text{M}$ ) and equal volume of transfection reagent were separately diluted in 125  $\mu\text{L}$  of Opti-MEM I Medium (Gibco, USA). Diluted siRNA and diluted transfection reagent were mixed (1:1 ratio) and incubated for 5 min at room temperature. Then, the siRNA/Lipofectamine solution mixture was added to the cells.

### **ROS detection**

Intracellular ROS level was detected using DCFDA/H<sub>2</sub>DCFDA Cellular ROS Assay Kit (Abcam, UK) according to the manufacturer's protocols. In brief, cells were stained with 25  $\mu\text{M}$  2,7-Dichlorodihydrofluorescein diacetate (DCFDA) for 45 min at 37°C, and stimulated with TNF- $\alpha$  (25 ng/mL) for 0.5, 1, 2, or 3 h, or tert-butyl hydroperoxide (TBHP) (50  $\mu\text{M}$ ) for 3 h as positive control. Signals were captured at 488 nm excitation wavelength with a fluorescence microscope (Olympus, Japan) at 20  $\times$  10 magnification with equal exposure time.

### **Cell proliferation analysis**

Cell proliferation assay was performed using the Enhanced Cell Counting Kit-8 (CCK-8) (Beyotime Biotechnology, China). RA-FLS were seeded in a 96-well plate at a density of 10<sup>3</sup> cells per well and incubated with or without indicated reagents. Two or four days later, cell culture medium was removed and diluted CCK-8 reagent were added to the cells. The optical density (OD) value at 450 nm was read on a microplate reader (Thermo Fisher Scientific, USA) after incubating for 2 h. The proliferation rate was expressed as the relative OD value at 450 nm.

### **Cell invasion analysis**

The invasion assay was performed in a 24-well plate using 6.5 mm diameter inserts with membranes containing 8  $\mu\text{m}$  pores (BD Bioscience, USA). The inserts were coated with Corning Matrigel matrix (Corning, USA) at 300  $\mu\text{g}/\text{mm}^2$  according to the manufacturer's instructions. Cells were digested and seeded into the upper chamber of the transwell insert in 100  $\mu\text{L}$  DMEM without FBS at a density of 10<sup>4</sup> cells/well. DMEM (650  $\mu\text{L}$ ) containing 10% FBS was placed in the lower chamber. The transwell plate was incubated at 37°C for 2 days. Then, the chamber membrane was fixed with 4% PFA for 10 min, and washed with double distilled water. Then the membrane was stained with crystal violet solution (Beyotime Biotechnology, China). Cells remaining on the upper surface of the chamber were wiped using a cotton swab. Pictures of the membrane were taken under an optical microscope (Olympus, Japan) at 20  $\times$  10 magnification. Density of cells that migrated to the lower side of the membrane represented the cell invasion ability. The cell density for each assay was calculated as the mean density of 5 random fields.

### **Statistical analysis**

Data analysis was conducted using IBM SPSS Statistics 22.0, and graphs were plotted using GraphPad Prism 8 software. The statistical comparisons between two groups were calculated using Student's t-test.

Multiple comparisons were calculated using one-way analysis of variance (ANOVA) or two-way ANOVA for repeated measurements with Dunnett's T3 post hoc test. P-values <0.05 were considered statistically significant.

## Results

### Upregulation of *nrf2* in the synovial tissues of RA patients

Immunohistochemistry staining was performed to determine the *nrf2* protein expression level in the synovial tissues from RA and OA patients using anti-*nrf2* antibodies and rabbit IgG (isotype). *Nrf2* expression could barely be detected in the OA synovium, while there was distinct positive stain in the RA synovium compared with OA; synovial hyperplasia was also observed in RA with HE staining (Fig. 1).

### Induction of *nrf2* expression and nuclear translocation by TNF- $\alpha$ through inducing ROS.

TNF- $\alpha$  is one of the most important mediators of RA synovitis. To investigate whether TNF- $\alpha$  could cause *nrf2* upregulation and activation, RA-FLS were treated with TNF- $\alpha$  (25 ng/mL). The upregulated expression of *nrf2* mRNA was demonstrated after 3, 6, and 24 h of TNF- $\alpha$  treatment with real-time PCR analysis (Fig. 2A). The nuclear protein level of *nrf2* was increased in RA-FLS after stimulation with TNF- $\alpha$  (25 ng/mL) at different time points (Fig. 2B, C), indicating that TNF- $\alpha$  could promote *nrf2* nuclear translocation. Furthermore, the nuclear translocation of *nrf2* after TNF- $\alpha$  (25 ng/mL) treatment was confirmed with immunofluorescence staining (Fig. 2D).

To illustrate whether ROS mediates TNF- $\alpha$ -induced *nrf2* activation, intracellular ROS level of RA-FLS was detected. TBHP was used as the positive control. It was found that TNF- $\alpha$  (25 ng/mL) enhanced cellular ROS levels in a time-dependent manner (Fig. 2E). Further treatment with ROS inhibitor N-acetylcysteine (NAC) blocked TNF- $\alpha$ -induced *nrf2* nuclear translocation (Fig. 2F).

### Enhancement of proliferation and invasion by knockdown of *nrf2* in RA-FLS

Next, the biological effects of *nrf2* in RA-FLS were explored. Successful *nrf2* knockdown by siRNA was confirmed by real-time PCR and Western blot analysis (Fig. 3A-C). The effects of knockdown of *nrf2* on the TNF- $\alpha$ -induced proliferation rate and invasion in RA-FLS were analyzed with CCK-8 and transwell assay. As a result, TNF- $\alpha$  (25 ng/mL) promoted the proliferation and invasion of RA-FLS, which were further enhanced by knockdown of *nrf2* (Fig. 3D-F).

### Effects of *nrf2* activator SFN and inhibitor ML385 on proliferation and invasion of RA-FLS

RA-FLS were incubated with *nrf2* activator SFN or inhibitor ML385 in the presence or absence of TNF- $\alpha$  (25 ng/mL), to verify the influence of *nrf2* on the TNF- $\alpha$ -induced proliferation and invasion of RA-FLS. SFN (10  $\mu$ M) exacerbated the TNF- $\alpha$ -induced *nrf2* nuclear translocation (Fig. 4A). Both proliferation and invasion of RA-FLS were inhibited by SFN (10  $\mu$ M) (Fig. 4B, D, and E). ML385 is a *nrf2* inhibitor that leads

to repressed transcriptional activation function of nrf2(23). ML385 (1.25  $\mu$ M) profoundly accelerated TNF- $\alpha$ -induced invasion and proliferation of RA-FLS (Fig. 4 C, F, and G).

### **Promotion of the TNF- $\alpha$ -induced MMPs expression by knockdown of nrf2.**

Proteins of the MMP family degrade extracellular matrix and mediate cell invasion. The mRNA expression of MMP1, MMP2, MMP3, MMP9 and MMP14 in RA-FLS after treatment of TNF- $\alpha$  (25 ng/mL) and knockdown of nrf2 were analyzed with real-time PCR. TNF- $\alpha$  (25 ng/mL) upregulated the expression of MMP1, MMP2, MMP3 and MMP9 in RA-FLS; and knockdown of nrf2 further enhanced the TNF- $\alpha$ -induced expression of MMP1, MMP3 and MMP9, without affecting the expression of MMP2 (Fig. 5A-D). TNF- $\alpha$  (25 ng/mL) did not influence the expression of MMP14, but knockdown of nrf2 would promote the MMP14 expression (Fig. 5E).

### **JNK mediating the modulation of proliferation, invasion and MMP9 expression in RA-FLS by knockdown of nrf2.**

To elucidate whether MAPK pathways mediate the enhanced proliferation, invasion and MMPs expression by knockdown of nrf2 in RA-FLS. Neither TNF- $\alpha$  (25 ng/mL) treatment nor nrf2 siRNA changed the phosphorylation level of ERK (Fig. 6A). The phosphorylation of JNK was upregulated by TNF- $\alpha$  (25 ng/mL) treatment, and knockdown of nrf2 facilitated the TNF- $\alpha$ -induced JNK phosphorylation (Fig. 6B). JNK pathway inhibitor SP600125 was applied to block the phosphorylation of JNK, and SP600125 (20  $\mu$ M) was found to be effective to remarkably suppress the proliferation and invasion of RA-FLS incubated with TNF- $\alpha$  and nrf2 siRNA (Fig. 6C-E). The mRNA expression of MMP9 induced by knockdown of nrf2 and TNF- $\alpha$  (25 ng/mL) was also significantly inhibited by SP600125 treatment (Fig. 6F). The mRNA expression levels of MMP1, MMP2, MMP3, and MMP14 were not changed (data not shown).

## **Discussion**

The present study demonstrated that TNF- $\alpha$  promoted the nrf2 expression and nuclear translocation by upregulating intracellular ROS level in RA-FLS, and silencing nrf2 further facilitated the TNF- $\alpha$ -induced RA-FLS proliferation and invasion by promoting expression of MMPs partly through activating the JNK pathway. These results indicate that nrf2 acts as a negative-feedback regulator in the RA-FLS against the TNF- $\alpha$ -induced proliferation, invasion and MMPs expression, and plays a protective role to relieve the severity of synovitis in RA.

First, nrf2 protein expression was verified in the human synovial tissues. Nrf2 expression in RA synovial tissues was much higher in comparison to OA, which was consistent with the previous study where Wruck et al(11) demonstrated that nrf2 was activated in RA synovium compared with healthy donors. Pro-inflammatory cytokines, such as TNF- $\alpha$ , IL-1 and IL-6, play pivotal roles in the pathogenesis of RA. TNF- $\alpha$  is one of the most established effectors that mediate the synovial inflammation and imbalanced systemic immune status(24). In monocytes, TNF- $\alpha$  induced sustained nrf2 activation and increased expression of nrf2 targeted genes(25). Besides, low concentration of TNF- $\alpha$  evoked significant nuclear

translocation of nrf2 and transactivation of nrf2 targets(26). Here, it was found that TNF- $\alpha$  not only upregulated nrf2 expression but also triggered nrf2 nuclear translocation, indicating that increased TNF- $\alpha$  level in RA patients might account for the nrf2 overexpression in the RA synovium. Then, the mechanisms that mediated the TNF- $\alpha$ -induced nrf2 activation were explored. It has been well recognized that intracellular ROS and electrophiles are potent nrf2 activators by modulating the conformation of the nrf2 adaptor keap1 and stop the nrf2 from going through keap1-dependent ubiquitination and proteasomal degradation(9). TNF- $\alpha$  could increase the production of ROS in mouse embryonic fibroblasts, Hela cells, and murine hepatocytes(27–29). As expected, in the present study TNF- $\alpha$  induced ROS production in RA-FLS, and pretreatment with ROS scavenger NAC inhibited the TNF- $\alpha$ -induced nrf2 nuclear translocation, indicating that ROS mediated the TNF- $\alpha$ -induced nrf2 activation in RA-FLS.

Despite the widely studied powerful antioxidant ability, nrf2 exerts multiple non-antioxidant effects. Recent studies addressed that nrf2 modulated cell proliferation, survival, and invasion, although the modulating direction depended on different cellular micro-environments. Nrf2 deficiency disrupted self-renewal of the airway basal stem cells(30). Knockout of nrf2 delayed proliferation of hepatocytes after hepatectomy(31). Nevertheless, type 2 innate lymphoid cells from nrf2-deficient mice underwent hyperproliferation in response to the stimulation of IL-33 combined with IL-2, IL-7, or thymic stromal lymphoprotein(32). ML385 is a probe molecule that binds to the Neh1 domain of nrf2 and interferes the binding of V-maf avian musculoaponeurotic fibrosarcoma oncogene homologue G with nrf2 and blocks the nrf2 targeted genes expression(33). Here, both knockdown of nrf2 by siRNA and inhibiting nrf2 by ML385 significantly promoted the TNF- $\alpha$ -induced proliferation of RA-FLS. SFN is the bioactive form of glucoraphanin that are mostly found in cruciferous vegetables; it effectively activates nrf2 by modifying the sensor cysteine in keap1(34). The present study further demonstrated that activating nrf2 by SFN profoundly inhibited the TNF- $\alpha$ -induced proliferation of RA-FLS. All these data suggest that increased and activated nrf2 secondary to inflammatory micro-environment may counteract the hypertrophy of RA synovium.

The invasive characteristic of RA-FLS is one of the major causes of bone and cartilage destruction in RA. So far, how nrf2 modulates the invasion of RA-FLS has not been elucidated. Most studies regarding the effect of nrf2 on cell invasion were conducted in cancer cells, and the results were paradoxical. Overexpression of nrf2 in oral squamous cell carcinoma cells promoted cell migration and invasion(35). However, stabilizing nrf2 in breast cancer cells inhibited invasion and expression of *recepteur d'origine nantais* which is a tyrosine kinase receptor that promotes cancer cell invasion(36). The present study showed that both silencing nrf2 and inhibiting nrf2 by ML385 promoted the invasion of RA-FLS. On the contrary, activating nrf2 by SFN profoundly inhibited the TNF- $\alpha$ -induced invasion of RA-FLS. These results further suggest nrf2, being a negative feedback in RA, might exert a protective effect by down-regulating the invasiveness of RA-FLS.

Rheumatoid synoviocytes express a battery of MMPs that are the primary effectors responsible for pathologic ECM remodeling and cell invasion in RA(37). The expression levels of MMP1, MMP2, MMP3, MMP9, and MMP14 were tested, which are among the major types contributing to the process of joint

destruction in RA(38, 39). It was found that knockdown of nrf2 could enhance the expression of these MMPs. Consistent with our results, Choi et al(40) revealed that nrf2 activator SFN inhibited IL-1 $\beta$ -induced expression of MMP-1, MMP-3 in RA-FLS. Activation of nrf2 reduced UVA-mediated MMP1 upregulation in human keratinocyte cell line(41). Moreover, disruption of nrf2 enhanced the MMP9 expression after spinal cord injury in mice(42). Taken together, the above data suggest that nrf2 may down-regulate the invasiveness of RA-FLS by inhibition of MMP production.

The promoter regions of many MMPs contain the binding site of c-Fos/activator protein-1 (AP-1), which is critical for MMPs expression and locates downstream of the JNK MAPK and ERK MAPK kinases(43). It has been well recognized that TNF- $\alpha$  regulates the expression of MMPs via the MAPK signaling cascades in RA-FLS(44). Knockdown of nrf2 could also activate the MAPK pathways in UV-radiated human keratinocytes(45, 46). Thus, the pathways involving JNK and ERK were analyzed in the present study. As a result, the phosphorylation of JNK was enhanced in the RA-FLS treated with TNF- $\alpha$  alone, and the up-regulated phosphorylation of JNK was further enhanced by nrf2 siRNA. In addition, blocking JNK with its inhibitor SP600125 reduced not only the invasion but also the MMP9 expression mediated by knockdown of nrf2 in RA-FLS. In addition, several studies noted the pro-proliferative effect of JNK pathway(47, 48), which was in line with the finding that the proliferation of RA-FLS incubated with TNF- $\alpha$  and nrf2 siRNA was inhibited by pre-treatment with JNK inhibitor SP600125. These data suggest that nrf2 may inhibit the activation of JNK pathway to down-regulate MMP expression, invasion, and proliferation of RA-FLS.

## Conclusions

In conclusion, nrf2 is over-expressed in synovial tissues of RA patients, which may be induced by TNF- $\alpha$  and ROS levels. Increased nrf2 may suppress TNF- $\alpha$ -induced proliferation, invasion, and MMPs expression in RA-FLS through inhibiting JNK activation. Our findings offer a novel insight into the protective role that nrf2 played in RA.

## Abbreviations

### **ANOVA**

One-way analysis of variance

### **BSA**

Bovine serum albumin

### **CCK-8**

Cell Counting Kit-8

### **DAB**

Diaminobenzidine

### **DCFDA**

2,7-Dichlorodi-hydrofluorescein diacetate

### **DMEM**

Dulbecco's Modified Eagle Medium

**ECM**

Extracellular matrix

**FBS**

Fetal bovine serum

**FLS**

Fibroblast-like synoviocytes

**HE**

Hematoxylin and eosin

**HRP**

Horseradish peroxidase

**Keap1**

Kelch-like ECH-associated protein 1

**MMPs**

Matrix metalloproteinases

**NAC**

N-acetylcysteine

**nrf2**

Nuclear factor erythroid 2-related factor 2

**OA**

Osteoarthritis

**OD**

Optical density

**PBS**

Phosphate buffered saline

**PFA**

Paraformaldehyde

**PVDF**

Polyvinylidene fluoride

**RA**

Rheumatoid arthritis

**ROS**

Reactive oxygen species

**SFN**

Sulforaphane

**TBHP**

Tert-butyl hydroperoxide

**TBS**

Tris buffered saline

**Declarations**

## **Ethics approval and consent to participate**

This study complied with the Declaration of Helsinki, and the research protocol was approved by the Ethic Committee of Shanghai Jiao Tong University Affiliated Sixth People's Hospital. Informed consent was obtained from all the patients for the tissue donation.

## **Consent for publication**

Not applicable.

## **Availability of data and materials**

All data are available from the corresponding author on reasonable request.

## **Competing interests**

The authors declare that they have no competing interests.

## **Funding**

This work was supported by the National Natural Science Foundation of China [81771746].

## **Authors' contributions**

YD, XZ and SD designed the research. YD, QW and NT performed the research. YD, NT, QW, and ML analyzed and interpreted the data. YD and SD wrote the manuscript. All authors read and approved the final manuscript.

## **Acknowledgements**

We would like to thank Prof. Si-feng Chen (Department of Physiology and Pathophysiology, School of Basic Medical Sciences, Fudan University, Shanghai, China) for scientific advice and technical assistance.

## **References**

1. McInnes IB, Schett G. The Pathogenesis of Rheumatoid Arthritis. *N Engl J Med*. 2011;365(23):2205–19.
2. Falconer J, Murphy AN, Young SP, Clark AR, Tiziani S, Guma M, et al. Review: Synovial Cell Metabolism and Chronic Inflammation in Rheumatoid Arthritis. *Arthritis & rheumatology (Hoboken, NJ)*. 2018;70(7):984–99.
3. Bustamante MF, Garcia-Carbonell R, Whisenant KD, Guma M. Fibroblast-like synoviocyte metabolism in the pathogenesis of rheumatoid arthritis. *Arthritis research therapy*. 2017;19(1):110-.


4. Bartok B, Firestein GS. Fibroblast-like synoviocytes: key effector cells in rheumatoid arthritis. *Immunol Rev.* 2010;233(1):233–55.
5. Smallwood MJ, Nissim A, Knight AR, Whiteman M, Haigh R, Winyard PG. Oxidative stress in autoimmune rheumatic diseases. *Free Radic Biol Med.* 2018;125:3–14.
6. Ferrándiz ML, Nacher-Juan J, Alcaraz MJ. Nrf2 as a therapeutic target for rheumatic diseases. *Biochem Pharmacol.* 2018;152:338–46.
7. Villeneuve NF, Lau A, Zhang DD. Regulation of the Nrf2-Keap1 antioxidant response by the ubiquitin proteasome system: an insight into cullin-ring ubiquitin ligases. *Antioxid Redox Signal.* 2010;13(11):1699–712.
8. Silva-Islas CA, Maldonado PD. Canonical and non-canonical mechanisms of Nrf2 activation. *Pharmacological research.* 2018;134:92–9.
9. Bellezza I, Giambanco I, Minelli A, Donato R. Nrf2-Keap1 signaling in oxidative and reductive stress. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research.* 2018;1865(5):721–33.
10. Ahmed SMU, Luo L, Namani A, Wang XJ, Tang X. Nrf2 signaling pathway: Pivotal roles in inflammation. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease.* 2017;1863(2):585–97.
11. Wruck CJ, Fragoulis A, Gurzynski A, Brandenburg LO, Kan YW, Chan K, et al. Role of oxidative stress in rheumatoid arthritis: insights from the Nrf2-knockout mice. *Ann Rheum Dis.* 2011;70(5):844–50.
12. Maicas N, Ferrándiz ML, Brines R, Ibáñez L, Cuadrado A, Koenders MI, et al. Deficiency of Nrf2 accelerates the effector phase of arthritis and aggravates joint disease. *Antioxid Redox Signal.* 2011;15(4):889–901.
13. Ferrándiz ML, Nacher-Juan J, Alcaraz MJ. Nrf2 as a therapeutic target for rheumatic diseases. *Biochem Pharmacol.* 2018;152:338–46.
14. Kong JS, Yoo SA, Kim HS, Kim HA, Yea K, Ryu SH, et al. Inhibition of synovial hyperplasia, rheumatoid T cell activation, and experimental arthritis in mice by sulforaphane, a naturally occurring isothiocyanate. *Arthritis rheumatism.* 2010;62(1):159–70.
15. Chu J, Wang X, Bi H, Li L, Ren M, Wang J. Dihydromyricetin relieves rheumatoid arthritis symptoms and suppresses expression of pro-inflammatory cytokines via the activation of Nrf2 pathway in rheumatoid arthritis model. *Int Immunopharmacol.* 2018;59:174–80.
16. Su X, Li T, Liu Z, Huang Q, Liao K, Ren R, et al. Licochalcone A activates Keap1-Nrf2 signaling to suppress arthritis via phosphorylation of p62 at serine 349. *Free Radic Biol Med.* 2018;115:471–83.
17. Zhang Y, Wang G, Wang T, Cao W, Zhang L, Chen X. Nrf2-Keap1 pathway-mediated effects of resveratrol on oxidative stress and apoptosis in hydrogen peroxide-treated rheumatoid arthritis fibroblast-like synoviocytes. *Ann N Y Acad Sci.* 2019;1457(1):166–78.
18. Zhang M, Zhang C, Zhang L, Yang Q, Zhou S, Wen Q, et al. Nrf2 is a potential prognostic marker and promotes proliferation and invasion in human hepatocellular carcinoma. *BMC Cancer.* 2015;15:531.

19. Pölönen P, Jawahar Deen A, Leinonen HM, Jyrkkänen HK, Kuosmanen S, Mononen M, et al. Nrf2 and SQSTM1/p62 jointly contribute to mesenchymal transition and invasion in glioblastoma. *Oncogene*. 2019;38(50):7473–90.
20. Rojo de la Vega M, Chapman E, Zhang DD. NRF2 and the Hallmarks of Cancer. *Cancer Cell*. 2018;34(1):21–43.
21. Kay J, Upchurch KS. ACR/EULAR 2010 rheumatoid arthritis classification criteria. *Rheumatology*. 2012;51(Suppl 6):vi5–9.
22. Altman R, Asch E, Bloch D, Bole G, Borenstein D, Brandt K, et al. Development of criteria for the classification and reporting of osteoarthritis. Classification of osteoarthritis of the knee. Diagnostic and Therapeutic Criteria Committee of the American Rheumatism Association. *Arthritis rheumatism*. 1986;29(8):1039–49.
23. Singh A, Venkannagari S, Oh KH, Zhang Y-Q, Rohde JM, Liu L, et al. Small Molecule Inhibitor of NRF2 Selectively Intervenes Therapeutic Resistance in KEAP1-Deficient NSCLC Tumors. *ACS Chem Biol*. 2016;11(11):3214–25.
24. Alam J, Jantan I, Bukhari SNA. Rheumatoid arthritis: Recent advances on its etiology, role of cytokines and pharmacotherapy. *Biomedicine pharmacotherapy = Biomedecine pharmacotherapie*. 2017;92:615–33.
25. Rushworth SA, Shah S, MacEwan DJ. TNF mediates the sustained activation of Nrf2 in human monocytes. *J Immunol*. 2011;187(2):702–7.
26. Shanmugam G, Narasimhan M, Sakthivel R, Kumar RR, Davidson C, Palaniappan S, et al. A biphasic effect of TNF- $\alpha$  in regulation of the Keap1/Nrf2 pathway in cardiomyocytes. *Redox Biol*. 2016;9:77–89.
27. Ventura JJ, Cogswell P, Flavell RA, Baldwin AS Jr, Davis RJ. JNK potentiates TNF-stimulated necrosis by increasing the production of cytotoxic reactive oxygen species. *Genes Dev*. 2004;18(23):2905–15.
28. Kim JJ, Lee SB, Park JK, Yoo YD. TNF- $\alpha$ -induced ROS production triggering apoptosis is directly linked to Romo1 and Bcl-XL. *Cell Death Differentiation*. 2010;17(9):1420–34.
29. Kastl L, Sauer S, Beissbarth T, Becker M, Krammer P, Gülow K. TNF- $\alpha$  stimulation enhances ROS-dependent cell migration via NF- $\kappa$ B activation in liver cells. *Free Radic Biol Med*. 2014;75(Suppl 1):32.
30. Paul MK, Bisht B, Darmawan DO, Chiou R, Ha VL, Wallace WD, et al. Dynamic changes in intracellular ROS levels regulate airway basal stem cell homeostasis through Nrf2-dependent Notch signaling. *Cell Stem Cell*. 2014;15(2):199–214.
31. Beyer TA, Xu W, Teupser D, auf dem Keller U, Bugnon P, Hildt E, et al. Impaired liver regeneration in Nrf2 knockout mice: role of ROS-mediated insulin/IGF-1 resistance. *EMBO J*. 2008;27(1):212–23.
32. Nagashima R, Kosai H, Masuo M, Izumiyama K, Noshikawaji T, Morimoto M, et al. Nrf2 Suppresses Allergic Lung Inflammation by Attenuating the Type 2 Innate Lymphoid Cell Response. *J Immunol*. 2019;202(5):1331–9.
33. Singh A, Venkannagari S, Oh KH, Zhang YQ, Rohde JM, Liu L, et al. Small Molecule Inhibitor of NRF2 Selectively Intervenes Therapeutic Resistance in KEAP1-Deficient NSCLC Tumors. *ACS Chem Biol*.

- 2016;11(11):3214–25.
34. Russo M, Spagnuolo C, Russo GL, Skalicka-Woźniak K, Daglia M, Sobarzo-Sánchez E, et al. Nrf2 targeting by sulforaphane: A potential therapy for cancer treatment. *Critical reviews in food science nutrition*. 2018;58(8):1391–405.
  35. Fan H, Paiboonrungruan C, Zhang X, Prigge JR, Schmidt EE, Sun Z, et al. Nrf2 regulates cellular behaviors and Notch signaling in oral squamous cell carcinoma cells. *Biochem Biophys Res Commun*. 2017;493(1):833–9.
  36. Thangasamy A, Rogge J, Krishnegowda NK, Freeman JW, Ammanamanchi S. Novel function of transcription factor Nrf2 as an inhibitor of RON tyrosine kinase receptor-mediated cancer cell invasion. *J Biol Chem*. 2011;286(37):32115–22.
  37. Sabeh F, Fox D, Weiss SJ. Membrane-type I matrix metalloproteinase-dependent regulation of rheumatoid arthritis synoviocyte function. *J Immunol*. 2010;184(11):6396–406.
  38. Jackson C, Nguyen M, Arkell J, Sambrook P. Selective matrix metalloproteinase (MMP) inhibition in rheumatoid arthritis—targetting gelatinase A activation. *Inflammation research: official journal of the European Histamine Research Society [et al]*. 2001;50(4):183–6.
  39. Burrage PS, Mix KS, Brinckerhoff CE. Matrix metalloproteinases: role in arthritis. *Frontiers in bioscience: a journal virtual library*. 2006;11:529–43.
  40. Choi YJ, Lee WS, Lee EG, Sung MS, Yoo WH. Sulforaphane inhibits IL-1 $\beta$ -induced proliferation of rheumatoid arthritis synovial fibroblasts and the production of MMPs, COX-2, and PGE2. *Inflammation*. 2014;37(5):1496–503.
  41. Chaiprasongsuk A, Lohakul J, Soontrapa K, Sampattavanich S, Akarasereenont P, Panich U. Activation of Nrf2 Reduces UVA-Mediated MMP-1 Upregulation via MAPK/AP-1 Signaling Cascades: The Photoprotective Effects of Sulforaphane and Hispidulin. *J Pharmacol Exp Ther*. 2017;360(3):388–98.
  42. Mao L, Wang H, Qiao L, Wang X. Disruption of Nrf2 enhances the upregulation of nuclear factor-kappaB activity, tumor necrosis factor- $\alpha$ , and matrix metalloproteinase-9 after spinal cord injury in mice. *Mediators Inflamm*. 2010;2010:238321.
  43. Brinckerhoff CE, Matrisian LM. Matrix metalloproteinases: a tail of a frog that became a prince. *Nature reviews Molecular cell biology*. 2002;3(3):207–14.
  44. Mengshol JA, Mix KS, Brinckerhoff CE. Matrix metalloproteinases as therapeutic targets in arthritic diseases: bull's-eye or missing the mark? *Arthritis rheumatism*. 2002;46(1):13–20.
  45. Chaiprasongsuk A, Lohakul J, Soontrapa K, Sampattavanich S, Akarasereenont P, Panich U. Activation of Nrf2 Reduces UVA-Mediated MMP-1 Upregulation via MAPK/AP-1 Signaling Cascades: The Photoprotective Effects of Sulforaphane and Hispidulin. *J Pharmacol Exp Ther*. 2017;360(3):388–98.
  46. Jeayeng S, Wongkajornsilp A, Slominski AT, Jirawatnotai S, Sampattavanich S, Panich U. Nrf2 in keratinocytes modulates UVB-induced DNA damage and apoptosis in melanocytes through MAPK signaling. *Free Radic Biol Med*. 2017;108:918–28.


47. Pinal N, Calleja M, Morata G. Pro-apoptotic and pro-proliferation functions of the JNK pathway of *Drosophila*: roles in cell competition, tumorigenesis and regeneration. *Open biology*. 2019;9(3):180256.
48. Guo J, Qiu X, Zhang L, Wei R. Smurf1 regulates macrophage proliferation, apoptosis and migration via JNK and p38 MAPK signaling pathways. *Molecular immunology*. 2018;97:20–6.

## Figures




**Figure 1**

Nuclear factor erythroid 2-related factor 2 (nrf2) expression in rheumatoid arthritis (RA) and osteoarthritis (OA). (A) Immunohistochemistry staining using anti-nrf2 antibodies and rabbit IgG (isotype), and hematoxylin and eosin (HE) staining of RA and OA synovial tissues. (B) Nrf2-positive cell densities in the synovial tissues of RA and OA. Scale bars represented 50  $\mu$ m. N=4. Data were shown as mean  $\pm$  standard deviation (SD), \*  $p < 0.05$ .




**Figure 2**

Induction of nrf2 expression and nuclear translocation by TNF- $\alpha$  through inducing ROS. (A-C) The mRNA expression (A) and nuclear protein level (B, C) of nrf2 in RA-FLS were detected by real-time PCR and Western blot after TNF- $\alpha$  (25 ng/mL) stimulation for 0, 3, 6, and 24 h. Histone-3 was used as the housekeeping gene in Western blot. N=3. (D) Nrf2 nuclear translocation (as shown by white arrows) after TNF- $\alpha$  (25 ng/mL) stimulation for 0, 3, 6, and 24 h was examined by immunofluorescence staining. Representative pictures (magnification  $\times 200$ ) from different groups and partially enlarged pictures (magnification  $\times 600$ ) were shown. Scale bar represented 50  $\mu$ m. (E) Intracellular ROS levels in RA-FLS stimulated with TNF- $\alpha$  (25 ng/mL) for 0, 0.5, 1, 2, and 3 h, and tert-butyl hydroperoxide (TBHP, 50  $\mu$ M) (positive control) for 3 h were detected using DCFDA probe. Scale bar represented 100  $\mu$ m. (F) The nuclear protein level of nrf2 in RA-FLS after TNF- $\alpha$  (25 ng/mL) stimulation for 3 h with or without N-acetylcysteine (NAC, 5  $\mu$ M) pretreatment for 1 h was detected by Western blot. Histone-3 was used as the housekeeping gene. Data were shown as mean  $\pm$  SD, \*  $p < 0.05$ , \*\*  $p < 0.01$ .



**Figure 3**


Knockdown of nrf2 promoted TNF- $\alpha$ -induced proliferation and invasion of RA-FLS. (A) Nrf2 mRNA expression was reduced by nrf2 siRNA transfection for 48 h, as revealed by real-time PCR. (B, C) Protein expression level of nrf2 was examined after nrf2 siRNA transfection for 72 h by Western blot analysis.  $\beta$ -actin was used as the housekeeping gene. (D) Proliferation rate of RA-FLS was analyzed by CCK-8 assay after transfection with scramble siRNA or nrf2 siRNA for 72 h. Cells were examined before and after 4 days of culture in the presence or absence of TNF- $\alpha$  (25 ng/mL). The proliferation rate was expressed as the relative optical density (OD) value at 450 nm between day 4 and day 0. (E, F) Invasion of RA-FLS was analyzed with transwell assay after transfection with scramble siRNA or nrf2 siRNA for 72 h. Cells were cultured in the transwell chamber coated with Matrigel in the presence or absence of TNF- $\alpha$  (25 ng/mL). Density of cells that migrated to the lower side of the membrane represented the cell invasion ability, as visualized by crystal violet staining. Pictures were taken at  $\times 200$  magnification. Scale bar represented 200  $\mu$ m. N=3. Data were shown as mean  $\pm$  SD, \*  $p < 0.05$ , \*\*  $p < 0.01$ .



**Figure 4**


Effects of nrf2 activator and inhibitor on proliferation and invasion of RA-FLS. (A) Nrf2 nuclear translocation (as shown by white arrows) was tested with immunofluorescence staining in RA-FLS after pre-treatment with nrf2 activator sulforaphane (SFN) at a concentration of 10  $\mu\text{M}$  for 1 h and subsequently stimulating with/without TNF- $\alpha$  (25 ng/mL) for 3 h. (B, C) Proliferation of RA-FLS treated with SFN or ML385 was analyzed by CCK-8 assay. Cells seeded in the 96-well plate were pre-treated with SFN (10  $\mu\text{M}$ ) (B) or ML385 (1.25  $\mu\text{M}$ ) (C) for 1 h, and subsequently stimulated with/without TNF- $\alpha$  (25 ng/mL) for 2 days. (D-G) Invasion analysis of RA-FLS treated by SFN or ML385 was performed with transwell assay. Cells seeded in the transwell chamber were pre-treated with SFN (10  $\mu\text{M}$ ) (D, E) or ML385

(1.25  $\mu$ M) (F, G) for 1 h, and subsequently stimulated with/without TNF- $\alpha$  (25 ng/mL) for 2 days. Scale bar represented 200  $\mu$ m. N=3. Data were shown as mean  $\pm$  SD, \*  $p < 0.05$ , \*\*  $p < 0.01$ .



**Figure 5**

Knockdown of nrf2 promoted TNF- $\alpha$ -induced matrix metalloproteinases (MMPs) expression. After transfection with scramble siRNA or nrf2 siRNA for 3 days, RA-FLS were stimulated with/without TNF- $\alpha$  (25 ng/mL) for 24 h. The expression of MMP1 (A), MMP2 (B), MMP3 (C), MMP9 (D), and MMP14 (E) were examined with real-time PCR. N=3. Data were shown as mean  $\pm$  SD, \*  $p < 0.05$ , \*\*  $p < 0.01$ .



**Figure 6**

JNK pathway mediates the modulation of proliferation, invasion and MMP9 expression in RA-FLS by nrf2. (A) After transfection with scramble siRNA or nrf2 siRNA for 3 days, RA-FLS were further stimulated with/without TNF- $\alpha$  (25 ng/mL) for 24 h. Representative figures showed the phosphorylation of ERK examined by Western blot.  $\beta$ -Tubulin was used as the housekeeping gene. (B) Representative figures showed the phosphorylation of JNK examined by Western blot. GAPDH was used as the housekeeping gene. (C) After transfection with nrf2 siRNA for 3 days, proliferation of RA-FLS was analysis by CCK-8 assay. Cells were cultured in the 96-well plate and stimulated with TNF- $\alpha$  (25 ng/mL) in the presence or absence of SP600125 (20  $\mu$ M) for 4 days. (D, E) After transfection with nrf2 siRNA for 3 days, invasion of RA-FLS was analyzed with transwell assay. Cells were cultured in the transwell chamber and stimulated with TNF- $\alpha$  (25 ng/mL) in the presence or absence of SP600125 (20  $\mu$ M) for 2 days. (F) After transfection with nrf2 siRNA for 3 days, RA-FLS were further stimulated with TNF- $\alpha$  (25 ng/mL) in the presence or

absence of SP600125 (20  $\mu$ M) for 24 h. The MMP9 mRNA expression was examined by real-time PCR. Scale bar represented 200  $\mu$ m. N=3. Data were shown as mean  $\pm$  SD, \*\*  $p < 0.01$ .