Assessment of Asymptomatic Bacteriuria and Sterile Pyuria among Antenatal Attendants in hospitals in Northern Ghana

Akosua Bonsu Karikari (✉ asbuks@yahoo.co.uk)

University for Development Studies <https://orcid.org/0000-0001-8163-1311>

Courage Kosi Setsoafia Saba

University for Development Studies

David Yembilla Yamik

University for Development Studies

Research article

Keywords: Asymptomatic bacteriuria, Sterile pyuria, Antibiotic susceptibility, Tamale, Ghana

Posted Date: October 7th, 2019

DOI: <https://doi.org/10.21203/rs.2.15813/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Pregnancy and Childbirth on April 22nd, 2020. See the published version at <https://doi.org/10.1186/s12884-020-02936-6>.

Abstract

Background Asymptomatic bacteriuria (ASB) and sterile pyuria (SP) are complexities of UTI whose prevalence are not known in the northern sector of Ghana. Our aim was to determine the occurrence of sterile pyuria and asymptomatic bacteriuria among pregnant women accessing antenatal care at a secondary and tertiary care hospitals in Tamale, northern Ghana.

Methods A cross sectional study was conducted by screening 530 pregnant women with no signs of acute urinary tract infection attending antenatal clinic for a period of 6 months. Midstream urine was collected for microscopy, quantitative urine culture and antibiotic susceptibility testing. Data analysis was carried out using the Statistical Package for Social Sciences version 20.

Results Of the 390 and 90 women screened for ASB, prevalence was 20% and 35.5% respectively at Tamale Central and Tamale Teaching Hospital. Sterile pyuria was found among 66% of the 50 women presenting at Tamale Central Hospital. More than 64% of isolates recovered from ASB patients were *S. aureus* and coagulase negative Staph. (CoNS). *Escherichia coli* was the dominant species among members of the enterobacteriaceae isolated. Highest susceptibility was recorded against gentamicin and amikacin while most resistance was to Ampicillin, cotrimoxazole, chloramphenicol and nitrofurantoin. Resistance to imipenem and vancomycin were 28.8% and 52%, with strains showing multiple drug resistance of between 81-92%.

Conclusion The prevalence of asymptomatic bacteriuria is appreciably higher (20-35.5%) than documented rates in the southern sector of the country. The presence of sterile pyuria which may be an indication of asymptomatic renal impairment and most often overlooked in antenatal management is 66%. Empirical treatment of UTIs at the Tamale Central and Teaching Hospital without confirmation of susceptibility may result in treatment failure. It is necessary to screen and treat pregnant women for ASB and SP due to the complications associated with these conditions.

Background

Urinary Tract Infection (UTI) is the invasion of microbes and the ensuing proliferation on part or the entire urinary tract [1]. It is the most common disorder caused by bacterial agents in pregnancy, which may lead to complications in neonates of such mothers in case of inappropriate diagnosis and treatment. As one of the most frequent acquired infections, UTI has an evident role in raising the number of stillbirth deliveries [2, 3]. Yearly, UTI and its related complications are the cause of nearly 150 million deaths worldwide [4]. Urinary tract infections in pregnancy is categorised into symptomatic and asymptomatic infections [5]. Asymptomatic bacteriuria is the commonest cause of UTI in pregnancy and mainly involves the lower urinary tract while the upper urinary tract engagement may result in symptomatic bacteriuria characterised by pyelonephritis [6].

Asymptomatic bacteriuria (ASB) is described as a significant bacteriuria without symptoms of acute UTI. Pregnant women with ASB have a higher risk to deliver premature or low-birth weight infants, develop pre-eclampsia and polyhydramnios [7, 8, 9]. Other conditions associated with ASB include transient renal failure, acute respiratory distress syndrome, shock and haematological abnormalities which occur in untreated or inadequately managed cases [9, 10]. Urinary tract infections and pyelonephritis in pregnancy

has also been linked to morbidity in both mother and foetus [8]. Culturable bacteria species usually recovered from ASB infections fall under fourteen genera although non-culturable pathogens have also been implicated [11]. The Enterobacteriaceae are responsible for nearly (90%) all cases of asymptomatic bacteriuria with *E. coli* being dominant. *Enterococcus* spp., *Staphylococcus aureus* and coagulase-negative *Staphylococci* may also cause ASB [12, 13].

Another urinary tract condition which is also not rare in pregnancy is Sterile Pyuria (SP). Although no current definition exists [14], SP is considered if a mid-stream urine specimen has 10 or more white blood cells per cubic millimeter or a urinary dipstick test is leukocyte esterase positive with no associated positive urinary culture [15]. Sterile pyuria has vast aetiological spectrum including sexually transmitted diseases, tuberculosis, interstitial cystitis, chlamydia and cystitis [15]. Population-based studies recount it as a highly prevalent condition with 13.9% of women and 2.6% of men being affected [16], with its preponderance in women attributable to pelvic infections [17].

Various studies have consistently revealed that early diagnosis and treatment of UTI in pregnancy exorbitantly reduce sequelae associated with the condition. Treatment of ASB in pregnancy with antibiotics has been found to decrease pyelonephritis from 35% to 4% [18], improve fetal outcomes and prevent preterm deliveries. Several institutions prescribe screening of pregnant women for ASB due to the benefits of treatment [19, 20]. But, in many developing countries such as Ghana, ASB screening in pregnancy is normally overlooked in antenatal management. The prevalence of asymptomatic bacteriuria and sterile pyuria in the Northern regions of Ghana are not known. The aim of this assessment was to screen mid-stream urine of antenatal attendants at the Tamale Teaching Hospital and Tamale Central Hospital for sterile pyuria and asymptomatic bacteriuria with associated agents and their susceptibility profiles. This was to contribute to the literature bank of urinary tract infections at these secondary and tertiary care hospitals.

Methods

Study design

A cross sectional study was conducted from 16th April 2018 to 10th September 2018 at the Tamale Teaching Hospital (TTH) and Tamale Central hospital (TCH). The Tamale Teaching Hospital is an 800 bed capacity tertiary care facility which provides referral services to three Regions in the northern sector of Ghana and affiliated to the Medical School of the University for Development Studies. The Tamale Central Hospital is a secondary care facility which supports the teaching hospital in providing healthcare services to the populace in Tamale and its environs.

Recruitment of study participants

A total of 530 pregnant women of all ages attending these hospitals for antenatal care within the study period were recruited. Informed consent was sought from all the women before enrollment. Attendees who

refused consent were excluded from the study. Socio-demographic information and diagnosis were obtained from the laboratory report form and their medical folders.

Specimen collection and processing

Pregnant women were provided with sterile urine containers and tutored on how to collect midstream urine. Approximately 10mls of urine were collected from each participant. The urine samples were transported to the Spanish laboratory of the University for Development Studies for processing and analysis, within four hours of collection.

Urine microscopy

About 5 mL of adequately mixed urine sample was centrifuged at 3000 rpm for 10 min. A drop of the sediment was placed on a glass slide, cover slipped and examined under the microscope to detect pus cells, red blood cells, casts and crystals. The presence of 10 or more pus cells/high-power field (HPF) was indicative of pyuria. Sterile pyuria in this study was defined as specimens scoring 10 or more pus cells but recorded no growth on CLED plates after adequate incubation period of 24hrs.

Urine culture and identification

Urine samples were cultured on cysteine lactose electrolyte deficient agar (CLED) using a standardized (0.01mL) wire loop. The plates were incubated at 37°C and read after 24 hours for significant bacteriuria. Significant bacteriuria was defined as a quantitative count of $\geq 10^5$ CFU/ml. Pathogens were identified using standardized biochemical tests, Mannitol salt agar and sugar fermentation using Triple Sugar Iron agar from 24h pure culture colonies. Asymptomatic bacteriuria in this study was considered when the bacterial value was $\geq 10^5$ but participants had no symptoms of acute urinary tract infections.

Antibiotic susceptibility test

Antibiotic susceptibility tests were conducted using Kirby Bauer disc diffusion method. Mueller-Hinton agar plates were inoculated with 0.5 McFarland standard saline suspension and incubated at 37°C for 24hours [21]. The antibiotics analysed included, ciprofloxacin 10µg, gentamicin 10 µg, erythromycin 15 µg, ceftriaxone 30 µg, chloramphenicol 30 µg, nitrofurantoin 50 µg, tetracycline 30 µg, ampicillin 10 µg, clindamycin 10 µg, vancomycin 30 µg, ceftiofur 30 µg, amikacin 30 µg, trimethoprim-sulfamethoxazole 25 µg, imipenem 10 µg, amoxicillin clavulanic acid 30 µg and norfloxacin 10 µg. The recorded inhibition zones were interpreted using CLSI breakpoints. Quality control strains of *Escherichia coli* (ATCC 25922) and *Staphylococcus aureus* (ATCC 25923) were used. Multidrug resistance in this study was defined as resistance of isolates to three (3) or more classes of antibiotics.

Data management

Collected data was entered into SPSS version 20. Descriptive statistics such as frequencies and percentages were used. Data was presented in tables. Associations between categorical outcome

variables were conducted using the Pearson- Chi square test at the 95% significant level. A two tailed p-value of <0.05 was considered statistically significant.

Ethical consideration

Ethics approval was obtained from the Ethical Review Committee of the Tamale Teaching Hospital (THERC/25/06/19/14). Verbal informed consent was sought from the pregnant women after sufficient information regarding the study was provided. This is because while some of the women were unable to read and write others felt reluctant to write and saw the process as a bother.

Results

Asymptomatic bacteriuria was present among 20% and 35.5% of pregnant women seeking antenatal care at the Tamale Central Hospital and Tamale Teaching Hospital respectively. More than 60% of these women were between the ages of 20-29 years but less frequent among ages less than 20 years and more than 40 year groups. The difference in prevalence at these hospitals was not significant, $p > 0.05$, Table 1.

Table 1. Asymptomatic bacteriuria among pregnant women at two hospitals in Tamale

	Tamale Central Hospital		Tamale Teaching Hospital	
	N=390		N=90	
Age/yrs.	Frequency (%)		Frequency (%)	P-value
< 20	3 (3.8)		1 (3.1)	0.157
20-29	50 (64.1)		21 (65.6)	
30-39	23 (29.5)		10 (31.3)	
40-49	2 (2.6)		0	
Total	78 (20)		32 (35.5)	

Organisms isolated from pregnant women with ASB at the Tamale Central and Teaching hospitals were similar and there was no significant difference in the rate of pathogen recovery ($p = 0.423$). At both hospitals, Gram positives mainly *S. aureus* and coagulase negative Staph. (C₀NS) were the dominant species identified as 64.1% and 75.0% were respectively found at Central and the Teaching hospital. These pathogens were rife among age group 20-29 with a rate of 42.3% at Central and 50% at Teaching hospital. This was followed by age group 30-39 with respective prevalence of 17.9% and 25%. *Escherichia coli* was more commonly isolated among the Gram negatives, followed by *Enterobacter* and *Klebsiella sp.*, Table 2.

Table 2. Recovered isolates from pregnant women presenting with asymptomatic bacteriuria

		Tamale Hospital			Central				
		Gram			Gram				
		Positives			Negatives				
Age/Yrs	Frequency (%)	<i>S. aureus</i>	<i>CoNS</i>	<i>Strept.</i>	<i>E. coli</i>	<i>Kleb.sp</i>	<i>Entero.</i>	<i>Serratia</i>	<i>Pseud.</i>
< 20	3 (3.8)	2	1	0	0	0	0	0	0
20-29	50 (64.1)	14	19	1	6	4	4	2	0
30-39	23 (29.5)	7	7	2	5	0	0	0	0
40-49	2 (2.6)	0	0	0	0	1	6	0	1
Total	78 (100)	23	27	3	11	4	7	2	1

		Tamale Teaching Hospital							
		Gram			Gram				
		Positives			Negatives				
Age/Yrs	Frequency (%)	<i>S. aureus</i>	<i>CoNS</i>	<i>Strept.</i>	<i>E. coli</i>	<i>Kleb.sp</i>	<i>Entero.</i>	<i>Serratia</i>	
< 20	1 (3.1)	0	0	1	0	0	0	0	
20-29	21 (65.6)	5	11	0	1	1	1	2	
30-39	10 (31.3)	6	2	0	0	1	1	0	
40-49	0	0	0	0	0	0	0	0	
Total	32 (100)	11	13	1	1	2	2	2	

Strept.-Streptococcus sp, Kleb.-klebsiella sp, Entero.-Enterobacter sp, pseud.-pseudomonas sp, CoNS-coagulase negative Staphylococcus

Generally resistance was commonly observed among the Gram negatives. The isolates were fairly susceptible to gentamicin and amikacin as resistance of below 20% was recorded at both hospitals. Resistance to the fluoroquinolones ranged from 0-26.6% to ciprofloxacin and 12-42.9% to norfloxacin. Among the broad spectrum antibiotics, isolates showed highest resistance to Ampicillin (76-96%) followed by erythromycin (32-85.7%), tetracycline (30-76%), chloramphenicol (39-71.4%) and amoxicillin

clavulanic acid (28-57%). Resistance to nitrofurantoin ranged from 33-71% and between 43-71% was recorded against trimethoprim sulphamethoxazole. Isolates resistance to ceftriaxone was less than 50% (22-42.9%). About 41-52% of the Gram positives showed resistance to vancomycin as Gram negative resistance to imipenem ranged from 8-28.6%, Tables 3 and 4. Isolates from Central Hospital showed less resistance to ceftriaxone (22-28%) and amoxicillin clavulanic acid (28-39.6%) as against 32-42.9% and 52-57.1% respectively recorded at the Tamale Teaching Hospital. But the Central Hospital strains showed higher resistance (30-76%) to tetracycline than isolates from Tamale Teaching Hospital (28-32%). The difference in resistance rates in the two hospitals was found to be statistically significant, $p=0.000$. Multidrug resistance of 92.3% and 81.3% were recorded among isolates recovered respectively from Central and the Teaching hospital, Table 5.

Table 3. Susceptibility profile of isolates recovered from pregnant women with asymptomatic bacteriuria

Antibiotic	Tamale Central Hospital						Tamale Teaching Hospital					
	N=53			N=25			N=25			N=7		
	Gram Positive			Gram Negative			Gram Positive			Gram Negative		
	S	I	R	S	I	R	S	I	R	S	I	R
Ciprofloxacin	41	4	8	18	1	6	25	0	0	5	0	2
Cotrimoxazole	27	3	23	8	2	15	13	1	11	2	0	5
Gentamicin	42	5	6	23	0	2	23	2	0	6	0	1
Amikacin	44	3	6	21	0	13	21	1	3	6	0	1
Ampicillin	5	0	48	4	2	19	1	0	24	1	0	6
Amoxicillin/clav.	31	1	21	12	6	7	12	0	13	2	1	4
Chloramphenicol	30	2	21	10	0	15	13	0	12	1	1	5
Nitrofurantoin	24	11	18	5	3	17	12	2	11	2	0	5
Ceftriaxone	13	28	12	12	6	7	5	12	8	2	2	3
Erythromycin	14	13	26	3	3	19	8	9	8	1	0	6
Tetracycline	31	6	16	5	1	19	15	2	8	4	1	2
Norfloxacin	34	1	18	18	0	7	22	0	3	4	0	3
Cefoxitin	34	8	11	NT			12	2	11	NT		
Clindamycin	21	4	28	NT			9	2	14	NT		
Vancomycin	30	1	22	NT			12	0	13	NT		
Imipenem	NT			23	0	2	NT			5	0	2

NT- Not tested, Cotrimoxazole- SXT, S- sensitive, I-intermediate, R-resistant

Table 4. Resistance patterns of recovered isolates from pregnant women with ASB

Antibiotic	Tamale Central Hospital		Tamale Teaching Hospital	
	Resistance %			
	Gram positive	Gram negative	Gram positive	Gram negative
Ciprofloxacin	15.1	24	0	26.6
Co-trimoxazole	43.4	60	44	71.4
Gentamicin	11.3	8	0	14.3
Amikacin	11.3	12	12	14.3
Ampicillin	90.6	76	96	85.7
Amoxicillin/clav.	39.6	28	52	57.1
Chloramphenicol	39.6	60	48	71.4
Nitrofurantoin	33.9	68	44	71.4
Ceftriaxone	22.6	28	32	42.9
Erythromycin	49.1	76	32	85.7
Tetracycline	30.2	76	32	28.6
Norfloxacin	33.9	28	12	42.9
Cefoxitin	20.8	NT	44	NT
Clindamycin	52.8	NT	56	NT
Vancomycin	41.5	NT	52	NT
Imipenem	NT	8	NT	28.6

NT- Not tested

Table 5. Multidrug resistance exhibited by isolates

Isolates	Tamale Central Hospital		Tamale Teaching Hospital	
	No. of isolates	MDR (%)	No. of isolates	MDR (%)
<i>S. aureus</i>	23	23(100.0)	11	11(100.0)
<i>CONS</i>	27	22(81.5)	13	8(61.5)
<i>Streptococcus sp.</i>	3	2(66.7)	1	1(100.0)
<i>E. coli</i>	11	11(100.0)	1	1(100.0)
<i>Klebsiella sp.</i>	4	4(100.0)	2	2(100.0)
<i>Enterobacter sp.</i>	7	7(100.0)	2	2(100.0)
<i>Serratia</i>	2	2(100.0)	2	1(50.0)
<i>Pseudomonas sp.</i>	1	1(100.0)	0	0
Total	78	72(92.3)	32	26(81.3)
Gram positives	53	47 (88.7)	25	20(80)
Gram negatives	25	25 (100.0)	7	6(85.7)

Of the 50 attendants screened for sterile pyuria at the Tamale Central Hospital, 66% were positive with more than 70% of the women within the ages of 20-29 years and a mean age of 25.9 was recorded, Table 6.

Table 6. Sterile pyuria among 50 pregnant women at Tamale Central Hospital

N=50		
Age/yrs.	Frequency	%
< 20	3	9.1
20-29	24	72.7
30-39	5	15.2
40-49	1	3
Total	33 (66%)	100

Discussion

The study revealed prevalence of asymptomatic bacteriuria among pregnant women seeking antenatal care at Tamale Central Hospital and Tamale Teaching Hospital at 20% and 35.5%; a rate much higher than 5.5% and 7.3% respectively reported from Korle-Bu Teaching Hospital in Accra and the Komfo Anokye Teaching Hospital in Kumasi both in the southern sector of Ghana [22, 23]. Other similar studies have shown varying prevalence rates; 10% in Egypt [24]; 4-7% in Canada [25]; 13.3% in Uganda [26] and 7% in Ethiopia [27]. But comparable to our finding is the 28.8% documented in Ibadan Nigeria although higher rates of 63.3% and 86.6% have been recorded in two studies in the same country [28, 29]. Disparities in ASB prevalence within and between countries are assignable to sexual contact, socioeconomic levels, cultural and religious behaviours concomitant to personal hygiene [24]. In Ghana, socioeconomic levels and cultural practices differ from the south to the north and this could have accounted for the sharp difference in prevalence.

Asymptomatic bacteriuria seem to be predominant in women aged 20 and 30 years and less in age group < 20 years [24, 30]. This was in agreement with our results and others reported in other sectors of the country [22, 23]. The explanation to the susceptibility of these age groups could be early and intensive sexual intercourse which may result in minor urethral trauma and transfer organisms from the perineum into the bladder [31].

Contrary to most studies which have Gram negatives especially *E. coli* as the frequently isolated bacteria, this investigation found Gram positive *S. aureus* and coagulase negative *Staphylococcus* (CoNS) as the commonly recovered pathogens at both hospitals. Among the isolated Gram negatives, *E. coli* was dominant followed by *Enterobacter* sp. and *Klebsiella* sp. This could be imputed to the fact that CoNS and *S. aureus* are normally encountered in UTIs in the 20-30 year group and more than 64% of our study population came from this group. In Ghana, Gram positive bacteria appears to be conventional in pregnant women presenting with ASB as *S. aureus* and *Enterococcus* dominated in similar analysis in the country [23, 32]

Susceptibility of isolates to gentamicin and amikacin were highest which could possibly be the infrequent prescription of these drugs in the treatment of UTI infections at both hospitals as these drugs are only used in the management of severe and vulnerable cases (Personal communication). Low susceptibilities were found against ampicillin, cotrimoxazole, chloramphenicol, erythromycin and nitrofurantoin. Antibiotic pressure resulting from persistent use and abuse could have accounted for the high resistance levels to these drugs. A worrying observation was the 28.8% and 52% resistance recorded against imipenem and vancomycin as these drugs (carbapenem and glycopeptides) used to be highly effective against members of the enterobacteriaceae and gram positives and assessed as the last line of treatment drugs in managing difficult to treat MDR pathogens in these hospitals. Considering these outcomes empirical treatment will not be safe without laboratory confirmation of the susceptibility of all classes of antibiotics.

Sterile pyuria is a pervasive condition in both primary and secondary care settings but limited literature exist to provide estimated prevalence in the community or hospital setting. This situation has led to

inconsistent management ranging from absolute neglect of the finding to over investigation and unwarranted referrals [33]. At the Tamale Central hospital (secondary care facility), 66% of the 50 pregnant women screened had sterile pyuria which relates to the study of Shipman et al. [34] who reported 74% in women presenting to an emergency department in the United States.

This study had some limitations which are that, detailed information such as gestational ages of pregnancy, sexual activity which is considered a risk factor for developing UTI in women could not be captured. This is because most women in the study area do not feel comfortable giving out information on such details. Any attempt to persuade them normally led to their withdrawal from the study. But this lapse did not affect the purpose of the study which was to establish the prevalence of asymptomatic bacteriuria and sterile pyuria among pregnant women in secondary and tertiary care facility.

Conclusion

The prevalence of asymptomatic bacteriuria at the Tamale Central and the Tamale Teaching Hospital is 20% and 35.5% which is considerably higher than what has been found in the southern sector of Ghana. *Staphylococcus aureus* and coagulase negative Staph. are the commonest causative organisms. Due to the complications associated with asymptomatic bacteriuria, the study recommends that screening is done for all pregnant women, and the appropriate antibiotics prescribed for treatment. Sterile pyuria; a common but neglected entity was frequently encountered (66%) which may be a presentation of asymptomatic renal disease and therefore must be considered for all patients.

List Of Abbreviations

ASB-Asymptomatic bacteriuria

CLED- Cysteine lactose electrolyte deficient agar

CFU- Colony forming unit

TCH- Tamale Central Hospital

TTH- Tamale Teaching Hospital

CoNS- Coagulase negative *Staphylococcus*

MDR-Multidrug resistance

SP- Sterile pyuria

RPM-Revolution per minute

Declarations

Ethics approval and consent to participate

Ethics approval was obtained from the Ethical Review Committee of the Tamale Teaching Hospital (TTHERC/25/06/19/14) and verbal informed consent was sought from the pregnant women after sufficient information regarding the study was provided.

Consent for publication

Not applicable

Availability of data and material

We consider our data private but the corresponding author will make it available upon reasonable request.

Competing interests

The authors declare that they have no competing interests

Funding

None

Authors' contributions

All authors read and consented to the final draft of the manuscript. ABK was involved in the conception, study design and drafting of the manuscript, CKSS contributed in conception and drafting of the manuscript, DYY was involved in sample collection and processing as well as data analysis

Acknowledgements

Our thanks go to the laboratory staff of the Tamale Teaching Hospital and the Tamale Central Hospital for their support in the sample and data collection.

Author Affiliation

¹Department of Clinical Microbiology, School of Medicine and Health Sciences, University for Development Studies.

References

1. Najar MS, Saldanha CL, Banday KA. Approach to urinary tract infections. *Indian J Nephrol* 2009; 19:129–39.
2. Masinde A, Gumodoka B, Kilonzo A, Mshana SE. Prevalence of urinary tract infection among pregnant women at Bugando Medical Centre, Mwanza, Tanzania. *Tanzan J Health Res.* 2009; 11:154–9.

3. Litza JA, Brill JR. Urinary tract infections. *Prim Care*. 2010; 37:491–507.
4. Amiri M, [Lavasani Z](#), [Norouzirad R](#), [Najibpour R](#), et al. * Prevalence of Urinary Tract Infection Among Pregnant Women and its Complications in Their Newborns During the Birth in the Hospitals of Dezful City, Iran, 2012 – 2013. *Iran Red Crescent Med J*. 2015; doi: 10.5812/ircmj.26946
5. Alemu A, Moges F, Shiferaw Y, Tafess K, Kassu A, Anagaw B, et al. Bacterial profile and drug susceptibility pattern of urinary tract infection in pregnant women at University of Gondar Teaching Hospital, Northwest Ethiopia. *BMC Res Notes*. 2012; 5:197.
6. Emamghorashi F, Mahmoodi N, Tagarod Z, Heydari ST. Maternal urinary tract infection as a risk factor for neonatal urinary tract infection. *Iran J Kidney Dis*. 2012; 6:178–80.
7. Schults R, Read AW, Straton JAY, Stanley FJ, Morich P. Genitourinary Tract Infection in Pregnancy and Low Birth Weight: Case Control Study in Australian Aboriginal Women. *Brit Med J*. 1991; 73: 576-582.
8. Smaill F. Asymptomatic bacteriuria in pregnancy. *Best Pract Res Clin Obstet Gynaecol*. 2007; 21:439–50.
9. Duarte G, Marcolin AC, Quintana SM, Cavalli RC. Urinary tract infection in pregnancy. *Braz J Gynecol Obstet*. 2008; 30:93–100.
10. Leigh DA, Groneberg RN, Brumfitt W. Long term follow-up of Bacteriuria in Pregnancy. *Lancet*, 1968; 1: 603-605.
11. Wolfe AJ, Toh E, Shibata N, et al. Evidence of uncultivated bacteria in the adult female bladder. *J Clin Microbiol*. 2012; 50: 1376–1383.
12. Ronald RA. Urinary tract infections in adults. *Experts guide to the management of common infectious diseases*. 2002:229-50.
13. Nicolle LE, Bradley S, Colgan R, Rice JC, Schaeffer A, Hooton TM. Infectious Diseases Society of America; American Society of Nephrology; American Geriatric Society. Infectious Diseases Society of America guidelines for the diagnosis and treatment of asymptomatic bacteriuria in adults. *Clin Infect Dis*. 2005; 40:643-54.
14. Glen P, Prashar A, Hawary A. Sterile pyuria: a practical management guide. *Br J Gen Pract*. 2016; 66:225-7.
15. Wise GJ, Schlegel PN. Sterile Pyuria. *N Engl J Med*. 2015; 372:1048-1054
16. Alwall N, Lohi A. A population study on renal and urinary tract diseases. II. Urinary deposits, bacteriuria and ESR on screening and medical examination of selected cases. *Acta Med Scand*. 1973;194:529-35
17. Hooker JB, Mold JW, Kumar S. *J Am Bd Fam Med*. 2014; 27:97-103.
18. Smaill F. Antibiotics for asymptomatic bacteriuria in pregnancy. *Cochrane Database Syst Rev* 2001; CD000490.
19. Mittendorf R, Williams MA, Kass EH. Prevention of preterm delivery and low birth weight associated with asymptomatic bacteriuria. *Clin Infect Dis*. 1992; 14: 927-32.

20. Romero R, Oyarzun E, Mazor M, Sirtori M, Hobbins JC, Bracken M. Meta-analysis of the relationship between asymptomatic bacteriuria and preterm delivery/low birth weight. *Obstet Gynecol.* 1989; 73:576-82.
21. Cavalieri SJ, et al. Manual of antimicrobial susceptibility testing. American Society of Microbiology. 2005; 39–52.
22. Turpin C, Minkah B, Danso K, Frimpong E. Asymptomatic bacteriuria in pregnant women attending antenatal clinic at Komfo Anokye Teaching hospital, Kumasi, Ghana. *Ghana Med J.* 2007; 41: 26-9.
23. Labi AK, Yawson AE, Ganyaglo GY, et al. Prevalence and associated risk factors of asymptomatic bacteriuria in ante-natal clients in a large teaching hospital in Ghana. *Ghana Med J.* 2015; 49:154–8.
24. Abdel-Aziz Elzayat M, Barnett-Vanes A, Dabour MFE, et al. Prevalence of undiagnosed asymptomatic bacteriuria and associated risk factors during pregnancy: a cross-sectional study at two tertiary centres in Cairo, Egypt. *BMJ Open* 2017; doi:10.1136/bmjopen-2016- 013198
25. Nicolle LE. Screening for Asymptomatic Bacteriuria in Pregnancy. *Canadian Guide on Preventive Health Care*, Ottawa Health, Canada. 1994; 100-106.
26. Andabati G, Byamugisha J. Microbial aetiology and sensitivity of asymptomatic bacteriuria among ante-natal mothers in Mulago hospital, Uganda. *Afr Health Sci.* 2010; 10:349–52.
27. Gabre-Selassie S. Asymptomatic Bacteriuria in Pregnancy; Epidemiological clinical and microbiological approach: *Ethiopia Med J.* 1998; 36: 185-192.
28. Akerele P, Abbulire F, Okonofua J. Prevalence of Asymptomatic Bacteriuria among Pregnant Women in Benin City, Nigeria. *J. Obstetric Gynaecol.* 2001; 21: 141-144.
29. Okon K, Nkwaku L, Balogun ST, Usman H, Adesina OO, Akuhwa RT, Uba A, Shidali NN. Antimicrobial Susceptibility Profile of Bacterial Pathogens Isolated From Pregnant Women with Asymptomatic Bacteriuria at Tertiary Hospital in Northeastern Nigeria. *Sierra Leone Journal of Biomedical Research.* 2012; 4:32-42.
30. Sujatha R, Nawani M. Prevalence of asymptomatic bacteriuria and its antibacterial susceptibility pattern among pregnant women attending the antenatal clinic at Kanpur, India. *J Clin Diagn Res.* 2014; 8:2–4.
31. Jalali M, Shamsi M, Roozbehani N, et al. Prevalence of urinary tract infection and some factors affected in pregnant women in Iran Karaj City 2013. *Middle East J Sci Res.* 2014; 20:781–5.
32. Addo VN. Asymptomatic bacteriuria and drug susceptibility patterns of mid-stream urine specimens among pregnant women at booking in a private hospital in Kumasi, Ghana. *Postgraduate Medical Journal of Ghana.* 2014; 3:82-86.
33. Rees J, Manley J. Assessment of sterile pyuria in primary care. *British Journal of Family Medicine.* 2015; 3:3.
34. Shipman SB, Risinger CR, Evans CM, Gilbertson CD, Hogan DE. [High Prevalence of Sterile Pyuria in the Setting of Sexually Transmitted Infection in Women Presenting to an Emergency Department.](#) *The Western Journal of Emergency Medicine.* 2018; doi: 10.5811/westjem.2017.12.35605.