Experience of Nutritional Counselling in a Nutritional Program in HIV Care in the Tigray Region of Ethiopia Using the Socio-ecological Model

Fisaha Haile Tesfay (✉ fisaha.1999@gmail.com)

Mekelle University College of Health Sciences <https://orcid.org/0000-0003-0399-1711>

Anna Ziersch

Royal Society for Public Health

Lillian Mwanri

Flinders University Faculty of Medicine Nursing and Health Sciences

Sara Javanparast

Flinders University Faculty of Medicine Nursing and Health Sciences

Research article

Keywords: Nutritional counselling, Nutritional program, Undernutrition, HIV, Tigray, Ethiopia

Posted Date: September 4th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-64812/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Journal of Health, Population and Nutrition on July 28th, 2021. See the published version at <https://doi.org/10.1186/s41043-021-00256-9>.

Abstract

Background: In many resource-poor settings, nutritional counselling is one component of nutrition support programs aiming to improve nutritional and health outcomes amongst people living with HIV. Counselling methods, contents, and recommendations that are culturally appropriate, locally tailored, and economically affordable are essential to ensure desired health and nutritional outcomes are achieved. However, there is little evidence showing the effectiveness of counselling in nutritional programs in HIV care, and the extent to which counselling policies and guidelines are translated into practice and utilised by people with HIV suffering from undernutrition. This study aimed to explore these gaps in the Tigray region of Ethiopia.

Methods and participants: A qualitative study was conducted in Tigray Region Ethiopia between May and August 2016. Forty-eight individual interviews were conducted with 20 undernourished adults living with HIV and 15 caregivers of children living with HIV enrolled in a nutritional program in three hospitals, as well as 11 health providers, and 2 program managers. Data analysis was undertaken using the Framework approach and guided by the socio-ecological model. Qualitative data analysis software (QSR NVivo 11) was used to assist data analysis. The study findings are presented using the consolidated criteria for the reporting of qualitative research (COREQ).

Result: The study highlighted that nutritional counselling as a key element of the nutritional program in HIV care varied in scope, content, and length. While the findings clearly demonstrated the acceptability of the nutritional counselling for participants, a range of challenges hindered the application of counselling recommendations in participants' everyday lives. Identified challenges included the lack of comprehensiveness of the counselling in terms of providing advice about the nutritional support and dietary practice, participants' poor understanding of multiple issues related to nutrition counselling and the nutrition program, lack of consistency in the content, duration and mode of delivery of nutritional counselling, inadequate refresher training for providers, and the absence of socioeconomic considerations in nutritional program planning and implementation. Evidence from this study suggests that counselling in nutritional programs in HIV care was not adequately structured and lacked a holistic and comprehensive approach.

Conclusion: Nutritional counselling provided to people living with HIV lacks comprehensiveness, consistency, and varies in scope, content, and duration. To achieve program goal of improved nutritional status, counselling guidelines and practices should be structured in a way that takes a holistic view of patient's life and considers cultural and socioeconomic situations. Additionally, capacity development of nutritional counsellors and health providers is highly recommended to ensure counselling provides assistance to improve the nutritional wellbeing of people living with HIV.

Introduction

There is strong evidence on the interaction between undernutrition and HIV infection, with impacts at individual and community levels. HIV infection and AIDS increases people's predisposition to undernutrition [1]. Due to clinical, social, and economic factors, people living with HIV are more likely to suffer from undernutrition and its consequences [1-3]. Estimates of the prevalence of undernutrition among people living with HIV in sub-Saharan Africa varies – for example from 19.4% in Tanzania to [4] to 42.5% in Ethiopia [5]. International health agencies including the World Health Organization (WHO) recommend nutritional care as an important component of HIV care services, particularly in resource-poor settings [6, 7].

Evidence suggests that a combination of antiretroviral therapy (ART) and nutritional care can improve adherence to ART and nutritional wellbeing. Ivers and colleagues found that adults living with HIV had better adherence to ART at 6 and 12 months of using nutritional supplementation respectively than those who were not provided with food assistance [8]. In addition, there is strong evidence on the benefits of nutritional support in terms of improving the nutritional status of people living with HIV [9-11].

To address HIV-related undernutrition and to improve HIV care outcomes, nutritional programs have been incorporated into HIV care services in many countries, particularly in sub-Saharan Africa [11-14]. Nutritional programs mainly comprise three components: 1) nutritional assessment; 2) nutritional support; and 3) nutritional counselling. A nutritional assessment can involve a range of parameters but most commonly includes nutritional screening which measures key anthropometric indicators to guide the provision of different therapeutic and supplementary foods [11, 12, 15]. Food-based supplements such as high-energy ready-to-use therapeutic foods (RUTF) and corn-soy-blended (CSB) are the most common nutritional supports used in HIV care settings to treat protein-energy undernutrition [16]. The third component of nutritional programs is nutritional counselling. The general aim of nutritional counselling is to assist patients to: a) maintain weight through increased energy intake; b) practice safe infant feeding; c) practice safe food or water handling; and d), manage HIV-related illness [15, 17].

Nutritional counselling plays a critical role in improving individuals' nutritional knowledge and understanding of the importance of nutrition in their lives [18]. Such engagement through individual or group counselling sessions can empower patients to make informed decisions regarding their eating practice based on their financial circumstances and food availability and to take control over their nutritional circumstances [18, 19]. Culturally appropriate nutritional counselling involves tailored content to the local cultural context, incorporating local knowledge, expertise and lifestyle choices, distributing locally and culturally appropriate food and adapting the nutritional counselling to the local people's educational level [19, 20]

The evidence base regarding the availability and practice of nutritional counselling in HIV care is mixed. While a study in sub-Saharan African countries found the inclusion of nutritional counselling in 95% of the study sites [21], another study from Ghana reported a lack of nutritional counselling in HIV care programs [22].

Evidence on the role of nutritional counselling in improving the nutritional status of people living with HIV is varied [23-25]. Some studies support the importance of nutrition counselling [24] but others are inconclusive about the benefits of counselling for the management of nutrition in HIV care. For example, a study conducted in Nigeria found that people living with HIV who were offered regular nutritional assessment and counselling had better nutritional outcomes than those who were not [25]. In contrast, a study from Brazil had shown that the provision of nutritional counselling to undernourished people living with HIV who had tuberculosis showed no improvement in their nutritional status [23], though tuberculosis infection might have undermined the positive impact of nutritional counselling. In a study from Uganda, nutritional counselling was found to be effective in improving the overall quality of life of people living with HIV, when given as a package of nutritional and other interventions such as nutritional support to [26].

There is little evidence on the best techniques of counselling such as individual vs groups, frequency and duration of counselling, and the possible roles of counsellors such as peers, nurses, physicians, and other auxiliary health workers in HIV care settings [27].

The nutritional program in HIV care in Ethiopia

The nutritional program in HIV care settings in Ethiopia commenced in 2010 and involves three main components to address the nutritional challenges of people living with HIV [12]. The program involves nutritional assessment for all people living with HIV on ART and pre-ART, classification of nutritional status, and enrolment of those undernourished into a nutritional program. The nutritional support program provides therapeutic food (Plumpynut) for six months for severely acute undernourished (SAM) and supplementary food (Plumpysup) for three months in moderately or mildly undernourished people living with HIV. Nutritional counselling is also one of the key components of the nutritional programs in HIV care in Ethiopia, similar to other areas in sub-Saharan Africa [11].

In Ethiopia according to the national guidelines, nutritional counselling involves the provision of general information to people living with HIV who are enrolled in the nutritional program as stated in the nutritional guideline [28]. The information includes how to: 1. conduct a periodic nutritional assessment at a health facility; 2. increase intake and dietary diversification; 3. maintain a good level of hygiene and sanitation; 4. increase clean and safe water intake; 5. maintain a healthy lifestyle; 6. early treatment and diagnosis of illness and symptoms; and 7. adherence to advice by a health provider on how to take medications and manage dietary needs. Despite the potential benefits of nutritional counselling to people living with HIV, there is little evidence about the practice and experiences of nutritional counselling in HIV care settings from the perspectives of beneficiaries, service providers, and program managers. A study conducted in Ethiopia examined the availability of nutritional education in a nutritional program in HIV care but failed to examine the experience of various stakeholders of the nutritional counselling [29]. This paper presents the experiences of different stakeholders on nutritional counselling in HIV care settings in

Ethiopia, with a view to help inform policy on how to effectively address the challenges of nutritional counselling for people living with HIV.

The Socioecological model

This study draws on the socioecological model (SEM) that can assist an understanding of a problem at multiple levels [30]. The SEM is a theory-based framework employed to comprehend the interaction of individual and environmental factors that impact health behaviours and outcomes [31, 32]. By highlighting that are problems are not solely explained by individual behaviour, the SEM helps to understand the dynamics and interactions between individual, interindividual, institutional, community, and social policy level factors in influencing health outcomes and behaviours [33]. The SEM has been applied in studies involving nutritional programs in HIV care, sexual and reproductive health among people living with HIV, adherence to ART, and acceptability of infant feeding to prevent HIV [34-37].

This paper sought to examine experiences of the nutritional counselling program in nutritional programs in HIV care settings in the Tigray region, from the perspectives of program users, staff and managers, to identify strengths and challenges of the nutritional counselling that is offered in HIV care.

Methodology

Study design, Sampling, and recruitment:

This study is part of a larger study that examined the determinants of nutritional outcomes and challenges of a nutritional program in the Tigray region of Ethiopia [37, 38]. This paper presents findings from a qualitative study involving individual interviews with adults living with HIV, caregivers of children living with HIV, health providers, and health managers to explore their experiences and perspectives about nutritional counselling in the nutritional program in HIV care settings.

The study was conducted in three hospitals namely: Mekelle, Shul, and Lemlem Karl hospitals in the Tigray region, northern Ethiopia. Twenty percent of people living with HIV on ART and 22% of those enrolled in the nutritional program in the Tigray region accessed HIV and nutrition related services from these hospitals.

Study participants consisted 20 adults and 15 caregivers of children living with HIV enrolled in the nutritional program. In addition, 11 health providers and two program managers were interviewed. Details of the study setting are described elsewhere [38].

Data collection procedure:

The first author (FT) conducted face-to-face in-depth interviews with participants. There was no relationship between the interviewer (first author) and the study participants prior to the interview date

and participant information sheet and consent form were the only source of information for participants to know about the research interviewer (FT). A semi-structured interview guide was developed (in English) and the interview guide was field-tested in a similar setting and adjustment was made based on the pre-test. The interview guides are attached as a supplementary file (supplementary file 1). To induce open discussion on the topics and maintain the privacy of the study participants, the interview was conducted in a private room inside the health facility. All interviews were conducted in the local Tigrigna language by directly translating the interview guide to the local language during the interview process by the first author (FT). He is a native speaker of the local Tigrigna language and fluent in English with a good understanding of the study context and culture. All in-depth interviews were audio-recorded using a high definition audio recorder. The duration of the in-depth interview ranged from 30-66 minutes. In addition, observation and field notes were taken during the field stay.

Data analysis:

Interviews were translated and transcribed verbatim from the Tigrigna local language to English by the first author (FT). A translation and transcription accuracy test was done to improve the quality of translation from the local language to English using a person who was fluent in the Tigrigna local language and English, with accuracy rated as high.

Data were analysed using a thematic framework analysis approach [39]. Interview transcripts were read and re-read to gain a holistic understanding of the range of responses and scope of counselling experiences. A list of codes was developed based on four transcripts which served as a framework for analysis and 11 nutritional counselling related working codes were generated. English translated transcripts were imported to NVivo 11 software for analysis. In the second phase of coding, the number of codes that emerged from all transcripts was increased to 14. Finally, these codes were merged, categorised and re-categorised to develop four themes to answer the research question. Coding and data analysis were done inductively by one person, and three transcripts were double coded by co-authors and disagreements were discussed in a team meeting to reach consensus. The socioecological model was used to guide the overall data analysis, presentation, and interpretation of findings.

Results

Characteristics of study participants

Participants' mean ages with standard deviation was 37.2 ± 9.7 for adults living with HIV (hereafter 'adults'), 36 ± 7.3 for caregivers of children living with HIV (hereafter 'children'), and 35.3 ± 8 for program staff. All caregivers and 12(60%) of adult study participants interviewed were female.

The majority of adults (60%) and caregivers (73%) were urban residents. While four (20%) and five (33%) of adults and caregivers respectively attended no school, 10(50%) and five (33%) adults and caregivers respectively had attended primary school. The average income in adults and caregivers ranged from no

reliable income (N=9,4 caregiver and 5 adults) to 6,000 (N=3) (equivalent to 458 Australian Dollars) per month which is much lower than the average annual per capita income in Ethiopia (857 Australian dollars) for the year 2018. At the time of the interview, six (30%) adults were single and six (30%) married, and nine (60%) of caregivers were married. An equivalent number of males and females were represented when it comes to health care providers and program managers, and they all had achieved education level at BSc and above (**Table 1**).

Table 1
Sociodemographic characteristics of study participants

Variables		Participants		
		Adults, n=20	Caregivers, n=15	Program staff, n=13
Mean age	Mean age	37.2±9.7	36±7.3	35.3± 8
Hospital	Mekelle	8	6	5
	Lemlem Karl	6	5	3
	Shul	6	4	3
	TRHB	N/A*	N/A	2
Gender	Male	8	0	6
	Female	12	15	7
Residence	Urban	12	11	13
	Rural	8	4	0
Educational level	No education	4	5	0
	Primary	10	5	0
	Secondary	6	5	0
	BSc and above	0	0	13
Marital status	Single (never married)	6	1	2
	Married	6	9	9
	Widowed	5	3	0
	Divorced	3	2	1

Experience of nutritional counselling in the nutritional program.

Four main themes that emerged from the interview data in relation to experiences of nutritional counselling in the nutritional program are summarised in **table 2** and discussed in detail below.

Table 2
Themes emerged from the qualitative analysis

Theme	Theme description
Frequency and duration of the nutritional counselling	This relates to the frequency of nutritional counselling over the course of participants' enrolment in the nutritional program. Duration indicates the length of one nutritional counselling session.
Contents of the nutritional counselling	This refers to the actual topics discussed during the nutritional counselling
Acceptability of the nutritional counselling	This theme relates to the acceptability of the nutritional counselling and recommendations by participants
Applicability of the nutritional program	Refers to the actual implementation of the nutritional recommendations given by the health professional
Challenges/barriers in relation to nutritional counselling	This theme identified the challenges of nutritional counselling in HIV care from the perspectives of study participants including applicability

Frequency and duration of the nutritional counselling

Frequency refers to the number of times patients received nutritional counselling during enrolment in the nutritional program. Session duration refers to how long one specific nutritional counselling session lasted.

Comments received about the frequency of nutritional counselling ranged from 'no counselling at all to more than one counselling. The majority of study participants noted that the nutritional counselling was provided as a one-off session, commonly at the beginning of the nutritional program.

...At the beginning when they give you [the nutritional support], they [the health providers] tell you that it is a medication: that is all, nothing else after... (Adult male, #7).

A small number of adults and caregivers reported receiving no nutritional counselling at all and said that they had been sent home only with the nutritional support:

Maybe since I am coming from far away, I may not reach the teaching session. Otherwise, I have not yet come across any counselling or teaching session (Adult female, #6).

We did the counselling at each visit. For example, when we have many clients. We send them to the case manager [peer counsellors] for counselling about ART adherence, cleanliness, or personal hygiene and so on" (Health provider #4)

Contents of the nutritional counselling as experienced by participants

Even though the nutritional counselling is meant to cover the seven elements mentioned in the introduction [40], participants reported that the nutritional counselling focused on three elements : a) nutritional support (Plumpynut/Plumpysup); b) dietary practice; and c) sanitation and hygiene, with a particular emphasis on the nutritional support.

In general, the nutritional counselling mainly focused on the nutritional support (Plumpynut/Plumpysup); including why the nutritional support is given and its benefits as stated by a caregiver below:

They [health providers] told me that my child's weight has been decreased and told me to give him the Plumpynut properly. They said, "this [the plumpynut] will improve his weight as well as general health status and told me to follow him properly"(Caregiver #5).

An adult female also reiterated the focus of the nutritional counselling around the nutritional support, specifically on the benefits of the nutritional support in terms of improving weight and wellbeing.

They [health providers] counsel me that the Plumpynut contains important nutrients including Plumpynut and vitamins. They told me that it is very important and beneficial for your weight and overall health (Adult female #1).

Discouraging patients to share and sell the nutritional support were also stated as part of the nutritional counselling regarding the nutritional support.

The health providers also advised us to take it (the Plumpynut) properly and not to share it with others. Selling, sharing or giving it to other persons is not allowed (Adult male #6)

The second content focus of the nutritional counselling identified in HIV care related to dietary practice and increasing the frequency and diversity of foods to improve nutritional wellbeing. Even though the lack of access to adequate food was a problem in many participants, counselling to diversify dietary practice with whatever food available at home was another element of the nutritional counselling.

They counsel us about everything. For instance, they tell us 'it is not only meat that we [herself and her child] should take, we can also take other foods like grains and vegetables which are equally important (Caregiver #4).

We counsel them how their dietary pattern looks like and they are counselled to take balanced foods (Health provider #2).

The third component covered in the counselling sessions was on mechanisms to maintain hygiene and food safety including handwashing, hygienic handling of the nutritional support and other foods as well as consuming cooked foods:

They [health providers] tell us to maintain our personal hygiene and environmental sanitation, to wash our hands before use of the nutritional support and to use Woha Agar^[1] to treat water (Caregiver #2).

Hygiene, to drink clean and safe water and to use Woha Agarto treat the water or boil it (water) before use, otherwise I didn't remember any specific counselling session (Adult male #6).

The health providers advised us to eat balanced food, drink clean water, and seek care for any sickness (Adult female, #4).

In a small number of instances, during the nutritional counselling other topics such as minimizing stress to improve overall health, wellbeing, and quality of life and ART adherence were also covered.

About our dietary practice, they told us to take foods that go with the ART medication such as eggs, meat, and milk. As I told you about that, if you have enough you will eat, if not you will use whatever you have with the ART (Caregiver #16).

Acceptability of the nutritional counselling

"Acceptability refers to determining how well an intervention will be received by the target population and the extent to which the new intervention might meet the needs of the target population and organizational setting"[41].

The majority of caregivers and adults felt that the nutritional counselling they received was acceptable and benefited them in terms of providing information about the nutritional support.

Since I understand the benefits of the Plumpynut, I believe that the counselling service has benefited me to take the Plumpynut accordingly (Adult male #6).

... counselling is very important. It helps me to understand its [the nutritional support] benefits to him [the child]) and use it properly, on time and get the necessary next ration (Caregiver #1).

Most adult and caregiver participants reported improved knowledge about food and nutrition as the benefit of the nutritional counselling, as described by an adult male below:

If you are taking the ART medication or the Plumpynut accepting it will benefit you, you will get the intended benefits. All the counselling is good for me for my weight and health. I fully practice the

counselling provided because I understand that all the benefits of counselling are to improve my weight. When they counsel you, they [the health providers] are giving your life not to get sick and weaker (Adult male, #13).

Sharing the information received from nutritional counselling with other adults and caregivers was an indication of a high level of acceptability amongst people living with HIV. This was noted by an adult male:

As a volunteer, I also transfer information to people who are taking Plumpynut not to share it instead to consume it to themselves (Adult male #5).

I also teach a mother who have children who are taking the Plumpynut not to share it with others (Caregiver #3).

Challenges for the provision of nutritional counselling in HIV care setting

A number of challenges were identified for the provision and implementation of nutritional counselling including a lack of understanding by recipients, training issues and inconsistent delivery, and a lack of consideration of the context of people's lives.

Despite the reported acceptability of the nutritional counselling by most adults and caregivers, some health providers said that nutritional counselling was not well implemented by some patients, particularly by older adults. It was believed that this may be partly due to a limited understanding of the nutritional counselling:

This is because we may not provide them with adequate counselling as well most HIV patients are uneducated which may result in poor or slow understanding of the counselling service (Health providers #1).

We give them similar counselling service to all patients, but some people understand you quickly and others didn't understand you at all despite the intensity of counselling. Some of those who didn't accept and apply the counselling are those old and uneducated ones who may not pay it adequate attention (Health provider #2).

Health providers also noted a range of issues relating to the training of staff and training materials and inconsistent delivery of the nutritional counselling. The lack of training of health providers related broadly to the general nutritional program provided in the HIV care service, with implications also for nutritional counselling:

Issue related to the health system, one is the professional. In order to give good service; their training should also be good. The health professional should know the contents of the two food supplements.

The knowledge of the health provider is low (Health provider #10)

“There are problems related to the health provider especially if they didn’t take training about food by prescription (Health provider #4).

Adults further reported that there was no detailed assessment of household socioeconomic status and family situation during the nutritional counselling but underlined that they have been told to eat whatever food available at home. An adult female described the lack of socioeconomic consideration of the nutritional counselling below:

They [the health providers] didn’t say about this [household socioeconomic consideration]. They didn’t ask about what we have and haven’t. They [health providers] only tell us to diversify our dietary practice. There is nobody who asks about our household situation (Adult female #19).

A lack of detailed assessment of household socioeconomic status and socioeconomic consideration of the nutritional counselling constrained the acceptability, applicability, and adherence of the nutritional counselling in HIV care setting with poor access to adequate food, for some participants, a key barrier to fulfilling the key messages promoted in the nutritional counselling:

About my dietary practice, I think you have to eat what you get at your home; otherwise, you can’t live according to the counselling given here in this health facility (caregiver #16).

A small number of caregivers did report the experience of assessment of their household socioeconomic status during the nutritional counselling as stated below:

It [household socioeconomic status] is considered. They ask me and write it down on my card and they have done everything. They know about my income and my job as well as other things about my family (Caregiver #3).

These overall challenges associated with nutritional counselling were seen by health providers to have a significant impact on program outcomes:

Had it been that they apply all the counselling services given to them like the ART medication, they may get the necessary benefits from the nutritional program: as well we may not have default or loss to follow up from the nutritional program (Health provider #6).

[1] Woha Agar is a water treatment chemical used in Ethiopia mainly distributed by health providers to people living with HIV

Discussion

Nutritional counselling in HIV care settings seeks to improve the effectiveness of nutritional programs in HIV care via improving knowledge and nutritional program utilisation. Research has indicated that locally and culturally sound nutritional counselling is vital to enhance and maintain the nutritional status of

people living with HIV [42–44]. Engaging with participants and empowering them to take control of their nutrition and health is the fundamental principle of successful nutritional counselling in HIV care settings [14]. Evidence from other studies suggests that standard nutritional counselling should empower individuals with essential nutritional knowledge and assist them to understand the importance of nutrition to make informed decisions regarding their nutritional options [19, 20].

The importance of nutritional counselling is stipulated in the Ethiopian national nutritional program [45]. However, despite evidence highlighting the importance of comprehensive nutritional counselling, this study has shown that counselling as one of the key components of nutritional program had a narrow focus, largely limited to a simple one way provision of information. One previous study from Ethiopia has highlighted the benefits of nutritional counselling in HIV care but did not provide detailed information on the level of engagement of people living with HIV in the counselling sessions [46]. Generally, there was no evidence of true engagement with patients, empowerment and patient's involvement in decisions made/counselling provided to people living with HIV.

While the technical guidelines for nutritional counselling recommends covering seven elements [40], the current study found that the nutritional counselling in the HIV care settings examined covered three elements—counselling about the nutritional support, dietary diversification, and food and personal hygiene. Among the three elements, the main focus was on the nutritional support, which led people living with HIV to consider and relate the nutritional program in HIV care only to the nutritional support. This is not in line with the objectives of the nutritional counselling which is more holistic and comprehensive [14]. This finding is consistent with other studies reporting a focus of the nutritional counselling on the nutritional support among HIV positive children enrolled in a nutritional program [47]. The study found there was more limited coverage of dietary diversification and, in those cases where there was, no or limited consideration of household socioeconomic status. There was even less coverage of personal and food hygiene.

Disproportional emphasis on the nutritional support was mainly related to the medicalised orientation of the nutritional program in general and the nutritional counselling in particular. Medical orientation refers to a focus of the nutritional program on treating medical malnutrition [48, 49]. The medicalised orientation of the nutritional counselling may contribute to the lack of a holistic approach – limiting the nutritional counselling to clinical malnutrition. A similar study from Kenya demonstrated that education about foods that were suitable for health and nutritional wellbeing was not given enough emphasis during the counselling of people living with HIV [50]. This highlights the need for reorientation of the nutritional counselling in HIV care settings toward a holistic approach.

According to the findings of this study, the nutritional counselling on the management of nutritional requirements was well received by patients, providing them with information on the nutritional supplements. Health providers did identify less educated and older people as having potential difficulties understanding the counselling. A study by Tobi and Voge found a lack of understanding of the nutritional counselling service reported as a barrier. [51]. Other studies have found that nutritional counselling

improved understanding of the nutritional program and their nutritional management of people living with HIV [14, 24, 25]. For instance, peer-delivered nutritional counselling among people living with HIV was found to improve dietary practice and food security [24]. Furthermore, a study from Nigeria also reported that nutritional counselling was effective in improving the nutritional management of people living with HIV [25]. Despite contextual differences, individualised nutritional counselling about dietary diversification and improved consumption was also deemed to improve the nutritional management of people living with HIV enrolled in a nutritional program as highlighted by a study from India [52]

There is no evidence on the specific duration of nutritional counselling to bring the required behavioural change among people living with HIV. However, inadequate nutritional counselling session duration was one of the key challenges demonstrated in this study - with most participants reporting only one counselling session and health providers indicating that session times in some contexts are too short. Other studies have also reported a short duration of nutritional counselling sessions. For example, a study by Tafesea and his colleagues which was conducted in Ethiopia indicated that a health provider counselled about 38 clients in a day and the average time spent on counselling a client was 3.26 minutes, which is very short to cover the contents and deliver key nutrition information [29]. Another study conducted in Cambodia among caregivers of children living with HIV also highlighted that counselling about the nutritional support was too short in duration [47]. Other studies suggest that shorter sessions also means less focus on prevention [53, 54].

Health providers who provide nutritional counselling need specialised training and knowledge in relation to culturally sensitive foods, disease progression, ART medication, and complications [51, 55]. However, a lack of continuous standardised training of health providers on nutrition for people living with HIV was identified and this finding is consistent with the findings of studies conducted in Ethiopia [29, 56]. A similar study by Kolasa and Rickett indicated that health providers lacked the necessary knowledge and skill to deliver brief and evidence-based nutritional counselling for people living with HIV [57]. This was also reflected in a lack of consistency in providing the nutritional counselling between health providers.

While the national policy framework in Ethiopia indicates the need for nutritional counselling [45], [40], despite, the need for clear technical nutritional counselling guidelines with clear outcomes in the nutritional program there were no clear and specific operational guidelines that guide the delivery strategy, session duration, and other technical issues related to the nutritional counselling except a simple pamphlet which suggests some seven key nutritional information for people living with HIV (see the introduction) [40]. In addition to the absence of technical guidelines, there was no training guideline for health providers. Despite the contextual differences, similar problems in relation to the training of health providers regarding the nutritional counselling were reported from the US [59].

A key issue also identified as a challenge for nutrition counselling programs was a lack of focus on the context of people's lives. This study found that food insecurity and poverty were a key issue among people living with HIV enrolled in the nutritional program [38] that affected their capacity to implement some elements of the counselling. Other studies in Kenya [50], South Africa, and Ghana [51, 58] have

likewise demonstrated the importance of considering the individual patient's financial and socioeconomic circumstances when providing the nutritional counselling for better effectiveness. For nutritional counselling to be effective, it should be tailored around the needs of people living with HIV.

The findings were interpreted using the socioecological model [30] and this indicated that effective nutritional counselling can be influenced by multilevel factors that interact with each other. At an individual level [60, 61], this can be influenced by the knowledge, cognition, and acceptability of the nutritional counselling of the individual patient. For instance, experiences of the acceptability of the nutritional counselling operated as an individual level factor that impacts the effectiveness of the nutritional counselling in HIV care setting. Acceptability can be further explained by the interaction of individual-level issues such as lack of understanding and social policy levels factors such as food insecurity and poverty [35, 62].

Inter-individual level factors such as lack of socioeconomic consideration of the nutritional counselling was a community-level factor which highly interacts with social policy level factors such as poverty and food insecurity [63]. The nutritional counselling was implemented in a broader socioeconomic environment where poverty and food insecurity was common [60, 64].

Nutritional counselling can also be an institutional level factor [60], where institutional structure, culture, and operational issues were important determinants of the nutritional counselling and outcomes. For instance, the nutritional counselling in the HIV care setting lacked standardised training for health providers as reflected by inconsistencies in the delivery of the counselling among and within the health facilities included in the current study. This could be related to health providers' skills and knowledge about the nutritional counselling and the lack of goal-oriented and individual problem-based focus of the nutritional counselling. Institutional level factors such as offering counselling session on an one-off basis, short duration of each session, lack of nutritional counselling guidelines, and lack of ongoing training of health providers coupled with policy level factors such as the medicalised orientation of the nutritional counselling negatively impacted the effectiveness of the nutritional counselling.

There is also an interaction between the institutional level factors which include, absence of structured and goal-oriented counselling and policy gap due to the lack of counselling guideline which responds to the individual needs as stipulated in the socio-ecological model [30].

The study was able to canvass the views of the nutritional counselling of people living with HIV and carers of children with HIV, supplemented by interviews with service providers and nutritional program managers. However, there are a number of potential limitations to the study. There may be issues around maintaining the original meaning of the data during translation and transcription as narrated by the study participants, but the maximum effort was made to maintain accuracy. Although issues of confidentiality were emphasised prior to and during interviews, some participants were concerned about the use of an audio recording device. This may have had an impact on participant responses. The issue of a man interviewing a woman may also constrain some participants to detail some views. However, the interviewer was very mindful in the interviews of being sensitive to this.

Conclusion And Implications

Compared to national guidelines, we found that nutritional counselling in HIV care is characterised by limited scope and is oftentimes provided as one-off session usually at enrolment to the nutritional program. The other challenges of the nutritional counselling were lack of comprehensiveness, consistency, and regularity, difficulties of some recipients in understanding the recommendations, and a failure to respond to the circumstances of people's lives.

Comprehensive nutritional counselling that is locally tailored and takes individual patients' needs and their socio-economic factors into consideration is crucial to improve nutritional health and wellbeing amongst people living with HIV. The development and improvement of national guidelines on nutritional counselling should include strategies and mechanisms that facilitate their translation into practice. Health providers' training, and reorientation of the nutritional counselling towards a holistic approach that engage and empower patients is also crucial to enhance the nutritional wellbeing of people living with HIV.

Declarations

Ethics approval and consent to participate:

Ethical approval was obtained from the Social and Behavioural Science research committee (SBRC) with ethics Approval n^o 718 of Flinders University, Australia, and from Ethics approval committee of Mekelle University, Ethiopia with approval n^o ERC 0621/2016.

Consent for publication:

Not applicable.

Availability of data and material:

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests:

The authors declare that they have no competing interests.

Funding:

This is a manuscript taken from a PhD thesis of the first author (FT) and Flinders University of South Australia funded the stipend and field trip to Ethiopia. Flinders University has no role in the design, collection, analysis, interpretation, and writing of this manuscript.

Authors' contributions:

FT has designed the study, collect, analyse and interpreted the data as well as drafted the manuscript. SJ, AZ, and LM critically reviewed the design, analysis, and interpretation of data, and critically reviewed the manuscript.

Acknowledgements:

This manuscript is taken from PhD thesis, and we would like to acknowledge Flinders University of South Australia for financing our study (student support payment and fieldwork) under the Australian Government Research Training program during the PhD candidature of the first author (FT). We would also like to express our acknowledgments to Mekelle University, Tigray Regional Health Bureau, and the study hospitals (Mekelle, Shul, and Lemlem Karl) for providing all administrative support and facilitating the fieldwork. The qualitative study participants also deserve a special appreciation for providing valuable information used in the current study.

References

1. Piwoz EG, Preble EA. **HIV/AIDS and nutrition: a review of the literature and recommendations for nutritional care and support in sub-Saharan Africa**. 2000.
2. nshait' NSS, SanGiovanni JuP. Synergism of nutrition, infection, and immunity: an overview. *Am J C/ii Nuir*. 1997;66:464S–477S.
3. Villamor E, Manji K, Fawzi WW: **Human Immunodeficiency Virus Infection**. In: *Nutrition and Health in Developing counreis Volume Second Edition* edn. Edited by Semba R, Bloem M. USA: Humana Press; 2008.
4. Kabalimu TK, Sungwa E, Lwabukuna WC. **Malnutrition and associated factors among adults starting on antiretroviral therapy at pasada hospital in Temeke district, Tanzania**. *Tanzania Journal of Health Research* 2018, 20(2).
5. Berhe N, Tegabu D, Alemayehu M. Effect of nutritional factors on adherence to antiretroviral therapy among HIV-infected adults: a case control study in Northern Ethiopia. *BMC Infect Dis*. 2013;13:233.
6. WHO. **Nutrient requirements for people living with HIV/AIDS**. 2003.
7. World Food Program. Organization WH. UNAIDS.: **HIV/AIDS, Food security and Nutrition**. In.; 2008.
8. Ivers LC, Chang Y, Jerome JG, Freedberg KA. Food assistance is associated with improved body mass index, food security and attendance at clinic in an HIV program in central Haiti: a prospective

- observational cohort study. *AIDS research therapy*. 2010;7(1):33.
9. Bahwere P, Deconinck H, Banda T, Collins S. Effective therapeutic feeding with chickpea sesame based ready-to-use therapeutic food (CS-RUTF) in wasted adults with confirmed or suspected AIDS. *World Journal of AIDS*. 2011;1(04):169.
 10. Olsen MF, Abdissa A, Kæstel P, Tesfaye M, Yilma D, Girma T, Wells JC, Ritz C, Mølgaard C, Michaelsen KF. Effects of nutritional supplementation for HIV patients starting antiretroviral treatment: randomised controlled trial in Ethiopia. *BMJ*. 2014;348:g3187.
 11. Nagata JM, Cohen CR, Young SL, Wamuyu C, Armes MN, Otieno BO, Leslie HH, Dandu M, Stewart CC, Bukusi EA, et al: **Descriptive characteristics and health outcomes of the food by prescription nutrition supplementation program for adults living with HIV in Nyanza Province, Kenya**. *Plos One* 2014, 9(3).
 12. Sadler K, Bontrager E, Rogers B, Coates SG J, Kidane Y. Food by Prescription: Measuring the impact and cost-effectiveness of prescribed food on recovery from malnutrition and HIV disease progression among HIV + adult clients in Ethiopia. Boston: In.: Feinstein International Center, Friedman School of Nutrition Science and Policy, Tufts University; 2012.
 13. Ahoua L, Umutoni C, Hueriga H, Minetti A, Szumilin E, Balkan S, Olson DM, Nicholas S, Pujades-Rodríguez M. **Nutrition outcomes of HIV-infected malnourished adults treated with ready-to-use therapeutic food in sub-Saharan Africa: A longitudinal study**. *Journal of the International AIDS Society* 2011, 14(1).
 14. Tang AMP, Quick TP, Chung MP, Wanke CAMD. Nutrition Assessment, Counseling, and Support interventions to improve health-related outcomes in people living with HIV/AIDS: a systematic review of the literature. *JAIDS Journal of Acquired Immune Deficiency Syndromes*. 2015;68(Supplement(3):340–9.
 15. WHO. **Nutrition counselling, care and support for HIV-infected women: guidelines on HIV-related care, treatment and support for HIV-infected women and their children in resource-limited settings**. 2004.
 16. Grobler L, Siegfried N, Visser ME, Mahlangu SS, Volmink J. Nutritional interventions for reducing morbidity and mortality in people with HIV. *Cochrane Database Syst Rev*. 2013;2:Cd004536.
 17. Food and Nutrition Technical Assistance (FANTA). **HIV/AIDS: a guide for nutritional care and support**. In.; 2004.
 18. Van Niekerk C, Smego R Jr, Sanne I. Effect of nutritional education and dietary counselling on body weight in HIV-seropositive South Africans not receiving antiretroviral therapy. *Journal of Human Nutrition Dietetics*. 2000;13(6):407–12.
 19. Kaye HL, Moreno-Leguizamon CJ. Nutrition education and counselling as strategic interventions to improve health outcomes in adult outpatients with HIV: a literature review. *African Journal of AIDS Research*. 2010;9(3):271–83.
 20. Martinez H, Palar K, Linnemayr S, Smith A, Derosé KP, Ramírez B, Farías H, Wagner G. Tailored nutrition education and food assistance improve adherence to HIV antiretroviral therapy: evidence from Honduras. *AIDS Behav*. 2014;18(5):566–77.

21. Anema A, Zhang W, Wu Y, Elul B, Weiser SD, Hogg RS, Montaner JS, El Sadr W, Nash D. Availability of nutritional support services in HIV care and treatment sites in sub-Saharan African countries. *Public Health Nutr.* 2012;15(5):938–47.
22. Sackey J, Zhang FF, Rogers B, Aryeetey R, Wanke C. Implementation of a nutrition assessment, counseling and support program and its association with body mass index among people living with HIV in Accra, Ghana. *AIDS care.* 2018;30(5):586–90.
23. Bacelo AC, Ramalho A, Brasil PE, Cople-Rodrigues CdS, Georg I, Paiva E, Argolo SVL, Rolla VC. Nutritional Supplementation Is a Necessary Complement to Dietary Counseling among Tuberculosis and Tuberculosis-HIV Patients. *PLOS ONE.* 2015;10(8):e0134785.
24. Deroose KP, Felician M, Han B, Palar K, Ramírez B, Farías H, Martínez H. A pre-post pilot study of peer nutritional counseling and food insecurity and nutritional outcomes among antiretroviral therapy patients in Honduras. *BMC nutrition.* 2015;1:21.
25. Alo C, Ogbonnaya LU, Azuogu BN: **Effects of nutrition counseling and monitoring on the weight and hemoglobin of patients receiving antiretroviral therapy in Ebonyi State, Southeast Nigeria.** *HIV/AIDS (Auckland, NZ)* 2014, **6**:91–97.
26. Weiser SD, Tuller DM, Frongillo EA, Senkungu J, Mukiibi N, Bangsberg DR. Food Insecurity as a Barrier to Sustained Antiretroviral Therapy Adherence in Uganda. *PLOS ONE.* 2010;5(4):e10340.
27. Tang AM, Quick T, Chung M, Wanke CA. Nutrition Assessment, Counseling, and Support (NACS) interventions to improve health-related outcomes in people living with HIV/AIDS: a systematic review of the literature. *J Acquir Immune Defic Syndr.* 2015;68(0 3):340.
28. Penda CI, Moukoko ECE, Nolla NP, Evindi NOA, Ndombo PK. **Malnutrition among HIV infected children under 5 years of age at the Laquintinie hospital Douala, Cameroon.** *Pan African Medical Journal* 2018, 30.
29. Tafesea Z, Birhanb Y. A H: Nutritional care and support among adults living with HIV at Hawassa Referral Hospital, southern Ethiopia: A qualitative study. *African Journal of AIDS Research.* 2013;12(2):105–11.
30. Bronfenbrenner U. The ecology of human development experiments by nature and design. Cambridge: Cambridge, MA: Harvard University Press; 1996.
31. Gregson J, Foerster SB, Orr R, Jones L, Benedict J, Clarke B, Hersey J, Lewis J, Zotz KJJone: **System, environmental, and policy changes: using the social-ecological model as a framework for evaluating nutrition education and social marketing programs with low-income audiences.** 2001, **33**:S4-S15.
32. Elder JP, Lytle L, Sallis JF, Young DR, Steckler A, Simons-Morton D, Stone E, Jobe JB, Stevens J, Lohman T, et al. A description of the social–ecological framework used in the trial of activity for adolescent girls (TAAG). *Health Educ Res.* 2007;22(2):155–65.
33. Taruwinga P. **Living with HIVAIDS: A nutritional support model for rural women in Zimbabwe.** *D.PH.* Ann Arbor: Capella University; 2014.
34. Parker ME, Bentley ME, Chasela C, Adair L, Piwoz EG, Jamieson DJ, Ellington S, Kayira D, Soko A, Mkhomawanthu C, et al. The acceptance and feasibility of replacement feeding at 6 months as an

- HIV prevention method in Lilongwe, Malawi: Results from the BAN Study. AIDS education prevention: official publication of the International Society for AIDS Education. 2011;23(3):281–95.
35. Chimphamba Gombachika B, Fjeld H, Chirwa E, Sundby J, Malata A, Maluwa A. A Social Ecological approach to exploring barriers to accessing sexual and reproductive health services among couples living with HIV in Southern Malawi. *ISRN Public Health*. 2012;2012:13.
 36. Coetzee B, Kagee A, Bland R. Barriers and facilitators to paediatric adherence to antiretroviral therapy in rural South Africa: a multi-stakeholder perspective. *AIDS Care*. 2015;27(3):315–21.
 37. Tesfay F: **Determinants of nutritional outcomes and challenges of nutritional programs in HIV care in Tigray region, Ethiopia: A mixed methods study**. Doctor of Philosophy, Thesis (Doctorate); 2019.
 38. Tesfay F, Ziersch A, Mwanri L, Javanparast S. Contextual and individual level factors influencing nutritional program effectiveness in HIV care setting in Tigray region, northern Ethiopia: Mixed methods study. *Plos one*. 2020;15(4):e0231859.
 39. Gale NK, Heath G, Cameron E, Rashid S, Redwood S. Using the framework method for the analysis of qualitative data in multi-disciplinary health research. *BMC Med Res Methodol*. 2013;13:117–7.
 40. FHAPCO E. **Staying strong with HIV/AIDS, Seven ways to maintain Strength if you are HIV-positive**. In..
 41. Ayala GX, Elder JP. Qualitative methods to ensure acceptability of behavioral and social interventions to the target population. *J Public Health Dent*. 2011;71(Suppl 1(0 1):69–79.
 42. Deroose KP, Felician M, Han B, Palar K, Ramírez B, Farías H, Martínez H. A pre-post pilot study of peer nutritional counseling and food insecurity and nutritional outcomes among antiretroviral therapy patients in Honduras. *BMC Nutrition*. 2015;1(1):21.
 43. Scarcella P, Buonomo E, Zimba I, Doro Altan AM, Germano P, Palombi L, Marazzi MC. The impact of integrating food supplementation, nutritional education and HAART (Highly Active Antiretroviral Therapy) on the nutritional status of patients living with HIV/AIDS in Mozambique: results from the DREAM Programme. *Ig Sanita Pubbl*. 2011;67(1):41–52.
 44. Schwenk A, Bürger B, Ollenschläger G, Stützer H, Wessel D, Diehl V, Schrappe M. Evaluation of nutritional counselling in HIV-associated malnutrition. *Clin Nutr*. 1994;13(4):212–20.
 45. Government of the Federal Democratic Republic of Ethiopia. **National Nutrition Programme** In. 2013;–2015:22–3.
 46. Kebede MA, Haidar J. **Factors influencing adherence to the food by prescription program among adult HIV positive patients in Addis Ababa, Ethiopia: a facility-based, cross-sectional study**. *Infectious Diseases of Poverty* 2014, 3(20).
 47. Bourdier F. **Socio-anthropological investigation related to the acceptability of Plumpy’nut in Cambodia**. *Paris: institut de Recherche pour le Développement (IRD)* 2009.
 48. Gitome S, Njuguna S, Kwenza Z, Ombati E, Njoroge B, Bukusi EA. Perspectives paper: medicalisation of HIV and the African response. *Afr J Reprod Health*. 2014;18(3):25–33.

49. Greenaway K: **Food by Prescription: A landscape paper**. In: *GAIN Working paper series*. edn.: Global Alliance for Improved Nutrition; 2009.
50. Kuria EN. Food consumption and nutritional status of people living with HIV/AIDS (PLWHA): a case of Thika and Bungoma Districts, Kenya. *Public Health Nutr*. 2010;13(4):475–9.
51. Tabi M, Vogel RL. Nutritional counselling: an intervention for HIV-positive patients. *J Adv Nurs*. 2006;54(6):676–82.
52. Gaikwad S, Giri P, Suryawanshi S, Garg S, Singh M, Gupta V. Impact of nutritional counseling on dietary practices and body mass index among people living with HIV/AIDS at a tertiary care teaching hospital in Mumbai. *Journal of Medical Nutrition Nutraceuticals*. 2013;2(2):99–102.
53. Pollak KI, Krause KM, Yarnall KS, Gradison M, Michener JL, Østbye T. Estimated time spent on preventive services by primary care physicians. *BMC Health Services Research*. 2008;8(1):245.
54. Food and Nutrition Technical Assistance II Project (FANTA-2). **Review of Kenya's Food by Prescription Program**. In. Washington, DC; 2009.
55. Wynn K, Trudeau JD, Taunton K, Gowans M, Scott I. **Nutrition in primary care. Current practices, attitudes, and barriers** 2010, 56(3):e109-e116.
56. Hussien S, Worku A, Aklilu A, Abate K: **Ready-to-Use Therapeutic Food for Management of Wasting in HIV Infected Adults: A Qualitative Investigation of Views and Experiences of Patients in Ethiopia**. *International Journal of Nutrition and Food Sciences* 2015; 2015, 4(5):518–529.
57. Kolasa KM, Rickett K. Barriers to providing nutrition counseling cited by physicians: a survey of primary care practitioners. *Nutrition in Clinical Practice*. 2010;25(5):502–9.
58. Van Niekerk C, Smego RA, Sanne I. Effect of nutritional education and dietary counselling on body weight in HIV-seropositive South Africans not receiving antiretroviral therapy. *Journal of Human Nutrition Dietetics*. 2000;13(6):407–12.
59. Julie M. **Barriers to nutrition management among people living with hiv on antiretroviral therapy**. 2011.
60. McLeroy KR, Bibeau D, Steckler A, Glanz K. An ecological perspective on health promotion programs. *Health Educ Q*. 1988;15(4):351–77.
61. Stokols D. Translating social ecological theory into guidelines for community health promotion. *Am J Health Promot*. 1996;10(4):282–98.
62. Mehtälä MAK, Sääkslahti AK, Inkinen ME, Poskiparta MEH. A socio-ecological approach to physical activity interventions in childcare: a systematic review. *International Journal of Behavioral Nutrition Physical Activity*. 2014;11(1):22.
63. Stokols D. Establishing and maintaining healthy environments: toward a social ecology of health promotion. *American psychologist*. 1992;47(1):6.
64. Robinson T. Applying the Socio-ecological Model to improving fruit and vegetable intake among low-income African Americans. *J Community Health*. 2008;33(6):395–406.

Figures

Figure 1

Shows the interrelationships of various factors that impact the nutritional counselling in HIV care settings

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Interveiwguide.docx](#)