Levels and Distribution of Organophosphate Esters (OPEs) in Typical Megacity Wetland Park Landscape Water Bodies in Southwest China

Hongling Yin (✉ bellings15@126.com)

Chengdu University of Information Technology <https://orcid.org/0000-0002-1662-4673>

Liya Liu

Chengdu University of Information Technology

Qin Liu

Chengdu University of Information Technology

Jiaojiao Song

Chengdu University of Information Technology

Shuhong Fang

Chengdu University of Information Technology

Xiaowen Liu

Chengdu University of Information Technology

Research Article

Keywords: Human activities, organophosphate esters (OPEs), suspended particulate matter (SPM), tris-(2-chloroethyl)-phosphate (TCEP) and trichloropropyl phosphate (TCIPP)

Posted Date: June 30th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-649375/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Archives of Environmental Contamination and Toxicology on October 28th, 2021. See the published version at <https://doi.org/10.1007/s00244-021-00899-z>.

Levels and distribution of organophosphate esters (OPEs) in typical megacity wetland park landscape water bodies in Southwest China

Hongling Yin, Liya Liu, Qin Liu, Jiaojiao Song, Shuhong Fang, Xiaowen Liu

Abstract

Human activities have led to the release of organophosphate esters (OPEs) into the environment. This study aims to investigate the levels and partitioning of OPEs in surface water, suspended particulate matter (SPM) and sediments of landscape waters across eleven parks in the city of Chengdu, a megacity in Southwest China. The average concentration of Σ_6 OPEs in the SPM samples (median: 2.94×10^3 ng/L, 6.88×10^4 ng/g dw) was 1-3 orders of magnitude higher than that in the surface water (median: 359 ng/L) and sediment (median: 82.8 ng/g) samples. Tri-n-butyl phosphate (TnBP), tris-(2-chloroethyl)-phosphate (TCEP) and trichloropropyl phosphate (TCIPP) were the primary OPE pollutants in the surface water and SPM samples, while TnBP, tris-(2-butoxyethyl) phosphate (TBEP) and tris-(2-ethylhexyl) phosphate (TEHP) predominated the sediment samples. The higher log K_{oc} values of OPEs in park landscape water bodies than other studies in the present study could be explained by the OPE properties (f_{oc} , K_{ow} , degradability) and the environmental conditions (the input sources and the hydraulic retention time, etc.).

✉ Hongling Yin
yhl@cuit.edu.cn
College of Resources and Environment, Chengdu University of Information Technology,
Cheng Du, China

As substitutes for brominated flame retardants, organophosphate esters (OPEs) are important organophosphorus flame retardants. In recent years, due to the large-scale production and use of OPEs worldwide, the accompanying environmental issues have also become increasingly concerning. Some OPEs have been confirmed to have obvious neurotoxicity, reproductive toxicity, carcinogenicity and genotoxicity (Van and Boer, 2012; Du et al. 2015). At present, the occurrence of OPEs has been reported in the air (Takigami et al. 2009; Stapleton et al. 2009; Clark et al. 2017; Yin et al. 2020), wastewater and sludge (Bester et al. 2005; Gao et al. 2016), surface water (Reemtsma et al. 2008; Regnery and Püttmann, 2010; Guo et al. 2017a), sediments (Cao et al. 2012; Cheng et al. 2014; Giulivo et al. 2017), soils (Mihajlovic et al. 2011; Matsukami et al. 2015; Wan et al. 2016; Cui et al. 2017; Deng et al. 2019) and humans (Shah et al. 2006; Schindler et al. 2009), even in remote areas (McDonough et al. 2018). OPEs have become recognized global organic pollutants. As contaminants of emerging concern, OPEs are gradually becoming of research interest to the scientific community. In this paper, alkyl OPEs (TnBP, TBEP, TEHP), chlorinated OPEs (TCEP and TCIPP) and phenyl OPEs (TPhP) were selected because they are widely used in industry and previous studies have shown that their concentrations in all environmental media are relatively high (Bacaloni et al. 2008; Shi et al. 2016; van der Veen and de Boer, 2012; Zhang et al. 2018).

After entering aquatic environments, OPEs endure a series of migration, transformation and bioconcentration related processes. The structural differences among OPEs lead to a variety of chemical and physical properties that result in differences in environmental behaviors. OPEs have only anthropogenic sources as opposed to natural sources, so cities with large populations are potential high-risk areas for OPE pollution in which water bodies are an essential sink of OPEs. Previous studies have

focused on the occurrence of OPEs in rivers and sediments (Bacaloni et al. 2008; Rodil et al. 2012; Giulivo et al. 2017; Guo et al. 2017a; Hou et al. 2019; Wu et al. 2019; Zeng et al. 2021), but almost no attention has been paid to urban landscape water bodies. Compared with rivers, urban landscape water has a relatively low flow rate and long water exchange time. The residence time of a relatively open water body ranges from approximately minutes to hours, and the residence time of a semiclosed water body can be hours to days depending on the water temperature. Closed water bodies have limited water exchange, and its retention time is the longest, resulting in its OPE pollution characteristics can reflect emissions from local sources. Therefore, the OPE pollution characteristics and its partitioning behavior in different types of landscape water bodies is of great interest.

The city of Chengdu, located in the middle reaches of the Minjiang River Basin, is the capital of the Sichuan Province and the only megacity (16.33 million people in 2018) in Southwest China. However, there have been few studies on OPEs in this region. To date, there are 25 parks in Chengdu city, and the landscape water for urban parks covers an area of 6.50×10^5 m². This study aims to 1) investigate the pollution levels and characteristics of OPEs in water, suspended particulate matter (SPM) and sediments of representative park landscape waters; 2) analyze the distribution behavior of OPEs in different water bodies; and 3) reveal the relationship between the OPE distribution and its influencing factors. The results can fill knowledge gaps regarding OPE pollution in different phases in the park landscape waters of Southwest China and shed light on their partition behavior, which is of great significance for OPE pollution prevention and control.

Materials and Methods

Chemicals

The main reagents, including ethyl acetate, acetone, hexane and acetonitrile, were high-performance liquid chromatography (HPLC) grade (Kelon Chemical, China). The standard solutions including tri-n-butyl phosphate (TnBP, 97%) (Cas NO.: 126-73-8), tris-(2-ethylhexyl) phosphate (TEHP, 97%) (Cas NO.: 78-42-2), tris-(2-butoxyethyl) phosphate (TBEP, 94%) (Cas NO.: 78-51-3), triphenyl phosphate (TPhP) (Cas NO.: 115-86-6), tris-(2-chloroethyl)-phosphate (TCEP) (Cas NO.: 115-96-8) and Tris-(2-chloroisopropyl) phosphate (TCIPP) (Cas NO.: 13674-84-5) and internal standard (triphenyl phosphate-d₁₅ (TPhP-d₁₅)) (Cas NO.:1173020-30-8) were all purchased from Sigma-aldrich Corp., USA. Copper, aluminium oxide, silica gel, Na₂SO₄ and other chemicals were purchased from Kelon Chemical Corp., China. Deionized water was supplied from a Milli-Q equipment.

Sample Collection

In April of 2017, samples were collected from 11 parks in Chengdu city. There was no rainfall during sample collection period. Geographic information on all 11 sampling sites is shown in table 1 and the spatial distribution of sampling locations is shown in Fig. 1. Due to the layout of cities in different directions, their population density, industrial development and environmental conditions vary. The main roads in Chengdu include the 1st ring road (Yihuan road), 2nd ring road (Erhuan road), 3rd ring road (Sanhuan road), and fourth ring road (Raocheng road), spreading from the downtown center to the outskirts with a radial pattern. Site 6# (People's Park) was located in the first ring, two sites including 4# (Jinghu Park) and 5# (Huanhuaxi Park) were located in the second ring, three sampling sites including 3# (Shahe Park), 7# (East Lake Park) and 9# (Daguanyan) were located in the third ring, and five sampling sites including 1# (Lianghe City Park), 2# (Fenghuangshan Park), 8# (Jincheng

Park), 10# (Bailu Bay Wetland Park) and 11# (Qinglong Lake Wetland Park) were located in the fourth ring. The number of sampling sites is positively correlated with the area of the ring ($p < 0.05$). Moreover, the sampling sites covered different types of landscape water bodies (relatively open: 5#, 7#, 8#; semiclosed: 3#, 10# and closed water bodies: 1#, 2#, 4#, 6#, 9#, 11#). Therefore, all samples are representative and typical. The area of the 11 wetland parks ranges from 0.11 km² to 20 km² (Table 1).

The surface water was sampled at 0.5 m below the water surface. The water samples collected from multiple sampling sites were mixed into one water sample, and parallel samples were obtained. A shovel immersed in the water body was used to collect sediment samples near the shore. The sediment samples were wrapped with aluminum foil, sealed in a seal bag, transported to the laboratory as soon as possible and stored in a refrigerator at -20 °C.

Sample Preparation and Analysis

The procedures for extraction and clean-up of the water and sediment samples followed a previously described method with minor modifications (Wu et al. 2019; Yin et al. 2021). Each sample was pretreated after spiking with a surrogate standard (100 ng of triphenyl phosphate-d₁₅ (TPhP-d₁₅)). 1L of surface water sample was passed through a 0.45 µm quartz filter to obtain SPM and water samples. The water samples were clear and transparent, indicating that the algae content was low. Solid-phase extraction with a C₁₈ column was used to enrich the target substances in the water samples. The C₁₈ column was eluted with 10 mL hexane for removal of impurities and then with 10 mL ethyl acetate/acetone (v/v, 4/3) for elution of the target substances. The eluent was concentrated and the final volume was adjusted to 100 µL for gas chromatography-mass spectrometry (GC-MS) analysis.

The SPM samples were freeze-dried, soaked in 20 mL ethyl acetate/acetone (v: v, 3:2) for 12

hours, and ultrasonically extracted twice for 30 minutes. Then, they were concentrated to approximately 1 mL with vacuum-condensing equipment and loaded onto an activated aluminum oxide/silica gel (v: v, 3:1) column. The column was first eluted with 20 mL hexane to remove impurities, then with 20 mL ethyl acetate/acetone (v: v, 3:2), and the eluate (ethyl acetate/acetone) was collected. The solvent extracts were concentrated to 100 μ L for GC-MS analysis.

By measuring the content of OPEs in surface water, SPM and sediments, as well as the content of organic matter in SPM and sediments, the water-SPM and water-sediment distribution coefficients were calculated. The distribution coefficients were calculated as follows:

$$K_a = C_{sed} / C_w \times 1000 \quad (1)$$

$$K_b = C_{spm} / C_w \times 1000 \quad (2)$$

K_a : equilibrium coefficient between water and SPM (cm^3/g),

K_b : equilibrium coefficient between water and sediment (cm^3/g),

C_{sed} : the content of OPEs in sediment at equilibrium (ng/g),

C_{spm} : the content of OPEs in SPM at equilibrium (ng/g),

C_w : the content of OPEs in surface water at equilibrium (ng/L).

$$K_{oc} = K_{a(or\ b)} \times 100 / f_{oc} \quad (3)$$

K_{oc} : standardized partition coefficient of organic carbon (cm^3/g);

f_{oc} : content of organic carbon in sediment/suspended particulate matter (%)

Total organic carbon was determined by potassium dichromate titration.

Detailed analytical parameters were similar with those in Wu et al. 2019 and Yin et al. 2021. The

GC was equipped with SH-Rxi-5Sil MS capillary column (30 m $\times$ 0.25 μ m $\times$ 0.25 mm, Shimadzu,

Japan) and operated with a 280 °C inlet temperature using splitless injection. The MS source was electron impact (EI), and it was operated in selected ion monitoring (SIM) mode. The GC oven temperature was held at 50 °C for 1 minute, increased to 200 °C at 15 °C min⁻¹ and held for 1 minute, increased to 250 °C at 4.00 °C min⁻¹, and then increased to 300 °C at 20 °C min⁻¹ and held for 4 minutes. The interface temperature was 280 °C, and the ion source temperature was 200 °C.

Quality Assurance and Quality Control (QA/QC)

Thorough QA/QC procedures for OPEs analysis were conducted to ensure data quality. To evaluate the recovery efficiencies of analytical procedures, internal standard (TPhP-d₁₅) was added in all samples, and the accuracy was evaluated via their recoveries. The concentrations of the 6 OPEs were determined by an external standard method. The correlation coefficients of the standard curves of the six OPE compounds (0.05-2.00 mg/L) were all higher than 0.99. The recoveries of the 6 OPEs were determined, ranged from 75% to 126% (Table 2). Procedural contamination from the analytical steps aforementioned was evaluated via running a matrix blank with every batch of 10 samples. Only TnBP and TCEP were detected in the blanks, and the levels were < 5% of the concentrations measured in all samples, which meant that they were negligible. The detection limit of the instrument (LODs) (S/N = 3) ranged from 0.02 mg/L to 0.04 mg/L and the instrument precision was 3%-7%.

Statistical Analysis

Data analysis was performed using the IBM SPSS 22.0. software. Before statistical analysis, data were tested for normality through Kolmogorov-Smirnoff and Shapiro-Wilk test which showed normal distribution of the data ($p > 0.05$ for all). Analysis of variance (ANOVA) was used to investigate

significant differences in individual OPEs concentrations in different matrix. Correlations between the three environmental phases were analyzed by Pearson's correlation coefficients. For the calculation of OPEs, non-detectable compounds and concentrations below the method detection limit (MDL) were treated as zero. Statistical significance was set as $p < 0.05$.

Results and Discussion

Occurrence and Levels of OPEs in Surface Water, SPM and Sediments

All OPE compounds had detection rates of 100% for all SPM samples, while all OPE compounds had an over 50% detection rate for all water and sediment samples. Although the TCIPP and TPhP had detection rates of 91% for all water samples, TCIPP (73%), TBEP (64%) and TPhP (55%) had moderate detection rates for sediment samples, and other OPE compounds had detection rates of 100%. High detection rates of OPEs in the aqueous environment indicated that OPE pollution was ubiquitous in the investigated area.

The concentrations of Σ_6 OPEs in the water samples were in the range of 213-658 ng/L, with a median value of 359 ng/L. Compared with OPEs in other lakes and reservoirs, the concentrations of OPEs in the present study were higher than those found in volcanic lakes (average: 165 ng/L) (Bacaloni et al. 2008) and the Great Lakes (7.3-96 ng/L) (Venier et al. 2014), but two times lower than those in the Pearl River Delta (median concentration: 837 ng/L) (Zhang et al. 2018) and one order of magnitude lower than those in found freshwater from Germany (average: 2.06×10^3 ng/L) (Ernst, 1988) and Spain (average: 1.98×10^3 ng/L) (Cristale et al. 2013).

The concentrations of Σ_6 OPEs in sediment samples were in the range of 49.2-482 ng/g, with a median value of 82.8 ng/g. The highest concentration of OPEs appeared at site 4#, followed by site 3#, which was highly related to the high density of human activities. The Σ_6 OPE level in sediment samples is comparable with those from the Pearl River Delta (8.3-470 ng/g) (Tan et al. 2016) in China and the Adige River basin in Italy (11.5-549 ng/g), higher than those in the Evrotas River basin in Greece (10.5-248 ng/g), the Sava River basin in Slovenia (0.31-310 ng/g) (Giulivo et al. 2017), Bui Dau in Vietnam (5-300 ng/g) (Giulivo et al. 2017), Austria (5.74 ng/g) (Matsukami et al. 2016) and Norway (4.98 ng/g) (Martínez-Carballo et al. 2007), but lower than those in the Flanders Rivers in Belgium (673 ng/g) (Green et al. 2008) and the Llobregat River basin in Spain (n.d.- 2.42×10^3 ng/g) (Santín et al. 2016).

The concentrations of Σ_6 OPEs in SPM varied from 468 to 6.74×10^3 ng/L (2.22×10^4 - 7.08×10^5 ng/g dw), with a median of 2.94×10^3 ng/L (6.88×10^4 ng/g dw). The highest concentration of OPEs in SPM was observed at site 8#. Site 8# is in a new wetland park with low population density. However, it is located in a high-tech zone with electronic industry and a large number of new company buildings and industrial parks, which was the cause of high emissions of TnBP and TBEP. Compared with the concentration of traditional brominated flame retardants, the levels of OPEs in SPM in our study were three to five orders of magnitude higher than those in PBDEs found in seawater near Hong Kong (Wurl et al. 2006). A similar trend was found in Dongjiang industrial water (Ruan et al. 2014). The concentration of brominated flame retardants in water was lower than that of OPEs, indicating that OPEs, as substitutes for brominated flame retardants, result in a higher concentration in some environmental media at present. Other researchers have found similar results in which the

concentrations of OPEs were higher than those of brominated flame retardants in the 40 rivers flowing into Bohai (Wang et al. 2015).

In general, the concentrations of OPEs in surface water and sediments of landscape waters in the present study were at the middle/low levels compared to other waters. However, the average concentration of OPEs in SPM was one to three orders of magnitude higher than that in surface water and sediments, indicating that SPM is more easily enriched with OPEs than water and sediments. Consequently, urgent attention should be given to OPE contamination in SPM.

In the surface water samples, TnBP (range: 104-227 ng/L, median: 146 ng/L, 39% of Σ_6 OPEs) and TCIPP (n.d.-285 ng/L, 125 ng/L, 35% of Σ_6 OPEs) were the predominant chemicals of OPEs. There were significant differences between the concentrations of individual OPEs in the aqueous phase ($p < 0.01$). The concentrations of TnBP and TCIPP were 1-2 orders of magnitude higher than those of TEHP and TPhP. The concentrations of the major pollutants in surface water was relatively low compared to the global lake waters (Bacaloni et al. 2008; Shi et al. 2006; Venier et al. 2014).

The main components of OPEs in sediments were TnBP (range: 16.2-439 ng/g, median: 61.7 ng/g, 67% of Σ_6 OPEs), TBEP (n.d.-128 ng/g, 11.5 ng/g, 13% of Σ_6 OPEs) and TEHP (8.6-32.6 ng/g, 8.64 ng/g, 8% of Σ_6 OPEs). Significant differences were also found between the concentrations of individual OPEs in the sediment samples ($p < 0.01$). The concentrations of dominant OPEs in the sediments were higher than those in TaiHu (TBEP: 1.03-5.00 ng/g) (Cao et al. 2012) and the Pearl River Delta (TBEP: 5.8-46 ng/g, TEHP: 6.2-44 ng/g) (Tan et al. 2016).

All OPE compounds were detected in the SPM samples, with TnBP (1.06×10^4 - 5.42×10^5 ng/g, median: 3.26×10^4 ng/g, 76% of Σ_6 OPEs), TBEP (1.82×10^3 - 7.74×10^4 ng/g, 7.72×10^3 ng/g, 7% of

205 Σ_6 OPEs), TCIPP (1.29×10^3 - 1.03×10^5 ng/g, 3.91×10^3 ng/g, 7% of Σ_6 OPEs) and TCEP (1.07×10^3 -
206 6.36×10^4 ng/g, 3.40×10^3 ng/g, 7% of Σ_6 OPEs) being the dominant components. TnBP has a
207 significantly higher concentration than the other compounds ($p < 0.01$). Compared with the main
208 pollutants of TnBP (median: 21.5 ng/g, 38% of Σ OPEs) and TCEP (median: 10.7 ng/g, 32% of Σ OPEs)
209 in the Pearl River Delta, the concentrations of individual OPEs in the SPM of the scenic water in this
210 study were approximately 2-4 orders of magnitude higher (Zhang et al. 2018).

211 In general, the distribution patterns of OPEs in the aqueous phase and SPM were similar, and
212 $\Sigma(\text{TnBP}+\text{TCIPP}+\text{TCEP})$ accounted for 82% and 91% of the Σ_6 OPEs in the aqueous phase and SPM,
213 respectively. Regarding the sediment samples, TnBP, TBEP and TEHP were the dominant compounds
214 found at all sampling sites, and $\Sigma(\text{TnBP}+\text{TBEP}+\text{TEHP})$ accounted for 86% of the Σ_6 OPEs. The
215 different patterns of OPEs could be attributed to differences in their accumulation features and
216 degradability due to varying physicochemical properties, as well as differences in their production and
217 usage (Wang et al. 2015).

218 To further elucidate the OPE patterns in surface water, sediments and SPM, OPEs were divided
219 into alkyl OPEs (TnBP, TBEP, TEHP), chlorinated OPEs (TCEP and TCIPP) and phenyl OPEs (TPhP).
220 Alkyl OPEs dominated the profile in three phases (water, sediments and SPM), and their percentage in
221 the latter two phases was greater than 85%. Chlorinated OPEs accounted for 43% of Σ_6 OPEs in the
222 water, but 10% and 14% of Σ_6 OPEs in the SPM and sediment samples, respectively.

223 Based on the specific water environment, the 11 sampling sites were divided into relatively open,
224 closed and semiclosed water body categories. 7# (Donghu Park) and 8# (Jincheng Park) are relatively
225 open water bodies that converge with the Fuhe River and Xiaojia River, respectively. Continuous

emission input, complex hydrodynamic conditions and frequent resuspension of OPEs in sediments results in high concentrations of OPEs in SPM and low concentrations in sediments. Sites 3# and 10# are typical semiclosed water bodies. Although these two sampling sites have inflows from other tributaries, they could be seen as closed water bodies as a whole. These sites showed higher OPE pollution in the water phases, whereas the concentrations of OPEs in SPM were low. The other sites (1-2#, 4#, 6#, 10-11#) are typical closed water bodies. As typical representative closed water body, sites 2# and 4# have the highest concentrations of OPEs in water and sediment samples respectively, which have almost no water exchange, coupled with a long history of pollution and a high density of human activities, resulting in the highest concentrations of OPEs in water and sediments and but low concentrations in SPM.

Distribution of OPEs Between Surface Water, SPM and Sediments

Distribution of OPEs Between Surface Water and SPM

Correlation analyses were performed between the concentration of Σ_6 OPEs in water, sediment and SPM samples and significant correlations were observed in different matrix (Table S1). A strong positive correlation was found between $\log K_a$ and $\log K_b$ ($R^2 = 0.80$, $p < 0.05$) (Fig. S1), which suggested that it was meaningful to study the partition behavior between different matrix (Wang et al. 2018). The calculated $\log K_{oc}$ value between water and SPM was shown in Table 3.

The $\log K_a$ values of TCIPP ($R = 0.654$, $p < 0.05$) and TPhP ($R = 0.662$, $p < 0.05$) were significantly and positively correlated with f_{oc} . The same results reported that only the $\log K_a$ of TnBP ($R = 0.677$, $p < 0.05$) and TCIPP ($R = 0.669$, $p < 0.05$) was significantly and positively correlated with

f_{oc} (Zhang et al. 2018). This suggests that the partitioning of most OPEs between the water and SPM was not mediated by f_{oc} .

As shown in Fig. 2, no significant correlation was observed between $\log K_{oc}$ and $\log K_{ow}$. These results differed from the research of Zhang et al (2018) who reported that the significant correlation was found between $\log K_{oc}$ and $\log K_{ow}$ in Pearl River Delta from China. Noticeably, the calculated $\log K_{oc}$ value of most of OPE compounds in the present study was higher than the data reported by Zhang et al (2018) and the predicted $\log K_{oc}$ using EPI Suite software (US EPA, 2012). Therefore, the partition behaviors of OPEs in wetland park landscape water bodies were different from lakes and rivers.

Distribution of OPEs Between Surface Water and Sediments

The correlation between $\log K_b$ and f_{oc} of the sediment samples showed that the $\log K_b$ of TnBP was highly correlated with f_{oc} ($R = 0.809$, $p < 0.01$), and TnBP was the most abundant component in the water (39% of Σ_6 OPEs) and sediment (67% of Σ_6 OPEs) samples. This result indicated that organic carbon was one of the most critical factors affecting the distribution of OPEs in water and sediments.

According to the correlation analysis between $\log K_{oc}$ and $\log K_{ow}$ of OPEs in the water and sediment samples, a significant positive linear correlation between $\log K_{oc}$ and $\log K_{ow}$ was only found at site 10# ($R = 0.839$, $p < 0.05$) (Fig. 3 and Fig. 4), suggesting that the partition was dominated by hydrophobic interactions at this site, which was similar to the results from Taihu Lake (Wang et al. 2018). Except for TEHP, the calculated $\log K_{oc}$ values between water and sediment samples were higher than the values reported by Wang et al (2018).

Overall, the calculated $\log K_{oc}$ value in the present study was markedly higher than those from previous studies both in water-SPM and water-sediment phase. The higher $\log K_{oc}$ values could be partly explained by the compositions of organic matters in SPM and the sorption kinetics of chemicals (Cao et al. 2017). Additionally, the landscape water bodies brought in the fresh discharges of OPEs. The spontaneously migration of OPEs in the sediment and SPM samples to the aqueous phase could not be ignored. These differed from the river or lakes in which the partitioning process was significantly influenced by the contributions of ongoing emissions from the water phase.

Conclusions

Generally, the concentration of OPEs in the surface water and sediment of landscape water in the megacity in Southwest China was at the middle/low level. The Σ_6 OPE concentration in the SPM was 1-3 orders of magnitude higher than that in the surface water and sediments. This phenomenon needs additional attention.

In terms of OPE profiles, TnBP, TCIPP and TCEP were the main OPE pollutants in surface water and SPM, while TnBP, TBEP and TEHP were dominant in sediments. Alkyl phosphates were the dominant pollutants in the three phases.

The calculated $\log K_{oc}$ in the park landscape water bodies were higher than the rivers reported by the previous studies. Differed from the lakes and rivers, the OPE properties (f_{oc} , K_{ow} , degradability) and the environmental conditions (the input sources and the hydraulic retention time, etc.) mainly influence the partitioning processes of OPEs in different matrix in the park landscape water bodies. Especially, K_{ow} can partly explain the partitioning process of OPEs between water and sediment samples. The OPE

286 pollution prevention should be based on the consideration on the changing of the main influencing
287 environmental factors except for the control of the source emissions.

288

289 **Declaration of Interests** The authors declare that they have no known competing financial interests or
290 personal relationships that could have appeared to influence the work reported in this paper.

291

292 **Acknowledgements** This work was financially supported by the National Natural Science Fund of
293 China (No.41773072, No.21407014)

294 **References**

- 295 Bacaloni A, Cucci F, Guarino C, Nazzari M, Samperi R, Laganà A (2008) Occurrence of
 296 organophosphorus flame retardant and plasticizers in three volcanic lakes of central Italy. *Environ*
 297 *Sci Technol* 42(6):1898–1903. <https://doi.org/10.1021/es702549g>
- 298 Bester K (2005) Comparison of TCIPP concentrations in sludge and wastewater in a typical German
 299 sewage treatment plant-comparison of sewage sludge from 20 plants. *J Environ Monit* 7(5):509–
 300 513. <https://pubs.rsc.org/>. <https://doi.org/10.1039/b502318a>
- 301 Cao S, Zeng X, Song H, Li H, Yu Z, Sheng G, Fu J (2012) Levels and distributions of organophosphate
 302 flame retardants and plasticizers in sediment from Taihu Lake, China. *Environ Toxicol Chem*
 303 31(7):1478–1484. <https://doi.org/10.1002/etc.1872>
- 304 Cao D D, Guo J H, Wang Y W, Li Z N, Liang K, Corcoran M B, Hosseini S, Bonina S M C, Rockne K
 305 J, Sturchio N C, Giesy J P, Liu J F, Li A (2017) Organophosphate Esters in Sediment of the Great
 306 Lakes. *Environ Sci Technol* 51(3):1441–1449. <https://doi.org/10.1021/acs.est.6b05484>
- 307 Cheng D, Liu X, Wang L, Gong W, Liu G, Fu W, Cheng M (2014) Seasonal variation and sediment-
 308 water exchange of antibiotics in a shallower large lake in North China. *Sci Total Environ*
 309 476(1):266–275. <https://doi.org/10.1016/j.scitotenv.2014.01.010>
- 310 Clark A E, Yoon S, Sheesley R J, Usenko S (2017) Spatial and temporal distributions of organophosph
 311 -ate ester concentrations from atmospheric particulate matter samples collected across Houston, T
 312 X. *Environ Sci Technol* 51(8):4239–4247. <https://doi.org/10.1021/acs.est.7b00115>
- 313 Cristale J, García V A, Barata C, Lacorte S (2013) Priority and emerging flame retardants in rivers:
 314 occurrence in water and sediment, *Daphnia magna* toxicity and risk assessment. *Environ Int*
 315 59:232–243. <https://doi.org/10.1016/j.envint.2013.06.011>
- 316 Cui K, Wen J, Zeng F, Li S, Zhou X, Zeng Z (2017) Occurrence and distribution of organophosphate
 317 esters in urban soils of the subtropical city, Guangzhou, China. *Chemosphere* 175:514–520.
 318 <https://doi.org/10.1016/j.chemosphere.2017.02.070>
- 319 Deng X, Yin H L, He W L, Luo Y, Wu D, Luo L, Chen J (2019) Distribution and migration of
 320 organophosphate in soil and crops in Chengdu City/suburb profile. *Environ Chem* 38(03):679–685.
 321 <https://doi.org/10.7524/j.issn.0254-6108.2018042803>
- 322 Du Z, Wang G, Gao S, Wang Z (2015) Aryl organophosphate flame retardants induced cardiotoxicity
 323 during zebrafish embryogenesis: by disturbing expression of the transcriptional regulators. *Aquat*
 324 *Toxicol* 161:25–32. <https://doi.org/10.1016/j.aquatox.2015.01.027>
- 325 Gao L, Shi Y, Li W, Liu J, Cai Y (2016) Occurrence and distribution of organophosphate triesters and
 326 diesters in sludge from sewage treatment plants of Beijing, China. *Sci Total Environ* 544:143–149.
 327 <https://doi.org/10.1016/j.scitotenv.2015.11.094>
- 328 Giulivo M, Capri E, Kalogianni E, Milacic R, Majone B, Ferrari F, Barceló D (2017) Occurrence of hal
 329 -ogenated and organophosphate flame retardants in sediment and fish samples from three Europea
 330 n river basins. *Sci Total Environ* 586:782–791. <https://doi.org/10.1016/j.scitotenv.2017.02.056>
- 331 Green N, Schlabach M, Bakke T, Brevik E M, Dye C, Herzke D, Huber S, Plosz B, Remberger M,
 332 Schoyen M, Uggerud H T, Vogelsang C (2008) Screening of selected metals and new organic
 333 contaminants. Norwegian Pollution Control Agency. <https://www.researchgate.net/publication>

- Guo J, Romank K, Westenbroek S, Hites R A, Venier M (2017a) Current use flame retardants in the water of Lake Michigan tributaries. *Environ Sci Technol* 51(17):9960–9969. <https://doi.org/10.1021/acs.est.7b01294>
- Hou L, Jiang J, Gan Z, Dai Y Y, Yang P, Yan Y, Ding S, Su S J, Bao X M (2019) Spatial distribution of organophosphorus and brominated flame retardants in surface water, sediment, groundwater, and wild fish in Chengdu, China. *Arch Environ Contam Toxicol* 77:279–290. <https://doi.org/10.1007/s00244-019-00624-x>
- Martínez-Carballo E, González-Barreiro C, Sitka A, Scharf S, Gans O (2007) Determination of selected organophosphate esters in the aquatic environment of Austria. *Sci Total Environ* 388(1):290–299. <https://doi.org/10.1016/j.scitotenv.2007.08.005>
- Matsukami H, Tue N, Suzuki G, Someya M, Tuyen L H, Viet P H, Takahashi S, Tanabe S, Takigami H (2015) Flame retardant emission from e-waste recycling operation in northern Vietnam: environmental occurrence of emerging organophosphorus esters used as alternatives for PBDEs. *Sci Total Environ* 514:492–499. <https://doi.org/10.1016/j.scitotenv.2015.02.008>
- Matsukami H, Suzuki G, Tue N M, Tuyenle H, Viet P H, Takahashi S, Tanabe S, Takigami H (2016) Analysis of monomeric and oligomeric organophosphorus flame retardants in fish muscle tissues using liquid chromatograph electrospray ionization tandem mass spectrometry: application to Nile tilapia (*Oreochromis niloticus*) from an e-waste processing area in northern Vietnam. *Emerg Contam* 2(2):89–97. <https://doi.org/10.1016/j.emcon.2016.03.004>
- McDonough C A, De Silva A O, Sun C, Cabrerizo A, Adelman D, Soltwedel T, Bauerfeind E, Muir D C G, Lohmann R (2018) Dissolved organophosphate esters and polybrominated diphenyl ethers in remote marine environments: Arctic surface water distributions and net transport through Fram Strait. *Environ Sci Technol* 52(11):6208–6216. <https://doi.org/10.1021/acs.est.8b01127>
- Mihajlovic T, Miloradov M V, Fries E (2011) Application of Twisselmann extraction, SPME, and GC-MS to assess input sources for organophosphate esters into soil. *Environ Sci Technol* 45(6):2264–2269. <https://doi.org/10.1021/es103870f>
- Reemtsma T, Quintana J B, Rodil R, García-Lo'pez M, Rodríguez I (2008) Organophosphorus flame retardants and plasticizers in water and air I. Occurrence and fate. *Trends Anal Chem* 27(9):727–737. <https://doi.org/10.1016/j.trac.2008.07.002>
- Regnery J, Püttmann W (2010) Seasonal fluctuations of organophosphate concentrations in precipitation and storm water runoff. *Chemosphere* 78(8):958–964. <https://doi.org/10.1016/j.chemosphere.2009.12.027>
- Rodil R, Quintana J B, Concha-Graña E, López-Mahía P, Muniategui-Lorenzo S, Prada-Rodríguez D (2012) Emerging pollutants in sewage, surface and drinking water in Galicia (NW Spain). *Chemosphere* 86(10):1040–1049. <https://doi.org/10.1016/j.chemosphere.2011.11.053>
- Ruan W, Tan X, Luo X (2014) Organophosphorus flame retardants in surface sediments of Dongjiang River. *China environmental science* 34:2394–2400. <https://doi.org/10.3969/j.issn.1000-6923.2014.09.042>
- Santín G, Eljarrat E, Barceló D (2016) Simultaneous determination of 16 organophosphorus flame retardants and plasticizers in fish by liquid chromatography-tandem mass spectrometry. *J Chromatogr A* 1441:34–43. <https://doi.org/10.1016/j.chroma.2016.02.058>

- Schindler B K, Förster K, Angerer J (2009) Determination of human urinary organophosphate flame retardant metabolites by solid-phase extraction and gas chromatography-tandem mass spectrometry. *J Chromatogr B* 877(4):375–381. <https://doi.org/10.1016/j.jchromb.2008.12.030>
- Shah M, Meija J, Cabovska B, Caruso J A (2006) Determination of phosphoric acid triesters in human plasma using solid-phase microextraction and gas chromatography coupled to inductively coupled plasma mass spectrometry. *J Chromatogr A* 1103(2):329–336. <https://doi.org/10.1016/j.chroma.2005.11.042>
- Shi Y, Gao L, Li W (2016) Occurrence distribution and seasonal variation of organophosphate flame retardants and plasticizers in urban surface water in Beijing, China. *Environ Pollut* 209:1–10. <https://doi.org/10.1016/j.envpol.2015.11.008>
- Stapleton H M, Klosterhaus S, Eagle S, Fuh J, Meeker J D, Blum A, Webster T F (2009) Detection of organophosphate flame retardants in furniture foam and U.S. house dust. *Environ Sci Technol* 43(19):7490–7495. <https://doi.org/10.1021/es9014019>
- Takigami H, Suzuki G, Hirai Y, Ishikawa Y, Sunami M, Sakai S I (2009) Flame retardants in indoor dust and air of a hotel in Japan. *Environ Int* 35(4):688–693. <https://doi.org/10.1016/j.envint.2008.12.007>
- Tan X X, Luo X J, Zheng X B, Li Z R, Sun R X, Mai B X (2016) Distribution of organophosphorus flame retardants in sediments from the Pearl River Delta in South China. *Sci Total Environ* 544:77–84. <https://doi.org/10.1016/j.scitotenv.2015.11.089>
- US EPA (2012) Estimation Programs Interface Suite for Microsoft Windows, v. U. S. E. P. A., Washington, DC, USA.
- Van der Veen I, de Boer D (2012) Phosphorus flame retardants: properties, production, environmental occurrence, toxicity and analysis. *Chemosphere* 88(10):1119–1153. <https://doi.org/10.1016/j.chemosphere.2012.03.067>
- Venier M, Dove A, Romanak K, Backus S, Hites R (2014) Flame retardants and legacy chemicals in Great Lakes' water. *Environ Sci Technol* 48(16):9563–9572. <https://doi.org/10.1021/es501509r>
- Wan W, Zhang S, Huang H, Wu T (2016) Occurrence and distribution of organophosphorus esters in soils and wheat plants in a plastic waste treatment area in China. *Environ Pollut* 214:349–353. <https://doi.org/10.1016/j.envpol.2016.04.038>
- Wang R, Tang J, Xie Z, Mi W, Chen Y, Wolschke H, Ebinghaus R (2015) Occurrence and spatial distribution of organophosphate ester flame retardants and plasticizers in 40 rivers draining into the Bohai Sea, north China. *Environ Pollut* 198:172–178. <https://doi.org/10.1016/j.envpol.2014.12.037>
- Wang T, Ding N, Wang T, Chen S J, Luo X J, Mai B X (2018) Organophosphorus esters (OPEs) in PM_{2.5} in urban and e-waste recycling regions in southern China: concentrations, sources, and emissions. *Environ Res* 167:437–444. <https://doi.org/10.1016/j.envres.2018.08.015>
- Wang X, Zhu L, Zhong W, Yang L (2018) Partition and source identification of organophosphate esters in the water and sediment of Taihu Lake, China. *J Hazard Mater* 360:43–50. <https://doi.org/10.1016/j.jhazmat.2018.07.082>
- Wu D, Yin H L, Li S P, Wang Z W, Deng X, Luo Y, Luo L (2019) Pollution characteristics of organophosphates in surface water and sediment of Jinjiang River in Chengdu. *Environ Sci* 40(3):1245–1251. <https://doi.org/10.13227/j.hjcx.201808038>

- Wurl O, Lam P K S, Obbard J P (2006) Occurrence and distribution of polybrominated diphenyl ethers (PBDEs) in the dissolved and suspended phases of the sea-surface microlayer and seawater in Hong Kong, China. *Chemosphere* 65(9):1660–1666. <https://doi.org/10.1016/j.chemosphere.2006.02.024>
- Wurl O, Obbard J P, Lam P K S (2006) Distribution of organochlorines in the dissolved and suspended phase of the sea-surface microlayer and seawater in Hong Kong, China. *Mar Pollut Bull* 52(7):768–777. <https://doi.org/10.1016/j.marpolbul.2005.11.024>
- Yin H L, Liang J F, Wu D, Li S P, Luo Y, Deng X (2020) Measurement report: seasonal, distribution and sources of organophosphate esters in PM_{2.5} from an inland urban city in southwest China. *Atmos Chem Phys* 20(23):14933–14945. <https://doi.org/10.5194/acp-20-14933-2020>
- Yin H L, Liu Q, Deng X, Liu X W, Fang S H, Xiong Y M, Song J J (2021) Organophosphates (OPEs) in water, suspended particulate matter (SPM) and sediments of the Minjiang River. *China Chemical Letters*. <https://doi.org/10.1016/j.cclet.2021.02.023>
- Zeng J M, Zhong S H, Qian W, Yuan S W, Zhu X S (2021) Pollution status and biological toxicity of organophosphate in water environment. *Environ Sci*. <https://doi.org/10.19674/j.cnki.issn1000-6923.20210324.004>
- Zhang Y, Zheng X, Wei L, Sun R, Guo H, Liu X, Mai B (2018) The distribution and accumulation of phosphate flame retardants (PFRs) in water environment. *Sci Total Environ* 630:164–170. <https://doi.org/10.1016/j.scitotenv.2018.02.215>

435 **Table captions:**

436 **Table 1** Information about the sampling sites.

437 **Table 2** Recoveries of OPEs in each medium.

438 **Table 3** The obtained log K_{oc} between water-SPM and water-sediment.

439

440 **Table 1** Information about the sampling sites.

Number	Location	Latitude and longitude	Position	Water type	Aera(km ²)
1#	Lianghe City Park	30°41'55"N 104°02'54"E	Northwest	Closed	4.44
2#	Fenghuangshan Park	30°75'20"N 104°08'72"E	North	Closed	10
3#	Shahe Park	30°42'58"N 104°04'10"E	North	Semiclosed	--
4#	Jinghu, Southwest Jiaotong University	30°39'35"N 104°03'10"E	North	Closed	--
5#	Huanhuaxi Park	30°39'38"N 104°01'43"E	Center	Relatively open	0.32
6#	The people's Park	30°39'31"N 104°03'19"E	Center	Closed	0.11
7#	East Lake Park	30°37'14"N 104°05'02"E	Southeast	Relatively open	1.52
8#	Jincheng Park	30°34'15"N 104°03'10"E	South	Relatively open	1.6
9#	Daguanyan	30°37'07"N 104°07'38"E	Southeast	Closed	--
10#	Bailu Bay Wetland Park	30°34'05"N 104°07'45"E	Southeast	Semiclosed	2
11#	Qinglong Lake Wetland Park	30°38'36"N 104°11'10"E	Southeast	Closed	20

441

442 **Table 2** Recoveries of OPEs in each medium.

compounds	Water	SPM	Sediments
TnBP	90~111%	82~130%	89~112%
TCEP	81~102%	90~112%	83~108%
TCIPP	75~91%	81~115%	84~99%
TDCPP	90~116%	85~120%	103~119%
TPhP	83~99%	96~110%	76~117%
TBEP	92~108%	82~109%	80~103%
TEHP	91~114%	75~98%	89~126%

443

444 **Table 3** The obtained log K_{oc} between water-SPM and water-sediment.

Log K_{oc} between water and SPM											
	1#	2#	3#	4#	5#	6#	7#	8#	9#	10#	11#
TnBP	6.21	5.80	5.74	6.70	6.41	6.07	6.74	7.11	5.17	5.00	5.34
TCEP	5.69	5.57	5.23	6.45	5.95	5.44	5.99	6.78	4.72	4.66	5.79
TCPP	5.07	4.55	4.81	5.33	4.55	4.69	6.19	5.83	--	4.35	4.92

TPhP	5.59	5.12	5.30	5.69	5.26	5.27	5.57	6.21	--	5.31	5.35
TBEP	5.57	5.86	5.73	5.96	5.65	5.49	5.68	7.03	5.28	5.74	5.78
TEHP	5.56	5.40	5.44	5.80	5.21	5.27	5.72	6.53	5.24	5.33	5.40
mean	5.62	5.38	5.38	5.99	5.51	5.37	5.98	6.58	5.10	5.07	5.43
Log K_{oc} between water and sediment											
	1#	2#	3#	4#	5#	6#	7#	8#	9#	10#	11#
TnBP	4.61	4.98	4.41	4.80	4.42	4.50	4.22	4.69	4.15	4.55	5.04
TCEP	3.70	4.08	3.92	3.84	3.48	3.92	4.27	5.18	4.42	4.32	4.46
TCPP	3.77	3.89	--	3.13	3.49	4.22	4.28	--	--	4.34	4.10
TPhP	4.51	--	3.99	4.08	--	5.00	4.90	--	--	5.14	--
TBEP	--	--	5.08	3.91	--	4.96	4.88	--	4.15	5.07	4.71
TEHP	4.54	4.82	4.51	3.90	4.36	4.94	4.61	5.42	4.23	5.42	4.68
mean	4.23	4.44	4.38	3.94	3.94	4.59	4.53	5.10	4.24	4.81	4.60

Figure captions:

Fig. 1 Map of sampling sites.

Fig. 2 Correlations between the log K_{oc} values of OPEs and their log K_{ow} in water and SPM (black square: the present study; blue square: the study of Zhang et al. 2018).

Fig. 3 Correlations between the log K_{oc} values of OPEs and their log K_{ow} in water and sediments (black square: the present study; pink square: the study of Wang et al. 2018).

Fig. 4 Correlations between the log K_{oc} values of OPEs and their log K_{ow} in water and sediments (only for site 10#).

Fig. 1 Map of sampling sites.

Fig. 2 Correlations between the log K_{oc} values of OPEs and their log K_{ow} in water and SPM (black square: the present study; blue square: the study of Zhang et al. 2018).

Fig. 3 Correlations between the log K_{oc} values of OPEs and their log K_{ow} in water and sediments (black square: the present study; pink square: the study of Wang et al. 2018).

Fig. 4 Correlations between the log K_{oc} values of OPEs and their log K_{ow} in water and sediments (only for site 10#).

Supplementary Table and Figure

Table S1 Correlations between water, sediments and SPM.

Fig. S1 Correlation between the log K_a and log K_b .

Table S1 Correlations between water, sediments and SPM.

		water	sediment	SPM			water	sediment	SPM
1#	water	1	.873*	.880*	4#	water	1	.689	.690
	sediment	.873*	1	.990**		sediment	.689	1	.998**
	SPM	.880*	.990**	1		SPM	.690	.998**	1
2#	water	1	.549	.459	5#	water	1	.633	.552
	sediment	.549	1	.975**		sediment	.633	1	.983**
	SPM	.459	.975**	1		SPM	.552	.983**	1
3#	water	1	.197	.597	6#	water	1	.881*	.877*
	sediment	.197	1	.778		sediment	.881*	1	.948**
	SPM	.597	.778	1		SPM	.877*	.948**	1
		water	sediment	SPM			water	sediment	SPM
7#	water	1	.551	.895*	10#	water	1	.823*	.250
	sediment	.551	1	.572		sediment	.823*	1	.659
	SPM	.895*	.572	1		SPM	.250	.659	1
8#	water	1	.241	.775	11#	water	1	.822*	.679
	sediment	.241	1	.457		sediment	.822*	1	.875*
	SPM	.775	.457	1		SPM	.679	.875*	1
9#	water	1	.981**	.894*	Total	water	1	.685	.735

sediment	.981**	1	.850*	sediment	.685	1	.991**
SPM	.894*	.850*	1	SPM	.735	.991**	1

473 **. $p < 0.01$ (2-tailed)

474 *. $p < 0.05$ (2-tailed)

475

476

477 Fig. S1 Correlation between the $\log K_a$ and $\log K_b$.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryTableandFigure.docx](#)
- [Supplementaldata.xlsx](#)
- [1.jpg](#)