

# The effects of protective lung ventilation on regional cerebral oxygen saturation in intracranial tumor operation during dura opening—study protocol for a randomized controlled trial.

**Jinlu Li**

The Second Affiliated Hospital of Soochow University <https://orcid.org/0000-0003-3248-2342>

**Xuemei Wu**

The Second Affiliated Hospital of Soochow University

**Hairui Liu** (✉ [hhr218@163.com](mailto:hhr218@163.com))

<https://orcid.org/0000-0002-3784-646X>

**Ying Huang**

The Second Affiliated Hospital of Soochow University

**Yueqin Liu**

The Second Affiliated Hospital of Soochow University

**Hong Xie**

The Second Affiliated Hospital of Soochow University

**Jun Dong**

The Second Affiliated Hospital of Soochow University

---

## Study protocol

**Keywords:** Randomized controlled trial, Regional cerebral oxygen saturation, Lung ultrasound, Brain relaxation, Craniotomy

**Posted Date:** December 19th, 2019

**DOI:** <https://doi.org/10.21203/rs.2.15841/v3>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published on February 7th, 2020. See the published version at <https://doi.org/10.1186/s13063-019-4025-9>.

# Abstract

**Objective:** To investigate the effects of protective lung ventilation on regional cerebral oxygen saturation during dura opening, that is from Ta (after dura opening) to Tb (before dura closing), in patients undergoing intracranial tumor surgery. **Methods:** This is a randomized, controlled trial which will be carried out at the second affiliated hospital of Soochow University. Fifty-four patients undergoing intracranial tumor surgery will be randomly allocated to the control group (C group) or the protective lung ventilation group (P group). In the C group, tidal volume (VT) will be set at 8 ml/kg of predicted body weight, but positive end-expiratory pressure (PEEP) and recruitment maneuvers will not be used. In the P group, VT will be set at 6 ml/kg of predicted body weight combined with individualized PEEP during dura opening, while in other periods of general anesthesia, VT will be set at 8 ml/kg of predicted body weight. Regional cerebral oxygen saturation (rSO<sub>2</sub>), partial pressure of oxygen (PaO<sub>2</sub>) and carbon dioxide (PaCO<sub>2</sub>), oxygenation index (OI), lactic acid level (Lac) in arterial blood, and mean arterial pressure (MAP) will be compared before anesthesia (T<sub>0</sub>), before dura opening (T<sub>1</sub>), after dura closing (T<sub>2</sub>) and 24 h after surgery (T<sub>3</sub>). Lung ultrasound scores (LUS) will be performed at T<sub>0</sub> and T<sub>3</sub>. The degree of brain relaxation at T<sub>1</sub> and T<sub>2</sub> will be evaluated by the surgeon using the brain relaxation scale. Amount of vasoactive drugs used and blood loss will be recorded during surgery. The duration of operation and reoperation rate will be recorded. The primary outcome of this study is the changes in rSO<sub>2</sub> within 24 hours postoperatively. **Discussion:** This study aims to determine whether protective lung ventilation during dura opening can improve regional cerebral oxygen saturation and the state of pulmonary ventilation in patients undergoing intracranial tumor surgery, and to investigate whether this strategy does not affect the degree of brain tissue swelling and the reoperation rate after operation. If our results are positive, this study will show that protective lung ventilation during dura opening can be used effectively and safely in neurosurgical patients undergoing craniotomy for tumor resection. Trial registration: [chictr.org.cn](http://chictr.org.cn), ID: ChiCTR1900025632. Registered on 3 September 2019. Study protocol version 3.0. **Keywords:** Randomized controlled trial, Regional cerebral oxygen saturation, Lung ultrasound, Brain relaxation, Craniotomy

## Background

With the development of society and the progress of science and technology, more and more people receive delicate and complex procedures such as neurosurgery. Almost all these patients are under general anesthesia which is basically inseparable from mechanical ventilation. 15%-20% of patients had different degrees of alveolar collapse at the bottom of the lung before operation, and this phenomenon could persist for several days after operation due to the influence of mechanical ventilation of endotracheal intubation. The pulmonary complications play an important role in death and disability in patients with general anesthesia[1-3]. Craniotomy always needs a long time for general anesthesia and prolonged mechanical ventilation, which leads to a higher risk of postoperative atelectasis and pulmonary infection[4,5]. Atelectasis and pulmonary infection can seriously affect pulmonary ventilation, even lead to severe hypoxemia. Moreover, the long period of brain operation is more likely to cause the

imbalance of brain oxygen supply and consumption. The unbalance of brain oxygen supply and consumption may lead to deterioration of brain function, such as postoperative cognitive function[6]. Postoperative cognitive dysfunction (POCD) will lower the quality of life, increase mortality and aggravate the financial and mental burden of patients.

Protective lung ventilation (PLV) strategies have been recognized by many anesthesiologists and widely used in clinical anesthesia[7,8]. Relevant studies suggest that low-tide volume combined with positive end-expiratory pressure (PEEP) ventilation and alveolar recruitment maneuver (ARM) is the most widely used lung protective ventilation strategy, which can reduce lung volume damage and pulmonary barotrauma, improve pulmonary function and decrease postoperative pulmonary complications[9]. Theoretically, low tide volume prevents excessive alveolar expansion[10] and higher PEEP prevents pulmonary atelectasis[11]. However, gradually increased PEEP to the level of 20 cm H<sub>2</sub>O or even higher is often needed in traditional protective lung ventilation strategies[5,9] which will obviously affect the circulation and intracranial pressure of patients[12], and may increase airway pressure, reduce cerebral venous reflux and intraoperative operating space, thus limiting its application in patients with craniotomy. In addition, anesthesiologists often use a single PEEP or pulmonary retention mode, ignoring individual differences among patients, thus affecting the effect of protective lung ventilation[13,14].

In recent years, with the development of medical monitoring equipment, regional cerebral oxygen saturation monitoring technology[15,16] has been gradually developed and used in clinical anesthesia. It provides a condition for real-time monitoring the perfusion level of brain tissue in patients undergoing craniotomy, and provides technical support for carrying out clinical research on protective pulmonary ventilation during craniotomy.

Near infrared spectrometer (NIRS) using near infrared technology is similar to the pulse oxygen monitor which is commonly used. Near-infrared light with wavelength of 650~1100nm has a good penetrability to human tissues such as scalp, skull and brain, even up to several centimeters. The major color base (Hemoglobin, Hb) attenuated in the intracranial area of NIR light result in the changes in the light intensity of penetrating human tissues. The oxygenation of brain tissues was evaluated by measuring the changes in the absorption spectrum, which was accompanied with changes in oxygenation state[17].

Up to present, in consideration of the risks associated with PEEP and recruitment maneuvers, there are no correlative randomized controlled trials to explore the efficacy and safety of intraoperative pulmonary protective ventilation strategies in patients undergoing craniotomy. However, due to the disappearance of intracranial pressure after the dura opening during craniotomy, individualized protective pulmonary ventilation strategy may avoid the adverse effects on cerebral perfusion. The purpose of this study is to evaluate the effects of protective lung ventilation strategies with individualized PEEP during dura opening on regional cerebral oxygen saturation in patients undergoing intracranial tumor surgery. Other outcomes include intraoperative brain relaxation, the lung ultrasound scores 24h after surgery, the reoperation rate within one week after operation, the amount of blood loss and dosage of vasoactive drugs during surgery.

# Methods

## Study design

This is a single-center, randomized, controlled trial which is being conducted at the second affiliated hospital of Soochow University. Recruitment began in 3 September 2019. All patients eligible for inclusion will be collected continuously till the recruitment completion. Schedule of enrolment, interventions, and assessments will be shown in figure 1. Basic information of patients will be shown in table 1.

Patients will be randomized after signed informed consent and confirm inclusion and exclusion criteria. Randomization will be conducted via a computer-generated age stratification randomized controlled table. Age stratification: a. 18≤younger≤40; b. 40≤elder≤65. Patients who meet the enrollment criteria will be randomly allocated to control group (C group) or protective lung ventilation group (P group) within 24 h before surgery. The allocation ratio is 1:1. Permuted randomization will be used and stratified by age. The designated staff will perform the allocation sequence. The designated staff assistants will assign participants to interventions. This research staff will implement the allocation sequence through sealed, opaque and stapled envelopes. Corresponding envelopes will not be opened until the enrolled participants complete the trial. The anesthesiologist who is responsible for the anesthesia implementation will know the grouping but not participate in the follow-up visit. However, the neurosurgeon who evaluates brain relaxation will be blinded to the group allocation. The patients and the outcome assessor are all blinded to the grouping.

Tab 1. Patient Characteristics and Baseline Data

Characteristic	C Group	P Group
Male/female		
Age, yr		
BMI, kg/m <sup>2</sup>		
Predicted body weight, kg		
ASA physical status, I/II		

## Selection and withdrawal of participants

### Recruitment

Participants will be recruited from the neurosurgical wards and identified by their presence on surgical lists. The investigator informs the participant or the participant's legal representative of all aspects. The study intervention will be completed immediately after the surgery, but follow-up visits will extend to 1 week after surgery. The medical records will be reviewed following hospital discharge for in-hospital complications and medication usage.

## Inclusion criteria

1. Patients scheduled to receive elective intracranial primary tumor resection who aged between 18 and 65 years old
2. The maximum diameter of the tumor is 2~5 cm (Magnetic Resonance Imaging, MRI)
3. ASA class forⅡ-Ⅲ
4.  $18.5 \leq \text{BMI} \leq 28$
5. Glasgow Coma Scale score of more than 8 points

## Exclusion criteria

1. Patients with chronic lung disease, pulmonary infection and other severe pulmonary complications such as acute respiratory failure
2. Patients with a history of pulmonary surgery
3. Patients with severe brain, heart, liver and kidney diseases
4. Patients with nerve injury affecting breathing preoperative
5. Pregnant women
6. Refuse to participate in the researcher

## Termination criteria

1. Duration of anesthesia  $<4$  h or  $>8$  h; or duration of operation  $<2$  h or  $>6$  h
2. Patients with significantly increased intraoperative intracranial pressure or swelling of brain tissue
3. Patients with intraoperative endotracheal catheter after surgery
4. Repeat intubation or operation within 24 h after operation

## Study intervention

### Related parameter setting during operation

All patients will be randomly allocated to the C group and the P group according to the computer-generated random number table. In the C group, tidal volume (VT) will be set at 8 ml/kg of predicted body weight, with PEEP=0, and recruitment maneuver will not be used. The predicted body weight is calculated as follows: for men,  $50+0.91(\text{height in centimeters}-152.4)$ ; and for women,  $45.5+0.91(\text{height in centimeters}-152.4)$ [18]. In the P group, VT will be set at 6 ml/kg of predicted body weight combined with individualized PEEP during intraoperative dura mater opening[6,7], but in other periods of general anesthesia, VT will be set at 8 ml/kg of predicted body weight. Titration method of individualized PEEP[19]: VT and respiratory rate will be fixed at 6ml/kg and 15 beats per minute during PEEP trial. Titration can only begin once the dura is opened. The titration for the individual PEEP can then be initiated by increasing PEEP from 0 to 10 cm H<sub>2</sub>O incrementally. Each PEEP level (0, 1, 2, 3, 4, 5, 6, 7, 8, 9,

10 cmH<sub>2</sub>O) will be maintained for 1 minute, and the pulmonary compliance of the last cycle will be recorded at each PEEP level. At last, the PEEP value at the highest compliance will be selected as the individual PEEP of patient.

Peripheral venous access will be established after the patient enters the operating room. If necessary, central venous access will be established. Noninvasive blood pressure (NBP), electrocardiogram (ECG), heart rate (HR), Oxygen saturation (SpO<sub>2</sub>) and bispectral index (BIS) will be routinely monitored. Radial artery catheterization under local anesthesia will be used to monitor invasive arterial pressure and collect blood samples. All of the above data will be collected completely. Fentanyl 5ug/kg, etomidate 0.3mg /kg, rocuronium 0.6mg/kg will be used for induction which will be started after oxygen flow of 0.1 L/kg/min was given by mask for 2 min. Volume controlled mechanical ventilation will be conducted by Primus anesthesia machine (Drager, Germany) after endotracheal catheter is inserted to a correct position. The VT will be set 8 ml/kg of predicted body weight, the inhalation oxygen fraction (FiO<sub>2</sub>) will be set at 0.5, the inhalation-expiration ratio (I: E)= 1:2, and the fresh gas flow will be set at 1 L/min. The respiration rate will be adjusted according to the result of end-expiratory carbon dioxide (ETCO<sub>2</sub>), and the end-tidal CO<sub>2</sub> pressures (PetCO<sub>2</sub>) will be maintained between 30-35 mmHg. 1% sevoflurane combined with propofol and remifentanyl to maintain anesthesia, and BIS value will be maintained at 45~55. During the operation, intermittent injection of fentanyl and rocuronium will be used to deepen the anesthesia. Extubation indications: patients were awake and cooperated, and muscle relaxation monitoring train of four stimulation (TOF)>90%[20]. Intraoperative fluid intake and urine volume will be monitored closely. Regional cerebral oxygen saturation (rSO<sub>2</sub>) will be recorded in the tumor surgery area of the patients before anesthesia (T0), before dura opening (T1), after dura closing (T2) and 24 h after surgery (T3). Arterial blood of the patients was collected for blood gas analysis. Lung ultrasound scores (LUS) will be performed at T0 and T3. The degree of brain relaxation at T1 and T2 will be evaluated by the surgeon using the brain relaxation scale. Amount of vasoactive drugs used and blood loss will be recorded during surgery. The duration of operation and reoperation rate will be recorded.

## **Study objective**

### Primary outcome

The primary outcome of this study is the changes in rSO<sub>2</sub> within 24 hours postoperatively.

The secondary outcomes are as follows:

1. Changes between preoperative and postoperative in Lung ultrasound in patients. Lung ultrasound scores will be used to evaluate postoperative atelectasis. The patient's chest is divided into 12 quadrants. Each of the 12 quadrants is assigned a score of 0 to 3 according to a modified grading system (Table 2). The LUS score (0-36) is then calculated by adding up the 12 individual quadrant scores with higher scores indicating more severe atelectasis[21].
2. The mean arterial pressure changes during intraoperative pulmonary protective ventilation.

3. The partial pressure of oxygen and carbon dioxide, oxygenation index, lactic acid level in arterial blood changes during intraoperative pulmonary protective ventilation.
4. The amount of vasoactive drugs and blood loss will be compared in two groups during surgery.
5. Intraoperative brain relaxation. Brain relaxation will be scored by the neurosurgeon after opening the cranium and before opening the dura. They will use a 4-point scale[22]: 1, completely relaxed; 2, satisfactorily relaxed; 3, firm brain; 4, bulging brain
6. The secondary operation rate in one week after surgery (Reoperation rate).

Table 2. Modified Lung Ultrasound Scores

	Normal Aeration	Small Loss of Aeration	Moderate Loss of Aeration	Severe Loss of Aeration
Quotation	0	1	2	3
Modified lung ultrasound score	0-2 B lines	≥3 B lines OR 1 or multiple small subpleural consolidations separated by a normal pleural line	Multiple coalescent B lines OR Multiple small subpleural consolidations separated by a thickened or irregular pleural line	Consolidation OR Small subpleural consolidation of >1 × 2 cm in diameter

Lung ultrasound scores can be calculated by adding up the 12 individual pulmonary quadrant scores yielding a score between 0 (no aeration loss) and 36 (complete aeration loss).

Tab 3. Perioperative parameters

	C Group	P Group
Tidal volume, ml		
Individual PEEP, cmH <sub>2</sub> O		
Duration of anesthesia, min		
Duration of operation, min		
Duration of dura opening, min		
Tumor size (maximum diameter)		
Brain relaxation scale (T1)		
Brain relaxation scale (T2)		
Lung ultrasound score (T0)		
Lung ultrasound score (T3)		
The volume of total fluid, ml		
The amount of bleeding, ml		
The volume of urine, ml		
Dosage of vasoactive agent, mg		
Reoperation rates		

Tab 4. Comparison of rSO<sub>2</sub>, PaO<sub>2</sub>, PaCO<sub>2</sub>, OI, Lac and MAP

		T0	T1	T2	T3
rSO <sub>2</sub>	C Group				
	P Group				
PaO <sub>2</sub>	C Group				
	P Group				
PaCO <sub>2</sub>	C Group				
	P Group				
OI	C Group				
	P Group				
Lac	C Group				
	P Group				
MAP	C Group				
	P Group				

T0:Before anesthesia; T1: Before dura opening; T2:After dura closing; T3:24h after surgery; rSO<sub>2</sub>: regional cerebral oxygen saturation; PaO<sub>2</sub>: partial pressure of oxygen; PaCO<sub>2</sub>: partial pressure of carbon dioxide; OI: oxygenation index; Lac: lactic acid level; MAP: mean arterial pressure.

### Reporting of adverse events

All adverse events will be recorded and closely monitored until resolution or stabilization. In the event of any serious adverse event ( $\geq$ grade 3)[23], it will be immediately reported to the Endpoint Adjudication Committee, which will determine the severity and causality of the adverse events. The chief investigator will be responsible for all adverse event reporting.

### Withdrawal from the trial

We will consider patient withdrawal from the trial if the following conditions occur: (1) severe brain swelling during the operation; (2) the patient has a cough during surgery; (3) the patient has persistent hypotension and circulatory instability.

### Data collect and management

All the patient information will be obtained through the electronic medical record system. The consent of the treating neurosurgeon, who will help us make the neurological diagnosis, has also been obtained. All personal information will be collected through the hospitalized medical records by a member of the research team and be kept strictly confidential for research purposes only. The research team members will be responsible for maintaining personal data. Only the primary investigator and the designated researcher can obtain interim results and final test data.

## Data Monitoring Committee (DMC)

The project will be monitored by a Data Monitoring Committee (DMC) composed of specialists in anesthesiology, ethics, statistics and methodology. The DMC will audit through regular interviews or telephone calls.

## Sample size and justification

We calculated the sample size through the website <http://www.sample-size.net/sample-size-proportions/>.

The difference of brain oxygen saturation before and after surgery was  $3.6 \pm 4.1$ ,  $\alpha=0.05$ ,  $\beta=0.2$ [7]. Based on this, it can be calculated that the sample size required for our study is 44 cases, plus 20% shedding rate, a total of 53 cases ( $44+44*20\%$ ) need to be recruited. Due to the 1:1 distribution ratio, a total of 54 cases will be recruited.

## Statistics

The SPSS 19.0 software package for Windows (SPSS, Inc. , Chicago, IL, USA) will be used for all statistical analyses. The quantitative variables will be expressed as mean $\pm$ SD or median (interquartile range (IQR)), and be analyzed by using the analysis of variance (ANOVA) or Mann-Whitney U test. Multiple comparison among continuous variables at different time points will be performed using S-N-K method for post hoc test. The incidence of reoperation rate will be expressed as the number of patients (percentage), and be analyzed by using the chi-square ( $\chi^2$ ). P value < 0.05 will be considered to be statistically significant. After the follow-up of half of the cases, the interim analysis will be conducted to evaluate the validity of the main results.

## Discussion

This study is a single-center, randomized, controlled trial of exploring whether protective lung ventilation during intraoperative dura opening can improve regional cerebral oxygen saturation in neurosurgical patients.

The incidence of postoperative pulmonary complications (PPCs) is high due to the long mechanical ventilation in neurosurgery. Qaseem et al.[24] reported that the risk of PPCs increased when operation time is more than 4 h. The incidence of PPCs was 28.4% (20.2–37.9%) in patients with neurosurgery lasting for longer than 300 min[25]. PEEP can lower the incidence of postoperative respiratory complications, prevent atelectasis and reduce the risk of ventilators associated with lung injury.

In this study, individual PEEP (< 10cmH<sub>2</sub>O) will be used to avoid the effect of high PEEP on intracranial pressure (ICP). It is a crucial issue that PEEP can be safely used in craniotomy. Therefore, pulmonary protective ventilation will be performed during dura opening and cerebral relaxation will be assessed before dura incision. If the intracranial pressure is elevated enough to affect the operation for using PEEP,

we will abandon the case and change ventilation parameters. The case will be reported to the principal investigator.

Regional cerebral oxygen saturation (rSO<sub>2</sub>) is actually the mixed oxygen saturation of local brain tissues, which can better reflect the change of brain oxygen supply and consumption balance during perioperative period. Samra et al. [26] studied 100 patients who underwent carotid endarterectomy, then found that if the rSO<sub>2</sub> value decreased by 20% compared with the baseline value after internal carotid artery occlusion, it predicted the possibility of neurological complications, and indicated that its sensitivity was 80% and specificity was 82%. Since the ratio of cerebral blood volume to arterial/venous blood flow is approximately 20:80, NIRS mainly represents cerebral venous oxygen saturation, which is completely unaffected by hypoxemia and hypocarbonemia, and better reflects the changes in the balance of oxygen supply and consumption in the brain[27]. Near infrared spectroscopy (NIRS) as a brain oxygen monitoring method has following characters: continuous and noninvasive, convenient, high degree of sensitivity and specificity[28]. Monitoring regional cerebral oxygen saturation can detect changes of the cerebral blood flow and oxygen supply and consumption balance in the brain area as early as possible, and judge the degree of cerebral ischemia and hypoxia and the changes of brain function. Timely adjustment of anesthesia plan is helpful to guide perioperative anesthesia management, so as to prevent POCD, shorten hospitalization period and improve quality of life.

We focus on whether pulmonary protective ventilation strategy can affect cerebral venous reflux and brain tissue oxygenation, and ultimately, the prognosis of patients. Protective lung ventilation after incision of dura can reduce the returned blood volume that results in exposing potential bleeding spots, which is beneficial for the surgeon to stop bleeding. Due to the opening dura, the intracranial pressure disappeared, and the decreased cerebral perfusion pressure caused by the expansion of the lung would be improved.

This study is a prospective, randomized controlled trial. This study aims to investigate the effect of intraoperative pulmonary protective ventilation in neurosurgical craniotomy. If we are able to demonstrate the safety and effectiveness of intraoperative pulmonary protective ventilation with individualized PEEP during dura opening in neurosurgical craniotomy, it will improve the prognosis of neurosurgical patients and reduce the medical costs.

## **Trial Status**

The study was also registered on the registry website <http://chictr.org.cn/> with the registration number ChiCTR1900025632 on 3 September 2019. Protocol version 3.0, 3/9/2019. The study began on 3 September 2019, and the planned completion date will be September 2020. Trial status was currently recruiting. The recruitment began on 3 September 2019, and the planned completion date will be June 2020.

## **List Of Abbreviations**

VT: tidal volume

PEEP: positive end-expiratory pressure

rSO<sub>2</sub>: regional cerebral oxygen saturation

PaO<sub>2</sub>: partial pressure of oxygen

PaCO<sub>2</sub>: carbon dioxide

OI: oxygenation index

Lac: lactic acid level

MAP: mean arterial pressure

PLV: protective lung ventilation

ARM: alveolar recruitment maneuver

NIRS: near infrared spectrometer

PPCs: postoperative pulmonary complications

ICP: intracranial pressure

FiO<sub>2</sub>: oxygen fraction

NBP: noninvasive blood pressure

ECG: electrocardiogram

HR: heart rate

SpO<sub>2</sub>: oxygen saturation

BIS: bispectral index

ETCO<sub>2</sub>: end-expiratory carbon dioxide

PetCO<sub>2</sub>: end-tidal CO<sub>2</sub> pressures

## **Declarations**

### **Ethics approval and consent to participate**

Ethics Committee of the Second Affiliated Hospital of Soochow University approved the study on November 23, 2018 (File number EC-AF (JD)-06/6.1, study protocol version 3.0). The study is designed in accordance to the principles of the Declaration of Helsinki. Written informed consent was obtained from every enrolled patient upon request by the review board.

### **Consent for publication**

Not applicable.

### **Availability of data and materials**

The datasets analyzed during the current study are available from the corresponding author on reasonable request.

### **Competing interests**

All authors declare that they have no competing non-financial/financial interests.

### **Funding**

This study was not funded.

### **Authors' contributions**

Hairui Liu conceived and designed the study, coordinates the overall study, and contributed to the final manuscript. Jinlu Li conceived and designed the study, coordinates the overall study, and contributed to the final manuscript. Xuemei Wu participated in the design of the study and drafted the manuscript. Ying Huang and Yueqin Liu performed the sample size calculation, drafted the statistical analysis plan and collected the data. Hong Xie participated in the design of the study. Jun Dong conceived the study, and guided the calculation of the sample size and the plans of the statistical analysis. All authors read and approved the final manuscript.

### **Corresponding author**

Correspondence to Hairui Liu.

### **Acknowledgements**

Not applicable.

### **Authors' information**

Not applicable.

## **References**

- [1]. Sutherasan Y, Vargas M, Pelosi P. Protective mechanical ventilation in the non-injured lung: review and meta-analysis. *Critical Care*, 2014,18(2):1-12.
- [2]. Fernandez-Bustamante, A., et al. Postoperative Pulmonary Complications, Early Mortality, and Hospital Stay Following Noncardiothoracic Surgery: A Multicenter Study by the Perioperative Research Network Investigators. *JAMA Surg*, 2017,152(2):157-166.
- [3]. Finks, J. F. , N. H. Osborne and J. D. Birkmeyer, Trends in hospital volume and operative mortality for high-risk surgery. *N Engl J Med*, 2011,364(22):2128-37.
- [4]. [Dasenbrock HH](#), [Liu KX](#), [Devine CA](#), et al. Length of hospital stay after craniotomy for tumor a National Surgical Quality Improvement Program analysis. *Neurosurg Focus*, 2015,39(6):E12.
- [5]. [Kourbeti IS1](#), [Vakis AF](#), [Ziakas P](#), et al. Infections in patients undergoing craniotomy risk factors associated with post-craniotomy meningitis. *J Neurosurg*, 2015,122(5):1113-9.
- [6]. Kim, J. , et al. Postoperative Cognitive Dysfunction and the Change of Regional Cerebral Oxygen Saturation in Elderly Patients Undergoing Spinal Surgery. *Anesthesia & Analgesia*, 2016,123(2):436-444.
- [7]. [Damian D](#), [Esquenazi J](#), et al. Incidence, outcome, and risk factors for postoperative pulmonary complications in head and neck cancer surgery patients with free flap reconstructions. *J Clin Anesth*, 2016,28(2):12-8.
- [8]. [Mercado P](#), [Maizel J](#), [Kontar L](#), et al. Moderate and Severe Acute Respiratory Distress Syndrome: Hemodynamic and Cardiac Effects of an Open Lung Strategy With Recruitment Maneuver Analyzed Using Echocardiography. *Crit Care Med*, 2018,46(10):1608-1616.
- [9]. Hew M, Tay TR. The efficacy of bedside chest ultrasound: from accuracy to outcomes. *Eur Respir Rev*, 2016,25(141):230-46.
- [10]. Tojo, K. , et al. Atelectasis causes alveolar hypoxia-induced inflammation during uneven mechanical ventilation in rats. *Intensive Care Med Exp*, 2015,3(1):56.
- [11]. Chikhani, M. , et al. High PEEP in acute respiratory distress syndrome: quantitative evaluation between improved arterial oxygenation and decreased oxygen delivery. *Br J Anaesth*, 2016,117(5):650-658.
- [12]. [Flexman A M](#), [Gooderham P A](#), [Griesdale D E](#), et al. Effects of an alveolar recruitment maneuver on subdural pressure, brain swelling, and mean arterial pressure in patients undergoing supratentorial tumour resection: a randomized crossover study. *Can J Anaesth*, 2017,64(3):626-633.
- [13]. [Pereira S M](#), [Tucci M R](#), [Morais C C](#), et al. Individual Positive End-expiratory Pressure Settings Optimize Intraoperative Mechanical Ventilation and Reduce Postoperative Atelectasis. *Anesthesiology*, 2018,129(6):1070-1081.

- [14]. Costa Leme, A. , et al. Effect of Intensive vs Moderate Alveolar Recruitment Strategies Added to Lung-Protective Ventilation on Postoperative Pulmonary Complications. *JAMA*, 2017,317(14):1422.
- [15]. Venkateswaran P, Sriganesh K, et al. Regional Cerebral Oxygen Saturation Changes After Decompressive Craniectomy for Malignant Cerebral Venous Thrombosis. *J Neurosurg Anesthesiol*, 2019,31(2):241-246.
- [16]. Jia Z, Teng Y, Liu Y, et al. Influence of high-flow modified ultrafiltration on brain oxygenation. *Perfusion*, 2018,33(3):203-208.
- [17]. Zheng, F. , et al. Cerebral near-infrared spectroscopy monitoring and neurologic outcomes in adult cardiac surgery patients: a systematic review. *Anesth Analg*, 2013,116(3):663-76.
- [18]. Treschan TA, Kaisers W, Schaefer MS, et al. Ventilation with low tidal volumes during upper abdominal surgery does not improve postoperative lung function. *Br J Anaesth*, 2012,109:263-271.
- [19]. Park M1, Ahn HJ, Kim JA, et al. Driving Pressure during Thoracic Surgery: A Randomized Clinical Trial. *Anesthesiology*, 2019,130(3):385-393.
- [20]. Eriksson, L. I., et al. Functional assessment of the pharynx at rest and during swallowing in partially paralyzed humans: simultaneous video manometry and mechanomyography of awake human volunteers. *Anesthesiology*, 1997,87(5):1035-43.
- [21]. Monastesse A, Girard F, Massicotte N, et al. Lung Ultrasonography for the Assessment of Perioperative Atelectasis: A Pilot Feasibility Study. *Anesth Analg*, 2017,124:494-504.
- [22]. Prabhakar H, Singh GP, Mahajan C, et al. Intravenous versus inhalational techniques for rapid emergence from anaesthesia in patients undergoing brain tumour surgery. *Cochrane Database Syst Rev*, 2016,9:CD010467.
- [23]. National Cancer Institute. Common Terminology Criteria for Adverse Events. Version 4.0. [https://evs.nci.nih.gov/ftp1/CTCAE/CTCAE\\_4.03/CTCAE\\_4.03\\_2010-06-14\\_QuickReference\\_8.5x11.pdf](https://evs.nci.nih.gov/ftp1/CTCAE/CTCAE_4.03/CTCAE_4.03_2010-06-14_QuickReference_8.5x11.pdf)
- [24]. Qaseem, A. , et al. Risk assessment for and strategies to reduce perioperative pulmonary complications for patients undergoing noncardiothoracic surgery: a guideline from the American College of Physicians. *Ann Intern Med*, 2006,144(8):575-80.
- [25]. Sogame, L. C. , et al. Incidence and risk factors for postoperative pulmonary complications in elective intracranial surgery. *J Neurosurg*, 2008,109(2):222-227.
- [26]. Samra, S. K. , et al. Evaluation of a Cerebral Oximeter as a Monitor of Cerebral Ischemia during Carotid Endarterectomy. *Anesthesiology*, 2000,93(4): 964-970.

[27]. Plachky, J. , et al. Regional cerebral oxygen saturation is a sensitive marker of cerebral hypoperfusion during orthotopic liver transplantation. *Anesth Analg*, 2004,99(2): 344-9, table of contents.

[28]. Brogan RJ, Kontojannis V, Garara B, et al. Near-infrared spectroscopy(NIRS) to detect traumatic intracranial haematoma: A systematic review and meta-analysis. *Brain Inj*, 2017,(31):581-588.

## Figures

TIMEPOINT**	STUDY PERIOD								
	Enrolment	Allocation	Post-allocation						
	-1day	Surgery day	T0	T1	Ta	Tb	T2	T3	
<b>ENROLMENT:</b>									
Eligibility screen	X								
Informed consent	X								
Random allocation		X							
<b>INTERVENTIONS:</b>									
<i>the control group</i>									
<i>the protective ventilation group</i>									
<b>ASSESSMENTS:</b>									
<i>Regional cerebral oxygen saturation</i>			X	X			X	X	
<i>Partial pressure of oxygen</i>			X	X			X	X	
<i>Partial pressure of carbon dioxide</i>			X	X			X	X	
<i>Oxygenation index</i>			X	X			X	X	
<i>Lactic acid</i>			X	X			X	X	
<i>Mean arterial pressure</i>			X	X			X	X	
<i>Lung ultrasound score</i>			X					X	
<i>The amount of vasoactive drugs</i>							X		
<i>Blood loss</i>							X		
<i>Brain relaxation</i>				X			X		
<i>The secondary operation rate</i>									

Figure 1

Schedule of enrolment, interventions, and assessments T0:Before anesthesia;T1:Before dura opening;Ta:After dura opening; Tb:Before dura closing;T2:After dura closing;T3:24h after surgery