

Characterization of genetic determinants of the resistance to phylloxera, *Daktulosphaira vitifoliae*, and the dagger nematode *Xiphinema index* from muscadine background

Bernadette Rubio

INRA

Guillaume Lalanne-Tisné

INRA

Roger Voisin

INRA

Jean-Pascal Tandonnet

INRA

Ulysse Portier

INRA

Cyril Van Ghelder

INRA

Maria Lafargue

INRA

Jean-Pierre Petit

INRA

Martine Donnart

INRA

Benjamin Joubard

INRA

Pierre-Francois Bert

Universite de Bordeaux <https://orcid.org/0000-0003-3209-952X>

Daciana Papura

INRA

Loïc Le Cunff

IFV

Nathalie Ollat (✉ nathalie.ollat@inrae.fr)

<https://orcid.org/0000-0002-6182-9686>

Daniel Esmenjaud

INRA

Research article

Keywords: ,

Posted Date: February 18th, 2020

DOI: <https://doi.org/10.21203/rs.2.15851/v3>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Muscadine (*Muscadinia rotundifolia*) is known as a resistance source to many pests and diseases in grapevine. The genetics of its resistance to two major grapevine pests, the phylloxera *D. vitifoliae* and the dagger nematode *X. index*, vector of the Grapevine fanleaf virus (GFLV) was investigated in a backcross progeny between the F1 resistant hybrid material VRH8771 (*Muscadinia-Vitis*) and *V. vinifera* cv. 'Cabernet-Sauvignon'. Phylloxera resistance was assessed by monitoring root nodosity number in an "in planta" experiment and larval development in a root "in vitro" assay. Nematode resistance was studied using 10-12 month long tests for the selection of durable resistance and rating criteria based on nematode reproduction factor and gall index. A major QTL for phylloxera larval development, explaining more than 70% of the total variance and co-localizing with a QTL for nodosity number, was identified on LG 7. Additional QTLs were detected on LG 3 and LG 10, depending on the "in planta" or "in vitro" experiments, suggesting that various loci may influence or modulate nodosity formation and larval development. Using a Bulk Segregant Analysis approach and a proportion test, markers clustered in three regions on LG 9, LG 10 and LG 18 were shown to be associated to the nematode resistant. All these results open the way to the development of marker-assisted selection in grapevine rootstock breeding programs based on muscadine derived resistance to phylloxera and to *X. index* in order to delay GFLV transmission.

Background

Domesticated grapevine, *Vitis vinifera* sub-sp. *vinifera* (This et al. 2006), is grown grafted in most countries worldwide. Grafting has been successfully used to cope mainly with phylloxera (*Daktulosphaira vitifoliae*), an insect-pest that destroyed the European vineyard after its introduction from North America in the middle of the 19th century (Pouget, 1990; Thiery & Vogelweith, 2016) and participate to the control of other soil-borne pests and adaptation to abiotic stresses.

In addition to Europe, grapevine phylloxera spread quickly to most wine grape growing regions of the world including South Africa, Middle East, Asia and Australia (Powell 2012). This insect has two different forms, the radicicoles and the gallicoles, which affect roots and leaves, respectively. Radicicoles are the most destructive in *V. vinifera* due to severe root damage, while gallicole forms are more common in most other species without lethal effects for the plants. Root feeding induces the formation of galls called nodosities (Powell 2012; Smith et al. 2018). Phylloxera root infection causes vine decline and finally plant death, partly explained by secondary fungal infections (Pouget 1990). The use of resistant rootstocks from *Vitis* species other than *V. vinifera* is advocated as the main method of radicicole phylloxera management and may be considered as the most sustainable example of biological control for a pest ever used (Powell 2008). Several sources of grape phylloxera resistance for rootstock breeding have been identified. Heritability studies have shown that a variable number of loci control the resistance trait (Davidis and Olmo 1964; Boubals 1966; Bouquet 1983; Roush et al. 2007; Ramming 2010). The first genetic mapping of a QTL for phylloxera resistance identified the *RDV1* (*RESISTANCE DAKTULOSPHEIRA VITIFOLIAE 1*) locus on chromosome 13 in the Börner (*V. riparia* x *V. cinerea*) rootstock (Zhang et al.

2009). The *RESISTANCE DAKTULOSPHEIRA VITIFOLIAE 2 (RDV2)* locus was mapped to chromosome 14 in *V. cinerea* cv. 'C2-50' (Smith et al. 2018). A leaf-specific phylloxera resistance locus overlapped the location of *RDV2* on LG 14 in a F1 family from a cross containing at least six *Vitis* species in the ancestry (*V. vinifera*, *V. riparia*, *V. rupestris*, *V. labrusca*, *V. aestivalis* and *V. berlandieri*). In contrast, from the same plant material, a locus for resistance to the root form of phylloxera mapped to chromosomes 5 and 10 (Clark et al. 2018).

In addition to phylloxera, rootstocks may contribute to the control of other soil-borne pests such as root-knot and dagger nematodes. Resistance to the dagger nematode *Xiphinema index* is an important objective in grape rootstock breeding programs. *X. index* is a migratory ectoparasite that primarily feeds on the root tips of grapevines and causes severe damage to their root system. More significantly, *X. index* is recognized as the vector of *Grapevine fanleaf virus (GFLV)*, the causal agent of the fanleaf degeneration disease which is considered to be one of the major threats to the grapevine industry (Hewitt et al. 1958; Andret-Link et al. 2004). *X. index* can survive in soils of ancient vineyards and retain GFLV for many years without the presence of host plant (Demangeat et al. 2005). Nematicides and fumigants failed to control the dagger nematode because of their poor penetration in deep soil layers where *X. index* mainly survives (Esmenjaud et al. 1992; Villate et al. 2008) and their use is now banned in most countries because of their high toxicity. In order to overcome these limitations, using nematode-resistant grapevine rootstocks appears as one of the most promising control method to significantly delay viral transmission. The highest level of resistance to *X. index* was found in *V. arizonica*, *V. candicans*, *V. rufotomentosa*, *V. smalliana* and *V. solonis* (Esmenjaud and Bouquet, 2009; Ollat et al. 2016). Resistance in *V. arizonica* is controlled by a single major locus, *XIPHINEMA INDEX RESISTANCE 1 (XiR1)* located on chromosome 19 (Xu et al. 2008; Riaz et al. 2012).

In the next decades, grape growing will have to face several challenges, such as climate change, decrease in pesticides application, water availability and competition for arable lands with food crops. In this context, it is important to consider the rootstock as a key component of adaptation and consequently, breeding new rootstocks is urgently needed. Rootstock breeding programs conducted throughout the world share similar goals, such as resistance to phylloxera and nematodes (dagger and root-knot nematodes), vigour management, mineral uptake efficiency, together with drought, salinity and limestone tolerances (Töpfer et al. 2011; Ollat et al. 2016). The world's existing rootstocks have a very narrow genetic base derived from only a few American grape species accessions (Riaz et al, 2019). Despite the diversity available among the genus *Vitis*, only 4 to 5 species are commonly used in crosses, i.e. *V. berlandieri*, *V. riparia*, *V. rupestris*, *V. champini* and *V. vinifera*. *Muscadinia rotundifolia*, the muscadine vine, was also considered as a highly interesting species for rootstock improvement because of its resistance to many pests and diseases (Bouquet 1983). Thus, hybrids between this American species and *V. vinifera* were created in order to be used as rootstocks (Walker et al. 1994) or as parents or elite materials for further crosses (Bouquet 1980). However, muscadines present major defaults such as poor rooting, and grafting incompatibility or mineral deficiencies, advanced crosses are necessary to breed highly performing rootstocks.

Grapevine breeding takes generally decades to release an advanced cultivar. Considering the difficulty to screen for most traits of interest in rootstocks, the development of molecular-assisted selection is a priority. This relies on the identification of one or few loci closely associated to the trait expression and explaining a high percentage of phenotypic variance. As previously mentioned, only four loci have been identified so far in response to phylloxera in various *Vitis* species and a single one in response to *X. index*. Therefore it is crucial to improve our knowledge about the resistance response of *Vitis* and *Muscadinia* spp. resources to major grapevine pests and to analyse their genetic determinism. Recent progress in the development and application of molecular markers, genetic mapping and whole genome sequencing combined with high throughput technologies will help to better characterize the genetic bases of the traits of interest (Varshney et al. 2009). For grapevine, the development and application of SSR (Simple Sequence Repeat) markers are considered as a key step of the construction of molecular maps (Doligez et al. 2006; Troglio et al. 2007; Riaz et al. 2012). Next-generation sequencing (NGS) has then facilitated the development of methods to genotype very large numbers of SNP (Single Nucleotide Polymorphisms) markers (Ray and Satya 2014; Vlček and Řepková 2017). Genotyping-by-sequencing (GBS) has been developed as a rapid and robust approach to sequencing of multiplexed samples (Elshire et al. 2011; Poland and Rife 2012). The work by Barba and collaborators (2014) demonstrated the first application of the GBS procedure in generating SNPs to construct a high resolution map for QTL mapping in grapevine.

In order to identify additional sources of resistance to the phylloxera and to the dagger nematode, a backcross 1 (BC1) population from the cross between a muscadine hybrid (VRH8771 = *V. vinifera* x *M. rotundifolia*) and *V. vinifera* cv. 'Cabernet-Sauvignon' has been characterized in this study. Through this cross, one seeks to expand the existing knowledge about the genetic determinism of *M. rotundifolia* natural resistances towards the pathogens and pests. The main objectives of our study were i) to build SSR and SNP-based linkage maps using pseudo-testcross strategy and ii) to exploit them for QTL mapping of the genetic bases of the resistance responses to these major grapevine pests. These goals are the first steps toward the development of marker-assisted breeding for resistance to phylloxera and *X. index*.

Results

Genotyping: sequencing, SNP calling and SNP selection

A total of 84 million cleaned reads were obtained, and the number of reads per sample ranged from 36,000 to 9.5 million, with an average of 3.05 million. This number of reads was equivalent to ~0.91-fold coverage of other *Vitis* genomes which are estimated to have a size of approximately 500 Mb. The SNP calling performed with GATK pipeline and VCFtools produced a total of 324,183 SNPs among which 52,625 have been identified as pseudotestcross markers. By excluding sites based on missing data or segregation distortion, two sets of 2,285 and 738 SNPs were obtained for the female (VRH8771) and the male (CS) maps, respectively. These sets were complemented by 135 and 148 SSRs genotyped on 90 BC1 individuals for the construction of the female and male maps, respectively. Because of the missing data threshold per individual and the genotype frequencies expected, the final dataset included 92 BC1

individuals. In the end, 75 BC1 individuals were genotyped with both SSRs and SNPs markers, 19 BC1 individuals with SNPs markers only and 17 BC1 individuals with SSRs markers only (Table S1).

High density maternal and paternal genetic maps

Maternal (VRH8771) and paternal (CS) maps were constructed using 2420 and 886 sets of markers (SSR and SNP markers), respectively (Fig. S1 and Fig S2). For each set of markers, 19 LGs were produced and the final sizes for the VRH8771 and CS maps were 2351 and 1982 cM, respectively. The map density or average distance between markers for VRH8771 and CS maps was 0.98 and 2.24 cM, respectively (Table 1). The comparison of marker order with their physical position in VRH8771 and CS maps showed that most plots were on the diagonal or adjacent position (Fig. S3).

Phenotypic evaluation of resistance to *D. vitifoliae*

The phenotypic scores of phylloxera root resistance were obtained in both *in planta* and *in vitro* experiments using 89 and 37 BC1 individuals, respectively. In the *in planta* experiment, results were expressed by a number of nodosities per plant whereas in the *in vitro* experiment they corresponded to a number of larvae counted on the five root pieces of each individual. A Spearman correlation between *in vitro* and *in planta* data was performed revealing a positive and significant correlation ($r = 0.47$, $p\text{-value} = 0.003$). The nodosity and larvae numbers varied considerably within the three replicates of each BC1 individual plant tested as illustrated by an average value of 73 % and 53 % for the variation coefficients of *in planta* and *in vitro* experiments, respectively. As this experimental system used the inoculation of insect eggs, the physiological variations in larval development, their successful attachment to the rootlets and the establishment of the galls cannot be avoided. Low values in nodosity or larvae numbers may result from less successful inoculations and not reflect the level of resistance of the host plant. To minimize this kind of effect, the maximal numbers of nodosities and larvae per individual were considered as the most reliable indicators of the quantitative resistance phenotype in *in planta* and *in vitro* experiments, respectively (Zhang et al. 2009). The negative control genotype, 'Börner', exhibited the expected results with a maximum number of 2 nodosities *in planta* and 9 larvae *in vitro*, respectively. Phenotypic responses of both parental genotypes were in agreement with what was expected, as VRH8771 resulted resistant and CS susceptible (Fig. 1). Over the population, the number of nodosities ranged from 0 to 37 (Fig. 1a) and the number of larvae ranged from 0 to 39 (Fig. 1b) in *in planta* and *in vitro* experiments, respectively. In both experiments, there were extensive variations of the phylloxera responses, with values of broad-sense heritabilities being 0.48 and 0.74 for nodosity and larvae numbers, respectively.

Phenotypic evaluation of resistance to *X. index*

In the first two experiments (2010-2011 and 2011-2012), resistance to *X. index* was screened from 35 BC1 individuals, using roots characteristics (RD and RW), the nematode reproduction factor (RF) and the gall index (GI). The data obtained from these two experiments were analyzed together since no year effect was identified (Fig. S4 and Table S2). A principal component analysis (PCA) using those 35 individuals and the RD, RW, RF and GI criteria showed that the two first axes explained 94.25 % of total variation (Fig.

2). Based on both the RF and GI criteria, resistant (R) and susceptible (S) groups were clearly identified. These two groups were separated along the first dimension of the PCA which is represented by these nematode development criteria. PCA demonstrated that susceptibility criteria vary independently from root characteristics. Statistical analyzes confirmed this result with significant differences in GI and RF values between resistant and susceptible BC1 individuals (Wilcoxon signed-rank test – *p-values* of 0.00015 and 0.00013 for RF and GI, respectively). On the other hand, no statistically significant differences were revealed between resistant and susceptible BC1 individuals for the criteria (RD and RW) related to the roots characteristics (Wilcoxon signed-rank test – *p-values* of 0.335 and 0.535 for RD and GW, respectively).

Spearman correlations were calculated among RD, RW, RF and GI criteria. There was no significant correlation between RD or RW on the one hand and RF or GI on the other hand, which confirmed the independence of the root and nematode developmental characteristics. The nematode reproduction factor (RF) was highly correlated with the gall index (GI) ($r = 0.59$, *p-value* = 0.00019) (Table S3). Such a high significant correlation made it possible to rely on the gall index rating in order to score the response of the 60 total BC1 individuals to *X. index* within the independent experiments. An individual is classified as resistant when its replicates have a mean GI value lower than 1 and as susceptible when its replicates have a mean GI value equal or above 1. Thus, on all five experiments, there were 50 and 10 BC1 individuals characterized as resistant and susceptible, respectively. This segregation ratio 50R:10S fits the ratio of 7:1 ($\chi^2_{0.05} = 0.5978$) expected when three dominant and independent genes control the resistance.

Genetic determinism and mapping of the response to *D. vitifoliae*

QTLs were detected for all traits scored and for their associated BLUP values in *in planta* and *in vitro* experiments. On the high-density genetic map of VRH8771 (Table 2), seven total QTLs were identified for both the maximum numbers of nodosities and the maximum numbers of larvae. These QTLs were localized on three LGs i.e. LG 3, LG 7 and LG 10. LOD scores and the explained phenotypic variances had higher values under *in vitro* conditions. This might be due to the lower environmental variations occurring under *in vitro* than under greenhouse conditions. On LG 7, several QTLs were found whatever the conditions and criteria within a confidence interval of 60 cM. In this group, the two QTLs detected for the criteria 'maximum number of nodosities' and 'BLUP nodosities' in *in planta* experiment explained each approximately 20 % of the variability while the two QTLs detected in *in vitro* conditions for the criteria 'number of larvae' and 'BLUP larvae' explained variances of 87 and 70 %, respectively. Two QTLs were identified on LG 10 in *in planta* and *in vitro* experiments, respectively with the maximal number of nodosities and the BLUP values associated to the maximal number of larvae. Finally, analysis of BLUP values associated to the maximal number of nodosities in *in planta* conditions, showed a QTL on LG 3.

Genetic determinism and mapping of the response to *X. index*

The 7R:1S segregation ratio observed in our tests might suggest the presence of three dominant and independent resistance (R) factors controlling the response to *X. index*. With this hypothesis in mind for the detection of markers linked to resistance, we first used a method derived from the bulked-segregant analysis (BSA) (Michelmore et al. 1991). As a double pseudo-testcross strategy was developed in the construction of the genetic maps, access to information on the different allelic forms was available only for SSR markers and, in a first step, the BSA-type analysis was performed using only these markers. From the 2 (ab), 3 (abc) or 4 (abcd) putative allelic forms of SSRs, we retained the markers for which an allele was detected in the resistant parent (VRH8771) and in a part of the resistant BC1 individuals but was lacking in all the susceptible BC1 individuals and the susceptible parent (CS). Markers meeting this requirement were located on the three linkage groups LG 9, LG 10 and LG 18 (Table 3), which is in line with the hypothesis of three dominant and independent R factors.

In order to use all the available genetic information related to the SSR and SNP markers, a proportion test was performed. This test aimed at comparing the proportion of resistant and susceptible individuals at each marker. Markers for which a statistically significant difference between resistant and susceptible proportions have been obtained, have been considered as potentially involved in the *X. index* resistance response. Such groups of markers with significant differences in their R/S proportions were found on LG 9, LG 10 and LG 18, corroborating the results obtained by BSA-type analyses from SSR markers (Table S4).

We finally performed a QTL analysis with a binary mapping model using the R/S phenotypic information from 60 BC1 individuals and the VRH8771 high-density genetic map. A first QTL explaining 22.73 % of the phenotypic variance was detected on LG 9 with a LOD score of 3.66 (Table 4 – Fig. S5). VVBX-A-06, the marker identified at the QTL peak on this group, corresponds to the marker identified previously through both the BSA-type analysis (Table 4) and the proportion test (Table S4). A second QTL explaining approximately the same proportion of phenotypic variance (21.19 %) was detected on LG 10 with an equivalent LOD score (3.59) (Fig. S5). The marker SC8-03 positioned at this QTL was the same as the marker detected by the proportion test on this linkage group but it was located distantly from the three SSR markers identified in the BSA-type analysis (MRBX-27, VVIN78-SEC and VVIN85). The third QTL with the highest LOD score (13.43% of the total phenotypic variance) was located on LG 18, i.e. in the same third chromosomal location as in the BSA-type analysis and the proportion test. Marker UDV-108 identified at this QTL was also the marker obtained from both other analyses. Nevertheless, this later QTL had a LOD score of ~2.5 that did not reach the significant threshold value of 3.12 (Table 4). In descending order, other loci were detected on LG 7, LG 19 and LG 11 but with much lower values of LOD scores (2.0 to 1.5) (Fig. 3).

Discussion

Genetic mapping of a *V. vinifera* x *M. rotundifolia* cross

Vitis x Muscadinia crosses have a very low fertility due to the differences in chromosome number that generate an incomplete homology between the genomes at meiosis (Riaz et al. 2012). Despite this limiting factor, a few F1 hybrids were obtained and have then been successfully backcrossed with *Vitis* species. Although reduced the size of the progeny, QTL analyses are reliable at the sight of the reproducibility of results obtained through different experimental conditions (*in planta* and *in vitro*) and different statistical methods (*in silico* BSA and QTL detection). This paper first reports the construction of high-density grapevine genetic maps for the BC1 progeny involving the *V. vinifera* x *M. rotundifolia* accession 'VRH8771' and the *V. vinifera* cv. 'Cabernet-Sauvignon' using both SSR and SNP types of markers. The framework maps established using SSR markers from other grapevine crosses were completed with SNP markers generated by GBS (Adam-Blondon et al. 2004; Doligez et al. 2006; Troggio et al. 2007; Wang et al. 2012; Barba et al. 2014). Among the whole genome and next generation sequencing techniques, GBS offers an inexpensive and robust solution for simultaneous SNP discovery and genotyping (Elshire et al. 2011; Poland and Rife 2012). In grapevine, GBS has already been used in the construction of high-resolution maps for the detection of QTLs linked to powdery and downy mildew resistance (Barba et al. 2014, Teh et al. 2017, Sapkota et al. 2019). In the present map, the number of SNP markers was lower in the paternal parent, CS, than in the maternal parent, VRH8771. This difference is linked to the use of the *V. vinifera* cv 'Pinot Noir' (PN40024) reference genome in our analyses (The French–Italian Public Consortium for Grapevine Genome Characterization 2007; Adam-Blondon et al. 2011). Actually, this genome is genetically closer to CS than to the *Vitis x Muscadinia* hybrid VRH8771, which explains a higher number of polymorphic markers in the maternal accession (Delame et al. 2019). As already shown in other studies, the comparison of the marker order with their physical position in VRH8771 map confirmed the high level of macrosynteny between the *V. vinifera* and *M. rotundifolia* genomes (Riaz et al. 2012; Blanc et al. 2012; Delame et al. 2019). The average distance between adjacent markers in all linkage groups in the maternal map was 0.98 cM, which is the same range of density reported by Teh et al. (2017) and Saptoka et al. (2019). The saturation level of the maternal genetic map is close to the map established by Delame et al (2019) for a muscadine derived progeny.

Identification of original QTLs for nodosity formation and larval development in response to root infection by *D. vitifoliae*

The response of the BC1 plants to phylloxera has been explored through both *in vitro* and *in planta* experiments. As expected in each experiment a wide variability of the response has been observed among replicates of a same plant individual due in particular to the vigour of the insect population. To minimize this effect, we considered the maximal values of the criteria studied as the most reliable indicators of the quantitative resistance phenotype (Zhang et al. 2009). Moreover, a linear mixed model was used to access the block effect in relation to the experimental design in randomized blocks. Thus, BLUP values were estimated and accounted for environmental effects (Piepho et al. 2008). This method has been used to optimize the statistical power for the detection of significant QTLs (Segura et al. 2009; Sadok et al. 2013). A QTL was detected on LG 7 for nodosity and larvae numbers with an explained phenotypic variance that reached 87 % in the *in vitro* experiment. The physiological status of the roots (which insects fed from) in each experiment, i.e. either connected to their plant (*in planta*) or completely detached from it

(*in vitro*), was quite different. Despite this, the strongest QTLs in both experimental conditions were located on the same chromosome (LG 7). Such results are in agreement with those obtained by Bouquet (1983) who reported also a good correlation of the data between the experiments conducted *in vitro* and in the greenhouse and/or the vineyard. Moreover, this author recorded a low ratio of phylloxera resistant plants in such *V. vinifera* x *M. rotundifolia* BC1 progenies. He suggested that this low ratio can be explained if the R factor controlling this trait is carried by a *Muscadinia* chromosome which has a low probability of pairing with its homologous *V. vinifera* chromosome in the F1 hybrids. By locating our major QTL in LG 7, the *Vitis* chromosome which had been split into the LG 7 (upper arm) and LG 20 (lower arm) in *Muscadinia*, our current data are in line with this previous hypothesis of a location of this R factor in the lower arm of chromosome 7.

Nevertheless, two other QTLs were also identified on LG 3 and LG 10 in our results, suggesting that other loci may influence or modulate nodosity formation and larval development.

The putative involvement of several QTLs in response to phylloxera has already been reported in the literature. Davidis and Olmo (1964) first hypothesized that the resistance from a *M. rotundifolia* accession was controlled by more than one locus. Then Bouquet (1983) suggested that grape phylloxera resistance in *M. rotundifolia* might be mediated by a semi-dominant locus regulated by three genetic modifiers. Similarly, in interspecific crosses involving different American *Vitis* species (*V. berlandieri*, *V. cinerea* and *V. rubra*) and *M. rotundifolia*, Boubals (1966) concluded that the resistance appears to be controlled by multiple loci. Four root grape phylloxera resistance QTLs have already been identified in *Vitis* spp. In *V. cinerea*, two QTLs have been mapped in two different accessions: *RDV1* located on LG 13 in cv. 'Arnold' (Zhang et al. 2009) and *RDV2* located on LG 14 in cv. 'C2-50' (Smith et al. 2018). Two other QTLs have been detected on LG 5 and LG 10 in an F1 cross containing at least six *Vitis* species in the ancestry (Clark et al. 2018). In our study, three new QTLs were identified from *M. rotundifolia* background. Interestingly, these seven QTLs have been mapped on six different LGs.

Resistance response to the dagger nematode *X. index* maps on three different LGs

The individuals were tested for their response to the dagger nematode *X. index* among successive experiments and no significant differences were found between the years of experiment, which underlines the reproducibility of the experimental protocol and the reliability of phenotyping.

Our evaluation of *X. index* resistance used a nematode reproduction factor and a root gall index assessed 10 to 12 months after inoculation. The root weight and a visual index of root development were also considered in order to study their relationships with the nematode development criteria previously mentioned. Gall index was shown to be highly correlated with the nematode reproduction factor but not with root characteristics. Such gall index rating method has been previously used in a genetic study for resistance to *X. index* in *V. arizonica*. It allowed the characterization of the early resistance conferred by the locus *XiR1* 4 to 8 weeks after inoculation in a fast greenhouse-based screening system (Xu et al. 2008; Hwang et al. 2010). Our current data demonstrate that gal index is also a reliable criterion for rating

resistance to *X. index* in longer tests (10-12 months) in order to identify R factors that should confer a more durable effect.

In our study, the distribution of the resistance levels to *X. index* was evaluated from 60 BC1 individuals. The approximate 7R:1S segregation ratio obtained suggested that three dominant and independent resistance factors might be involved in the response to the nematode. To cope with this hypothesis, a method derived from the bulked-segregant analysis (Michelmore et al. 1991) and performed on SSRs revealed markers on LG 9, LG 10 and LG 18. A proportion test using both SSRs and SNPs identified markers in the same three LGs and therefore confirmed previous data.

The QTL analysis performed on the same individuals supported these findings since two QTLs were identified on LG 9 and LG 10. QTL location on LG10 was not clearly estimated on this chromosome since the LOD score distribution curve showed 3 significant peaks, A third QTL was detected on LG 18 and, even though it was not significant, its LOD score value was higher than those observed on other linkage groups. These results provide the first location of resistance factors against the vector nematode *X. index* in muscadine. On the basis of segregation patterns observed in the progenies from *Vitis* crosses involving 13 species, one-gene and two-gene modes of tolerance to *X. index* were previously reported (Meredith et al. 1982), and Xu et al. (2008) identified a single resistance locus, *XiR1*, on the LG 19 of *V. arizonica* using a map from 185 plants. Interestingly, some hybrids, in particular the BC1 rootstock 'Nemadex Alain Bouquet', derived from the *M. rotundifolia* accession 'NC184-4' presently studied, were shown to decrease both *X. index* numbers and GFLV infection under greenhouse and field conditions (Bouquet et al. 2000; Ollat et al. 2011; Claverie et al. 2016; Nguyen 2018). Consequently, we hypothesize that resistance to *X. index* conferred by this muscadine accession will be associated with a delay in GFLV transmission by the nematode to the grapevine.

QTLs identified for response to *D. vitifoliae* and *X. index* map to chromosomal regions enriched in resistance gene analogs

While *V. vinifera* is highly sensitive to the most critical pests and diseases for grapevine, wild *Vitis* and *Muscadinia* materials are characterized by various levels of resistance. Among all, powdery and downy mildew have been studied in details, and numerous loci controlling partial or total resistance to both diseases have been identified (Meridinoglu et al. 2018). Interestingly, some of the chromosomal regions identified in our study as carrying R factors to *D. vitifoliae* and *X. index* already encompass QTLs identified in response to other major grapevine diseases. LG 18 contains many loci detected for resistance to downy mildew *Plasmopara viticola* and powdery mildew *Erysiphe necator* in muscadine (Wiedemann-Meridinoglu et al. 2006; Riaz et al. 2011) or *Vitis* spp. (Welter et al. 2007; Bellin et al. 2009; Zyprian et al. 2016; Sapkota et al. 2019). Thus, among the markers flanking the *Rpv3* (resistance to *P. viticola*) locus identified from 'Regent' genotype by van Heerden and collaborators (2014) on LG 18, is the UDV-108 marker which has been identified in our study with the maximum LOD score in one of the three QTLs highlighted in response to *X. index*. Additionally, in *Vitis* spp., the loci *rpv7* (resistance to downy mildew) and *rpv5* and *Ren6* (downy and powdery mildew) have been reported on LG 7 and LG9

respectively (Bellin et al. 2009; Marguerit et al. 2009; Pap et al. 2016). These three chromosomes have been previously demonstrated to be enriched in NBS-LRR genes (Di Gaspero et al. 2007). The current development of sequencing technologies should contribute to the comparison of *Vitis* genomes and to gather more precise knowledge about these resistance genes in order to use them in breeding new cultivars with multi-pest resistances.

Conclusions

With the objective of grapevine rootstock breeding, our data open the way to the use of muscadine as a source for the obligate phylloxera R feature and as an R source to *X. index* in order to delay GFLV transmission to grapevine by this nematode. In this work, we could successfully identify root resistances against phylloxera and *X. index* and map the first cognate QTLs from the *M. rotundifolia* source 'NC184-4'. Thus QTLs were identified on LG 7 completed by LG 3 and LG 10 in response to phylloxera and on LG 9, LG 10 and LG 18 in response to *X. index*. These newly identified trait-linked loci should now be tested in alternative crosses with *M. rotundifolia*. In the future, validated markers at those loci may allow their use in a marker-assisted breeding approach for producing new rootstocks with durable resistances.

Methods

Plant material

We used the backcross 1 (BC1) segregating population of 135 individuals (Table S1) from the cross between the hybrid VRH8771 [(‘Cabernet-Sauvignon’ x ‘Alicante Bouschet’) x *M. rotundifolia*] and *V. vinifera* cv. ‘Cabernet-Sauvignon’ (VRH8771 x CS). This cross has been initiated by Alain Bouquet at INRA Montpellier since 2005 and continued at INRA UMR EGFV (Bordeaux, France) after 2008. VRH8771, the female parent, is resistant to *D. vitifoliae* and *X. index* whereas *V. vinifera* ‘Cabernet-Sauvignon’ (CS), the male parent, is susceptible (Bouquet et al. 2000). Parental genotypes and BC1 individuals are conserved at the INRA germplasm repository at INRA UMR EGFV (Bordeaux, France).

Insect and nematode material

For *D. vitifoliae*, all experiments were conducted with the isofemale clone ‘Pcf7’. The original population was sampled in 2010 at Pineuilh (France) on *V. vinifera* cv. ‘Cabernet franc’ scions grafted on SO4 rootstock (*V. berlandieri* x *V. riparia*) and maintained at INRA UMR SAVE (Bordeaux, France) on leaves of the American variety ‘Harmony’, a complex hybrid between Dog-Ridge (*V. champinii*) and ‘1613C’ (*V. labrusca* x *V. riparia* x *V. vinifera*) and on root pieces of *V. vinifera* cv. ‘Cabernet Sauvignon’.

For *X. index*, all experiments were conducted with the isofemale line ‘Frejus’. The original population has been sampled in a GFLV-infected grapevine field in Frejus (Provence, France). Using the population grown on grapevine in the greenhouse, the line had been created from a single female inoculated on a fig plant previously grown from *in vitro*.

Experimental designs and phenotyping

Phylloxera assays

An *in planta* assay was performed with 89 BC1 individuals. Two control genotypes were also tested: *V. vinifera* cv. 'Pinot Noir' and the rootstock 'Börner' (*V. riparia* x *V. cinerea*) as susceptible and resistant genotypes respectively. This experiment was organized according to a randomized complete block design with three experimental blocks, each block being an independent randomization of one replicate of each BC1 individual. Plants were grown in individual pots of 1L in a soil substrate composed of at least 50 % clay with pebbles at the bottom to improve drainage. The soil was sterilized by autoclaving in order to prevent cross contamination with other phylloxera population sources. Each pot was covered with an insect-proof transparent plastic bell and an automatic watering system was adapted (Fig. S6 A). One hundred phylloxera eggs of 'Pcf7' clone, previously grown on root pieces of *V. vinifera* cv. 'Cabernet Sauvignon' for two generations, were deposited on a moistened and sterilized filter paper near the root (~3 cm depth) of each pot. Three months after inoculation, the plants were uprooted and the nodosity number was counted (Fig. S6 B). The maximal number of nodosities scored among the three replicates of each BC1 individual was considered as the most reliable indicator of the quantitative resistance phenotype.

An *in vitro* assay was also performed according to Pouget (1975) to assess the larval development of phylloxera on 37 BC1 individuals. Five woody root pieces per individual, 6 to 7 cm long, were arranged in small bundles (a contact between all the roots is required) on a disk of dampened blotting paper in a Petri dish sealed with parafilm. Three replicates were realized for each individual. A fungicide treatment (Ridomil Gold, Syngenta® - 2.3g/L) was carried out on the roots to prevent *Botrytis cinerea* infection. In each Petri dish, 50 phylloxera eggs were deposited on the roots pieces. Then Petri dishes were incubated at 25°C in the dark. The number of larvae that have developed on the five root pieces was counted one month after inoculation. The maximal number of larvae scored among the three replicates was considered as the best indicator of the quantitative resistance phenotype.

1. *index assays*

A total of 60 BC1 individuals were evaluated during five successive experiments conducted between 2010 and 2017 (Table S4). Three reference genotypes were also tested: the susceptible *V. rupestris* cv. 'du Lot', the resistant BC1 rootstock 'Nemadex Alain Bouquet' from the cross 'VRH8773 (*V. vinifera* x *M. rotundifolia*) x 140 Ru (*V. berlandieri* x *V. rupestris*)' and the susceptible to intermediate genotype 'VRH8624'. VRH8773 is a brother clone of VRH8771, and VRH8624 in an F1 hybrid (*V. vinifera* x *M. rotundifolia*) whose muscadine parent is the accession 'Trayshed' (Esmenjaud et al. 2010).

For each evaluation, homogenous hardwood cuttings of the individuals were rooted annually in alveolated plates in the nursery at INRA UMR EGFV (Bordeaux, France) in February. In May, plants were

delivered to INRA UMR ISA (Sophia-Antipolis, France), planted individually into 2-L pots with six replicates per individual and grown in a greenhouse. At the end of June, each pot was inoculated with a fixed number of nematodes that ranged from 300 to 900 depending on the year of experiment (Table S5). The plants were grown for ten to twelve months, which is the time that allows approximately three to four nematode developmental cycles over two successive calendar years.

At harvest, the aerial part of the plant was cut at the collar level and removed and each pot was hermitically placed into a plastic bag and stored in a cold chamber at 6°C. This stopped plant and nematode development simultaneously in all individual replicates until plant and nematode ratings. Ratings were done sequentially, i.e. replicate after replicate. Total soil of each 2-L pot was recovered in a 10-L bucket, over which plant roots were washed individually with caution under tap water. The entire root system of each plant was rated for its root development (RD) based on a 0-5 scale and its fresh root weight (RW) was also measured. Root galling was rated for each plant using a 0-5 gall index (GI) scale derived from studies for resistance to the root-knot nematode *Meloidogyne* spp. : 0 = no gall; 1 = 1-10 %; 2 = 11-30 %; 3 = 31-70 %; 4 = 71-90 %; 5 > 90 % of root system galled (Barker 1985). Nematodes of each plant were extracted from the total soil suspended in the bucket using an adapted Oostenbrink method (De Goede and Verschoor 2000). In the two first experiments (2010-2011 and 2011-2012), final nematode numbers were counted under a binocular microscope. Then the ratio between nematode final and initial numbers was calculated to evaluate the mean nematode reproduction factor (RF) for each BC1 individual and the parental and reference genotypes. Individuals were classified as resistant (R) when their RF value was lower than 1 and susceptible (S) when their RF value was equal or above 1. As the two first experiments showed that RF ratings were significantly correlated with GI ratings (see Results section), no nematode extraction was performed for the three last experiments (2012-2013, 2015-2016 and 2016-2017) and individuals were classified directly as R or S from their visual symptoms.

Treatment of phenotypic data sets

Phylloxera assays

The number of nodosities and the number of larvae were explored using the following generalized mixed model:

$$P_{ij} = \mu + \text{individual} + \text{block}_j + \varepsilon_{ij}$$

where P_{ij} is the observed phenotype, μ is the overall mean of the phenotypic data, 'individual' corresponds to the genetic differences among the BC1 individuals, 'block' accounts for the differences in microenvironmental conditions among the three blocks, and ε_{ij} is the residual term (R_{LME4} package). The factor 'individual' was treated as a random factor, whereas the factor 'block' was treated as a fixed factor. The 'Best linear unbiased predictions' (BLUP) of random effects were extracted from the selected generalized mixed model. The BLUP values were noted by the name of the trait preceded by the word BLUP.

For each quantitative phenotypic trait, broad sense heritability was estimated using the formula $h^2 = \sigma_g^2 / [\sigma_g^2 + \sigma_e^2/n]$, where σ_g^2 is the genetic variance, σ_e^2 is the environmental variance and n is the number of plants per accession.

Nematode assays

All principal component analyses (PCA) and parametric and non-parametric statistical tests were performed using R version 3.4.0. Statistical significance was set at $P < 0.05$.

Mapping of the resistance traits

Simple Sequence Repeats (SSR) markers

DNA was extracted from 50 to 100 mg of young leaves grown in a greenhouse from 90 BC1 individuals (Table S1). Leaves were ground in 5 mL of a first buffer extraction containing sodium metabisulfite [sodium metabisulfite 20 mM, Tris-HCl pH8 0.2 M, EDTA pH8 70 mM, NaCl 2 M]. 500 μ L of this homogenate was incubated with 450 μ L of a second buffer with CTAB [CTAB 2 %, NaCl 1.4 M, EDTA pH8 20 mM, Tris-HCl pH8 0.1 M] during one hour at 65°C. Then the solution was centrifuged for 30 minutes at 13,000 rpm at 4°C. 500 μ L of supernatant was sampled and cleaned with the same volume of a solvent chloroforme:octanol (24:1). After 20 min of spin at 13,000 rpm at 4°C, we recovered 300 μ L of the aqueous phase in 450 μ L of a solution of isopropanol: ammonium acetate (2:1). We left one hour at 4°C and then we centrifuged at 13,000 rpm for 20 min at 4°C. We discarded the solution and washed the pellet with 70% ethanol v/v (two times). After centrifugation (13,000 rpm for 20 min at 4°C), we discarded the ethanol, and precipitated the pellet in 100 to 200 μ L of TE 0.1 X [Tris HCl 10mM, EDTA 1mM]. DNAs were quantified by Nanodrop™.

A total of 217 primers pairs were used: 61 VMC (Vitis Microsatellite Consortium, managed through AGROGENE, Moissy Cramayel, France), 7 VVMD (Bowers et al. 1999), 1 VVS (Thomas and Scott 1993), 6 SC08 (Cipriani et al. 2011), 1 VrZAG (Sefc et al. 1999), 7 VVC (Thomas and Scott 1993), 59 VVI (Merdinoglu et al. 2005), 11 MRBX (Zah-Bi et al. 2010), 21 UDV (Di Gaspero et al. 2005) and marker Gf13-9 (Zhang et al. 2009). Two new series of SSRs, VVBX and VVBX-A, were designed from the genome 12X of *V. vinifera* cv. 'Pinot Noir' (PN40024) using Primer3 software (Untergasser et al. 2012). Primers characteristics of VVBX and VVBX-A markers are reported in Table S6.

PCRs were performed by a single reaction with M13-tailed forward primer (Oetting et al. 1995) conjugated with four different dyes (6FAM™, VIC®, NED™ and PET®) in 15 μ L reaction volume containing : 5 ng of DNA template, 1.5 μ L of 10xPCR reaction buffer, 2 mM of MgCl₂, 0.2 mM of each dNTP, 0.05 μ M of M13 tailed SSR forward primer, 0.2 μ M of reverse primer, 0.2 μ M of dye conjugated with M13 primer, and 0.2 U of JumpStart™ Taq DNA Polymerase (Sigma-Aldrich). Amplification conditions were as follows: 5 min initial denaturation step at 95°C followed by 2 cycles (30 s denaturation at 95°C, 1.5 min annealing at 60°C or 56°C and 1 min extension at 72°C) followed by 35 cycles (30 s denaturation at 95°C, 30 s annealing at 60°C or 56°C and 1 min extension at 72°C) then followed by 10 min final extension at 72°C.

Visualization was performed by a 3730 DNA analyzer (Applied Biosystem™). Eight to sixteen PCR products were pooled, according to the size of SSRs and the dyes, and analyzed in a single run. Electropherograms were analyzed using free software STRand. Any ambiguous genotypes were re-run, re-amplified or left as unknown.

Single Nucleotide Polymorphisms (SNP) markers

Two foliar disks (1.5 cm of diameter) of 128 BC1 individuals grown in a greenhouse were sampled (Table S1). DNA extractions were realized on dried leaf tissues using the same protocol as described by Cormier et al. (2019). Genotyping-by-sequencing (GBS) was performed as described by Elshire et al. (2011) (Keygene N.V. owns patents and patent applications protecting its Sequence Based Genotyping technologies) integrating two 96-well plates across 96 barcodes for library preparation. The genomic library was prepared using ApeKI restriction enzyme. Paired-end sequencing of 150 bp reads was performed on an Illumina HiSeq3000 system (at the GeT-PlaGe platform in Toulouse, France).

Raw reads were checked with FastQC (Andrews 2016), demultiplexed with a custom script (<https://github.com/timflutre>) and cleaned with CutAdapt (Martin 2011). Cleaned reads were then mapped to the *V. vinifera* cv 'Pinot Noir' (PN40024) genome assemblies for SNP calling. Alignment on this genome was performed using Burrows-Wheeler Aligner maximal exact match (BWA-MEM) with default parameters (Li and Durbin 2009), SAMtools and Picard (<http://broadinstitute.github.io/picard/>). SNP calling was performed with GATK using the hardfilter parameters (McKenna et al. 2010; DePristo et al. 2011; Van der Auwera et al. 2013).

VCFtools was used to remove SNPs with a quality score < 200 and with depth values < 10 (Danecek et al. 2011). Quality-filtered SNPs were analyzed with the *major_minor* and *get_pseudo_test_cross* scripts from HetMappS pipeline to identify pseudo-testcross markers (Hyma et al. 2015). In the variant call format (VCF) output file only sites with less than 10 % missing data were retained. Individuals with more than 50% missing data and those with genotype frequencies different from expected 1:1 marker segregation were discarded. Two sets of markers were obtained corresponding to the two parental genotypes.

Map construction

Individual maps were constructed for each parental genotype following a double pseudo-testcross strategy (Grattapaglia and Sederoff 1994). Marker segregation was analysed with regard to goodness-of-fit to the expected Mendelian ratio using the Chi-square test ($P < 0.05$). Marker types of *Im x ll* and *nn x np* were retained for construction of maternal and paternal maps, respectively. Genetic maps and marker order were determined using the maximum likelihood (ML) algorithm with Haldane function and default parameters of JoinMap®4.1 software (Van Ooijen 2006). Linkage groups (LGs) were constructed with a minimum threshold logarithm of odds (LOD) score of 6.0. LGs were grouped and numbered based on their corresponding physical chromosome numbers (The French–Italian Public Consortium for Grapevine Genome Characterization 2007).

Methods for resistance mapping

Phylloxera assays

The maximum number of nodosities (*in planta* assay), the maximum number of larvae (*in vitro* assay) and the BLUP values associated were used as the quantitative scores of susceptibility/resistance response.

Detection of quantitative trait loci (QTLs) was performed with the one-dimension scan function, *scanone*, of R/qtl software using a normal model but also a non-parametric analysis and the expectation-maximization (EM) algorithm method depending on the normality of the data (Dempster et al. 1977; Broman et al. 2003). Multipoint genotype probabilities were calculated beforehand using *calc.genoprob* with step = 1 and default parameters. Logarithm of odd score (LOD) significance threshold was estimated with 1000 permutations and for a significant level α of 0.05. An interval estimate of the location of each QTL was calculated using the 1.5-LOD intervals method of Rqtl (Broman and Sen 2009). The percentage of the phenotypic variation explained by a QTL corresponds to the regression value R^2 taken at its peak LOD score.

Nematode assays

Based on SSR markers, an *in silico* method derived from the Bulk Segregant Analysis (BSA) was performed (Michelmore et al. 1991). In the VRH8771 x CS progeny, polymorphic markers were distributed into either 2 (ab), 3 (abc) or 4 (abcd) allelic forms. From each of the markers screened, markers retained were those for which an allele was detected in the resistant parent (VRH8771) and in part of the resistant BC1 individuals but in none of all susceptible BC1 individuals.

The resistant and susceptible phenotypes were converted to values equal to 0 and 1, respectively. The one-dimension scan function, *scanone*, of R/qtl software was performed with the argument *model = 'binary'* (Broman and Sen 2009). LOD significance threshold, the QTL interval and the R^2 value were obtained with the procedure described in the previous paragraph. A number of 47 phenotyped individuals were used for the BSA analysis with SSR markers whereas the 60 total individuals phenotyped were used for QTL analysis with SSR and SNP markers.

Abbreviations

BLUP: Best linear unbiased predictions

BSA: Bulk segregant analysis

EM: expectation-maximization

GBS: Genotyping-by-sequencing

LG: Linkage group

LOD: Logarithm of odds ratio

ML: maximum likelihood

NGS: Next-generation sequencing

PCA: principal component analysis

QTL: Quantitative trait locus

SNP: Single nucleotide polymorphism

VCF: variant call format

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and material

The data sets during the current study are available from the corresponding author.

Competing interests

The authors declare there are no conflicts of interest.

Funding

This study has received funding from the European Union's Seventh framework program for research, technological development and demonstration under grant agreement n°311775 (Innovine, 2013-2016) which supported the SSR genotyping of the population, the establishment of the SSR map, the phylloxera assays and GLT initial contract. It was also partly funded by the French Ministry of Agriculture through the CASDAR contracts 'RemuneX' (2015-2018) (CTPS Project n°C-2014-09), for the nematode assays, the completion of the SSR map, the BSA analysis and GLT contract. 'PIVERT I and II' (2016-2019) (France AgriMer EDP 09 16 00 2775, EDP 09 17 00 3392, EDP 09 18 00 3764) contributed to the GBS for the population, the establishment of the SNP map, the QTL detection, GLT final contract and BR contract. A complementary funding has been provided by the contracts INRA-AgriObtentions IVD3 (2011-2015) and IVD4 (2016-2020) for phenotyping activities, SSR and SNP genotyping.

Authors' Contributions

NO, DE and DP designed the study. ML and J-PT performed the crosses and participated to the maintenance of the progeny. ML, J-PP, BJ and DP performed the phylloxera tests and ML, MD, BJ, DP, GLT and BR analyzed the data for phylloxera. DE, RV, UP and CVG performed the *X. index* tests and DE, GLT and BR analyzed the data for *X. index*. GLT and MD collected the samples for DNA extraction and performed microsatellite genotyping. GLT, LLC and BR constructed the genetic maps and performed the statistical analysis. GLT and BR contributed to the QTL analyses. P-FB aided in interpreting the results and contributed to the correction of the manuscript. DE, GLT and BR wrote the first version of the manuscript. DE, BR, LLC and NO generated the last version of the manuscript. All authors read and approved the manuscript

Acknowledgements

This manuscript is dedicated to Alain Bouquet who initiated numerous crosses involving muscadine in particular the cross used in this study. Without him, nothing would have been possible. The authors thank A. El Msayryb and A. Bakhkhouch for their help in nematological analyses, B. Douens, J-P Robert and C. Hévin for their contribution to plant material multiplication and maintenance, and N. Girollet for his wise advices and help in bioinformatic processing of data.

References

- Adam-Blondon AF, Martinez-Zapater JM, Kole C (2011) Genetics, Genomics, and Breeding of Grapes. Science Publishers and CRC Press Book
- Adam-Blondon A-F, Roux C, Claux D, et al (2004) Mapping 245 SSR markers on the *Vitis vinifera* genome: a tool for grape genetics. Theor Appl Genet 109:1017–1027
- Andret-Link P, Laporte C, Valat L, et al (2004) *Grapevine fanleaf virus*: still a major threat to the grapevine industry. J Plant Pathol 86:183-195
- Barba P, Cadle-Davidson L, Harriman J, et al (2014) Grapevine powdery mildew resistance and susceptibility loci identified on a high-resolution SNP map. Theor Appl Genet 127:73–84
- Barker KR (1985) Design of greenhouse and microplot experiments for evaluation of plant resistance to nematodes. In: Zuckerman BM, MAi WS, Harrison MB eds. *Plant Nematology Laboratory Manual*. Amherst: University of Massachussets Agriculture Experiment Station, 103-113
- Bellin D, Peressotti E, Merdinoglu D, et al (2009) Resistance to *Plasmopara viticola* in grapevine 'Bianca' is controlled by a major dominant gene causing localised necrosis at the infection site. Theor Appl Genet 120:163-176
- Blanc S, Wiedemann-Merdinoglu S, Dumas V, et al (2012) A reference genetic map of *Muscadinia rotundifolia* and identification of Ren5, a new major locus for resistance to grapevine powdery mildew. Theor Appl Genet 125:1663–1675

- Bordenave L, Tandonnet JP, Decroocq S, et al (2014) Wild vitis as a germplasm resource for rootstocks. In: Exploitation of autochthonous and more used vines varieties - Oenoviti International Network I, Geisenheim, Germany.
- Boubals D (1966) Hérité de la résistance au phylloxéra radicole chez la vigne. Ann Amélioration Plantes 16:327–347
- Bouquet A (1980) *Vitis x Muscadinia* hybridization: A new way in grape breeding for disease resistance in France Proc Third Intl Symp on Grape breeding pp. 42-61
- Bouquet A (1983) Etude de la résistance au phylloxéra radicole des hybrides *Vitis vinifera x Muscadinia rotundifolia*. Vitis 22:311-323
- Bouquet A, Danglot Y, Bongiovanni M, et al (2000) Breeding rootstocks resistance to *Grape fanleaf virus* spread, using *Vitis x Muscadinia* hybridization. Acta Hort 528: 517-526
- Bowers JE, Dangl GS, Meredith CP (1999) Development and characterization of additional microsatellite DNA markers for grape. Am J Enol Vitic 50:243–246
- Broman KW, Sen S (2009) A guide to QTL mapping with R/qtl. Springer, Dordrecht
- Broman KW, Wu H, Sen S, Churchill GA (2003) R/qtl: QTL mapping in experimental crosses. Bioinformatics 19:889–890
- Cipriani G, Di Gaspero G, Canaguier A, et al (2011) Molecular linkage maps: strategies, resources and achievements. In AF Adam-Blondon JM Martinez-Zapater C Kole (eds) Genet Genomics Breed Grapes pp. 111–136
- Clark MD, Teh SL, Burkness E, et al (2018) Quantitative trait loci identified for foliar phylloxera resistance in a hybrid grape population: Foliar phylloxera resistance. Aust J Grape Wine Res 24:292–300.
- Claverie M, Audeguin L, Barbeau G, et al (2016) Neadex AB : Bilan des réseaux d'expérimentation en France. Progrès Agricole et Viticole pp.17-29.
- Cormier F, Lawac F, Maledon E, et al (2019) A reference high-density genetic map of greater yam (*Dioscorea alata* L.). Theor Appl Genet <https://doi.org/10.1007/s00122-019-03311-6>
- Danecek P, Auton A, Abecasis G, et al (2011) The variant call format and VCFtools. Bioinformatics 27:2156–2158.
- Davidis UX, Olmo HP (1964) The *Vitis vinifera x V. rotundifolia* Hybrids as phylloxera resistant rootstocks. Vitis 4:129–143
- De Goede RGM, Verschoor B (2000) The nematode extraction efficiency of the Oostenbrink elutriator-cottonwool filter method with special reference to nematode body size and life strategy. Nematology

2:325–342.

Delame M, Prado E, Blanc S, et al (2019) Introgression reshapes recombination distribution in grapevine interspecific hybrids. *Theor Appl Genet* 132:1073-1087

Demangeat G, Esmenjaud D, Voisin R, et al (2005) Le court-noué de la vigne. *Phytoma - La défense des végétaux* 587:38–42

Dempster AP, Laird NM, Rubin DB (1977) Maximum Likelihood from Incomplete Data Via the *EM* Algorithm. *J R Stat Soc Ser B Methodol* 39:1–22.

DePristo MA, Banks E, Poplin R, et al (2011) A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nat Genet* 43:491–498.

Di Gaspero G, Cipriani G, Marrazzo MT, et al (2005) Isolation of (AC) n-microsatellites in *Vitis vinifera* L. and analysis of genetic background in grapevines under marker assisted selection. *Mol Breed* 15:11–20

Doligez A, Adam-Blondon AF, Cipriani G, et al (2006) An integrated SSR map of grapevine based on five mapping populations. *Theor Appl Genet* 113:369–382.

Duval H, Hoerter M, Polidori J, et al (2014) High-resolution mapping of the RMia gene for resistance to root-knot nematodes in peach. *Tree Genet Genomes* 10:297–306

Elshire RJ, Glaubitz JC, Sun Q, et al (2011) A Robust, Simple Genotyping-by-Sequencing (GBS) Approach for High Diversity Species. *PLoS ONE* 6(5): e19379

Esmenjaud D, Walter B, Valentin G, et al (1992) Vertical distribution and infectious potential of *Xiphinema index* (Thorne et Allen, 1950) (*Nematoda: Longidoridae*) in fields affected by grapevine fanleaf virus in vineyards in the Champagne region of France. *Agronomie* 12:395–399

Esmenjaud D. and Bouquet A. (2009) Selection and application of resistant germplasm for grapevine nematodes

management. In: Ciancio A. and Mukerji K.G. (eds.). *Integrated Management of Fruit Crops and Forest Nematodes*. Springer Science+Business Media B.V. pp. 195-214

Esmenjaud D, Decroocq S, Bouquet A, Ollat N (2010) Host Suitability of *Vitis* and *Vitis-Muscadinia* material to the nematode *Xiphinema index* over one to four years *Am J Enol Vitic* 61:96-101

Esmenjaud D, Bouquet A, Demangeat G, et al (2011) Nematode-resistant rootstocks as a major component of the management alternative for *Grapevine fanleaf virus* control in grape. *Acta Hort* 904:111-115

Ferris H, Zheng L, Walker MA (2012) Resistance of grape rootstocks to plant-parasitic nematodes. *J. Nematol.* 44:337-86

- Grattapaglia D, Sederoff R (1994) Genetic linkage maps of *Eucalyptus grandis* and *Eucalyptus urophylla* using a pseudo-testcross: mapping strategy and RAPD markers. *Genetics* 137:1121–1137
- Hewitt WB, Raski DJ, Goheen AC (1958) Nematode vector of soil-borne fanleaf virus of grapevines. *Phytopathology* 48:586–595
- Hwang C-F, Xu K, Hu R, et al (2010) Cloning and characterization of *XiR1*, a locus responsible for dagger nematode resistance in grape. *Theor Appl Genet* 121:789–799
- Hyma KE, Barba P, Wang M, et al (2015) Heterozygous Mapping Strategy (HetMappS) for High Resolution Genotyping-By-Sequencing Markers: A Case Study in Grapevine. *PLOS ONE* 10:e0134880
- Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25:1754–1760
- Marguerit E, Boury C, Manicki A, et al (2009) Genetic dissection of sex determinism, inflorescence morphology and downy mildew resistance in grapevine. *Theor Appl Genet* 118:1261–1278
- Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet Journal* 17:10-12
- McKenna A, Hanna M, Banks E, et al (2010) The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res* 20:1297–1303
- Merdinoglu D, Butterlin G, Bevilacqua L, et al (2005) Development and characterization of a large set of microsatellite markers in grapevine (*Vitis vinifera* L.) suitable for multiplex PCR. *Mol Breed* 15:349–366
- Merdinoglu D, Schneider C, Prado E, et al (2018) Breeding for durable resistance to downy and powdery mildew in grapevine. *OENO one* 52:203-209
- Meredith CP, Lider LA, Raski DJ, Ferrari NL (1982) Inheritance of tolerance to *Xiphinema index* in *Vitis* species. *Am J Enol Vitic* 33:154-158
- Michelmore RW, Paran I, Kesseli RV (1991) Identification of markers linked to disease-resistance genes by bulked segregant analysis: a rapid method to detect markers in specific genomic regions by using segregating populations. *Proc Natl Acad Sci* 88:9828–9832
- Nguyen V. C. 2018. Genetic diversity of the grapevine vector nematode *Xiphinema index* and application to optimize

the resistance strategy. Doctoral dissertation, Université Côte d'Azur. 179 pp.

Oetting WS, Lee HK, Flanders DJ, et al (1995) Linkage analysis with multiplexed short tandem repeat polymorphisms using infrared fluorescence and M13 tailed primers. *Genomics* 30:450–458

Ollat N, Claverie M, Esmenjaud D, et al (2011). Nemadex Alain Bouquet : un nouveau porte-greffe contrôlant la

recontamination des vignes par la maladie du court-noué. *Phytoma. La Défense des Végétaux* 649:29-33.

Ollat N, Peccoux A, Papura D, et al (2016). Rootstocks as a component of adaptation to environment. In: Gerós

H.V., Chaves M.M., Gil H.M. and Delrot S. (eds). *Grapevine in a Changing Environment: A Molecular and Ecophysiological Perspective*, Wiley Blackwell, Chichester, UK. pp.68-108

Pap D, Riaz S, Dry I B, et al (2016) Identification of two novel powdery mildew resistance loci, *Ren6* and *Ren7*,

from the wild Chinese grape species *Vitis piasezkii*. *BMC Plant Biol* 16:170

Patel G, Olmo HP (1955) Cytogenetics of *Vitis*: The hybrid *V. vinifera* x *V. rotundifolia*. *Am J Bot* 42:141-159

Piepho HP, Möhring J, Melchinger AE, Büchse A (2008) BLUP for phenotypic selection in plant breeding and variety testing. *Euphytica* 161:209–228

Poland JA, Rife TW (2012) Genotyping-by-Sequencing for Plant Breeding and Genetics. *The Plant Genome* 5:92-102

Pouget R (1975) Méthode de contamination de racines de vigne *in vitro* par le phylloxéra radicicole: application à la recherche de porte-greffes résistants. *Connaiss. Vigne Vin* 9: 165-176

Pouget R (1990) Histoire de la lutte contre le phylloxéra de la vigne en France. INRA/OIV, Paris. 157 pp.

Powell KS (2012) A holistic approach to future management of grapevine phylloxera. In: N J. Bostanian, C. Vincent, R. Isaacs *Arthropods Management in Vineyards: Pests, approaches, and future directions* pp. 219–251

Powell KS (2008) Grape Phylloxera : An Overview. In: S.N. Johnson and G.M. Murray (eds) *Root feeders: An ecosystem perspective*. CAB International pp. 96-114

Ramming DW (2010) Greenhouse screening of grape rootstock populations to determine inheritance of resistance to phylloxera *Am J Enol Vitic* 61:234-239

- Ray S, Satya P (2014) Next generation sequencing technologies for next generation plant breeding. *Front Plant Sci* 5:1-4
- Riaz S, Tenschler A.C, Ramming D.W, Walker M.A (2011) Using a limited mapping strategy to identify major QTLs for resistance to grapevine powdery mildew (*Erysiphe necator*) and their use in marker-assisted breeding. *Theor Appl Genet* 122:1059-1073
- Riaz S, Hu R, Walker MA (2012) A framework genetic map of *Muscadinia rotundifolia*. *Theor Appl Genet* 125:1195–1210
- Riaz S, Pap D, Uretsky J, Laucou V, Boursiquot J-M, Kocsis L, Andrew Walker M (2019) Genetic diversity and parentage analysis of grape rootstocks. *Theor Appl Genet* [https://doi: 10.1007/s00122-019-03320-5](https://doi.org/10.1007/s00122-019-03320-5)
- Roush TL, Granett J, Walker MA (2007) Inheritance of gall formation relative to phylloxera resistance levels in hybrid grapevines *Am J Enol Vitic* 58:234-241
- Sadok IB, Celton J-M, Essalouh L, et al (2013) QTL mapping of flowering and fruiting traits in olive. *PLoS ONE* 8:e62831
- Sapkota S, Chen L, Yang S, et al (2019) Construction of a high-density linkage map and QTL detection of downy mildew resistance in *Vitis aestivalis*-derived 'Norton'. *Theor Appl Genet* 132:137-147
- Saucet S, Van Ghelder C, Abad P, et al (2016) Resistance to root-knot nematodes *Meloidogyne* spp. in woody plants. *New Phytol* 211:41–56
- Sefc KM, Regner F, Turetschek E, et al (1999) Identification of microsatellite sequences in *Vitis riparia* and their applicability for genotyping of different *Vitis* species. *Genome* 42:367–373
- Segura V, Durel C-E, Costes E (2009) Dissecting apple tree architecture into genetic, ontogenetic and environmental effects: QTL mapping. *Tree Genet Genomes* 5:165–179
- Smith BP, Morales NB, Thomas MR, et al (2017) Grapevine rootstocks resistant to the root-knot nematode *Meloidogyne javanica*: Grapevine resistance to root-knot nematodes. *Aust J Grape Wine Res* 23:125–131
- Smith HM, Clarke CW, Smith BP, et al (2018) Genetic identification of SNP markers linked to a new grape phylloxera resistant locus in *Vitis cinerea* for marker-assisted selection. *BMC Plant Biol* 18:360
- Teh S-L, Fresnedo-Ramirez J, Clark M D, et al (2017) Genetic dissection of powdery mildew resistance in

interspecific half-sib grapevine families using SNP-based maps. *Mol Breed* 37:1

<https://doi.org/10.1007/s11032-016-0586-4>

The French–Italian Public Consortium for Grapevine Genome Characterization (2007) The grapevine genome sequence suggests ancestral hexaploidization in major angiosperm phyla. *Nature* 449:463–467

Thiéry D, Vogelweith F (2016). La colocation en vigne : interactions entre plante, les insectes ravageurs et leurs

parasitoïdes. *La Garance Voyageuse* 113:19-25

This P, Lacombe T, Thomas MR (2006) Historical origins and genetic diversity of wine grapes. *Trends Genet* 22:511–519

Thomas MR, Scott NS (1993) Microsatellite repeats in grapevine reveal DNA polymorphisms when analyzed as sequence-tagged sites (STSs). *Theor Appl Genet* 86:985–990

Töpfer R, Hausmann L, Harst M, et al (2011) New horizons for grapevine breeding. In: Flachowsky H., Hanke M.-V (eds) *Methods Temperate Fruit Breed* pp 79–100

Troggio M, Malacarne G, Coppola G, et al (2007) A dense single-nucleotide polymorphism-based genetic linkage map of grapevine (*Vitis vinifera* L.) anchoring Pinot Noir bacterial artificial chromosome contigs. *Genetics* 176:2637–2650

Untergasser A, Cutcutache I, Koressaar T, et al (2012) Primer3—new capabilities and interfaces. *Nucleic Acids Res* 40:e115

van der Auwera GA, Carneiro MO, Hartl C, et al (2013) From FastQ Data to High-Confidence Variant Calls: The Genome Analysis Toolkit Best Practices Pipeline: The Genome Analysis Toolkit Best Practices Pipeline. In: Bateman A, Pearson WR, Stein LD, et al. (eds) *Current Protocols in Bioinformatics*. John Wiley & Sons, pp 11.10.1-11.10.33

van Ghelder C, Lafargue B, Dirlewanger E, et al (2010) Characterization of the *RMja* gene for resistance to root-knot nematodes in almond: spectrum, location, and interest for *Prunus* breeding. *Tree Genet Genomes* 6:503–511

van Heerden CJ, Burger P, Vermeulen A, Prins R (2014) Detection of downy and powdery mildew resistance QTL

in a 'Regent' x 'RedGlobe' population. *Euphytica* 200:281-295

van Ooijen (2006) JoinMap4, software for the calculation of genetic linkage maps in experimental populations Wageningen, Kyazma BV

- Varshney RK, Nayak SN, May GD, Jackson SA (2009) Next-generation sequencing technologies and their implications for crop genetics and breeding. *Trends Biotechnol* 27:522–530
- Villate L, Fievet V, Hanse B, et al (2008) Spatial distribution of the dagger nematode *Xiphinema index* and its associated *Grapevine fanleaf virus* in French vineyard. *Phytopathology* 98:942–948
- Vlk D, Řepková J (2017) Application of next-generation sequencing in plant breeding. *Czech J Genet Plant Breed* 53:89–96
- Walker MA, Wolpert JA, Weber E. (1994) Viticultural characteristics of VR hybrid rootstocks in a vineyard site infected with *Grapevine fanleaf virus*. *Vitis* 33: 19-23.
- Wang N, Fang L, Xin H, et al (2012) Construction of a high-density genetic map for grape using next generation restriction-site associated DNA sequencing. *BMC Plant Biol* 12:148
- Welter LJ, Göktürk-Baydar N, Akkurt M, et al (2007) Genetic mapping and localization of quantitative trait loci affecting fungal disease resistance and leaf morphology in grapevine (*Vitis vinifera* L). *Mol Breed* 20:359-374
- Wiedemann-Merdinoglu S, Prado E, Coste P, et al (2006) Genetic analysis of resistance to downy mildew from *Muscadinia rotundifolia*. In: 9th International conference on Grape Genetics and Breeding, Udine - Italy
- Xu K, Riaz S, Roncoroni NC, et al (2008) Genetic and QTL analysis of resistance to *Xiphinema index* in a grapevine cross. *Theor Appl Genet* 116:305–311
- Zah-Bi I, Blanc S, Bras M, et al (2010) Development of resources for comparative physical mapping between *Muscadinia rotundifolia* and *Vitis vinifera*. In: Proceedings of the 6th International Workshop of Grapevine Downy and Powdery Mildew. GDPM 2010. Villenave d'Ornon, France
- Zhang J, Hausmann L, Eibach R, et al (2009) A framework map from grapevine V3125 (*Vitis vinifera* 'Schiava grossa' × 'Riesling') × rootstock cultivar 'Börner' (*Vitis riparia* × *Vitis cinerea*) to localize genetic determinants of phylloxera root resistance. *Theor Appl Genet* 119:1039–1051
- Zyprian E., Ochßner I., Schwander F., et al (2016) Quantitative trait loci affecting pathogen resistance and ripening of grapevines. *Mol Genet Genomics* 291:1573–1594

Tables

Table 1 Characteristics of the maternal (VRH8771) and the paternal (*V. vinifera* cv. 'Cabernet-Sauvignon', CS) genetic maps of the BC1 cross VRH8771 x CS.

Linkage Group	Map of female parent VRH8771				Map of male parent CS			
	Number of markers		Length (cM)	Average marker interval (cM)	Number of markers		Length (cM)	Average marker interval (cM)
	SNPs	SSRs			SNPs	SSRs		
1	68	9	90.36	1.17	46	7	123.13	2.32
2	84	8	107.66	1.17	41	6	106.75	2.27
3	140	8	117.91	0.80	21	9	96.94	3.23
4	106	5	127	1.14	24	6	69.84	2.33
5	176	7	173.84	0.95	67	7	115.5	1.56
6	154	6	112.04	0.70	28	7	90.03	2.57
7	47	7	87.7	1.62	36	10	121.33	2.64
8	115	9	123.22	0.99	50	7	121.89	2.14
9	120	7	119.21	0.94	25	7	73.25	2.29
10	107	9	91.8	0.79	45	8	92.07	1.74
11	120	4	121.07	0.98	42	6	92.56	1.93
12	126	5	125.13	0.96	21	7	81.32	2.90
13	108	7	212.89	1.85	42	7	98.58	2.01
14	172	8	160.33	0.89	48	8	135.63	2.42
15	17	5	57.74	2.62	14	7	107.16	5.10
16	105	6	90.35	0.81	36	9	92.91	2.06
17	158	6	154.09	0.94	63	10	96.87	1.33
18	207	11	151.66	0.70	49	11	165.52	2.76
19	141	8	126.69	0.85	40	9	101.18	2.06
TOTAL	2271	135	2350.69	0.98	738	148	1982.46	2.24

Table 2 Location and characteristics of the QTLs identified in *in planta* and *in vitro* experiments in response to *D. vitifoliae*.

Experiment	Trait	Linkage group (nearest marker)	LOD score	LOD threshold = 5%	Confidence interval (in cM)	Total variance explained by the QTL (%)
<i>In planta</i>	Max. number of nodosities	7 (VVIQ67)	3.06	2.59	35 - 87.70	19.02
		10 (VMC3D7)	2.59		0 - 63	11.50
<i>In planta</i>	BLUP nodosities	7 (VVIQ67)	4.33	2.94	55 - 87.70	20.13
		3 (3_5494608)	3.20		27 - 117.91	15.27
<i>In vitro</i>	Max. number of larvae	7 (7_4261424)	4.31	3.25	22 - 87.70	87.74
<i>In vitro</i>	BLUP larvae	7 (7_13408919)	10.93	3.44	29 - 77.16	70.11
		10 (VVIN85)	3.83		29 - 50.21	37.94

Table 3 Distribution of the resistance alleles identified in the resistant parent VRH8771 using SSRs markers among resistant BC1 individuals in response to *X. index*.

Linkage Group	SSR Marker	Number of the 47 resistant BC1 individuals displaying the resistance allelic form at the marker
9	VVBX-A-06	30
10	MRBX-27	26
10	VVIN78-SEC	26
10	VVIN85	26
18	UDV-108	24

Table 4 QTLs identified using a binary model in response to *X. index* performed on 60 BC1 individuals.

Linkage group (nearest marker)	LOD score	LOD threshold = 5%	LOD threshold = 10%	Confidence interval (in cM)	Total variance explained by the QTL (%)
9 (VVBX-A-06)	3.658	3.12	2.94	28.99 - 52.19	22.73
10 (SC8-03)	3.592	3.12	2.94	41.79 - 91.80	21.19
18 (UDV-108)	2.499	3.12	2.94	133.27 - 151.66	13.43

Figures




Figure 1

Distribution of the BC1 individuals according to (A) the maximal number of nodosities (in planta experiment) and (B) the maximal number of larvae (in vitro experiment). Parental genotypes, VRH8771

and CS, are represented by a green and pink bar, respectively.




Figure 2

Principal component analysis (PCA) of the 35 BC1 individuals tested in 2010-2011 and 2011-2012 experiments with root system development (RD), root weight (RW), nematode reproduction factor (RF) and gall index (GI). The two major principal components that accounted for 94.25 % of the variance have been plotted. The rating criteria factor map (left) and the individuals factor map (right) are reported. In the individual factor map, the BC1 individuals were assigned to two distinct groups, the resistant group in green and the susceptible group in pink, based on their RF and GI criteria.




Figure 3

QTL analysis of the resistance to X. index performed on 60 BC1 individuals. The y-axis represents the LOD score obtained by the binary mapping and the x-axis represents the 19 linkage groups related to the maternal genetic map (VRH8771). Curves in plot indicate the genetic coordinate (x-axis) and LOD score (y-axis). The red dotted line represents the LOD significant threshold estimated with 1000 permutations

for a level α of 0.05. The blue arrow corresponds to the genomic region below the significant threshold but with a high LOD score value.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Figures.docx](#)
- [Table.docx](#)