

Do the human gut metagenomic species possess the minimal set of core functionalities necessary for life?

Matteo Soverini (✉ matteo.soverini5@unibo.it)

University of Bologna <https://orcid.org/0000-0002-3026-9460>

Simone Rampelli

University of Bologna

Silvia Turrone

University of Bologna

Patrizia Brigidi

University of Bologna

Elena Biagi

University of Bologna

Marco Candela

University of Bologna

Research article

Keywords: Gut microbiome, metagenomic assembled genomes, uncultured metagenomic species, minimal bacterial genome

Posted Date: March 24th, 2020

DOI: <https://doi.org/10.21203/rs.2.15864/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on September 30th, 2020. See the published version at <https://doi.org/10.1186/s12864-020-07087-8>.

Abstract

Background: Advances in bioinformatics recently allowed for the recovery of 'metagenomes assembled genomes' from human microbiome studies carried on with shotgun sequencing techniques. Such approach is used as a mean to discover new unclassified metagenomic species, putative biological entities having distinct metabolic traits. **Results:** In the present analysis we compare 400 genomes from isolates available on NCBI database and 10,000 human gut metagenomic species, screening all of them for the presence of a minimal set of core functionalities necessary, but not sufficient, for life. As a result, the metagenome-assembled genomes resulted systematically depleted in genes encoding for essential functions apparently needed to support autonomous bacterial life. **Conclusions :** The relevant degree of lacking core functionalities that we observed in metagenome-assembled genomes raises some concerns about the effective completeness of metagenome-assembled genomes, suggesting caution in extrapolating information about their metabolic propensity and ecological role in a complex environment like the human gastrointestinal tract.

Background

Integral to the human biology, the Gut Microbiome (GM) is a key determinant of our health and its dysbiotic variations have been associated with several inflammatory diseases (1). Species-level variation in GM has been indicated as an emergent factor to be considered both for a better understanding of the biology of the GM-host mutualism (2) and for a refined evaluation of the individual health risk (3). However, even if it is perceived as strategic in GM study, the capability of shotgun metagenomics to infer species-level taxonomic and functional information is traditionally limited by the relative paucity of reference genomes. Indeed, despite the important progresses in culturomics, the degree of unclassified GM diversity at the species level is still very high. An important step forward in this direction has been recently provided by genome-resolved metagenomics, which involves the simultaneous recovery of draft and complete genomes directly from sequenced metagenomes (4). In particular, this approach consists in a de novo assembly of shotgun metagenomic reads into contigs, which are binned on the basis of coverage and tetranucleotide frequency (5, 6). This strategy allows the recovery of thousands of new genomes, i.e. the so called 'Metagenomes Assembled Genomes' (MAGs), directly from metagenomic reads, considerably expanding the tree of life beyond the limits of cultivability (7). Recently, Almeida et al. (8) provided a first extensive discovery campaign of MAGs from 13,133 human metagenomic samples. In particular, the Authors successfully characterized 1,175 nearly complete MetaGenomic Species (MGS), estimating a median completeness of 96.5% and 0.8% of contamination. Further, additional 893 medium quality MGS were also detected, with median completeness of 77.8% and 1.1% of contamination. The 94% of these MAG (1,952) did not match any bacterial isolate genome included in the Human-specific Reference (HR) (9) and RefSeq databases and thus indicated as new Unclassified MetaGenomic Species (UMGS). 74% of UMGS correspond to entirely novel genomes. 26% of the UMGS belonged to potential new families and 40% to new genera, thus expanding our current knowledge of human bacterial lineage by 281%. The Authors also performed an in-depth functional characterization of 2,505 human gut

species, 1,952 UMGS and 553 isolates from the Human Gut Reference (HGR) database, i.e. gut-specific species from the HR database (8). Interestingly, UMGS resulted depleted in genes involved in antioxidant activities and redox functions, being conversely enriched in iron-sulfur and iron binding genes. Thus, the Authors concluded that the recovered UMGS corresponded to strict anaerobes, with a distinctive metabolic propensity, well adapted to specific niches of the gastrointestinal tract with particularly low oxygen tension and high iron concentration.

Several research projects have been carried out with the specific purpose to define a 'minimal genome' as a model for understanding the basic functions of life (10). This resulted in the identification of a set of core functionalities necessary for a bacteria to survive and reproduce, as a universal minimal gene set represented in all living systems (11). In order to explore the efficacy of genome resolved metagenomics in providing comprehensive biological information on the uncultured members of the human microbiome, here we wondered if UMGS, which now remain bioinformatic entities, possesses the minimal set of core genes necessary – even if not sufficient – for life. To this aim, two publicly available minimal genomes were used as reference to generate a Core gene set of Minimal Functions (CMF), apparently necessary – but not sufficient – for life. Then we attempted to screen both UMGS and isolated NCBI genomes for the presence of genes included in CMF, showing that a remarkable number of UMGS were depleted in essential functionalities generally necessary for autonomous life.

Results And Discussion

The aim of the present study was to provide a first screening of UMGS and isolates genomes for a minimal subset of genetic functions (CMF) necessary – but not sufficient - to sustain bacterial life. In order to generate the CMF, two publicly available minimal genomes were downloaded from NCBI website: JCVI-syn 3.0 genome generated by Hutchison et al. (11) and *C. Eth-2.0* genome generated by Venetz et al. (12). The two genomes were annotated and only the genes assigned with certainty (not being prefixed by putative or hypothetical) and present in both genomes were retained and used as a reference set for the CMF. The CMF mostly includes genes involved in genetic information processing and cytosolic metabolism (Additional file 1). In particular, of the 190 genes included in CMF (Additional file 2), 143 were assigned by KEGG orthology to the genetic information processing pathways, with the functions involved in translation highly represented, including 115 genes among which 44 encode for ribosomal subunits, 20 for aminoacids-tRNA ligases, and 24 for tRNA. Replication and repair are other groups of functions highly represented in the CMF list, including 16 genes encoding for DNA polymerases, gyrases, and topoisomerases among others. Conversely, 35 out of 190 genes are devoted to metabolic functions, including especially carbohydrate metabolic pathways (e.g. glycolysis and gluconeogenesis, galactose metabolism, starch and sucrose metabolism, etc), energy metabolism (including all subunits of ATP synthase), and metabolism of nucleotides. Only two genes included in CMF are exclusively devoted to environmental information processes, and other two to cellular processes. However, 7 out of 190 genes showed multiple functionalities according to their KEGG orthology; for instance, Enolase is involved in metabolism, genetic information processes and environmental information processes, Phosphoglycerate kinase is involved in both metabolism and environmental information process, and two Protein

translocase subunits (SecA and SecY) are involved in genetic information processes, environmental information processes and cellular processes.

Next, we scanned both UMGS and isolated NCBI genomes for the presence of genes included in CMF. To this aim, a total of 10,400 genomes were selected and downloaded, covering all the most represented phyla in the gastrointestinal tract both for UMGS and NCBI genomes (see Additional file 3 for the distributions of the genomes at phylum level). In particular, 400 genomes of isolated species were obtained from NCBI, covering a wide array of bacterial species, and 10,000 genomes were selected from the UMGS in the database generated by Almeida et al. (8). For more information about the genomes included in this study, and the species included in the selected genomes, see Additional file 4. Each genome set was then annotated, and for both the NCBI and UMGS genomes, the presence or the absence of each gene included in CMF was verified, generating a binary matrix of CMF presence/absence profiles. For each tested genome, the percentages of adherence to the CMF and the absolute amounts of missing entries were also computed. Our analysis revealed that the NCBI and the UMGS genomes are characterized by a significant different presence of the CMF ($P < 0.001$, Kruskal-Wallis test), with the NCBI genomes showing a higher average representativeness value and a lower standard deviation when compared to UMGS ($93.2 \% \pm 2.9$ and $67.9 \% \pm 9.5$ Standard Deviation for NCBI and UMGS genomes, respectively) (Figure 1A). In Figure 1B the overall profile of the missing CMF in NCBI and UMGS genomes is reported. The CMF were found generally less represented in UMGS, with a total of 45 genes lacking in more than 50% analyzed genomes, with respect to the NCBI isolates. Among the missing genes, the 16S rRNA, fundamental for bacterial life, has not been retrieved in 8,034 out of 10,000 UMGS. This was due by the nature of the 16S rRNA gene, consisting in conserved and variable region and an overall similarity that can be up to 97% between two different bacterial species, making the assembly very hard in metagenomic data from microbial community. By analyzing the data at a higher level of detail, it has been possible to see how the genes encoding for ATP-dependent metalloprotease FtsH, DNA topoisomerase subunit 4, Dihydrolipolysine-residue acetyltransferase and Histidine biosynthesis protein HisB were the most absent within the UMGS genomes analyzed – i.e. systematically missing in more than 85% of the UMGS genomes. This group of 4 genes represent a set of various and biologically diverse functions: ATP-dependent metalloprotease FtsH gene encodes for a metalloprotease that plays a crucial role in the control of membrane protein integrity and regulates LPS biosynthesis (13)"; DNA topoisomerase subunit 4 gene encodes for a protein crucial in the chromosome segregation process, decatenating newly replicated chromosomes (14). A lack of this gene can lead to the impossibility for a bacterium to resolve DNA supercoilings, determining the lack of viability of the organism (15). On the other hand, the lack of Dihydrolipolysine-residue acetyltransferase, a component of the pyruvate dehydrogenase complex, can confirm the strict anaerobic propensity of the novel-identified bacteria species. Finally, Histidine biosynthesis protein HisB is a protein involved in step 6 and 8 of the sub-pathway that synthesizes L-histidine from 5-phospho-alpha-D-ribose 1-diphosphate. The function of this protein is crucial for bacterial life, since histidine is required for multiple biological processes (16).

Clustering analysis and PCA of the presence/absence profiles of CMF genes in NCBI and UMGS genomes showed a segregation between the two groups of genomes (Figure 2). Clustering analysis clearly

separates the genome batch in two parts, demarking a difference between the two types of genomes ($P < 0.001$, Fisher's exact test), with UMGS grouped on the left and right sides of the heatmap, flanking the NCBI genomes (Figure 2A). In the same graphics, it is possible to notice how UMGS genomes systematically lack in more genes when compared to NCBI genomes ($32\% \pm 34.5$ and $11.5\% \pm 24.2$ genomes missing for a single CMF gene in UMGS and NCBI set, respectively). Finally, the PCA analysis carried out using the binary Euclidean metric showed a separation of the genomes in the two-dimensional plan ($P < 0.001$, permutation test with pseudo-F ratio), with NCBI genomes less disperse if compared to UMGS, indicating a greater homogeneity in the representativeness of the CMF genes inside the NCBI group (Figure 2A).

Conclusions

The present report provides a first attempt at screening both newly-proposed Unknown metagenomic species and isolated genomes for the presence of a minimal set of core functionalities necessary but not sufficient for life. Our results showed that UMGS were substantially depleted in such essential genetic functions, including a recurrent depletion of 4 essential genes in more than the 85% of the UMGS, encoding for ATP-dependent metalloprotease FtsH, DNA topoisomerase subunit 4, Dihydrolipolysine-residue acetyltransferase and Histidine biosynthesis protein HisB. Even if a median completeness of up to 96.5% has been reported for UMGS (8), our data suggest that a relevant fraction of these genomes is missing genes encoding for essential functionalities to support life, thus raising concerns about their extrapolated biology. The recurrence and number of missed functionalities in at least some of the UMGS suggest caution when interpreting their metabolic and ecological propensity on the basis on their peculiar profile of gene relative abundance. Our analysis also points out the need to confirm the recovery of UMGS on metagenomic datasets obtained using third generation sequencing platforms providing longer reads, which have shown to aid genome completeness in *de novo* assembly and preserve more genomic information useful for species-level taxonomic assignment, such as operon structures (17). Further studies also comprising different UMAGs from different niches, are needed to verify and refine the accuracy of our results.

Methods

The minimal genomes JCVI-syn 3.0 (11) and *C. Eth-2.0* (12) were downloaded from NCBI website and annotated using prokka 1.13.3 standard pipeline and `-addgenes` flag (18) in a Unix CentOS environment. Genes assigned with certainty (not flagged with 'hypothetical' or 'putative' and manually screened) were retained for the creation of the CMF. A total of 400 NCBI and 10,000 UMGS genomes have been downloaded from the "Assembly" page of NCBI (the parameters "Complete" and "Representative" were selected) and from the European Nucleotide Archive under study ID PRJEB26432, respectively, and annotated as reported above for the synthetic genomes. For each, the presence/absence profile of CMF was obtained comparing the detected functions among each genome and the CMF, retaining only the matching hits. Clustering of the CMF presence/absence profiles of NCBI and UMGS genomes was

performed in R studio (version 1.2.1355 - R version 3.5.1 (19)), using the binary distance and the Ward's minimal variance clustering method (packages 'stats' V3.6.0 (15) and 'gplot' V3.0.1.1 (20)). Finally, the Euclidean distances between the CMF presence/absence profiles of UMGS and NCBI genomes were calculated and a multivariate analysis was carried out using the vegan package (V2.5-5 (21)). The separation between the NCBI genomes and the UMGS in the two-dimensional space was verified using a permutation test with pseudo-F ratio ('adonis' function of the 'vegan' package).

Abbreviations

CMF: Core gene set of Minimal Functions

GM: Gut Microbiota

HGR: Human Gut Reference

HR: Human-specific Reference

MAGs: Metagenome Assembled Genomes

PCA: Principal Components Analysis

UMGS: Unknown MetaGenomic Species

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

Availability of data and material

The datasets analyzed during the current study are available in the ENA repository, https://www.ebi.ac.uk/ena/data/view/PRJEB26432&portal=wgs_set, and in the NCBI repository, <https://www.ncbi.nlm.nih.gov/assembly/?term>.

Competing interests

The authors declare that they have no competing interests.

Funding

Not applicable.

Authors' contributions

MS and MC conceived the concept. MS developed the different scripts and performed the bioinformatics and statistical analysis. MS, MC, EB, and SR wrote the manuscript. PB, ST and EB revised and edited the draft. All authors discussed the results and have read and approved the manuscript.

Acknowledgements

Not applicable.

References

1. Integrative HMP (iHMP) Research Network Consortium. The Integrative Human Microbiome Project. *Nature*. 2019;569:641-8.
2. Zhao S, Lieberman TD, Poyet M, Kauffman KM, Gibbons SM, Groussin M, et al. Adaptive Evolution within Gut Microbiomes of Healthy People. *Cell Host Microbe*. 2019;25:656-67.
3. Zeevi D, Korem T, Godneva A, Bar N, Kurilshikov A, Lotan-Pompan M, et al. Structural variation in the gut microbiome associates with host health. *Nature*. 2019;568:43-8.
4. Devoto AE, Santini JM, Olm MR, Anantharaman K, Munk P, Tung J, et al. Megaphages infect *Prevotella* and variants are widespread in gut microbiomes. *Nat Microbiol*. 2019;4:693-700.
5. Anantharaman K, Brown CT, Hug LA, Sharon I, Castelle CJ, Probst AJ, et al. Thousands of microbial genomes shed light on interconnected biogeochemical processes in an aquifer system. *Nat Commun*. 2016;7:13219.
6. Delmont TO, Quince C, Shaiber A, Esen ÖC, Lee ST, Rappé MS, et al. Nitrogen-fixing populations of Planctomycetes and Proteobacteria are abundant in surface ocean metagenomes. *Nat Microbiol*. 2018;3:804-13.
7. Parks DH, Rinke C, Chuvochina M, Chaumeil PA, Woodcroft BJ, Evans PN, et al. Recovery of nearly 8,000 metagenome-assembled genomes substantially expands the tree of life. *Nat Microbiol*. 2017;2:1533-42.
8. Almeida A, Mitchell AL, Boland M, Forster SC, Gloor GB, Tarkowska A, et al. A new genomic blueprint of the human gut microbiota. *Nature*. 2019;568:499-504.
9. Forster SC, Kumar N, Anonye BO, Almeida A, Viciani E, Stares MD, et al. A human gut bacterial genome and culture collection for improved metagenomic analyses. *Nat Biotechnol*. 2019;37:186-92.

10. Martínez-García E, de Lorenzo V. The quest for the minimal bacterial genome. *Curr Opin Biotechnol*. 2016;42:216-24.
11. Hutchison CA 3rd, Chuang RY, Noskov VN, Assad-Garcia N, Deerinck TJ, Ellisman MH, et al. Design and synthesis of a minimal bacterial genome. *Science*. 2016;351:aad6253.
12. Venetz JE, Del Medico L, Wölfle A, Schächle P, Bucher Y, Appert D, et al. Chemical synthesis rewriting of a bacterial genome to achieve design flexibility and biological functionality. *Proc Natl Acad Sci U S A*. 2019;116:8070-9.
13. Katz C, Ron EZ. Dual role of FtsH in regulating lipopolysaccharide biosynthesis in *Escherichia coli*. *J Bacteriol*. 2008 Nov;190(21):7117-22.
14. Zechiedrich EL, Khodursky AB, Cozzarelli NR. Topoisomerase IV, not gyrase, decatenates products of site-specific recombination in *Escherichia coli*. *Genes Dev*. 1997 Oct 1;11(19):2580-92.
15. Usongo, V., & Drolet, M. (2014). Roles of type 1A topoisomerases in genome maintenance in *Escherichia coli*. *PLoS genetics*, 10(8), e1004543. <https://doi.org/10.1371/journal.pgen.1004543>
16. Alifano P, Fani R, Liò P, et al. Histidine biosynthetic pathway and genes: structure, regulation, and evolution. *Microbiol Rev*. 1996;60(1):44–69.
17. Nicholls SM, Quick JC, Tang S, Loman NJ. Ultra-deep, long-read nanopore sequencing of mock microbial community standards. *Gigascience*. 2019;8.
18. Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics*. 2014;30:2068-9.
19. R Core Team. R: A language and environment for statistical computing. 2017. R Foundation for Statistical Computing, Vienna, Austria. <https://www.r-project.org/>
20. Warnes GR, Bolker B, Bonebakker L, Gentleman R, Liaw WHA, Lumley T, et al. gplots: Various R Programming Tools for Plotting Data. R package version 3.0.1.1. <https://CRAN.R-project.org/package=gplots>.
21. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, et al. vegan: Community Ecology Package. R package version 2.5-5. <https://CRAN.R-project.org/package=vegan>.

Additional Files

Additional file 1 - file format: PDF. List of genes included in CMF. For each gene the KEGG orthology is detailed, with three functional levels reported whenever possible. CMF genes are grouped based on the lower level of KEGG orthology available.

Additional file 2 - file format: PDF. Detailed list of genomes included in the analysis. For the 400 NCBI isolates genomes species name is reported. For the 10,000 genomes retrieved from Almeida et al. study the values of completeness and contamination are reported.

Additional file 3 – file format: PDF. Pie charts representing the phyla distribution in the two cohorts of genomes analyzed (A for NCBI genomes, B for UMGS)

Figures




Figure 1

A) Percentage of genes in NCBI (skyblue) and UMGS (gold) genomes that were included in CMF. NCBI genomes show a significantly greater adherence to CMF ($P < 0.001$, Wilcoxon test). B) Superimposed distribution of missing CMF genes in NCBI (skyblue) and UMGS (gold) genomes. For each gene included in CMF, the number of genomes lacking the correspondent function is plotted. Genes in CMF are clustered according to the functional classes, as in Additional file 1.





Figure 2

A) Genomes clustering based on the presence/absence profile of CMF genes. The two generated clusters are highlighted by gold and skyblue underlying vectors for UMGS and NCBI genomes, respectively. The separation between the two groups is statistically significant ($P < 0.001$, Fisher's exact test). B) PCA based on Euclidean distances showing a significant separation between NCBI (skyblue) and UMGS (gold)

genomes according to the presence/absence profile of CMF genes ($P < 0.001$, permutation test with pseudo-F ratio).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile3.pdf](#)
- [Additionalfile1.pdf](#)
- [Additionalfile2.pdf](#)