

Enhanced production of falcarinol-type polyacetylenes in hairy roots of cultivated carrot (*Daucus carota* subsp. *sativus*)

Frank Dunemann (✉ frank.dunemann@julius-kuehn.de)

Julius Kühn-Institut (JKI), Institute for Breeding Research on Horticultural Crops, Quedlinburg, Germany
<https://orcid.org/0000-0002-3348-5177>

Christoph Böttcher

Julius Kühn-Institut (JKI), Institute for Ecological Chemistry, Plant Analysis and Stored Product Protection, Berlin, Germany

Short Report

Keywords: *Daucus carota*, *Rhizobium rhizogenes*, hairy root culture, HPLC/DAD, C17-polyacetylene, falcarinol, falcarindiol

Posted Date: December 1st, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-653311/v2>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Polyacetylenes (PAs) are a large group of bioactive phytochemicals, which are primarily produced by higher plants of the families Apiaceae and Araliaceae. Especially aliphatic C₁₇-polyacetylenes of the falcarinol-type such as falcarinol (FaOH) and falcarindiol (FaDOH) are known for their numerous biological functions including positive effects on human health. In this study we investigate the potential of carrot hairy root (HR) cultures for accumulation of PAs *in vitro*. Three individual plants of seven differently colored carrot cultivars were used for the development of HR cultures by transformation of root discs with the wild-type *Rhizobium rhizogenes* strain 15834. A total of 51 individual HR lines were established and quantitatively analysed by HPLC/DAD together with root, petiole and leaf tissue samples for their FaOH and FaDOH levels. Among the five tissues sampled from the donor plants, root periderm samples generally exhibited the highest PA levels with FaDOH as prevailing PA and large differences between cultivars. Compared with the periderm, FaOH levels were highly increased in HRs of all investigated carrot cultivars, whereas the FaDOH levels were not significantly increased. Considering the low to moderate PA concentration in root and leaf tissues of the orange cultivars there was an up to 30-fold increase of the FaOH concentration in HRs derived from orange genotypes. In this study, we demonstrate, that carrot HRs are able to produce large amounts of PAs. Hence, we suggest that HRs might be a suited tool to assess the function of candidate genes involved in the biosynthesis of major PAs.

Introduction

Polyacetylenes (PAs) are a large group of bioactive phytochemicals, which are primarily produced by higher plants of the families Apiaceae and Araliaceae both belonging to the order Apiales. They are also common in Asteraceae family members (Dawid et al. 2015). Among the more than 1.400 known PAs (Christensen and Brandt 2006), a subset of 12 structurally related PAs were isolated from *D. carota* (Dawid et al. 2015). These PAs are found in common vegetables and herbs of the Apiaceae family such as carrots, parsnip, fennel, celery, and parsley, with the C₁₇-PAs falcarinol (FaOH) and falcarindiol (FaDOH) as the main PAs (Czepa and Hoffmann 2003, Kidmose et al. 2004). Both FaOH and FaDOH are known to have a major impact on bitter off-taste in carrots, which is often the reason for consumer rejection of carrots or carrot products (Schmiech et al. 2008; Kreutzmann et al. 2008). On the other hand, several *in vitro* and *in vivo* studies showed that in particular the aliphatic C₁₇-PAs of the falcarinol-type exhibit potent anti-microbial, anti-inflammatory, and anti-cancer effects (Zidorn et al. 2005; Christensen and Brandt 2006; Christensen 2011; Dawid et al. 2015; Christensen 2020). For instance, falcarinol-type PAs may function as inhibitors of the breast cancer resistance protein BCRP/ABCG2, indicating their prospective use as agents for cancer chemotherapy (Tan et al. 2014). Rat feeding experiments with purified FaOH and FaDOH suggest preventing effects of these PAs on the development of colorectal cancer (Kobaek-Larsen et al. 2019). Despite their numerous biological functions the biosynthesis of PAs and its genetic basis are poorly understood. An efficient test system based on fast growing transformed root cultures might support the functional characterization of gene candidates involved in PA production.

Hairy root (HR) cultures obtained after transformation with the soil bacterium *Rhizobium rhizogenes* (*Rrhi*), formerly known as *Agrobacterium rhizogenes*, have often been used for tissue culture-based production of specialized plant metabolites (Malik 2018). Especially medicinal plants were transformed in the past with wild-type *Rrhi* strains to produce pharmaceutically relevant phytochemicals (Guillon et al. 2008). Since the advent of genome editing techniques such as CRISPR/Cas9 HR-based transformation systems are evolving to an efficient research tool to elucidate physiological processes and biosynthetic pathways (Shi et al. 2021). Examples for genome editing approaches based on HRs were reported for several plant species such as *Salvia* (Li et al. 2017), *Eucalyptus* (Dai et al. 2020), and *Medicago* (Zhang et al. 2020).

The cultivated carrot (*Daucus carota* ssp. *sativus*) is one of the most important root vegetable crops grown worldwide. Carrots provide not only one of the richest sources of the vitamin A precursor β -carotene, but contain many other bioactive and putatively health-relevant phytochemicals including anthocyanins, phenylpropanoids, terpenoids, and the above mentioned PAs. Several Apiaceae species were investigated in the past for production of plant natural products in HR culture systems (Baranski 2008). Starting with pioneer work in early 1980s (Chilton et al. 1982; Willmitzer et al. 1982) several HR-based studies were reported for *D. carota* (Kim and Yoo 1996; Ridgway et al. 2004; Baranski et al. 2006). Recently, HR cultures derived from dark-purple carrots were shown to be a suited platform for research on the *in vitro* production of anthocyanins and phenolic compounds (Barba-Espin et al. 2020).

To our knowledge, in carrots the production of PAs in HR cultures has not been described yet.

In this study we use a *Rrhi*-based transformation system for generation of HRs derived from orange, yellow and purple carrots and elucidate their potential to produce sufficient amounts of two major PAs, FaOH and FaDOH. We show that especially FaOH is strongly increased in HRs of orange carrots compared to peridermal root tissue and suggest that carrot HR cultures are well suited for functional studies of genes involved in carrot PA biosynthesis.

Material And Methods

Plant material

Seven carrot cultivars (*D. carota* subsp. *sativus*) with differently coloured roots were used in this study: 'Anthonina' (AN) - purple, 'Deep Purple' (DP) - purple, 'Yellowstone' (YS) - yellow, '710015' (71) - yellow, 'Rotin' (RO) - orange, 'Lange Rote Stumpfe ohne Herz' (LR) - orange, 'Shinsuu Senkou Oonaga' (SO) - orange. Seeds were obtained from the JKI carrot seed repository kept by Dr. Thomas Nothnagel. Three plants of each cultivar were sown and cultivated in 17 cm plastic pots in a standard soil mixture (sand/humus, 3/1 (v/v)) under controlled greenhouse conditions (16 h photoperiod, 20 – 25°C) until harvest about 15 weeks after sowing. Taproots were washed to remove adhering soil and blotted dry. Periderm samples were obtained from a 3-cm section from the middle of each carrot root using a peeler. Afterwards, roots were latitudinally halved and an 8 mm thick slice was removed from each root half. Using a scalpel phloem and xylem samples were prepared from both slices. The remaining root material

was sterilized and used for transformation. Petiole samples were obtained by cutting a 5-cm section from the middle of three petioles. Leaf tissue was sampled from younger leaflets. Plant material was shock frozen in liquid nitrogen immediately after harvest and stored at -80°C until freeze-drying.

Transformation of carrot with *Rhizobium rhizogenes*

Pieces of tap roots remaining after PA sampling were surface-sterilized via immersion in 70% ethanol for 5 min followed by a 20 min treatment with 4% sodium hypochlorite supplemented with 0.1% Tween 20. Each carrot root was sliced into 0.5 – 0.8 cm thick discs and dead tissue of the outer edge of the discs was removed. The apical (shoot-averted) side was labeled with a sterile toothpick, and the carrot discs were placed (apical side up) onto water agar (2% agar in water; Duchefa, Haarlem, Netherlands). Wild-type *Rrhi* strain ATCC 15834 (Lippincott et al. 1973) was used to grow bacterial inocula in dark conditions at 28°C for about 24 h in liquid CPY medium (bacto-yeast extract 0.1%, peptone 0.5%, saccharose 0.5%, bacto-agar 20 g L⁻¹, MgSO₄x7H₂O 200 mM) on a rotary shaker (180 rpm). About 100 µl of the bacterial suspension (OD₆₀₀ of 0.6 - 0.8) was spread along the cambial ring on the apical side of the root discs. After co-cultivation for seven days on 2% water agar medium the carrot discs were placed for up to five weeks on 2% water agar supplemented with 200 mg L⁻¹ cefotaxime (Duchefa, Haarlem, Netherlands) and 100 mg L⁻¹ carbenicillin (Roth, Karlsruhe, Germany) to eliminate the bacteria. HRs started to regenerate generally after two weeks and were harvested weekly alongside with a small piece of original root tissue. Regenerated HRs were placed in small petridishes (6 cm diameter) on hormone-free half-concentrated MS (Murashige & Skoog) medium with 200 mg L⁻¹ cefotaxime and 100 mg L⁻¹ carbenicillin (½ MS-CC) and cultured in the dark at 22°C. HR cultures developing from single excised roots that originated unambiguously from different regions of the root disc were considered to have an independent origin of regeneration. They were placed in a small single petridish and were designated as 'hairy-root line' (HR line). About four weeks later the propagating HRs were subcultured on the same culture medium. After further one to two subcultures in small plates, the HRs were placed on filter paper located on the surface of a ½ MS culture medium without antibiotics in 9 cm petri dishes. Four weeks later the HR lines were harvested and frozen immediately at -80°C until chemical analysis.

In order to verify the transgenic character of the HRs and to prove that they were free of contaminating *Rrhi* bacteria we performed PCR analyses. For DNA isolation, 50 - 100 mg fresh hairy roots were snap frozen in liquid N₂ and grinded using a swing mill to extract total genomic DNA with the innuPREP Plant DNA kit (Analytik Jena, Jena, Germany). The DNA content was measured via NanoDrop device (Thermo Fisher). To check the transgenic status of the HR lines, we used PCR primer for the *rolC* gene from the Ri plasmid of *Rrhi*, which are diagnostic for the integration of the T-DNA into the plant genome (for primer information and PCR conditions, see Medina-Bolivar et al. 2007). To prove that the presence of the *rolC* gene is due to its integration into the genome of *D. carota* and not due to contamination with *Rrhi* we performed PCR analysis with primer for the *Rhizobium virD2* gene which is not transmitted into the plant genome (Medina-Bolivar et al. 2007).

Sample preparation and analysis of polyacetylenes by HPLC/DAD

Deep-frozen plant material (leaf, petiole, periderm, phloem, xylem, hairy roots) was freeze-dried for four days (Christ Gamma 1-16 LSC, condenser temperature -50°C , pressure 0.04 mbar). Depending on the quantity, dried plant material was homogenized in 1.5-ml or 5-ml polypropylene centrifuge tubes or in 10-ml polyethylene vials using steel balls (\varnothing 3 - 10 mm) and a mixer mill (Retsch MM 400, 30 Hz, 1 - 2 min). Homogenized plant material was precisely weighed [leaf and petiole tissue: (40 ± 1) mg; periderm, phloem, xylem, hairy root tissue: (20 ± 1) mg] into a 1.5-mL polypropylene centrifuge tube. After addition of three steel balls (\varnothing 3 mm), 50 μL internal standard solution (0.2 g L^{-1} *N*-vanillylnonamide in MeOH) and 300 μL acetone the mixture was homogenized using a mixer mill (Retsch MM400, 30 Hz, 60 s, room temperature). Afterwards, the sample was sonicated (ElmaSonic P, 37 kHz, 100 W, 5 min, 20°C) and shaken (2400 min^{-1} , 10 min, room temperature). After centrifugation ($13000 g$, 5 min, 22°C) a 200- μL aliquot of the supernatant was transferred into an HPLC vial with micro-insert and stored at 6°C until analysis.

Polyacetylene analyses were performed on an 1100 Series HPLC system (Agilent Technologies) comprising a degasser (G1322A), a binary pump (G1312A), an autosampler (G1329A), an autosampler thermostat (G1330A), a column compartment (G1316A) and a diode array detector (G1315A). Extracts (injection volume 2.5 μL) were separated on a Zorbax Eclipse XDB-C18 column (3 mm \times 150 mm, 3.5 μm particle size, Agilent Technologies) using water and acetonitrile as eluent A and B, respectively. The following binary gradient program at a flow rate of 1 mL min^{-1} was used: 0-10 min, linear from 50 to 80 % B; 10-10.5 min, linear from 60 to 100 % B; 10.5-13 min, isocratic, 100 % B; 13-15 min, isocratic 50 % B. The column and autosampler temperature was maintained at 40°C and 6°C , respectively. The diode array detector response time was set at 0.2 s, the optical slit width at 4 nm. Polyacetylenes were detected at 196 nm with a spectral bandwidth of 4 nm, the internal standard *N*-vanillylnonamide at 204 nm with a spectral bandwidth of 4 nm. ChemStation software (version B.03.02) was applied for controlling the instrument, data acquisition and quantitative analysis. *N*-Vanillylnonanamide (t_{R} 2.53 min), FaDOH (t_{R} 5.09 min) and FaOH (t_{R} 9.48 min) were quantified based on peak area using external standard calibration method. Therefore, the following calibration curves were established : (i) *N*-vanillylnonamide: calibration range 1 – 600 ng, 14 points, linear regression model ($y = m x$), equal weighting, $R^2 = 0.99987$; (ii) FaDOH: calibration range 5 – 500 ng, 9 points, linear regression model ($y = m x$), equal weighting, $R^2 = 0.99996$; (iii) FaDOH: calibration range 500 – 1000 ng, 6 points, logarithmic regression model ($y = m \ln(x) + b$), equal weighting, $R^2 = 0.99919$; (iv) FaOH: calibration range 5 – 1000 ng, 14 points, linear regression model ($y = m x$), equal weighting, $R^2 = 0.99970$. Polyacetylene levels were corrected using the recovery rate of the internal standard. The average recovery rate of the internal standard was 101.3 % (standard deviation 4.6 %, $n = 156$).

Results And Discussion

Establishment of carrot hairy root lines

Hairy root induction in seven differently coloured *D. carota* cultivars was accomplished by co-culture of carrot root disks and *Rrho* wild strain 15834. Two weeks after inoculation first HRs protruded from the upper surface of the root disks. HR lines were established from single excised roots and, after selection for sufficient growth, used for PA analysis. To prove the presence of the T-region of the Ri plasmid and the integration of the *rol* (*root loci*) genes into the genome of *D. carota* we performed PCR analysis with *rolC* primer. The transgenic character of the HR lines was verified, and the *virD2* fragment indicating bacterial contamination was not detected in any HR line analyzed (PCR results not shown). A single transformation experiment based on three individual tap roots of each carrot cultivar used for PA analysis resulted in a total number of 51 well growing HR lines at the end of the propagation process (Table 1). Cultivars DP, YS, and LR totally yielded the desired three lines per plant (totally 9), and for the other cultivars the number of HR lines varied from 8 in RO to 4 in AN (Table 1). For graphical presentation and statistical evaluation of PA data from different plant tissues and HRs (Fig. 1) only the cultivars DP, YS, LR and RO were considered, for which HRs from all three original plants were available.

Rrho strain ATCC 15834 proved to be highly effective for transformation of *D. carota*. This wild strain, used first for carrot transformation by Willmitzer et al. (1982), has shown strong virulence in many plant studies. Among the about 100 plant species, which have been reported for a subsequent plant regeneration from HRs carrying a wild type Ri plasmid, about 20 were transformed by the 15834 strain (Desmet et al. 2020). John et al. (2017) examined four different *Rrho* strains, of which strain 15834 showed significantly higher transformation efficiency and was capable of inducing HRs from different explants of *Achyranthes aspera*. Transformation of 'Indian ginseng' (*Withania somnifera*) was only successful with strain 15834 (Saravanakumar et al. 2012).

Polyacetylene contents and distribution patterns in carrot donor plants

Two main PAs FaOH and FaDOH were quantified in carrot plant tissues and HRs (Fig. 1). Among the five leaf and root tissues we detected the highest concentration of both compounds in the periderm samples and the lowest in the leaf and xylem tissues, but there were large differences among the cultivars. Peridermal tissue of the purple cultivar DP accumulated approximately 600 µg/g DW (dry weight) FaOH, but in the orange cultivars LR and RO significantly lower FaOH levels of approximately 50 and 100 µg/g DW were registered, respectively. In DP the peridermal PA levels were significantly enhanced in comparison to the other plant samples (Fig. 1). However, due to the large variation of the PA data among the samples from individual plants most other differences were not significant. With regard to FaDOH, the purple cultivar DP also showed highest concentration in the periderm (1200 µg/g) whereas the peridermal FaDOH levels of the orange cultivars LR and RO were in the range of 450 - 650 µg/g DW (Fig. 1). The highest FaDOH contents in the leaf samples were measured in DP with about 100 µg/g and in YS with 240 µg/g.

PA accumulation patterns are influenced by numerous factors such as genotype, developmental stage, and abiotic and biotic stress factors (Dawid et al. 2015). Besides, the PA distribution among the different

plant organs and within the different root segments can vary considerably. Since the major aim of this study was not the analytical dissection of carrot plants, but to determine the potential of PA production in HRs, we analysed three selected individuals of each cultivar for HR induction and used leaf and root samples taken from each donor plant before *Rrhi* transformation as reference. Nevertheless, it was evident, that in all cultivars the periderm samples displayed the highest FaDOH levels. According to Czepa and Hofmann (2004) the most abundant PA in cultivated orange carrots is FaDOH, and the highest total PA levels were found in the periderm tissue (Baranska and Schulz 2005; Baranska et al. 2005). Busta et al. (2018) analysed tissue-specific accumulation of PAs in the orange cultivar 'Danvers' and also measured the highest total PA level in the peridermal tissues, with FaDOH as the dominating PA.

With regard to FaOH, the purple cultivars DP and AN (data not shown for AN) appeared to accumulate considerably higher contents in the periderm compared with the other tissue samples. One reason for this finding might be the ancestry of purple carrots. It is generally recognized, that the origin of the cultivated carrot was in Central Asia (Afghanistan), where purple-rooted wild carrots were domesticated and spread westward more than 1.000 years ago (Simon et al. 2008). Orange carrots occurred in the 15th and 16th centuries in Europe (Stolarczyk and Janick 2011), which indicates a secondary domestication process. Purple and red carrots persisted in Asia and the Middle East, and it can be assumed that current purple carrot cultivars are genetically closer to early domesticated purple carrots than to the modern orange forms, which were selected for less bitterness and likely for less PA contents.

Table 1 Levels of falcarinol (FaOH) and falcarindiol (FaDOH) in hairy root cultures and periderm samples obtained from seven carrot cultivars

Cv.	Plant	FaOH Level					FaDOH Level				
		[$\mu\text{g (g dry weight)}^{-1}$]					[$\mu\text{g (g dry weight)}^{-1}$]				
		HR Line#1	HR Line#2	HR Line#3	HR Mean ¹	Periderm	HR Line#1	HR Line#2	HR Line#3	HR Mean ¹	Periderm
AN	1	2027	na ²	na ²	2027	934	2405	na ²	na ²	2405	1010
	2	1967	1558	1642	1722	442	4248	1521	1553	2440	1481
	3	na ²	na ²	na ²	na ²	793	na ²	na ²	na ²	na ²	1108
DP	1	2173	2082	4162	2805	551	2640	2436	2679	2585	1134
	2	3766	2358	1955	2693	576	1321	1905	1638	1621	972
	3	2862	3203	2101	2722	623	1522	2154	1522	1733	1596
YS	1	2047	3481	3374	2967	87	1407	1417	1407	1411	722
	2	1142	1219	1513	1291	51	826	696	1458	994	565
	3	2021	1692	2401	2038	493	2323	1931	2011	2088	1673
71	1	1877	1995	1580	1818	493	849	739	1255	948	720
	2	na ²	na ²	na ²	na ²	127	na ²	na ²	na ²	na ²	750
	3	2050	1642	1701	1798	287	678	721	582	660	900
LR	1	1994	1284	1822	1700	74	832	403	763	666	827
	2	1534	1155	2207	1632	27	984	1243	1105	1111	85
	3	1726	884	977	1196	38	2431	1010	1034	1491	416
RO	1	1683	1287	na ²	1485	77	1164	1220	na ²	1192	729
	2	2270	2306	1384	1987	193	807	660	885	784	649
	3	1841	1289	1448	1526	68	1301	1288	839	1143	574
SO	1	711	1682	1370	1254	70	848	940	1136	975	627
	2	1188	1133	1334	1219	128	1354	1234	946	1178	747
	3	na ²	na ²	na ²	na ²	512	na ²	na ²	na ²	na ²	1391

¹mean level of HR lines developed from an individual plant, ²not available

Polyacetylene production in carrot hairy roots

Quantification of PAs in freeze-dried root samples of 51 individual HR lines revealed strong increases of the FaOH contents in all seven analysed cultivars ranging - on average of a single donor plant - from 1200 to 3000 $\mu\text{g/g DW}$ in the HRs (Table 1). In case of the four cultivars DP, YS, RO and LR a significant increase of the FaOH level was observed in comparison to the periderm. For FaDOH there was also an increase in the HR lines (range 660 to 2600 $\mu\text{g/g DW}$) but compared with the periderm values the increase was less pronounced and not significant in the four fully analyzed cultivars (Table 1, Fig. 1). The HR line with the highest FaOH content was 'DP 1-3' with 4162 $\mu\text{g/g DW}$, which is about 7 times more than the content in the periderm of the original taproot. The HR line 'AN 2-1' contained with 4247 $\mu\text{g/g DW}$ the highest FaDOH content (Table 1). Considering the low to moderate PA contents even in the periderm of the orange cultivars such as LR and RO, there was an approximately 30-fold increase of the FaOH concentration in HRs of LR. It might be possible, that especially the amount of FaOH is enhanced in HRs of cultivated (orange) carrot genotypes due to unknown genetic factors influenced by the domestication process. HRs derived from purple and orange carrots might be a suited system for further research to identify genes involved in the accumulation of FaOH and FaDOH.

The results of our study indicate, that field-grown carrots with relatively high PA contents, such as for instance purple cultivars like DP, might be used in future for the large-scale production of pharmaceutically relevant falcarinol-type PAs. Carrot HR cultures might be a suited alternative plant source, and bioreactor techniques seem to be a possible way to produce large quantities of PAs. However, for latter approach further research on the consistency and kinetics of PA production in HRs would be necessary. HR cultures ave been used in numerous plant species for the synthesis of a wide spectrum of phytochemicals (Malik 2018; Gutierrez-Valdes et al. 2020), but few reports are available for falcarinol-type PA production in HRs. FaOH was among the main components of oil samples from HR cultures of the Apiaceae species *Levisticum officinale* (Santos et al. 2005), and Washida et al. (2003) reported the production of FaOH and FaDOH in HRs of *Panax* hybrids. HRs of *Panax ginseng* were used to investigate the natural formation of FaOH and FaDOH by ¹³C-labeling experiments (Knispel et al. 2013). A major bottleneck in functional gene studies in non-model plants is often the lack of an efficient transformation protocol for recalcitrant genotypes or whole species. Since it is mostly easier to induce HRs using *R. rhizogenes* than to generate transformants using *A. tumefaciens*, HR-based functional gene studies might be a practicable alternative, especially for research on root-derived natural compounds.

Declarations

Acknowledgements

The authors wish to thank Jochen Kumlehn (IPK Gatersleben) for providing the *Rrhi* 15834 strain, and Vicky Bartels (JKI Quedlinburg) and Astrid Hansen (JKI Berlin) for their excellent technical assistance on HR culture development and PA analyses.

Authors' contributions

F.D. and C.B. designed the study. F.D. performed the transformation experiments, and C.B. performed and PA analyses. F.D. and C.B. wrote the manuscript. Both authors read and approved the final manuscript.

References

Barba-Espin G, Chen S-T, Agnolet S, Nymark Hegelund J, Stanstrup J, Christensen JH, Müller R, Lütken H (2020) Ethephon-induced changes in antioxidants and phenolic compounds in anthocyanin-producing black carrot hairy root cultures. *Journal of Experimental Botany* 71:7030-7045. <https://doi.org/10.1093/jxb/eraa376>

Baranska M, Schulz H, Baranski R, Nothnagel T, Christensen L (2005) In situ simultaneous analysis of polyacetylenes, carotenoids and polysaccharides in carrot roots. *Journal of Agricultural and Food*

Chemistry 53:6565–6571. <https://doi.org/10.1021/jf0510440>

Baranski R (2008) Genetic transformation of carrot (*Daucus carota*) and other Apiaceae species. *Transgenic Plant Journal* 2:18-38.

Baranski R, Klocke E, Schumann G (2006) Green fluorescent protein as an efficient selection marker for *Agrobacterium rhizogenes* mediated carrot transformation. *Plant Cell Reports* 25:190–197.

<https://doi.org/10.1007/s00299-005-0040-2>

Busta L, Yim WC, LaBrant EW, Wang P, Grimes L, Malyszka K, Cushman JC, Santos P, Kosma DK, Cahoon EB (2018) Identification of genes encoding enzymes catalyzing the early steps of carrot polyacetylene biosynthesis. *Plant Physiology* 178:1507–1521. <https://doi.org/10.1104/pp.18.01195>

Chilton MD, Tepfer D, Petit A, David C, Casse-Delbart F, Tempe J (1982) *Agrobacterium rhizogenes* inserts T-DNA into plant roots. *Nature* 295:432-434.

Christensen LP (2011) Aliphatic C(17)-polyacetylenes of the falcarinol type as potential health promoting compounds in food plants of the Apiaceae family. *Recent Patents on Food, Nutrition & Agriculture* 3:64–77. <https://doi.org/10.2174/2212798411103010064>

Christensen LP (2020) Bioactive C17 and C18 acetylenic oxylipins from terrestrial plants as potential lead compounds for anticancer drug development. *Molecules* 25:2568. <https://doi.org/10.3390/molecules25112568>

Christensen LP, Brandt K (2006) Bioactive polyacetylenes in food plants of the Apiaceae family: occurrence, bioactivity and analysis. *Journal of Pharmaceutical and Biomedical Analysis* 41:683–693. <https://doi.org/10.1016/j.jpba.2006.01.057>

Czepa A, Hofmann T (2003) Structural and sensory characterization of compounds contributing to the bitter off-taste of carrots (*Daucus carota* L.) and carrot puree. *Journal of Agricultural and Food Chemistry* 51:3865–3873. <https://doi.org/10.1021/jf034085>

Czepa A, Hofmann T (2004) Quantitative studies and sensory analyses on the influence of cultivar, spatial tissue distribution, and industrial processing on the bitter off-taste of carrots (*Daucus carota* L.) and carrot products. *Journal of Agricultural and Food Chemistry* 52:4508–4514. <https://doi.org/10.1021/jf0496393>

Dai Y, Hu G, Dupas A, Medina L, Blandels N, Clemente H, et al. (2020) Implementing the CRISPR/Cas9 technology in *Eucalyptus* hairy roots using wood-related genes. *International Journal of Molecular Sciences* 21(10). <https://doi.org/10.3390/ijms21103408>

Dawid C, Dunemann F, Schwab W, Nothnagel T, Hofmann T (2015) Bioactive C₁₅-polyacetylenes in carrots (*Daucus carota* L.): Current knowledge and future perspectives. *Journal of Agricultural and Food Chemistry* 63:9211–9222. <https://doi.org/10.1021/acs.jafc.5b04357>

Desmet S, Dhooghe E, De Keyser E, Van Huylenbroeck J, Müller R, Geelen D, Lütken H (2020) Rhizogenic agrobacteria as an innovative tool for plant breeding: current achievements and limitations. *Applied Microbiology and Biotechnology* 104:2435–2451. <https://doi.org/10.1007/s00253-020-10403-7>

Guillon S, Trémouillaux-Guiller J, Kumar PP, Gantet P (2008) Hairy Roots: a powerful tool for plant biotechnological advances. In: Ramawat K, Merillon J (eds) *Bioactive Molecules and Medicinal Plants*. Springer, Berlin, Heidelberg. https://doi.org/10.1007/978-3-540-74603-4_14

Gutierrez-Valdes N, Häkkinen S, Lemasson C, Guillet M, Oksman-Caldentey KM, Ritala A, Cardon F (2020) Hairy root cultures - a versatile tool with multiple applications. *Frontiers in Plant Science* 11:33. <https://doi.org/10.3389/fpls.2020.00033>

John R, Shajitha PP, Devassy A, Mathew L (2017) Hairy-root cultures of *Achyranthes aspera* Linn. as a novel route for the production of 20-hydroxyecdysone. *Acta Physiol Plant* 39:255.

<https://doi.org/10.1007/s11738-017-2555-x>

Kidmose U, Hansen SL, Christensen LP, Edelenbos M, Larsen E, Nørbaek R (2004) Effects of genotype, root size, storage, and processing on bioactive compounds in organically grown carrots (*Daucus carota* L.). *Journal of Food Science* 69:S388-S394.

Kim YH, Yoo YJ (1996) Peroxidase production from carrot hairy root cell culture. *Enzyme and Microbial Technology* 18:531-535.

Knispel N, Ostrozhenkova E, Schramek N, Huber C, Pena-Rodriguez LM, Bonfill M, Palazon J, Wischmann G, Cusido RM, Eisenreich W (2013) Biosynthesis of panaxynol and panaxydol in *Panax ginseng*. *Molecules* 18:7686-7698. <https://doi.org/10.3390/molecules18077686>

Kobaek-Larsen M, Baatrup G, Notabi MK, El-Houri RB, Pipó-Ollé E; Christensen-Arnspang E, Christensen LP (2019) Dietary polyacetylenic oxylipins falcarinol and falcarindiol prevent inflammation and colorectal neoplastic transformation: A mechanistic and dose-response study in a rat model. *Nutrients* 11:2223. <https://doi.org/10.3390/nu11092223>

Kreutzmann S, Christensen LP, Edelenbos M (2008) Investigation of bitterness in carrots (*Daucus carota* L.) based on quantitative chemical and sensory analyses. *LWT - Food Science and Technology* 41:193–205.

Li B, Cui G, Shen G, Zhan Z, Huang L, Chen J, Qi X (2017) Targeted mutagenesis in the medicinal plant *Salvia miltiorrhiza*. *Scientific Reports* 7:43320. <https://doi.org/10.1038/srep43320>

Lippincott JA, Beiderbeck R, Lippincott BB (1973) Utilization of octopine and nopaline by *Agrobacterium* sp. *J Bacteriol* 116:378-383.

Malik S (ed) (2018): *Production of Plant Derived Natural Compounds through Hairy Root Culture*. Springer International Publishing. 1st edition 2018. Springer International Publishing.

Medina-Bolivar F, Condori J, Rimando AM, Hubstenberger J, Shelton K, et al. (2007) Production and secretion of resveratrol in hairy root cultures of peanut. *Phytochemistry* 68:1992-2003. <https://doi.org/10.1016/j.phytochem.2007.04.039>

Ridgway HJ, Kandula J, Stewart A (2004) Optimising production of carrot hairy roots. *New Zealand Plant Protection* 57:77-80.

Santos PAG, Figueiredo AC, Oliveira MM, Barroso JG, Pedro LG, Deans SG, Scheffer JJC (2005) Growth and essential oil composition of hairy root cultures of *Levisticum officinale* W.D.J. Koch (lovage). *Plant Science* 168:1089-1096.

Saravanakumar A, Aslam A, Shajahan A (2012) Development and optimization of hairy root culture systems in *Withania somnifera* (L.) Dunal for withaferin-A production. *African Journal of Biotechnology* 11:16412-16420. <https://doi.org/10.5897/AJB11.3867>

Schmiech L, Uemura D, Hofmann T (2008) Reinvestigation of the bitter compounds in carrots (*Daucus carota* L.) by using a molecular sensory science approach. *J. Agric. Food Chem.* 56:10252-10260.

<https://doi.org/10.1021/jf8023358>

Shi M, Liao P, Nile SH, Georgiev MI, Kai G (2021) Biotechnological exploration of transformed root culture for value-added products. *Trends in Biotechnology* 39, No.2. <https://doi.org/10.1016/j.tibtech.2020.06.012>

Simon PW, Freeman RE, Vieira JV et al (2008) Carrot. In: Prohens J, Nuez F (eds) *Handbook of plant breeding*, vol 2. Springer, New York, NY, pp 327–357

Stolarczyk J, Janick J (2011) Carrot: History and Iconography. *Chronica Horticulturae* 2:13-18.

Tan KW, Killeen DP, Li Y, Paxton JW, Birch NP, Scheepens A (2013) Dietary polyacetylenes of the falcarinol type are inhibitors of breast cancer resistance protein (BRCP/ABCG2). *Eur J Pharmacology* 723:346–352. <https://doi.org/10.1016/j.ejphar.2013.11.005>

Washida D, Shimomura K, Kitanaka S (2003) Polyacetylenes in hairy roots of a *Panax* hybrid. *Planta Med* 69:1163-1165.

Willmitzer L, Sanchez-Serrano J, Buschfeld E, Schell J (1982) DNA from *Agrobacterium rhizogenes* is transferred to and expressed in axenic hairy root plant tissues. *Mol Gen Genet* 186:16-22.

Zhang H, Cao Y, Zhang H, Xu Y, Zhou C, Liu W, et al. (2020) Efficient generation of CRISPR/Cas9-mediated homozygous/biallelic *Medicago truncatula* mutants using a hairy root system. *Frontiers in Plant Science* 11:294. <https://doi.org/10.3389/fpls.2020.00294>

Zidorn C, Jöhner K, Ganzera M, Schubert B, Sigmund EM, Mader J, et al. (2005) Polyacetylenes from the Apiaceae vegetables carrot, celery, fennel, parsley, and parsnip and their cytotoxic activities. *Journal of Agricultural and Food Chemistry* 53:2518–2523. <https://doi.org/10.1021/jf048041s>

Figures




Figure 1

Falcarinol (FaOH) and falcarindiol (FaDOH) levels in different leaf and root tissues and hairy roots. Bars represent means and error bars standard errors of mean (leaf, petiole, periderm, phloem, xylem: n = 3; hairy root: n = 9 for DP, YS and LR; n = 8 for RO). Significance analyses were performed after log-transformation using Tukey's HSD test. Means with the same latin or greek letter are not significantly different ($P \geq 0.05$).