

Association between social capital and loneliness among older adults: A cross-sectional study in Anhui Province, China

Zhongliang Bai

Anhui Medical University

Zijing Wang

Anhui Medical University

Tiantai Shao

Anhui Medical University

Xia Qin

Anhui Medical University

Zhi Hu (✉ aywghz@ahmu.edu.cn)

Anhui Medical University

Research article

Keywords: Social Capital, Aging, Health, Loneliness, China

Posted Date: January 7th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-66051/v3>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on January 7th, 2021. See the published version at <https://doi.org/10.1186/s12877-020-01973-2>.

Abstract

Background: We aimed to examine the association between social capital and loneliness in Anhui Province, China.

Methods: Data were collected from a cross-sectional study using a multi-stage stratified cluster sampling strategy. Data on demographic characteristics, socioeconomic factors, social capital, and loneliness in 1810 older adults (aged 60 years and older) were used for analysis. Binary logistic regression models and a classification and regression tree model were performed to assess the association of social capital and loneliness.

Results: Our results indicated that social capital in terms of lower level of social participation (AOR = 1.38; 95% CI: 1.10-1.74), social connection (AOR = 1.51; 95% CI: 1.18-1.93), and reciprocity (AOR = 1.47; 95% CI: 1.13-1.90) were associated with higher odds of developing loneliness. We noted the interactive effect of different social capital dimensions on loneliness, suggesting that the risk for suffering loneliness was greatest in older people limited in functional ability, with less trust, less social connection, and less social participation.

Conclusions: Our findings show that social capital is associated with loneliness in older adults. This implies that social capital, especially in terms of trust, social connection, and social participation may be significant for alleviating loneliness in later life.

Background

Although the size of an aging society among the people has been soaring globally, yet a prolonged life is not accompanied by good health and well-being [1, 2]. Studies have found that loneliness, as a negative emotional status, can degenerate the health and well-being of older people during aging [3-5]. For example, loneliness has been found to prompt coronary heart disease and stroke [6], and dementia [7], consequently leading to poor quality of life [8, 9].

Loneliness is frequent among older people [5], yet the prevalence of loneliness varies across different countries according to different assessment tools and study samples. A national longitudinal study in Singapore found that about 23.0% of older people suffered from chronic loneliness [5]. A cross-sectional survey in Finland found that 27.3% of the older adults had frequent loneliness [10]. Prior findings from the Chinese Longitudinal Healthy Longevity Survey (CLHLS) revealed that about 22.9% of older men and 30.6% of older women experienced loneliness [11]. Therefore, how to prevent and reduce the incidence of loneliness is a pressing issue in public health research.

A previous study has evidenced that determining the risk factors of loneliness is of importance to prevent loneliness [12]. Some factors such as females [11], advanced age [13], low level of educational attainment, and socioeconomic status [14], residing in rural areas [15] have been identified as having a greater odds of developing loneliness in later life. Additional factors including physical inactivity, living

alone, retirement, and loss of companions or friends were associated with a higher risk of loneliness among older individuals [16-19]. In the last decades, an increasing volume of studies concluded that more engagement in activities and good relationships with social contact could benefit older adults and reduce loneliness [14, 20]. With the development of determinants of health, the significance and relevancy of social resources, including social capital, in maintaining the well-being and health of older people have been emphasized [13]. Social capital refers to a concept that describing social relations at individual or community levels that can be obtained through mutual interaction within a neighborhood, community, or society [10, 21, 22]. Growing studies have documented that social capital is related to the health and wellness of older people [23, 24], and many studies including from China have examined the association of social capital and loneliness [10, 14, 15].

However, existing researches exploring the relationship between social capital and loneliness among older populations assessed social capital with varied factors and obtained mixed findings. For example, a study in Western Finland found that low levels of social capital that measured by social contacts, participation, trust, and sense of belonging, while only lower trust was associated with loneliness among the oldest age group [10]. In a study conducted in 14 European countries, social capital, including regular social participation was linked to a decreased experience of loneliness and a reduction in the impact of loneliness in low-income families [14]. A prior study found that social capital (bonding social capital and bridging social capital) was associated with loneliness among rural widowed older people in China [15]. Similar conclusions have also been drawn from an intervention study in Spain, which indicated that the risk of experiencing loneliness decreased with an increase in social capital regarding social participation and social support [25]. Hence, to better explore and examine the linkage of social capital and loneliness among older adults, an accepted and defined method to measure social capital is highly needed.

As mentioned above, loneliness in later life was exposed to a myriad of factors, including a low level of social capital [11, 13-15, 19]. Despite that these factors had been well explored in many studies, whether these factors and social capital interact to impact loneliness remains to be confirmed by further studies. Collectively, a more accurate and comprehensive analytical approach exploring the interaction of different variables that prompt health outcomes in the older population is proposed. [26]. Exploring multiple interactions is fundamental in obtaining the most targeted and effectual interventions to reduce the onset of loneliness and these findings may be valuable in developing appropriate intervention measures or programs to prevent the incidence of loneliness.

Given this, the present study aims to examine the relationship between social capital (six dimensions) and loneliness and investigates whether there is an interactive effect of social capital and demographic factors and health-related factors on loneliness among community-dwelling older adults in Anhui Province, China.

Methods

Study design and data collection

This cross-sectional study was carried out from July to September 2017 in Anhui Province, which is located in the east of China (Additional file 1). As of December 2019, the population residing in Anhui is around 71.19 million, among which people aged 60 years and above had accounted for about 18.41% [27]. In brief, 1935 participants were interviewed, while 1810 were finally included for analysis. A detailed description of the study design and data collection has been fully described previously [28, 29].

Measures

Measurement of loneliness

Loneliness, the main outcome variable, was assessed by a single question in line with previous studies [10, 11, 30]. Participants were asked the question “Do you have the feeling of loneliness?” with answers ranging from “often, sometimes, and never”. We dichotomized this variable into two categories (yes, no) by combining “often and sometimes” as the presence of loneliness, while “never” as the absence of loneliness, which was in line with other studies [10, 11, 31]. Details about the measurement of loneliness please refer to Additional file 2.

Measurement of social capital

Social capital, the main independent variable, included six dimensions: social participation, social connection, social support, trust, cohesion, and reciprocity. Social capital was measured with a five-point Likert scale and 22 items. More information about the measurement of social capital has been published previously [28, 29] and can also be found in Additional file 2.

Subjects were asked to respond to each item with answers ranging from 1 = “never”, 2 = “seldom”, 3 = “usually”, 4 = “often”, and 5 = “more often”, where each number indicated the associated score accordingly. The score ranges for each dimension were obtained by summarizing answers to each item as follows; social participation (4-20), social support (4-20), social connection (3-15), trust (3-15), cohesion (5-25), reciprocity (3-15). A higher score denoted a better social capital. During analysis, binary variables (high and low level) were generated by dichotomizing each dimension into two categories according to their relative median value, which was in accordance with other studies [32, 33].

Measurement of other variables

Other variables including age, sex, body mass index (BMI, kg/m²), living status, residence (urban, rural), marital status, education level, functional ability, smoking status, drinking status, and the number of diseases were collected as well. In this study, we measured the functional ability of the participants using the work of Lawton and Brody [28, 34], meanwhile, the number of diseases was collected based on the physical health section of the Older Americans Resources and Services (ORAS) [35], which were ranked into four categories: 0, 1, 2, >2 [36]. More detailed information about these variables has been described in our previous papers [28, 29].

Statistical Analysis

First, to compare the difference between two groups (lonely versus not lonely), a Chi-square test was used with variables expressed as numbers and percentages.

Second, we applied logistic regression models to calculate the odds ratio (OR) of developing loneliness and its 95% confidence interval (95% CI). Based on the literature review [23, 37], related variables such as age, sex, BMI, living status, residence, marital status, education level, functional ability, number of diseases, smoking status, and drinking status, were adjusted in all regression models.

Third, we used a classification and regression tree (CART) model to explore the interactive relationship between some indicators of social capital and demographic factors and health-related factors associated with loneliness. This nonparametric model has been used to explore the different interactions among various variables in public health research [38, 39]. Several homogenous subgroups related to the development of loneliness were generated using this model. The variables included in this model were based on the unadjusted results from the logistic regression model. The tree model growing method has been described previously [28].

There no missing data in this study and all analyses were processed in the SPSS 23.0 statistical software (SPSS Inc., Chicago, IL, USA). The P-value that smaller than 0.05 indicated the statistical significance level. We used the ArcMap software (version 10.6) to make the map to show the location of study area. The map is available on request from the corresponding author.

Results

Results of descriptive analysis

The summary statistics of the general characteristics of the respondents are presented in Table 1. The study included 1810 participants (71.20 ± 7.51 years). There were differences in age, sex, BMI, living status, residence, marital status, education level, functional ability, number of diseases, drinking status, and social capital dimensions between the two groups (lonely versus not lonely). About 38.7% (700 out of 1,810) of the participants reported a sense of loneliness in this study, of which 39.7% (278/700) were between 60-69 years, 62.3% (436/700) were females, 48.7% (341/700) of subjects' BMI was between 18.5-22.9, 81.0% (567/700) were living with others, about 65.0% (455/700) lived in the rural area, 67.9% (475/700) belonged to married or cohabited group, and 82.0% (574/700) attended primary school and below. Meanwhile, above half the lonely participants 64.9% (454/700) reported limited functional ability. 38.1% (267/700) of them suffered at least one disease. Most of the participants reporting loneliness were non-smoker (80.3%) and non-drinker (84.9%).

Table 1 General characteristics of the respondents (N = 1810)

Results of logistic regression analysis

Table 2 shows the results of the logistic regression models. In model 1 (unadjusted), subjects aged ≥ 80 years (OR = 2.32; 95% CI: 1.76-3.06), who were females (OR = 1.38; 95% CI: 1.14-1.68), living alone (OR =

2.13; 95% CI: 1.62-2.80), residing in rural areas (OR = 1.86; 95% CI: 1.53-2.26), were single (OR = 2.40; 95% CI: 1.92-3.00), with low education level (OR = 2.58; 95% CI: 1.86-3.57), and limited functional ability (OR = 4.48; 95% CI: 3.66-5.48) were found to have a higher risk for experiencing loneliness. The odds for developing loneliness increased with the number of diseases and were the highest among respondents reporting ≥ 2 diseases (OR = 3.70; 95% CI: 2.66-5.15). Individuals with lower social participation (OR = 1.86; 95% CI: 1.54-2.26), social support (OR = 1.23; 95% CI: 1.02-1.49), social connection (OR = 2.36; 95% CI: 1.92-2.90), trust (OR = 1.81; 95% CI: 1.49 – 2.19), cohesion (OR = 1.82; 95% CI: 1.50-2.20), and reciprocity (OR = 2.03; 95% CI: 1.68-2.46) were at a greater risk of developing loneliness.

After adjusting all covariates in model 2, respondents with lower social participation (AOR = 1.38; 95% CI: 1.10-1.74), social connection (AOR = 1.51; 95% CI: 1.18-1.93), and reciprocity (AOR = 1.47; 95% CI: 1.13-1.90) were more likely to suffer from loneliness.

Table 2 Logistic regression analysis examining the association between social capital and loneliness (N = 1810)

Results of classification and regression tree model

The results of the CART model are shown in Fig. 1. The development of loneliness was associated with functional ability, marital status, trust, social connection, number of diseases, and social participation. The interactive effect among social capital dimensions and diverse variables were also observed.

Our analyses indicated that functional ability was the most determinant factor related to loneliness. The sample was therefore split into subsets based on this factor. Participants with a robust functional ability (Node 1), married or cohabited (Node 4), and reported no or less number of diseases (Node 9), were least likely to experience loneliness. However, participants who were single (Node 3) and with a low level of social connection (Node 7) were more prone to experience loneliness as compared to those with a higher level of social connection (Node 8).

Those who reported limitations in functional ability (Node 2) and had a lower level of trust (Node 6) and social participation (Node 14) were the most likely to experience loneliness. Meanwhile, subjects with a higher level of trust (Node 5) and were single (Node 11), inclined to suffer from loneliness than those who were married or cohabited (Node 12).

Discussion

In this study, we examined the association between social capital and loneliness among the older population and explored the interactive effect of social capital and demographic factors and health-related factors on loneliness in Anhui Province, China. Our results confirmed the association of social capital and loneliness and the combined influence of social capital and some other factors on the development of loneliness in later life, suggesting the relevance of social capital in preserving the emotional health of older people.

The results of the current study demonstrated the relationship between loneliness and social capital. Specifically, older people who lacked social capital concerning social participation, social connection, and reciprocity were more likely to experience loneliness, which was echoed by findings from previous studies [10, 20, 25, 41]. Similar to our results, Nyqvist et al. [10] found that infrequent social connection with neighbours had an increased likelihood of being lonely among older people aged 65-80 years in Western Finland. Results from the fifth wave of the Survey of Health, Ageing and Retirement in Europe (SHARE) also concluded that more social participation was a protective factor for loneliness and mitigated the impact of unfavorable socioeconomic status among older people [14]. Some studies also suggested that forming and building reciprocal connections or relationships with others could alleviate the impact of some mental health issues, including loneliness [20, 42].

According to the results of our adjusted logistic regression analysis, a non-significant association was observed between a lower level of social capital regarding social support, trust, and cohesion, and loneliness. Different from our results, a prior study revealed that less trust and a weak sense of belonging to the community were significantly linked to a higher risk for developing loneliness [10]. Meanwhile, a study also found that an increase of trust at the community-level also contributed to a reduction in loneliness among community-dwelling older people [43]. Also, a previous study showed that insufficient social support is significantly related to the onset of loneliness [44-46]. There are two possible reasons for this inconsistency. First, this may be due to different measurements that were used to assess social capital in the current study and earlier works. For example, in this paper, social support was measured by asking participants "how often can they get mental or material support when they are in need". But Chen et al. [44] measured social support by asking surveyed subjects the quantity of social support offered by friends. The lack of common and widely used social capital measurement tool has been recognized as a repeated issue, which contributes to some inconclusive results in this research field. Another reason may be that our adjusted model included several variables such as demographic and health-related as covariates, which were associated with social capital and loneliness. As a result, the association between social support, trust, and cohesion with loneliness was attenuated or even became non-significant. However, more research is still warranted to further verify our results in the future.

Previous studies demonstrated that limitation in functional ability, being single, and multimorbidity was related to a higher risk for developing loneliness among older people [19, 47]. In the present study, most importantly, an interacting relationship between social capital and functional ability, marital status, and the number of diseases was observed. That is, older people who reported limitations in functional ability and had a lower level of trust and social participation were the most likely to experience loneliness. This indicates that trust and social participation as a social capital dimension may be of relevance in loneliness prevention [10, 25]. Besides, single older people, who had less social connection were more prone to loneliness as compared to those who had a higher-level social connection, which further highlights the protective role of social connection in preventing the incidence of loneliness [48]. Interestingly, the significance of trust was not found in the adjusted logistic regression model, however, it was observed in the CART model. We suggest that the role of trust may depend on the appearance of other social capital dimensions and variables, which emphasizes the effectiveness of the CART model in

examining the complex interactions among multiple variables that may be overlooked in the conventional analytical approach [26]. The importance of this finding lies in adding a scientific explanation of using the CART model to help examine the association between social capital and loneliness while revealing how social capital interacts with other factors and produces an effect on the development of loneliness. Additionally, the CART model was used as a predictive model to estimate the subsets of older people that are more likely to become lonely.

Regarding the assessment of loneliness, different tools and methods were employed to identify loneliness as well. For instance, the UCLA loneliness scale was the most common and validated scale used to collect loneliness data in previous research [49, 50]. However, a single item by asking participants how often they feel lonely, which potential responses included often, sometimes, and never, etc., also had been validated a good predictive validity in many studies [10, 11, 31]. Moreover, the use of a single question to assess loneliness has been proved to have relevant edges such as succinctness, easily understood, and well accepted by subjects [30]. Besides, in the process of data analysis, many studies dichotomized the status of loneliness into two groups (lonely, not lonely) by combining often and sometimes as the presence of loneliness, while rarely or never as the absence of loneliness [10, 11, 31, 37]. Despite such dichotomization could result in some loss of information about the outcome, to obtain statistical power during statistical process and make comparisons with other research. Likewise, we also categorized loneliness into two kinds in this paper.

Loneliness is a subjective and negative feeling that may be the consequence of dissatisfaction with an individual's social relations and unsatisfied social needs [8]. Therefore, by looking at these findings from the present study, some relevant suggestions for the interventions of social capital to reduce the level and alleviate the impact of loneliness among community-dwelling older adults can be offered. First, we suggest older people should engage in more social activities, for instance, older adults are encouraged to participate the formal/informal groups (political/non-political parties, elections, hobby groups, etc.), voluntary activities, and services (health lecture, culture, and physical education activities) within the community, which provide opportunities for them to meet the social needs, share life experiences and exchange interests. Second, the children, relatives, friends, and neighbors are encouraged to have more frequent interactions and communications with the older people and care about them, which is beneficial to maintain a good social relationship, promote and improve the quality of social connections, in turn, reduce the chance of developing loneliness among older adults. Third, to have an ideal reciprocal relationship, programs, and activities designed to cultivate and escalate the willingness to actively provide help to each other, including their relatives, friends or neighbors, and strangers, should be introduced. Lastly, to make good use of the role of social capital in the prevention of loneliness and maintain good emotional health, more attention should be paid to those who reported limitations in functional ability, had a lower level of trust and insufficient social participation, and single older people who had a less social connection.

This study has several limitations. First, since it was a cross-sectional study, which limited to conclude the causal relations between social capital and loneliness. Future studies using a longitudinal or

randomized control trial design are warranted. Second, data in our study were based on self-report and might be subject to a recall or reporting bias. Nevertheless, to improve the data accuracy, we formulated clear and precise questions and carried out a pilot study before investigation. During data collection, forward or backward recall techniques were also used. Third, since our study was only conducted in Anhui Province, this might constrain the generalization of our findings to other regions or countries. Future studies that include extensive sites and larger samples are needed. Fourth, in the present study, we did not focus on the concept of emotional and social loneliness yet took loneliness as an umbrella concept instead. Besides, the variables included in this study were not broad enough; some other variables such as depression and other mental health data were not well considered. More attention should be given to these mental health factors in the future.

Despite the limitations, this study also has some strength. Findings from this study are reliable because the sample was representative and had high response rates from participants. We also used a validated and standardized social capital scale, which may facilitate the development of social capital theory. Moreover, to our knowledge, this study is the first using the CART model allowed for further exploring multi-variable interactions and yielded a straightforward and visible tree, which is suitable in devising more specific and accurate strategies to counteract the impact of loneliness.

Conclusions

In summary, we observed an association between social capital and loneliness among the older population. Specifically, social capital in terms of social participation, social connection, trust, and reciprocity may be significant in designing intervention programs and measures to prevent and reduce the incidence of loneliness.

Declarations

Ethics approval and consent to participate

In the present study, written informed consent with a signature was obtained from all participants. All participants were medically and ethically capable of providing their consent. For those who could not write, a fingerprint was replaced instead after the participants have fully understood the information mentioned in the informed consent and agreed to take part in this study as a study subject. The procedure was approved and Ethical approval for this study was obtained from the Biomedical Ethics Committee, Anhui Medical University (No. 20150297)

Consent for publication

Not applicable.

Availability of data and materials

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the National Natural Science Foundation of China (No.71573002 and 71673002). This organization had no role in the research design, data collection, data analysis, manuscript writing, and submission.

Authors' contributions

XQ and ZH conceived and designed the study. ZB undertook data collection, performed statistical analyses, and drafted the manuscript. ZW and TS undertook data collection. XQ and ZH revised the manuscript. All authors checked, interpreted results, and approved the final version.

Acknowledgments

The authors wish to express sincere appreciation to all who assisted and engaged in our study.

Authors' information

¹Department of Epidemiology and Biostatistics, School of Public Health, Anhui Medical University, Hefei, 230032, China. ²Department of Health Services Management, School of Health Services Management, Anhui Medical University, Hefei, 230032, China.

Abbreviations

CLHLS: The Chinese Longitudinal Healthy Longevity Survey;

BMI: Body Mass Index;

ADL: The Activities of Daily Living;

IADL: Instrumental Activities of Daily Living;

ORAS: The Older Americans Resources and Services;

CHD: Coronary Heart Disease;

OR: Odds Ratio;

95% CI: 95% confidence interval;

References

1. Partridge L, Deelen J, Slagboom PE: **Facing up to the global challenges of ageing.** *Nature* 2018, **561**(7721):45-56.
2. Beard JR, Officer A, de Carvalho IA, Sadana R, Pot AM, Michel J-P, Lloyd-Sherlock P, Epping-Jordan JE, Peeters GME, Mahanani WR *et al.*: **The World report on ageing and health: a policy framework for healthy ageing.** *The Lancet* 2016, **387**(10033):2145-2154.
3. Hawkley LC, Cacioppo JT: **Loneliness matters: a theoretical and empirical review of consequences and mechanisms.** *Annals of behavioral medicine : a publication of the Society of Behavioral Medicine* 2010, **40**(2):218-227.
4. Luhmann M, Hawkley LC: **Age differences in loneliness from late adolescence to oldest old age.** *Developmental psychology* 2016, **52**(6):943-959.
5. Lim KK, Chan A: **Association of loneliness and healthcare utilization among older adults in Singapore.** *Geriatrics & gerontology international* 2017, **17**(11):1789-1798.
6. Valtorta NK, Kanaan M, Gilbody S, Ronzi S, Hanratty B: **Loneliness and social isolation as risk factors for coronary heart disease and stroke: systematic review and meta-analysis of longitudinal observational studies.** *Heart* 2016, **102**(13):1009-1016.
7. Holwerda TJ, Deeg DJ, Beekman AT, van Tilburg TG, Stek ML, Jonker C, Schoevers RA: **Feelings of loneliness, but not social isolation, predict dementia onset: results from the Amsterdam Study of the Elderly (AMSTEL).** *J Neurol Neurosurg Psychiatry* 2014, **85**(2):135-142.
8. Arslantas H, Adana F, Abacigil Ergin F, Kayar D, Acar G: **Loneliness in Elderly People, Associated Factors and Its Correlation with Quality of Life: A Field Study from Western Turkey.** *Iranian journal of public health* 2015, **44**(1):43-50.
9. Luo Y, Hawkley LC, Waite LJ, Cacioppo JT: **Loneliness, health, and mortality in old age: a national longitudinal study.** *Social science & medicine (1982)* 2012, **74**(6):907-914.
10. Nyqvist F, Victor CR, Forsman AK, Cattan M: **The association between social capital and loneliness in different age groups: a population-based study in Western Finland.** *BMC public health* 2016, **16**:542.
11. Zhou Z, Mao F, Zhang W, Towne DS, Wang P, Fang Y: **The Association Between Loneliness and Cognitive Impairment among Older Men and Women in China: A Nationwide Longitudinal Study.** *International journal of environmental research and public health* 2019, **16**(16).
12. Valtorta N, Hanratty B: **Loneliness, isolation and the health of older adults: do we need a new research agenda?** *Journal of the Royal Society of Medicine* 2012, **105**(12):518-522.
13. Chen Y, Hicks A, While AE: **Loneliness and social support of older people in China: a systematic literature review.** *Health & social care in the community* 2014, **22**(2):113-123.
14. Niedzwiedz CL, Richardson EA, Tunstall H, Shortt NK, Mitchell RJ, Pearce JR: **The relationship between wealth and loneliness among older people across Europe: Is social participation protective?**

Preventive medicine 2016, **91**:24-31.

15. Jiang D, Hou Y, Hao J, Zhou J, Jiang J, Wang Q: **Association between Personal Social Capital and Loneliness among Widowed Older People.** *International journal of environmental research and public health* 2020, **17**(16).
16. Findlay RA: **Interventions to reduce social isolation amongst older people: where is the evidence?** *Ageing and Society* 2003, **23**(5):647-658.
17. Pinquart M, Sorensen S: **Influences on loneliness in older adults: A meta-analysis.** *Basic and applied social psychology* 2001, **23**(4):245-266.
18. Hawkey LC, Hughes ME, Waite LJ, Masi CM, Thisted RA, Cacioppo JT: **From social structural factors to perceptions of relationship quality and loneliness: the Chicago health, aging, and social relations study.** *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences* 2008, **63**(6):S375-S384.
19. Cohen-Mansfield J, Hazan H, Lerman Y, Shalom V: **Correlates and predictors of loneliness in older-adults: A review of quantitative results informed by qualitative insights.** *International psychogeriatrics* 2016, **28**(4):557-576.
20. Murayama Y, Murayama H, Hasebe M, Yamaguchi J, Fujiwara Y: **The impact of intergenerational programs on social capital in Japan: a randomized population-based cross-sectional study.** *BMC public health* 2019, **19**(1):156.
21. Putnam RD: **Bowling alone: The collapse and revival of American community:** Simon and schuster; 2000.
22. Coll-Planas L, Nyqvist F, Puig T, Urrútia G, Solà I, Monteserín R: **Social capital interventions targeting older people and their impact on health: a systematic review.** *Journal of epidemiology and community health* 2017, **71**(7):663-672.
23. Tomioka K, Kurumatani N, Hosoi H: **Age and gender differences in the association between social participation and instrumental activities of daily living among community-dwelling elderly.** *BMC geriatrics* 2017, **17**(1):99.
24. Imamura H, Hamano T, Michikawa T, Takeda-Imai F, Nakamura T, Takebayashi T, Nishiwaki Y: **Relationships of Community and Individual Level Social Capital with Activities of Daily Living and Death by Gender.** *International journal of environmental research and public health* 2016, **13**(9):860.
25. Coll-Planas L, Del Valle Gómez G, Bonilla P, Masat T, Puig T, Monteserín R: **Promoting social capital to alleviate loneliness and improve health among older people in Spain.** *Health & social care in the community* 2017, **25**(1):145-157.
26. Jung SY, Vitolins MZ, Fenton J, Frazier-Wood AC, Hursting SD, Chang S: **Risk profiles for weight gain among postmenopausal women: a classification and regression tree analysis approach.** *PloS one* 2015, **10**(3):e0121430-e0121430.
27. **Facts about Anhui** [<http://english.ah.gov.cn/AboutAnhui/StatisticalCommunique/8321041.html>]
28. Bai Z, Wang Z, Shao T, Qin X, Hu Z: **Relationship between Individual Social Capital and Functional Ability among Older People in Anhui Province, China.** *International journal of environmental research*

- and public health* 2020, **17**(8): 2775.
29. Bai Z, Xu Z, Xu X, Qin X, Hu W, Hu Z: **Association between social capital and depression among older people: evidence from Anhui Province, China.** *BMC public health* 2020, **20**(1):1560.
 30. Bartlett MY, Arpin SN: **Gratitude and Loneliness: Enhancing Health and Well-Being in Older Adults.** *Research on aging* 2019, **41**(8):772-793.
 31. Routasalo PE, Savikko N, Tilvis RS, Strandberg TE, Pitkälä KH: **Social contacts and their relationship to loneliness among aged people—a population-based study.** *Gerontology* 2006, **52**(3):181-187.
 32. Maselko J, Hughes C, Cheney R: **Religious social capital: Its measurement and utility in the study of the social determinants of health.** *Social Science & Medicine* 2011, **73**(5):759-767.
 33. Ito T, Okuyama K, Abe T, Takeda M, Hamano T, Nakano K, Nabika T: **Relationship between Individual Social Capital and Cognitive Function among Older Adults by Gender: A Cross-Sectional Study.** *International journal of environmental research and public health* 2019, **16**(12).
 34. Lawton MP, Brody EM: **Assessment of older people: self-maintaining and instrumental activities of daily living.** *The Gerontologist* 1969, **9**(3_Part_1):179-186.
 35. George LK, Fillenbaum GG: **OARS Methodology.** *Journal of the American Geriatrics Society* 1985, **33**(9):607-615.
 36. Domènech-Abella J, Mundó J, Lara E, Moneta MV, Haro JM, Olaya B: **The role of socio-economic status and neighborhood social capital on loneliness among older adults: evidence from the Sant Boi Aging Study.** *Social psychiatry and psychiatric epidemiology* 2017, **52**(10):1237-1246.
 37. Nyqvist F, Cattan M, Andersson L, Forsman AK, Gustafson Y: **Social capital and loneliness among the very old living at home and in institutional settings: a comparative study.** *J Aging Health* 2013, **25**(6):1013-1035.
 38. Farrell RT, Gamelli RL, Aleem RF, Sinacore JM: **The Relationship of Body Mass Index and Functional Outcomes in Patients With Acute Burns.** *Journal of Burn Care & Research* 2008, **29**(1):102-108.
 39. BeLue R, Francis LA, Rollins B, Colaco B: **One Size Does Not Fit All: Identifying Risk Profiles for Overweight in Adolescent Population Subsets.** *Journal of Adolescent Health* 2009, **45**(5):517-524.
 40. Nakabe T, Sasaki N, Uematsu H, Kunisawa S, Wimo A, Imanaka Y: **Classification tree model of the personal economic burden of dementia care by related factors of both people with dementia and caregivers in Japan: a cross-sectional online survey.** *BMJ open* 2019, **9**(7):e026733-e026733.
 41. Child ST, Lawton L: **Loneliness and social isolation among young and late middle-age adults: Associations with personal networks and social participation.** *Aging & mental health* 2019, **23**(2):196-204.
 42. Salehi A, Ehrlich C, Kendall E, Sav A: **Bonding and bridging social capital in the recovery of severe mental illness: a synthesis of qualitative research.** *Journal of mental health (Abingdon, England)* 2019, **28**(3):331-339.
 43. Yang J, Moorman SM: **Beyond the Individual: Evidence Linking Neighborhood Trust and Social Isolation Among Community-Dwelling Older Adults.** *International journal of aging & human*

development 2019:91415019871201.

44. Chen L, Alston M, Guo W: **The influence of social support on loneliness and depression among older elderly people in China: Coping styles as mediators.** *Journal of community psychology* 2019, **47**(5):1235-1245.
45. Bai X, Yang S, Knapp M: **Sources and directions of social support and life satisfaction among solitary Chinese older adults in Hong Kong: the mediating role of sense of loneliness.** *Clinical interventions in aging* 2018, **13**:63-71.
46. Chen Y, Hicks A, While AE: **Loneliness and social support of older people living alone in a county of Shanghai, China.** *Health & social care in the community* 2014, **22**(4):429-438.
47. Domènech-Abella J, Mundó J, Haro JM, Rubio-Valera M: **Anxiety, depression, loneliness and social network in the elderly: Longitudinal associations from The Irish Longitudinal Study on Ageing (TILDA).** *Journal of affective disorders* 2019, **246**:82-88.
48. Cacioppo JT, Fowler JH, Christakis NA: **Alone in the crowd: the structure and spread of loneliness in a large social network.** *Journal of personality and social psychology* 2009, **97**(6):977.
49. Wang J, Lloyd-Evans B, Marston L, Mann F, Ma R, Johnson S: **Loneliness as a predictor of outcomes in mental disorders among people who have experienced a mental health crisis: a 4-month prospective study.** *BMC psychiatry* 2020, **20**(1):249.
50. Wang G, Zhang X, Wang K, Li Y, Shen Q, Ge X, Hang W: **Loneliness among the rural older people in Anhui, China: prevalence and associated factors.** *International journal of geriatric psychiatry* 2011, **26**(11):1162-1168.

Tables

Table 1 General characteristics of the respondents (N = 1810)

Variables	Loneliness		χ^2	p-value
	No N = 1110	Yes N = 700		
Age (years)			37.365	<0.001
60-69	545(49.1%)	278(39.7%)		
70-79	435(39.2%)	268(38.3%)		
≥ 80	130(11.7%)	154(22.0%)		
Sex			10.881	0.001
Male	506(45.6%)	264(37.7%)		
Female	604(54.4%)	436(62.3%)		
BMI (kg/m²)			8.722	0.033
<18.5	108(9.7%)	81(11.6%)		
18.5-22.9	484(43.6%)	341(48.7%)		
23.0-27.4	417(37.6%)	227(32.4%)		
≥ 27.5	101(9.1%)	51(7.3%)		
Living status			30.518	<0.001
Not living alone	1000(90.1%)	567(81.0%)		
Living alone	110(9.9%)	133(19.0%)		
Residence				<0.001
Urban	556(50.1%)	245(35.0%)		
Rural	554(49.9%)	455(65.0%)		
Marital status			60.266	<0.001
Married/cohabited	927(83.5%)	475(67.9%)		
Single	183(16.5%)	225(32.1%)		
Education level			63.648	<0.001
Primary school and below	717(64.6%)	574(82.0%)		
Junior school	219(19.7%)	72(10.3%)		
High school and above	174(15.7%)	54(7.7%)		
Functional ability			222.842	<0.001
Robust	786(70.8%)	246(35.1%)		
Limited	324(29.2%)	454(64.9%)		
Number of diseases			67.128	<0.001
0	378(34.1%)	152(21.7%)		
1	447(40.3%)	267(38.1%)		
2	197(17.7%)	150(21.4%)		
>2	88(7.9%)	131(18.7%)		
Smoking status			3.848	0.146
Non-smoker	850(76.6%)	562(80.3%)		
Former smoker	62(5.6%)	37(5.3%)		
Smoker	198(17.8%)	101(14.4%)		
Drinking status			7.737	0.021
Non-drinker	890(80.2%)	594(84.9%)		
Former drinker	43(3.9%)	27(3.9%)		
Drinker	177(15.9%)	79(11.3%)		
Social participation			40.766	<0.001
High	705(63.5%)	338(48.3%)		
Low	405(36.5%)	362(51.7%)		
Social support			4.670	0.031

High	578(52.1%)	328(46.9%)		
Low	532(47.9%)	372(53.1%)		
Social connection			67.738	<0.001
High	862(77.7%)	417(59.6%)		
Low	248(22.3%)	283(40.4%)		
Trust			37.188	<0.001
High	690(62.2%)	333(47.6%)		
Low	420(37.8%)	367(52.4%)		
Cohesion			37.064	<0.001
High	723(65.1%)	355(50.7%)		
Low	387(34.9%)	345(49.3%)		
Reciprocity			53.083	<0.001
High	675(60.8%)	303(43.3%)		
Low	435(39.2%)	397(56.7%)		

Table 2 Logistic regression analysis examining the association between social capital and loneliness (N = 1810)

Variables	Model 1	Model 2
	OR (95% CI)	AOR (95% CI)
Age (years)		
60-69	1	1
70-79	1.21 (0.98-1.49)	0.96 (0.75-1.22)
≥ 80	2.32 (1.76-3.06) ***	1.25 (0.90-1.74)
Sex		
Male	1	1
Female	1.38 (1.14-1.68) **	1.05 (0.79-1.38)
Body mass index (kg/m²)		
Normal weight	1	1
Underweight	1.06 (0.77-1.47)	0.81 (0.56-1.17)
Overweight	0.77 (0.62-0.96) *	0.88 (0.69-1.12)
Obese	0.72 (0.50-1.03)	0.68 (0.45-1.03)
Living status		
Not living alone	1	1
Living alone	2.13 (1.62-2.80) ***	1.39 (0.96-2.00)
Residence		
Urban	1	1
Rural	1.86 (1.53-2.26) ***	1.26 (1.00-1.60)
Marital status		
Married/cohabited	1	1
Single	2.40 (1.92-3.00) ***	1.66 (1.22-2.25) **
Education level		
High school and above	1	1
Primary school and below	2.58 (1.86-3.57) ***	1.10 (0.76-1.60)
Junior school	1.06 (0.71-1.59)	0.80 (0.51-1.24)
Functional ability		
Robust	1	1
Limited	4.48 (3.66-5.48) ***	3.10 (2.45-3.93) ***
Number of diseases		
0	1	1
1	1.49 (1.17-1.89) **	1.30 (0.99-1.70)
2	1.89 (1.43-2.51) ***	1.48 (1.08-2.03) *
>2	3.70 (2.66-5.15) ***	2.38 (1.64-3.45) ***
Smoking status		
Non-smoking	1	1
Former smoking	0.90 (0.59-1.37)	1.23 (0.72-2.11)
Smoking	0.77 (0.59-1.00)	1.14 (0.80-1.62)
Drinking status		
Non-drinking	1	1
Former drinking	0.94 (0.58-1.54)	0.81 (0.44-1.49)
Drinking	0.67 (0.50-0.89) *	0.80 (0.56-1.13)
Social participation		
High	1	1
Low	1.86 (1.54-2.26) ***	1.38 (1.10-1.74) **
Social support		
High	1	1

Low	1.23 (1.02-1.49) *	0.79 (0.62-1.01)
Social connection		
High	1	1
Low	2.36 (1.92-2.90) ***	1.51 (1.18-1.93) **
Trust		
High	1	1
Low	1.81 (1.49-2.19) ***	1.21 (0.93-1.56)
Cohesion		
High	1	1
Low	1.82 (1.50-2.20) ***	1.15 (0.88-1.49)
Reciprocity		
High	1	1
Low	2.03 (1.68-2.46) ***	1.47 (1.13-1.90) **

Note: Model 2 adjusted age, sex, body mass index, living status, residence, marital status, education level, functional ability, number of diseases, smoking, and drinking status.

OR: odds ratio, 95% CI: 95% confidence interval

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

Figures




Figure 1

CART model analysis examining the interactive association between social capital and loneliness. (This classification and regression tree shows the variables related to the development of loneliness and the

likelihood of experiencing loneliness in the node boxes, which also displays the interactive relationship between social capital and demographic factors and health-related factors associated with loneliness.)

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile2.docx](#)
- [Additionalfile1.docx](#)