

Dissemination of Carbapenem-resistance and Plasmids-encoding Carbapenemases in Gram-negative Bacteria Isolated from India

Prasanth Manohar

VIT University

Sebastian Leptihn

ZJE Institute, Zhejiang University

Bruno S. Lopes (✉ bruno.lopes@abdn.ac.uk)

University of Aberdeen

Nachimuthu Ramesh (✉ drpnramesh@gmail.com)

VIT University <https://orcid.org/0000-0002-5546-1117>

Research article

Keywords: Carbapenem, Gram-negative bacteria, plasmid incompatibility grouping, conjugative plasmid, carbapenem-resistance genes.

Posted Date: December 18th, 2020

DOI: <https://doi.org/10.21203/rs.2.15952/v2>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Dissemination of Carbapenem-resistance and Plasmids-encoding Carbapenemases in Gram-negative Bacteria Isolated from India

Prasanth Manohar^{1#}, Sebastian Leptihn^{2,3}, Bruno S. Lopes^{4*}, Nachimuthu Ramesh^{1*}

¹*Antibiotic Resistance and Phage Therapy Laboratory, Department of Biomedical Sciences, School of Bioscience and Technology, Vellore Institute of Technology (VIT), Vellore- 632014, Tamil Nadu, India*

²*Zhejiang University-University of Edinburgh (ZJU-UoE) Institute, Zhejiang University, Haining, Zhejiang 314400, China and Department of Infectious Diseases, Sir Run Run Shaw Hospital, Zhejiang University School of Medicine, Hangzhou, China*

³*University of Edinburgh Medical School, Biomedical Sciences, College of Medicine & Veterinary Medicine, The University of Edinburgh, 1 George Square, Edinburgh, EH8 9JZ, United Kingdom*

⁴*School of Medicine, Medical Sciences and Nutrition, Medical Microbiology, University of Aberdeen, Aberdeen, AB25 2ZD UK*

#Present address: Zhejiang University-University of Edinburgh (ZJU-UoE) Institute, Zhejiang University International Campus, Haining, Zhejiang 314400, P.R.China

***Correspondence**

Dr. N. Ramesh, Assistant Professor, Antibiotic Resistance and Phage Therapy Laboratory, School of Biosciences and Technology, VIT, Vellore-632014, India.

Phone: +91 9842660673, E. mail: drpnramesh@gmail.com

Dr. B. S. Lopes, School of Medicine, Medical Sciences and Nutrition, Medical Microbiology, University of Aberdeen, AB25 2ZD UK.

E. mail: bruno.lopes@abdn.ac.uk

Abstract

Carbapenem resistance in Gram-negative bacteria is an ongoing public-health problem of global dimensions leaving very few treatment options for severely infected patients. This study focuses on the dissemination of plasmid-borne carbapenemase genes in Gram-negative bacteria in Tamil Nadu, India. A total of 151 non-repetitive isolates belonging to 11 genera were collected from a diagnostic center in Tamil Nadu. *E. coli* (n=57) isolates were classified as, Enteropathogenic (n=12), Enteroinvasive (n=9),

Enterohemorrhagic (n=8), Enterotoxigenic (n=3), Enteroinvasive (n=1) and unclassified *E. coli* (n=24). Of the 45 *Klebsiella* species, 14 were K1 whereas 11 were K2 serotype and in 20 *Klebsiella* serotype could not be determined. Other isolates (n=49) consisted of *P. aeruginosa*, *S. typhi*, *E. cloacae*, *A. baumannii*, *S. marcescens*, *A. xylosoxidans*, *P. mirabilis* and *E. meningoseptica*. Of the 151 isolates, 71% (n=107) and 68% (n=103) were found to be resistant to meropenem and imipenem respectively. The most prevalent beta-lactamase gene was *bla*_{NDM-1} (21%, 12/57) followed by *bla*_{OXA-181} (16%, 9/57), *bla*_{GES-9} (n=8), *bla*_{OXA-23} (n=7), *bla*_{IMP-1} (n=3), *bla*_{GES-1} (n=11) and *bla*_{OXA-51} (n=9). The unusual presence of *bla*_{OXA-23} was seen in *E. coli* (n=4), and *bla*_{OXA-23} and *bla*_{OXA-51} (IncA/C) in *K. pneumoniae* (n=3). Plasmid incompatibility (inc/rep) typing results showed that the plasmids carrying resistance genes (n=11) belonged to IncX, IncA/C, IncFIA-FIB and IncFIIA groups. *E. coli* and *K. pneumoniae* were able to transfer plasmid-borne carbapenemase via conjugation. This study highlights the prevalence of carbapenem resistance and the acquisition of plasmid-borne carbapenemase genes in Gram-negative bacteria highlighting the role of plasmid transfer in disseminating resistance.

Keywords: Carbapenem, Gram-negative bacteria, Plasmid incompatibility grouping, Conjugative plasmid, Carbapenem-resistance genes.

1.0 Introduction

Antibiotic resistance is an emerging global health problem due to the injudicious use of antibiotics [1]. It is considered a major clinical and public health problem because of increasing bacterial resistance to most of the available antibiotics including penicillin, cephalosporins, carbapenems, and colistin [1]. World Health Organization (WHO) recently listed carbapenem-resistant *Acinetobacter baumannii*, *Pseudomonas aeruginosa* and Extended Spectrum Beta-Lactamases (ESBL) -producing *Enterobacteriaceae* as pathogens that are of critical importance [2]. Gram-negative bacteria (GNB) especially *Enterobacteriaceae* have developed resistance towards a broad spectrum of antibiotics responsible for significant mortality around the globe [3]. Carbapenems are considered as one of the last resort antibiotics against infections caused by multi-drug resistant GNB [4]. The emergence of carbapenem resistance especially in *Enterobacteriaceae* is a threat to the patients, particularly with complex

infections, immunocompromised conditions and multiple diseases [5]. Because pathogens that are resistant to carbapenems often shows high resistance to other commonly used antibiotics that are often used for treatment, not only the mortality rates are high with increased hospital stay, but also huge medical expenditure placing emotional, economic and financial burden on families especially in resource limited countries [6].

The assessment of the rise in global antibiotic resistance has become very difficult due to the increasing rate of multi-drug resistance shown by pathogens with no proper harmonized surveillance systems in resource limiting countries [7]. Moreover, the co-existence of more than one carbapenem resistance gene with other genes like plasmid mediated AmpC, or plasmid mediated quinolone resistance has resulted in an increased acquisition of resistance among *Enterobacteriaceae* for community as well as hospital acquired infections [8,9]. The carbapenem-hydrolyzing oxacillinas (CHDL) are the major resistance mechanisms to carbapenems in *A. baumannii*. The first report of OXA-23 beta-lactamase in *A. baumannii* was from United Kingdom, in 1985 [10]. Later, OXA-23 was found to confer carbapenem-resistance in *A. nosocomialis* [11] and recently, it was reported in members of *Enterobacteriaceae* family [12-14]. In 1996, the first report of OXA-51 type beta-lactamase was from Argentina and at present, there are more than 150 variants of OXA-51 were reported globally [15]. These intrinsic enzymes in *A. baumannii* are naturally chromosomal-borne but rare cases of plasmid-borne genes are also reported [16]. Earlier, we reported the distribution of carbapenem and colistin resistance, and the role of integrons serving as the horizontal gene transfer agents in disseminating resistance among Gram-negative bacteria [17,18]. In the present study, molecular characterization of Gram-negative bacteria was performed and the role of interspecies plasmid transfer as evolutionary mechanism of carbapenem resistance was determined.

2.0 Materials and methods

2.1 Isolate collection and classification

During January 2015 and December 2016, a total of 151 Gram-negative bacterial isolates were collected from Hi-Tech diagnostic center in Chennai, Tamil Nadu, India. Bacteria were isolated from urine, blood, pus, bronchial secretion, cerebrospinal fluid, pulmonary secretion and bile fluid. The collected isolates were received at the Antibiotic Resistance and Phage Therapy Laboratory, VIT, Vellore, for further analyses. Genomic DNA was extracted from all the isolates using boiling lysis method [18]. Bacterial identification was carried out using VITEK identification system (bioMerieux) and 16S rRNA gene nucleotide sequence analysis using universal primers 27F and 1492R [18]. The PCR products were sequenced and identified to the species level using the BLASTN tool.

2.2 Antibiotic susceptibility testing and Minimal Inhibitory Concentration

Antibiotic resistance profiling was performed using the disk-diffusion method according to CLSI guidelines. The antibiotics used for this study were gentamicin (10 µg), amoxyclav (30 µg), cefotaxime (30 µg), ertapenem (10 µg), amikacin (30 µg), meropenem (10 µg), colistin (10 µg) and ceftazidime (30 µg). Minimum Inhibitory Concentration (MIC) was determined by broth micro-dilution method for meropenem and imipenem as described previously [18] and the results were interpreted according to CLSI guidelines [19].

2.3 Molecular analysis of *E. coli* pathotypes and *Klebsiella* serotypes

The *E. coli* pathotypes namely enteropathogenic *E. coli* (EPEC); enterohemorrhagic *E. coli* (EHEC); enterotoxigenic *E. coli* (ETEC); enteroaggregative *E. coli* (EAEC) and enteroinvasive *E. coli* (EIEC) were identified as described earlier [20]. The *Klebsiella* serotypes K1, K2 and K5 were determined using PCR primers and conditions as described earlier [21].

2.4 Molecular analysis of resistance-related genes

The isolates were screened for the presence of carbapenem resistance genes *bla*_{NDM}, *bla*_{OXA-48-like}, *bla*_{KPC}, *bla*_{IMP} and *bla*_{VIM} [18]. A second multiplex PCR was also performed for *bla*_{DIM}, *bla*_{BIC}, *bla*_{GIM}, *bla*_{SIM} and *bla*_{AIM} [22]. The *bla*_{OXA-1}, *bla*_{OXA-4}, *bla*_{OXA-30}, *bla*_{GES-1},

*bla*_{GES-9} and *bla*_{GES-11} were screened as described earlier [23]. The *bla*_{OXA-23-like}, *bla*_{OXA-24-like}, *bla*_{OXA-51-like}, *bla*_{OXA-58-like} were screened as described by Karunasagar et al. [24]. The PCR amplicons of the resistance genes were sequenced and genes were confirmed using NCBI BLASTN programme.

2.5 Plasmid isolation and plasmid incompatibility grouping

Plasmid isolation was performed for all the isolates harbouring resistance genes. The isolation of plasmid DNA was performed using HiPurA Plasmid DNA Miniprep Purification Kit (Himedia, India). Chromosomal DNA contamination was checked using the 16S rRNA primers as described earlier [25]. Plasmid incompatibility (*inc/rep*) typing (FIA, FIB, FIC, HI1, HI2, I1-Ig, L/M, N, P, W, T, A/C, K, B/O, X, Y, F, and FIIA replicons) was performed using multiplex PCR following the primers and PCR conditions as described by Carattoli et al. [26].

2.6 Conjugation studies

Representative carbapenem-resistant isolates harbouring plasmid-borne resistances (n=11) were subjected for conjugation using broth-mating method [18]. Briefly, the donor strain (strains carrying resistance genes) and the recipient strain (*E. coli* AB1157, Str^r) were grown overnight, and mixed in 9:1 ratio each of donor and recipient. The cells were kept undisturbed for 6 hours at 37°C and plated on to antibiotic containing medium. The isolates which grew on both meropenem and streptomycin were considered as transconjugants. All the transconjugants were confirmed for the presence of respective carbapenem resistance genes using PCR.

3.0 Results

3.1 Bacterial classification

In this cross-sectional study, a total of 151 non-duplicate, Gram-negative bacteria belonging to 11 genera were studied which include *Escherichia coli* (n=57, 37.7%), *Klebsiella pneumoniae* (n=40, 26.4%), *Klebsiella oxytoca* (n=5, 3.3%), *Pseudomonas aeruginosa* (n=10, 6.6%), *Salmonella typhi* (n=8, 5.2%), *Enterobacter cloacae* (n=8, 5.2%), *Acinetobacter baumannii* (n=7, 4.6%), *Serratia marcescens* (n=5, 3.3%),

Achromobacter xylosoxidans (n=5, 3.3%), *Proteus mirabilis* (n=5, 3.3%) and *Elizabethkingia meningoseptica* (n=1, 0.6%). Most of the isolates were isolated from urine 37% (56/151) and blood 28% (42/151) and from other sources such as pus (7%), bronchial secretion (2%), cerebrospinal fluid (1%), pulmonary secretion (1%), bile fluid (5%) and unknown (19%).

3.2 Antibiotic susceptibility studies

Table 1 summarizes the antibiotic susceptibility pattern of all the isolates tested against eight different antibiotics. MIC for meropenem showed that 107/151 (71%) isolates were resistant (fig.1), whereas 128 (84.7%) isolates were meropenem- resistant by the disk-diffusion method. For imipenem, 68% (n=103) were resistant by micro-broth dilution method whereas 83% (n=125) resistant by the disk-diffusion method. MIC₅₀ and MIC₉₀ values for meropenem were 16 mg/L and 8 mg/L respectively and for imipenem MIC₅₀ = 8 mg/L and MIC₉₀ = 8 mg/L.


Table 1: Antibiotic resistance pattern and the prevalence of multi-drug resistant isolates among 151 Gram-negative bacteria isolated from clinical samples.

Bacteria/antibiotic	GEN	AMY	IMP	ETP	AMK	MER	COL	CEF	Total MDR isolates (n=151)
<i>E. coli</i> (n=57)	51 (89)	45 (79)	46 (81)	38 (67)	49 (86)	43 (75)	35 (61)	45 (79)	54 (95)
<i>K. pneumoniae</i> (n=40)	33 (83)	31 (78)	32 (80)	28 (70)	36 (90)	32 (80)	29 (73)	30 (75)	32 (80)
<i>P. aeruginosa</i> (n=10)	10 (100)	10 (100)	9 (90)	6 (60)	8 (80)	10 (100)	7 (70)	8 (80)	10 (100)
<i>S. typhi</i> (n=8)	6 (75)	7 (88)	5 (63)	5 (63)	6 (75)	7 (88)	5 (63)	7 (88)	7 (88)
<i>E. cloacae</i> (n=8)	7 (88)	8 (100)	7 (88)	6 (75)	8 (100)	8 (100)	6 (75)	7 (88)	8 (100)
<i>A. baumannii</i> (n=7)	7 (100)	6 (86)	7 (100)	6 (86)	7 (100)	7 (100)	5 (71)	7 (100)	7 (100)
<i>S. marcescens</i> (n=5)	5 (100)	5 (100)	5 (100)	3 (60)	5 (100)	5 (100)	4 (80)	5 (100)	5 (100)
<i>A. xylosoxidans</i> (n=5)	5 (100)	5 (100)	4 (80)	2 (40)	5 (100)	5 (100)	4 (80)	5 (100)	5 (100)
<i>K. oxytoca</i> (n=5)	4 (80)	5 (100)	5 (100)	4 (80)	5 (100)	5 (100)	4 (80)	5 (100)	5 (100)

P. mirabilis (n=5)	5 (100)	5 (100)	4 (80)	4 (80)	5 (100)	5 (100)	4 (80)	5 (100)	5 (100)
E. meningoseptica (n=1)	1 (100)	1 (100)	1 (100)	0	1 (100)	1 (100)	1 (100)	1 (100)	1 (100)

Values represent the number of resistant isolates, % is listed in brackets. GEN-gentamicin, AMY-amoxyclav, IMP-imipenem, ETP-ertapenem, AMK-amikacin, MER-meropenem, COL-colistin, CEF-cefepime. Isolates were deemed as MDR only when the isolates are resistant to three or more antibiotics.

Figure 1: The distribution of Gram-negative bacteria and comparison of minimal inhibitory concentrations (MIC) of imipenem and meropenem resistance.



3.3 Distribution of carbapenemase resistance genes

The distribution of beta-lactamase resistance genes among 151 Gram-negative isolates is summarized in Table 2. Of the 57 *E. coli*, 32 isolates carried carbapenemase and five *E. coli* isolates carried more than one carbapenem resistance genes. Among the *K. pneumoniae*, 19/40 carried the studied genes and one isolate was positive for both

*bla*_{NDM} and *bla*_{OXA-48-like}. Carbapenem resistance genes were detected in 71/151 by PCR and 10 isolates had more than one gene. The most prevalent resistance gene was *bla*_{NDM-1} (n=22), *bla*_{OXA-48-like} (n=21), *bla*_{GES-1} (n=11), *bla*_{GES-9} (n=8), *bla*_{OXA-23-like} (n=7), *bla*_{OXA-51-like} (n=9) and *bla*_{IMP-1} (n=3). The beta-lactamase genes *bla*_{KPC}, *bla*_{VIM}, *bla*_{BIC}, *bla*_{GIM}, *bla*_{DIM}, *bla*_{SIM} and *bla*_{AIM} were not detected in the isolates. Sequencing of genes showed that all the amplified NDM genes were NDM-1, OXA-48-like genes were OXA-181 and IMP genes were IMP-1.

Table 2: The distribution of carbapenemase genes among Gram-negative bacteria isolated from the clinical isolates.

Bacteria/resistance gene	<i>bla</i> _{NDM-1}	<i>bla</i> _{OXA-181}	<i>bla</i> _{IMP-1}	<i>bla</i> _{GES-1}	<i>bla</i> _{GES-9}	<i>bla</i> _{OXA-23}	<i>bla</i> _{OXA-51}
<i>E. coli</i> (n=57)	12	9	1	6	5	4	-
<i>K. pneumoniae</i> (n=40)	5	7	-	3	2	1	2
<i>P. aeruginosa</i> (n=10)	1	-	-	2	-	-	-
<i>E. cloacae</i> (n=8)	2	1	1	-	1	-	-
<i>A. baumannii</i> (n=7)	-	2	-	-	-	2	7
<i>S. marcescens</i> (n=5)	1	-	-	-	-	-	-
<i>A. xylosoxidans</i> (n=5)	1	-	-	-	-	-	-
<i>K. oxytoca</i> (n=5)	-	1	-	-	-	-	-
<i>P. mirabilis</i> (n=5)	-	1	1	-	-	-	-
Total	22	21	3	11	8	7	9

-A total of 20 resistance genes were studied that include *bla*_{NDM}, *bla*_{OXA-48-like}, *bla*_{KPC}, *bla*_{IMP}, *bla*_{VIM}, *bla*_{DIM}, *bla*_{BIC}, *bla*_{GIM}, *bla*_{SIM}, *bla*_{AIM}, *bla*_{OXA-1}, *bla*_{OXA-4}, *bla*_{OXA-30}, *bla*_{GES-1}, *bla*_{GES-9}, *bla*_{GES-11}, *bla*_{OXA-23-like}, *bla*_{OXA-24-like}, *bla*_{OXA-51-like}, *bla*_{OXA-58-like}.

-The genes *bla*_{KPC}, *bla*_{VIM}, *bla*_{DIM}, *bla*_{BIC}, *bla*_{GIM}, *bla*_{SIM}, *bla*_{AIM}, *bla*_{OXA-1}, *bla*_{OXA-4}, *bla*_{OXA-30}, *bla*_{GES-11}, *bla*_{OXA-24-like} and *bla*_{OXA-58-like} were not observed in any of the isolates.

-GES-1 is an extended spectrum beta-lactamase which shows a significant degree of inhibition by imipenem indicating that it may be able to bind imipenem with strong affinity without being able to hydrolyze it. This enzyme was detected in 11 isolates.

-OXA-51 is a beta-lactamase which can only act as a carbapenemase if it is upregulated by insertion elements or has a Trp222 site mutation. This enzyme was detected in 9 isolates.

3.4 Identification of *E. coli* pathotypes and *Klebsiella* serotypes

The *E. coli* pathotyping results showed that, of the 57 *E. coli* isolates tested 12 were Enteropathogenic (EPEC), 9 Enteroaggregative (EAEC), 8 Enterohemorrhagic (EHEC), 3 Enterotoxigenic (ETEC), 1 Enteroinvasive (EIEC) and 24 unclassified *E. coli* (Table 3). Of the 12 EPEC isolates 3 were positive for NDM-1, 2 for OXA-181 and one for OXA-23; among 8 EHEC isolates NDM-1 was detected in 2 isolates, OXA-181 in one isolate and GES-1 plus GES-9 in one isolate; among 9 EAEC isolates GES-1 was found in 1 isolate, OXA-23 in 1 and OXA-23 along with GES-9 in one isolate; among 3 ETEC isolates 1 isolate carried NDM-1 and one EIEC isolate carried NDM-1 along with OXA-181. The virulence genes found in *E. coli* isolates included *eaeA* (n=20), *LT* (n=3), *aggR* (n=6), *astA* (n=5) and *VirA* (n=1). 24 *E. coli* isolates did not belong to any of the tested pathotypes.

Table 3: Number of *E. coli* pathotypes and *Klebsiella* serotypes among all the *E. coli* and *Klebsiella* species isolated from clinical samples.

Bacterial strain	<u>Pathotypes</u>					
	EPEC	EHEC	EIEC	ETEC	EAEC	Unknown
<i>E. coli</i> (57)	12	8	1	3	9	24
<u>Virulence gene</u>						
	<i>eaeA</i>	<i>LT</i>	<i>ST</i>	<i>aggR</i>	<i>astA</i>	<i>VirA</i>
<i>E. coli</i> (57)	20	3	0	6	5	1
<u>Serotypes</u>						
	K1	K2	K5	Unknown		
<i>Klebsiella</i> sp. (45)	14	11	0	20		
<u>Serotype gene</u>						
	<i>wzy</i>_{KPK1}	<i>wzy</i>_{KPK2}	<i>wzy</i>_{KPK5}			
<i>Klebsiella</i> sp. (45)	14	11	0			

Of the 45 *Klebsiella* species, 14 belonged to K1 serotype, 11 were K2, none of the isolates were of K5 serotypes and 20 were of unknown serotypes (Table 3). Of the 14 *K. pneumoniae* K1, NDM-1 (n=1), OXA-181 (n=4) and OXA-51 (n=1) was detected in

K1 serotypes. Among 11 *K. pneumoniae* K2, OXA-181 (n=1) and GES-9 (n=1) were detected in K2 serotypes.

3.5 Plasmid incompatibility typing and conjugation

Plasmid DNA was isolated from 70 isolates which carried resistance genes (Table 4). The plasmids ranged from 10-100 kb in size. In total, of the 151 isolates studied 70 isolates carried resistance genes of which 11 were plasmid-borne and 59 were chromosomal. Of the 37 *E. coli* isolates, 32 isolates carried resistance genes of which 6 were plasmid-borne. Among 40 *K. pneumoniae*, only 19 carried resistance genes of which 3 were associated with plasmids. In *E. cloacae*, one isolate carried *bla*_{NDM-1} on plasmid and in *P. mirabilis*; one isolate carried plasmid-borne *bla*_{IMP-1}. Plasmid incompatibility/replicon (inc/rep) typing results showed that the plasmids were belonging to Inc/rep types: IncX, IncA/C, IncFIA-FIB and IncFIIA (Table 4). *E. coli* isolates harboured *bla*_{NDM-1} genes in IncX (EC10), IncA/C (EC21) and IncFIA-FIB (EC29) type plasmids whereas *bla*_{OXA-48-like} genes were associated with IncFIIA (EC39) and IncFIA-FIB (EC29), and *bla*_{GES-1/9} genes with IncFIA-FIB (EC47) type plasmid.

K. pneumoniae isolates harboured *bla*_{NDM-1} genes in IncFIA-FIB (KP10) and *bla*_{GES-1}, *bla*_{OXA-23/51-like} genes in IncA/C (KP31 and KP39) type plasmids. One *E. cloacae* isolate harboured *bla*_{NDM-1} gene in IncFIIA (EL3) type plasmid and one *P. mirabilis* isolate harboured *bla*_{IMP-1} gene in IncFIA-FIB (PM5) type plasmid.

Overall, 6 *E. coli* (EC10, 21, 29, 39, 44, 47) isolates, 3 *K. pneumoniae* isolates (KP10, 31, 39), one *E. cloacae* isolate (EL3) and one *P. mirabilis* isolate (PM5) carried resistance genes on plasmid of the identified inc/rep types. All the 6 *E. coli* isolates (EC10, 21, 29, 39, 44, 47) were found to transfer resistance plasmids to susceptible *E. coli* AB1157. Inter-generic transfer of NDM-1 was observed in one *K. pneumoniae* isolate (KP10) in which *bla*_{NDM-1} harbouring plasmid IncFIA-FIB was transferrable to *E. coli* AB1157 (Table 4).

Table 4: Distribution of resistance genes, plasmid incompatibility grouping and transconjugation studies on Gram-negative isolates that were harbouring resistance genes.

Bacterial isolate	Source	Pathotype/ Serotype	Resistance gene	Plasmid <i>inc/rep</i> typing	Conjugative plasmid
<i>E. coli</i> EC1	Urine	EPEC	<i>bla</i> _{NDM-1}	-	-
<i>E. coli</i> EC2	Urine	EAEC	ND	-	-
<i>E. coli</i> EC3	Blood	EHEC	ND	-	-
<i>E. coli</i> EC4	Pus	Unknown	<i>bla</i> _{NDM-1}	-	-
<i>E. coli</i> EC5	Urine	Unknown	<i>bla</i> _{NDM-1}	-	-
<i>E. coli</i> EC6	Pus	EAEC	ND	-	-
<i>E. coli</i> EC7	Urine	EPEC	<i>bla</i> _{NDM-1}	-	-
<i>E. coli</i> EC8	Urine	Unknown	ND	-	-
<i>E. coli</i> EC9	Urine	EAEC	ND	-	-
<i>E. coli</i> EC10	Blood	EHEC	<i>bla</i> _{NDM-1}	*IncX	+
<i>E. coli</i> EC11	Unknown	Unknown	ND	-	-
<i>E. coli</i> EC12	Urine	Unknown	<i>bla</i> _{NDM-1}	-	-
<i>E. coli</i> EC13	Unknown	Unknown	ND	-	-
<i>E. coli</i> EC14	Unknown	EPEC	ND	-	-
<i>E. coli</i> EC15	Urine	ETEC	ND	-	-
<i>E. coli</i> EC16	Urine	Unknown	ND	-	-
<i>E. coli</i> EC17	Urine	Unknown	<i>bla</i> _{NDM-1}	-	-
<i>E. coli</i> EC18	Urine	EPEC	ND	-	-
<i>E. coli</i> EC19	Unknown	ETEC	ND	-	-
<i>E. coli</i> EC20	Urine	Unknown	ND	-	-
<i>E. coli</i> EC21	Blood	EHEC	<i>bla</i> _{NDM-1}	*IncA/C	+
<i>E. coli</i> EC22	Unknown	Unknown	<i>bla</i> _{NDM-1}	-	-
<i>E. coli</i> EC23	Urine	ETEC	<i>bla</i> _{NDM-1}	-	-
<i>E. coli</i> EC24	Unknown	EAEC	ND	-	-
<i>E. coli</i> EC25	Urine	EPEC	<i>bla</i> _{NDM-1}	-	-
<i>E. coli</i> EC26	Urine	Unknown	ND	-	-
<i>E. coli</i> EC27	Urine	EPEC	ND	-	-
<i>E. coli</i> EC28	Bile fluid	Unknown	ND	-	-
<i>E. coli</i> EC29	Unknown	EIEC	<i>bla</i> _{NDM-1} , <i>bla</i> _{OXA-181}	*IncFIA-FIB	+
<i>E. coli</i> EC30	Urine	Unknown	<i>bla</i> _{OXA-181}	-	-
<i>E. coli</i> EC31	Unknown	EPEC	<i>bla</i> _{OXA-181}	-	-
<i>E. coli</i> EC32	Blood	EAEC	ND	-	-
<i>E. coli</i> EC33	Blood	Unknown	<i>bla</i> _{OXA-181}	-	-
<i>E. coli</i> EC34	Bile fluid	EPEC	<i>bla</i> _{OXA-181}	-	-
<i>E. coli</i> EC35	Urine	Unknown	ND	-	-
<i>E. coli</i> EC36	Urine	Unknown	<i>bla</i> _{OXA-181}	-	-
<i>E. coli</i> EC37	Bile fluid	EPEC	ND	-	-

<i>E. coli</i> EC38	Urine	EPEC	ND	-	-
<i>E. coli</i> EC39	Blood	Unknown	<i>bla</i> _{OXA-181}	*IncFIIA	+
<i>E. coli</i> EC40	Blood	EHEC	<i>bla</i> _{OXA-181}	-	-
<i>E. coli</i> EC41	Blood	Unknown	<i>bla</i> _{OXA-181}	-	-
<i>E. coli</i> EC42	Blood	Unknown	<i>bla</i> _{IMP-1}	-	-
<i>E. coli</i> EC43	Urine	EPEC	ND	-	-
<i>E. coli</i> EC44	Pus	EAEC	<i>bla</i> _{GES-1}	*IncFIA-FIB	+
<i>E. coli</i> EC45	Unknown	Unknown	<i>bla</i> _{GES-1}	-	-
<i>E. coli</i> EC46	Urine	EAEC	<i>bla</i> _{GES-1}	-	-
<i>E. coli</i> EC47	Blood	EHEC	<i>bla</i> _{GES-1, bla} _{GES-9}	*IncFIA-FIB	+
<i>E. coli</i> EC48	Pus	Unknown	ND	-	-
<i>E. coli</i> EC49	Blood	EHEC	<i>bla</i> _{GES-1, bla} _{GES-9}	-	-
<i>E. coli</i> EC50	Pus	Unknown	<i>bla</i> _{GES-1, bla} _{GES-9}	-	-
<i>E. coli</i> EC51	Bile fluid	Unknown	<i>bla</i> _{GES-9}	-	-
<i>E. coli</i> EC52	Unknown	EAEC	<i>bla</i> _{GES-9,} <i>bla</i> _{OXA-23}	-	-
<i>E. coli</i> EC53	Blood	Unknown	<i>bla</i> _{OXA-23}	-	-
<i>E. coli</i> EC54	Urine	EPEC	<i>bla</i> _{OXA-23}	-	-
<i>E. coli</i> EC55	Unknown	EHEC	ND	-	-
<i>E. coli</i> EC56	Urine	EAEC	<i>bla</i> _{OXA-23}	-	-
<i>E. coli</i> EC57	Blood	EHEC	ND	-	-
<i>K. pneumoniae</i> KP1	Urine	K1	ND	-	-
<i>K. pneumoniae</i> KP2	Urine	K2	ND	-	-
<i>K. pneumoniae</i> KP3	Blood	Unknown	<i>bla</i> _{NDM-1}	-	-
<i>K. pneumoniae</i> KP4	Bile fluid	K1	ND	-	-
<i>K. pneumoniae</i> KP5	Urine	K2	ND	-	-
<i>K. pneumoniae</i> KP6	Blood	K1	ND	-	-
<i>K. pneumoniae</i> KP7	Blood	Unknown	<i>bla</i> _{NDM-1}	-	-
<i>K. pneumoniae</i> KP8	Blood	K1	ND	-	-
<i>K. pneumoniae</i> KP9	Urine	K1	<i>bla</i> _{NDM-1}	-	-
<i>K. pneumoniae</i> KP10	Blood	Unknown	<i>bla</i> _{NDM-1}	*IncFIA-FIB	+
<i>K. pneumoniae</i> KP11	Unknown	Unknown	<i>bla</i> _{NDM-1}	-	-
<i>K. pneumoniae</i> KP12	Bile fluid	K2	ND	-	-
<i>K. pneumoniae</i> KP13	Urine	K1	ND	-	-
<i>K. pneumoniae</i> KP14	Urine	Unknown	ND	-	-
<i>K. pneumoniae</i> KP15	Pulmonary secretion	K2	ND	-	-
<i>K. pneumoniae</i> KP16	Urine	K2	ND	-	-
<i>K. pneumoniae</i> KP17	Blood	K1	<i>bla</i> _{OXA-181}	-	-
<i>K. pneumoniae</i> KP18	Unknown	K2	ND	-	-

<i>K. pneumoniae</i> KP19	Blood	Unknown	<i>bla</i> _{OXA-181}	-	-
<i>K. pneumoniae</i> KP20	Unknown	K1	<i>bla</i> _{OXA-181}	-	-
<i>K. pneumoniae</i> KP21	Unknown	Unknown	<i>bla</i> _{OXA-181}	-	-
<i>K. pneumoniae</i> KP22	Unknown	Unknown	ND	-	-
<i>K. pneumoniae</i> KP23	Blood	K1	ND	-	-
<i>K. pneumoniae</i> KP24	Blood	K2	ND	-	-
<i>K. pneumoniae</i> KP25	Unknown	K2	<i>bla</i> _{OXA-181}	-	-
<i>K. pneumoniae</i> KP26	Blood	Unknown	ND	-	-
<i>K. pneumoniae</i> KP27	Unknown	K1	<i>bla</i> _{OXA-181}	-	-
<i>K. pneumoniae</i> KP28	Blood	K1	<i>bla</i> _{OXA-181}	-	-
<i>K. pneumoniae</i> KP29	Unknown	Unknown	ND	-	-
<i>K. pneumoniae</i> KP30	Urine	K1	ND	-	-
<i>K. pneumoniae</i> KP31	Blood	Unknown	<i>bla</i> _{GES-1}	*IncA/C	-
<i>K. pneumoniae</i> KP32	Unknown	Unknown	<i>bla</i> _{GES-1}	-	-
<i>K. pneumoniae</i> KP33	Urine	Unknown	<i>bla</i> _{GES-1}	-	-
<i>K. pneumoniae</i> KP34	Blood	K2	ND	-	-
<i>K. pneumoniae</i> KP35	Unknown	K1	ND	-	-
<i>K. pneumoniae</i> KP36	Urine	Unknown	<i>bla</i> _{GES-9}	-	-
<i>K. pneumoniae</i> KP37	Bile fluid	K2	<i>bla</i> _{GES-9}	-	-
<i>K. pneumoniae</i> KP38	Blood	K2	ND	-	-
<i>K. pneumoniae</i> KP39	Urine	Unknown	<i>bla</i> _{OXA-23} , <i>bla</i> _{OXA-51}	*IncA/C	-
<i>K. pneumoniae</i> KP40	Urine	K1	<i>bla</i> _{OXA-51}	-	-
<i>P. aeruginosa</i> PA1	Pus	NA	<i>bla</i> _{NDM-1}	-	-
<i>P. aeruginosa</i> PA2	Pus	NA	ND	-	-
<i>P. aeruginosa</i> PA3	Pus	NA	ND	-	-
<i>P. aeruginosa</i> PA4	Bronchial secretion	NA	ND	-	-
<i>P. aeruginosa</i> PA5	Urine	NA	<i>bla</i> _{GES-1}	-	-
<i>P. aeruginosa</i> PA6	Unknown	NA	ND	-	-
<i>P. aeruginosa</i> PA7	Blood	NA	<i>bla</i> _{GES-1}	-	-
<i>P. aeruginosa</i> PA8	Pus	NA	ND	-	-
<i>P. aeruginosa</i> PA10	Pus	NA	ND	-	-
<i>S. typhi</i> ST1	Blood	NA	ND	-	-
<i>S. typhi</i> ST2	Unknown	NA	ND	-	-
<i>S. typhi</i> ST3	Urine	NA	ND	-	-
<i>S. typhi</i> ST4	Blood	NA	ND	-	-
<i>S. typhi</i> ST5	Urine	NA	ND	-	-
<i>S. typhi</i> ST6	Blood	NA	ND	-	-
<i>S. typhi</i> ST7	Blood	NA	ND	-	-

<i>S. typhi</i> ST8	Unknown	NA	ND	-	-
<i>E. cloacae</i> EL1	Urine	NA	ND	-	-
<i>E. cloacae</i> EL2	Blood	NA	<i>bla</i> _{NDM-1}	-	-
<i>E. cloacae</i> EL3	Urine	NA	<i>bla</i> _{NDM-1}	*IncFIIA	-
<i>E. cloacae</i> EL4	Bronchial secretion	NA	ND	-	-
<i>E. cloacae</i> EL5	Blood	NA	<i>bla</i> _{OXA-181}	-	-
<i>E. cloacae</i> EL6	Urine	NA	-	-	-
<i>E. cloacae</i> EL7	Urine	NA	<i>bla</i> _{IMP-1}	-	-
<i>E. cloacae</i> EL8	Urine	NA	<i>bla</i> _{GES-9}	-	-
<i>A. baumannii</i> AB1	Cerebrospinal fluid	NA	<i>bla</i> _{OXA-181} , <i>bla</i> _{OXA-51}	-	-
<i>A. baumannii</i> AB2	Urine	NA	<i>bla</i> _{OXA-181} , <i>bla</i> _{OXA-51}	-	-
<i>A. baumannii</i> AB3	Unknown	NA	<i>bla</i> _{OXA-51}	-	-
<i>A. baumannii</i> AB4	Pus	NA	<i>bla</i> _{OXA-23} , <i>bla</i> _{OXA-51}	-	-
<i>A. baumannii</i> AB5	Blood	NA	<i>bla</i> _{OXA-23} , <i>bla</i> _{OXA-51}	-	-
<i>A. baumannii</i> AB6	Urine	NA	<i>bla</i> _{OXA-51}	-	-
<i>A. baumannii</i> AB7	Urine	NA	<i>bla</i> _{OXA-51}	-	-
<i>S. marcescens</i> SM1	Bronchial secretion	NA	ND	-	-
<i>S. marcescens</i> SM2	Blood	NA	<i>bla</i> _{NDM-1}	-	-
<i>S. marcescens</i> SM3	Unknown	NA	ND	-	-
<i>S. marcescens</i> SM4	Urine	NA	ND	-	-
<i>S. marcescens</i> SM5	Unknown	NA	ND	-	-
<i>A. xylosoxidans</i> AY1	Unknown	NA	ND	-	-
<i>A. xylosoxidans</i> AY2	Blood	NA	ND	-	-
<i>A. xylosoxidans</i> AY3	Urine	NA	ND	-	-
<i>A. xylosoxidans</i> AY4	Urine	NA	ND	-	-
<i>A. xylosoxidans</i> AY5	Urine	NA	<i>bla</i> _{NDM-1}	-	-
<i>K. oxytoca</i> KO1	Blood	NA	ND	-	-
<i>K. oxytoca</i> KO2	Urine	NA	ND	-	-
<i>K. oxytoca</i> KO3	Blood	NA	ND	-	-
<i>K. oxytoca</i> KO4	Blood	NA	<i>bla</i> _{OXA-181}	-	-
<i>K. oxytoca</i> KO5	Urine	NA	ND	-	-
<i>P. mirabilis</i> PM1	Unknown	NA	ND	-	-
<i>P. mirabilis</i> PM2	Blood	NA	<i>bla</i> _{OXA-181}	-	-
<i>P. mirabilis</i> PM3	Urine	NA	ND	-	-

<i>P. mirabilis</i> PM4	Blood	NA	ND	-	-
<i>P. mirabilis</i> PM5	Urine	NA	<i>bla</i> _{IMP-1}	*IncFIA-FIB	-
<i>E. meningoseptica</i> EM1	Cerebrospinal fluid	NA	ND	-	-

NA- serotyping/pathotypic not applicable; ND- resistance gene not detected; *Plasmids carrying resistance genes; ‘-’ – Absence; ‘+’ – Conjugation positive; highlighted in grey represents the isolates carrying resistance genes on conjugative plasmids

4.0 Discussion

In India, the prevalence of carbapenem-resistant Gram-negative bacteria has been reported with an increasing frequency [17,18]. In this study, the distribution of carbapenem-resistant isolates among 11 genera of Gram-negative bacteria isolated from diagnostic center in Tamil Nadu, India is reported. Previously, the increasing prevalence of ESBL and MBL producers among Gram-negative bacteria has been reported in India [27-30].

In this study, MIC results showed that 107/151 (71%) were resistant to meropenem in accordance with 128 isolates by the disk-diffusion method. All the 71 isolates harbouring carbapenem resistance genes were resistant with MIC and disc-diffusion method.–The studied *E. coli* pathotypes (EPEC, EHEC, EIEC, EAEC and ETEC) are associated with intestinal diseases, they are collectively called as diarrheagenic *E. coli* (DEC) or intestinal pathogenic *E. coli* (IPEC) [31,32]. All these pathotypes are linked directly to their virulence properties and severity of infections. Though there are studies that showed the prevalence of DEC in India [33,34], still the studies on *E. coli* pathotypes (virulence) interrelation to carbapenem resistance is not well established in India. In this study, all the five *E. coli* pathotypes were found to harbour carbapenem resistance genes, namely EPEC (NDM-1, OXA-181, and OXA-23), EHEC (NDM-1, OXA-181, GES-1, and GES-9), EIEC (NDM-1, OXA-181), EAEC (GES-9, OXA-23, and GES-1), ETEC (NDM-1) and some are unknown pathotypes (Table 4). Adding to their virulence, the presence of resistance genes makes these bacterial infections (mostly diarrhoea) more complicated due to unavailability of treatment options. The *Klebsiella* isolates can be grouped into serotypes using surface antigens or surface exposed lipopolysaccharides [35]. The *Klebsiella* belonging to K-serotypes have K-antigen that

relates to the capsule polysaccharide (CPS) [35]. Of the known capsular types (eight serotypes), the serotypes K1 and K2 are the most virulent among the hypervirulent *K. pneumoniae* (hvKP) [36]. In this study, of the 14 *Klebsiella* isolates belonging to K1 serotypes, six isolates carried carbapenem resistance genes, *bla*_{NDM-1}, *bla*_{OXA-181} and *bla*_{OXA-51}. Among the 11 *Klebsiella* isolates belonging to K2 serotype, two isolates were found to carry carbapenem resistance genes, *bla*_{OXA-181} and *bla*_{GES-9} (Table 4). In India, NDM-1 and OXA-48 genes were detected in *Klebsiella* belonging to K2 serotypes [36], but to the best of our knowledge, this is the first study to detect the presence of carbapenemase genes NDM-1, OXA-181 and OXA-51 among K1 serotypes.

As carbapenems are the last resort of antibiotics available to treat infections caused by Gram-negative bacteria, the prevalence of carbapenem resistance is given a global attention. Our previous studies had reported the dissemination of carbapenem-resistant bacteria and carbapenem resistance genes among Gram-negative bacteria [17,18]. Here, we report the prevalence (71%) of carbapenem resistant isolates among 11 genera of Gram-negative bacteria. Beta-lactamase resistance genes such as *bla*_{NDM-1} (n=22), *bla*_{OXA-181} (n=21), *bla*_{GES-1} (n=11), *bla*_{GES-9} (n=8), *bla*_{OXA-23} (n=7), *bla*_{OXA-51} (n=9) and *bla*_{IMP-1} (n=3) were found in 71 isolates (10 isolates carrying more than one genes), comparatively our earlier studies showed the low prevalence (27%) of *bla*_{NDM-1} and *bla*_{OXA-181} genes among carbapenem-resistant isolates [18]. The coexistence of *bla*_{NDM-1} and *bla*_{OXA-181} in *E. coli* is one of the serious concerns from healthcare prospective. All the *A. baumannii* isolates (n=7) were found to have either the class D carbapenem hydrolyzing oxacillinases (OXA-23, OXA-181) and OXA-51 is naturally existed in *Acinetobacter* spp. [37]. There were earlier reports in India showing the presence of OXA-23 and OXA-51 in carbapenem-resistant *Acinetobacter* causing serious health care problems [12]. *Enterobacteriaceae* are encoded by OXA-48-like genes as carbapenem-hydrolyzing class D β-lactamases [13,38]. But the unusual occurrence of *bla*_{OXA-23} in *E. coli*, and plasmid-borne (IncA/C) *bla*_{OXA-23} and *bla*_{OXA-51} in *K. pneumoniae* is one of the important findings of this study. There are very few earlier studies that reported the presence of *bla*_{OXA-23} gene in *E. coli* [39,40]. To the best of our knowledge, this is the first study to report the plasmid-borne (IncA/C) OXA-23 and OXA-51 in *K. pneumoniae*. The OXA-23-like genes in *Enterobacteriaceae* may be carried within a

transposon but was not characterized in this study. The resistance reports on *E. meningoseptica* are very rare in India [39,40] and in our study, it was found that one isolate of *E. meningoseptica* was resistant to imipenem and meropenem. Though earlier studies showed the presence of carbapenemase genes in *E. meningoseptica*, in this study no carbapenem resistance genes were amplified.

Carbapenem resistance among Gram-negative bacteria is becoming very common in India and the spread of carbapenem resistance genes are one of the troublesome problems. These resistance genes that are located adjacent to the mobile genetic elements (integrons and transposons), which facilitates the easy transposition between replicons [41]. The most common plasmid replicon types for carbapenem resistance genes are IncF, IncA/C₂, IncX3, IncL/M and IncH [42]. In this study, *bla*_{NDM-1} was found to be harboured in IncX, IncA/C, IncFIA-FIB and IncFIIA; *bla*_{OXA-181} in IncA/C, IncFIA-FIB and IncFIIA; *bla*_{GES-1/9} in IncFIA-FIB and IncA/C; *bla*_{IMP-1} in IncFIA-FIB and *bla*_{OXA-23/51} in IncA/C. The presence of plasmid-borne *bla*_{OXA-23/51} is very rare and important finding, considering the rapid spread of carbapenem-resistance among Gram-negative bacteria. Interestingly, the isolates such as *P. aeruginosa*, *Salmonella typhi*, *A. baumannii*, *S. marcescens*, *A. xylosoxidans*, *K. oxytoca*, and *E. meningoseptica* do not carry any plasmids harbouring resistance genes. This clearly showed that the beta-lactamase or carbapenemase resistance genes were present in the plasmids with different replicon types in the study region. Earlier, the *bla*_{NDM} IncFII plasmids were reported from India [42] and IncFIA-FIB plasmids carrying carbapenem resistance genes such as *bla*_{NDM} was report from India in the samples collected from river and sewage treatment plants [42,43]. This study also showed that some plasmids were carrying more than one resistance genes which are an alarming threat to the public health. Conjugative plasmids are known to spread their resistance characteristics among the bacteria from the same or different genus. This study showed that all the six *E. coli* isolates carrying plasmid-borne resistance genes (*bla*_{NDM-1}, *bla*_{OXA-181}, *bla*_{GES-1}, *bla*_{GES-9}) were conjugative and one *K. pneumoniae* isolate plasmid (IncFIA-FIB with *bla*_{NDM-1}) was transferable which clearly shows the way by which resistance genes can rapidly spread in clinical bacteria.

5.0 Conclusion

The emerging antibiotic resistance in bacteria is a worrisome problem. This study highlighted the distribution of carbapenem resistant isolates in the study region with the extra emphasis on the existence of *bla*_{NDM-1}, *bla*_{OXA-48-like}, *bla*_{IMP-1}, *bla*_{GES-1}, *bla*_{GES-9}, *bla*_{OXA-23-like}, *bla*_{OXA-51-like} among the clinical pathogens. Alternative therapeutic options should be undertaken immediately to combat the problem of resistance especially to treat infections caused by carbapenem resistant bacteria. Our study shows that the ‘conjugative plasmids’ can strongly contribute to the resistance transfer in pathogens leading to dissemination of resistance genes. Alternative approaches are necessary to combat the problem of resistance and concepts such as ‘one-health approach’ can be appreciated.

Authors' contribution:

Authors PM and NR, collected the isolates from the clinical samples. Authors PM and NR undertook the laboratory work, NR and BSL interpreted the data, and PM and NR wrote the initial manuscript. Authors NR and BSL revised and edited the manuscript. All the authors' have read and approved the manuscript.

Competing interest:

The authors declare that they have no competing interest.

Funding information:

This research work was not funded by any external agencies.

Ethics approval:

Ethical approval from Institutional Ethical Committee for studies on Human subjects (IECH), ref. no. VIT/IECH/004/Jan2015

Availability of data and materials:

All the datasets are presented in the main manuscript. The raw datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Acknowledgement:

The authors would like to thank Vellore Institute of Technology (VIT) for providing partial funding, ‘VIT Seed Grant’ and Council of Scientific and Industrial Research (CSIR) for providing financial assistance to PM in the form of senior research fellowship (SRF) to support this research.

References:

1. Roca I, Akova M, Baquero F, Carlet J, Cavalieri M *et al*. The global threat of antimicrobial resistance: science for intervention. *New Microbes New Infect* 2015;6:22-29. <https://doi.org/10.1016/j.nmni.2015.02.007>
2. Global priority list of antibiotic-resistant bacteria to guide research, discovery, and development of new antibiotics. WHO publishes list of bacteria for which new antibiotics are urgently needed. <http://www.who.int/mediacentre/news/releases/2017/bacteria-antibiotics-needed/en/>
3. Ventola CL. The antibiotic resistance crisis: part 1: causes and threats. *Pharmacy and Therapeutics* 2015;40(4):277.
4. McKenna M. Antibiotic resistance: the last resort. *Nat News* 2013;499(7459):394.
5. Ben-David D, Kordevani R, Keller N, Tal I, Marzel A *et al*. Outcome of carbapenem resistant *Klebsiella pneumoniae* bloodstream infections. *Clin Microbiol Infect* 2012;18(1):54-60. <https://doi.org/10.1111/j.1469-0691.2011.03478.x>
6. Xu Y, Gu B, Huang M, Liu H, Xu T *et al*. Epidemiology of carbapenem resistant Enterobacteriaceae (CRE) during 2000-2012 in Asia. *J Thorac Dis* 2015;7(3):376-385. <https://doi.org/10.3978/j.issn.2072-1439.2014.12.33>
7. Akiba M, Sekizuka T, Yamashita A, Kuroda M, Fujii Y *et al*. Distribution and relationships of antimicrobial resistance determinants among extended-spectrum-cephalosporin-resistant or carbapenem-resistant *Escherichia coli* isolates from rivers and sewage treatment plants in India. *Antimicrob Agents Chemother* 2016;60(5):2972-2980. <https://doi.org/10.1128/AAC.01950-15>
8. Chandran SP, Sarkar S, Diwan V, Pathak A, Shah H *et al*. Detection of virulence genes in ESBL producing, quinolone resistant commensal *Escherichia coli* from rural Indian children. *J Infect Dev Ctries* 2017;11(5):387-392. <https://doi.org/10.3855/jidc.8574>
9. Manohar P, Shanthini T, Marathe N, Jadhav S, Slathia S *et al*. Genetic characteristics of plasmid-mediated extended-spectrum β-lactamases (CTX-M) and its coexistence with

- carbapenemases (NDM-1) in clinical Gram negative bacteria. *Ind J Biotechnol* 2017;16(2):189-194.
10. Scaife W, Young HK, Paton RH and Amyes SG. Transferable imipenem-resistance in *Acinetobacter* species from a clinical source. *J Antimicrob Chemother* 1995;36(3):585-586. <https://doi.org/10.1093/jac/36.3.585>
 11. Teixeira AB, Martins AF, Barin J, Hermes DM, Pitt CP et al. First report of carbapenem-resistant *Acinetobacter nosocomialis* isolates harboring ISAbalbla OXA-23 genes in Latin America. *J Clin Microbiol* 2013;51(8):2739-2741. <https://doi.org/10.1128/JCM.00469-13>
 12. La MV, Jureen R, Lin RT and Teo JW. Unusual detection of an *Acinetobacter* class D carbapenemase gene, bla OXA-23, in a clinical *Escherichia coli* isolate. *J Clin Microbiol* 2014;52(10):3822-3823. <https://doi.org/10.1128/JCM.01566-14>
 13. Paul D, Ingati B, Bhattacharjee D, Maurya AP, Dhar D et al. An unusual occurrence of plasmid-mediated blaOXA-23 carbapenemase in clinical isolates of *Escherichia coli* from India. *Int J Antimicrob Agents* 2017;49(5):642-645. <https://doi.org/10.1016/j.ijantimicag.2017.01.012>
 14. Österblad M, Karah N, Halkilahti J, Sarkkinen H, Uhlin BE et al. Rare detection of the *Acinetobacter* class D carbapenemase blaOXA-23 gene in *Proteus mirabilis*. *Antimicrob Agents Chemother* 2016;60(5):3243-3245. <https://doi.org/10.1128/AAC.03119-15>
 15. Brown S, Young HK and Amyes SGB. Characterisation of OXA-51, a novel class D carbapenemase found in genetically unrelated clinical strains of *Acinetobacter baumannii* from Argentina. *Clin Microbiol Infect* 2005;11(1):15-23. <https://doi.org/10.1111/j.1469-0691.2004.01016.x>
 16. Chen TL, Lee YT, Kuo SC, Hsueh PR, Chang FY et al. Emergence and distribution of plasmids bearing the blaOXA-51-like gene with an upstream ISAbal in carbapenem-resistant *Acinetobacter baumannii* isolates in Taiwan. *Antimicrob Agents Chemother* 2010;54(11):4575-4581. <https://doi.org/10.1128/AAC.00764-10>
 17. Nachimuthu R, Subramani R, Maray S, Gothandam KM, Sivamangala K et al. Characterization of carbapenem-resistant Gram-negative bacteria from Tamil Nadu. *J Chemother* 2016;28(5):371-374. <https://doi.org/10.1179/1973947815Y.0000000056>
 18. Manohar P, Shanthini T, Ayyanar R, Bozdogan B, Wilson A et al. The distribution of carbapenem-and colistin-resistance in Gram-negative bacteria from the Tamil Nadu region in India. *J Med Microbiol* 2017;66(7):874-83. <https://doi.org/10.1099/jmm.0.000508>

19. Clinical and Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing; twenty-fifth informational supplement. Approved standard M100-S27. Wayne, PA: CLSI; 2017.
20. Bisi-Johnson MA, Obi CL, Vasaikar SD, Baba KA and Hattori T. Molecular basis of virulence in clinical isolates of *Escherichia coli* and *Salmonella* species from a tertiary hospital in the Eastern Cape, South Africa. *Gut pathogen* 2011;3(1):9. <https://doi.org/10.1186/1757-4749-3-9>
21. Siu LK, Fung CP, Chang FY, Lee N, Yeh KM et al. Molecular typing and virulence analysis of serotype K1 *Klebsiella pneumoniae* strains isolated from liver abscess patients and stool samples from noninfectious subjects in Hong Kong, Singapore, and Taiwan. *J Clin Microbiol* 2011;49(11):3761-3765. <https://doi.org/10.1128/JCM.00977-11>
22. Gheorghe I, Czobor I, Chifiriuc MC, Borcan E, Ghiță C et al. Molecular screening of carbapenemase-producing Gram-negative strains in Romanian intensive care units during a one year survey. *J Med Microbiol* 2014;63(10):1303-1310. <https://doi.org/10.1099/jmm.0.074039-0>
23. Dallenne C, Da Costa A, Decré D, Favier C and Arlet G. Development of a set of multiplex PCR assays for the detection of genes encoding important β -lactamases in Enterobacteriaceae. *J Antimicrob Chemother* 2010;65(3):490-495. <https://doi.org/10.1093/jac/dkp498>
24. Karunasagar A, Maiti B, Shekar M, Shenoy MS and Karunasagar I. Prevalence of OXA-type carbapenemase genes and genetic heterogeneity in clinical isolates of *Acinetobacter* spp. from Mangalore, India. *Microbiol Immunol* 2011;55(4):239-246. <https://doi.org/10.1111/j.1348-0421.2011.00313.x>
25. Lin L, Ling BD and Li XZ. Distribution of the multidrug efflux pump genes, adeABC, adeDE and adeIJK, and class 1 integron genes in multiple-antimicrobial-resistant clinical isolates of *Acinetobacter baumannii*-*Acinetobacter calcoaceticus* complex. *Int J Antimicrob Agents* 2009;33(1):27-32. <https://doi.org/10.1016/j.ijantimicag.2008.06.027>
26. Carattoli A, Bertini A, Villa L, Falbo V, Hopkins KL et al. Identification of plasmids by PCR-based replicon typing. *J Microbiol Methods* 2005;63(3):219-228. <https://doi.org/10.1016/j.mimet.2005.03.018>
27. Oberoi L, Singh N, Sharma P and Aggarwal A. ESBL, MBL and AmpC β lactamases producing superbugs—Havoc in the Intensive Care Units of Punjab India. *J Clin Diagn Res* 2013;7(1):70.

28. Azim A, Dwivedi M, Rao PB, Baronia AK, Singh RK *et al.* Epidemiology of bacterial colonization at intensive care unit admission with emphasis on extended-spectrum β -lactamase-and metallo- β -lactamase-producing Gram-negative bacteria—an Indian experience. *J Med Microbiol* 2010;59(8):955-960. <https://doi.org/10.1099/jmm.0.018085-0>
29. Nordmann P, Naas T and Poirel L. Global spread of carbapenemase-producing Enterobacteriaceae. *Emer Infect Dis* 2011;17(10):1791. <https://doi.org/10.3201/eid1710.110655>
30. Khurana S, Mathur P, Kapil A, Valsan C and Behera B. Molecular epidemiology of beta-lactamase producing nosocomial Gram-negative pathogens from North and South Indian hospitals. *J Med Microbiol* 2017;66(7):999-1004. <https://doi.org/10.1099/jmm.0.000513>
31. Robins-Browne RM, Holt KE, Ingle DJ, Hocking DM, Yang J *et al.* Are *Escherichia coli* pathotypes still relevant in the era of whole-genome sequencing?. *Front Cell Infect Microbiol* 2016;6:141. <https://doi.org/10.3389/fcimb.2016.00141>
32. Bajaj P, Singh NS and Virdi JS. *Escherichia coli* β -lactamases: what really matters. *Front Microbiol* 2016;7:417. <https://doi.org/10.3389/fmicb.2016.00417>
33. Singh T, Das S, Ramachandran VG, Dar SA, Snehaa K *et al.* Spectrum of diarrhoeagenic *Escherichia coli* in paediatric population suffering from diarrhoea and as commensals in healthy children. *Ind J Med Microbiol* 2017;35(2):204. https://doi.org/10.4103/ijmm.IJMM_16_21
34. Gupta D, Sharma M, Sarkar S, Thapa BR and Chakraborti A. Virulence determinants in enteroaggregative *Escherichia coli* from North India and their interaction in in vitro organ culture system. *FEMS Microbiol Lett* 2016;363(17). <https://doi.org/10.1093/femsle/fnw189>
35. Follador R, Heinz E, Wyres KL, Ellington MJ, Kowarik M *et al.* The diversity of *Klebsiella pneumoniae* surface polysaccharides. *Microb Genom* 2016;2(8). <https://doi.org/10.1099/mgen.0.000073>
36. Remya P, Shanthi M and Sekar U. Occurrence and characterization of hyperviscous K1 and K2 serotype in *Klebsiella pneumoniae*. *J Lab Physicians* 2018;10(3):283. https://doi.org/10.4103/JLP.JLP_48_18
37. Durante-Mangoni E and Zarrilli R. Global spread of drug-resistant *Acinetobacter baumannii*: molecular epidemiology and management of antimicrobial resistance. *Fut Microbiol* 2011;6(4):407-422. <https://doi.org/10.2217/fmb.11.23>

38. Bali NK, Fomda BA, Bashir H, Zahoor D, Lone S *et al.* Emergence of carbapenem-resistant *Acinetobacter* in a temperate north Indian State. *Br J Biomed Sci* 2013;70(4):156-160. <https://doi.org/10.1080/09674845.2013.11669950>
39. Shinha T and Ahuja R. Bacteremia due to *Elizabethkingia meningoseptica*. *IDCases* 2015;2(1):13-15. <https://doi.org/10.1016/j.idcr.2015.01.002>
40. Bhat KS, Priya R, Krishnan L and Kanungo R. *Elizabethkingia meningoseptica* bacteremia in a neonate: A case report and mini-review of the literature. *J Curr Res Scient Med* 2016;2(1):42. <https://doi.org/10.4103/2455-3069.184130>
41. Partridge SR and Iredell JR. Genetic contexts of blaNDM-1. *Antimicrob Agents Chemother* 2012;56(11):6065-6067. <https://doi.org/10.1128/AAC.00117-12>
42. Sugawara Y, Akeda Y, Sakamoto N, Takeuchi D, Motooka D *et al.* Genetic characterization of blaNDM-harboring plasmids in carbapenem-resistant *Escherichia coli* from Myanmar. *PLoS one* 2017;12(9):e0184720. <https://doi.org/10.1371/journal.pone.0184720>
43. Akiba M, Sekizuka T, Yamashita A, Kuroda M, Fujii Y *et al.* Distribution and relationships of antimicrobial resistance determinants among extended-spectrum-cephalosporin-resistant or carbapenem-resistant *Escherichia coli* isolates from rivers and sewage treatment plants in India. *Antimicrob Agents Chemother* 2016;60(5):2972-2980. <https://doi.org/10.1128/AAC.01950-15>

Figures




Figure 1

The distribution of Gram-negative bacteria and comparison of minimal inhibitory concentrations (MIC) of imipenem and meropenem resistance.